

Cite this: *Chem. Sci.*, 2018, 9, 2991Received 15th January 2018
Accepted 18th February 2018DOI: 10.1039/c8sc00232k
rsc.li/chemical-science

Gold-catalyzed [4+3]- and [4+2]-annulations of 3-en-1-ynamides with isoxazoles *via* novel 6π -electrocyclizations of 3-azahepta trienyl cations†



Sovan Sundar Giri and Rai-Shung Liu *

New gold-catalyzed [4+3]-annulations of 3-en-1-ynamides with isoxazoles afford 4*H*-azepines efficiently; this process involves 6π electrocyclizations of gold-stabilized 3-azaheptatrienyl cations. In the presence of $\text{Zn}(\text{OTf})_2$, the resulting 4*H*-azepines undergo skeletal rearrangement to furnish substituted pyridine derivatives. We subsequently develop new catalytic [4+2]-annulations between the same 3-en-1-ynamides and isoxazoles to deliver substituted pyridine products using $\text{Au}(\text{I})/\text{Zn}(\text{II})$ catalysts. This work reports the first success of the 6π electrocyclizations of heptatrienyl cations that are unprecedented in literature reports.

Introduction

Electrocyclizations of acyclic conjugated π -motifs are powerful tools to access five-, six- and seven-membered carbocycles;¹ prominent examples include Nazarov cyclizations of penta-dienyl cations² and 6π electrocyclizations of trienes,³ which have found widespread applications in organic synthesis.

In the context of seven-carbon π -motifs, heptatrienyl anions **I** undergo facile 8π electrocyclizations *via* rapid interconversions among various anion configurations (Scheme 1).⁴ In contrast, heptatrienyl cations **III**⁵ exclusively undergo Nazarov reactions because of the difficulties of forming all σ -*cis* configured cations **V** that have a high energy state.^{5b} 1-Aza- and 1-oxaheptatrienyl cations⁶ were also reported to follow Nazarov cyclizations. The realization of a 6π electrocyclization of conjugated seven-membered cations is formidable but challenging. This work reveals the first success of such seven-membered cyclizations of gold-stabilized 3-azaheptatrienyl cations **V'** to form azacyclic products **3–4** *via* a new C–C bond formation.

Scheme 1 Electrocyclizations of conjugated π -motifs.

Department of Chemistry, National Tsing-Hua University, Hsinchu, Taiwan, Republic of China. E-mail: rslu@mx.nthu.edu.tw

† Electronic supplementary information (ESI) available. CCDC 1589549, 1589562, 1589561, 1589558, 1589559 and 1589560. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8sc00232k

The advent of gold catalysis has inspired new annulations between alkynes and poor nucleophiles.⁷ N–O containing nucleophiles serve as useful building blocks to construct valuable azacyclic frameworks.⁷ Ye and Hashmi reported interesting [3+2]-annulations of isoxazoles or benzisoxazoles with electron-

rich ynamides, yielding substituted pyrrole derivatives through aza-Nazarov cyclizations of the key intermediate [eqn (1)].^{7,8} These [3+2]-annulations were extensively expanded to other N–O heterocycles including benzisoxazoles, 1,2,4-oxadiazoles, 1,4,2-dioxazoles and 4,5-dihydro-1,2,4-oxadiazoles, yielding additional five-membered azacycles as depicted in [eqn (1)].⁹ Here, we report two distinct [4+3]- and [4+2]-annulations between 3-en-1-ynamides and isoxazoles using varied catalysts. An Au(I) catalyst alone delivers 4*H*-azepines **3–4** through 6π electrocyclizations of intermediates **V'** [eqn (2)] whereas a combined action of Au(I)/Zn(II) on the same reactants furnishes highly functionalized pyridines **5** [eqn (3)]. With our convenient synthesis, the synthetic utility of new 4*H*-azepines **3–4** is also reported.¹⁰

Results and discussion

We examined the reactions of 3-methyl-3-en-1-ynamide **1a** with 3,5-dimethylisoxazole **2a** using various gold catalysts. Heating this mixture (**1a/2a** = 1 : 2 ratio) in hot DCE with 5 mol% LAuCl/AgNTf₂ [L = *p*-(*t*-Bu)₂(*o*-biphenyl)] and IPr afforded a [4+3]-annulation product, 4*H*-azepine **3a**, in 64% and 75% yields respectively (Table 1, entries 1–2). Under these conditions, a low loading (1.2 equiv.) of 3,5-dimethylisoxazole **2a** gave **3a** in a decreased yield, *ca.* 62% (entry 3). With a 10 mol% catalyst, IPrAuCl/AgNTf₂ gave a clean reaction, yielding desired **3a** up to 91% (entry 4). We tested other phosphine ligands such as PPh₃ and P(OPh)₃, yielding desired **3a** in satisfactory yields (78–81%, entries 5–6). Other counter anions such as OTf[−] and SbF₆[−] were also effective in producing **3a** in 85–88% yields (entries 7–8). AgNTf₂ alone was not active at all (entry 9).

Table 1 [4+3]-Annulations over various gold catalysts

Entry	Catalyst [mol%]	x	Time [h]	Yield ^b [%]		
				1a	3a	1a-H'/1a-H''
1 ^c	LAuCl/AgNTf ₂ [5]	2	3	20	64	—
2 ^d	IPrAuCl/AgNTf ₂ [5]	2	7	12	75	7 [2.5 : 1]
3	IPrAuCl/AgNTf ₂ [5]	1.2	7	23	62	5 [1 : 1]
4	IPrAuCl/AgNTf ₂ [10]	2	3	—	91	Trace
5	PPh ₃ AuCl/AgNTf ₂ [10]	2	3.5	—	81	5 [1.25 : 1]
6	[PhO] ₃ PAuCl/AgNTf ₂ [10]	2	3.5	—	78	13 [1.1 : 1]
7	IPrAuCl/AgSbF ₆ [10]	2	2.5	—	85	6 [1.4 : 1]
8	IPrAuCl/AgOTf [10]	2	2	—	88	Trace
9	AgNTf ₂ [10]	2	15	33	—	11

^a [1a] = 0.15 M. ^b Product yields are reported after separation from a silica column. ^c L = *p*-(*t*-Bu)₂(*o*-biphenyl). ^d IPr = 1,3-bis(diisopropylphenyl)-imidazol-2-ylidene. Ms = methanesulfonyl, DCE = 1,2-dichloroethane, and Tf = trifluoromethanesulfonyl.

Table 2 [4+3]-Annulations with various 3-en-1-ynamides

(1) R ⁴ , EWG = CH ₃ , Ts (3b , 3 h, 84%, x-ray)	(5) (3f , 22 h, 64%)
(2) R ⁴ , EWG = <i>c</i> -Pr, Ts (3c , 7 h, 86%)	
(3) R ⁴ , EWG = Bn, Ts (3d , 4.5 h, 87%)	
(4) R ⁴ , EWG = <i>n</i> -Bu, <i>n</i> -BuSO ₂ (3e , 6 h, 90%)	
(6) R ¹ = <i>i</i> -Pr (3g , 4 h, 74%)	(9) R ⁴ , EWG = <i>n</i> -Bu, Ms (3j , 5 h, 55%, 3j/3j' = 5:1)
(7) R ¹ = <i>c</i> -Pr (3h , 2 h, 79%)	(10) R ⁴ , EWG = CH ₃ , Ts (3k , 4 h, 68%, 3k/3k' = 11.1:1)
(8) R ¹ = Ph (3i , 2.5 h, 58%)	
(11) (3l , 2.5 h, 48%, x-ray) (6l , 43%, <i>E/Z</i> = 3.3:1)	(12) (3m , 14 h, 16%) (6m , 73%, x-ray)

^a [1] = 0.15 M. ^b Product yields are reported after separation from a silica column. EWG = electron withdrawing group.

Table 3 [4+3]-Annulations with various isoxazoles

Entry	(R ¹ , R ²)	2	Time [h]	Yield [%]	4
(1)	H, H	2b	4	84	4a (X-ray)
(2)	H, Me	2d	3	8	7a'
(3)	Me, H	2c	3	75	4b
(4)	Et, Et	2e	6	87	4c
(5)	<i>n</i> -Bu, <i>n</i> -Bu	2f	7	85	4d
(6)	Me, <i>n</i> -Bu	2g	3	81	4e
(7)	<i>n</i> -Bu, <i>c</i> -Pr	2h	3	82	4f
(8)	Ph, <i>n</i> -Bu	2i	2	77	4g
(9)	Ph, Ph	2j	4	69	4h
			6.5	61	4i
				30	5i (X-ray)
(10)	Me, Ph	2k	4	71	4j
				15	5j

^a [1b] = 0.15 M. ^b Product yields are reported after separation from a silica column.



Suitable substituents of 3-en-1-ynamides **1** are crucial to achieve 6π cyclizations of 3-azaheptatrienyl cations **V'** [eqn (2)]. We tested the reactions on 3-en-1-ynes **1b–1m** bearing a C(3)-substituent to circumvent aza-Nazarov cyclizations as reported in Ye's work.⁷ Herein, only entries 9 and 10 showed the presence of 3-azanorcaradienes **3'**. We examined these [4+3]-annulations on 3-methyl-3-en-1-ynamides **1b–1e** bearing various sulfonamides NTsR⁴ (R⁴ = Me, cyclopropyl, benzyl and N(*n*-C₄H₉) (–SO₂Bu)), affording the desired 4*H*-azepines **3b–3e** in high yields (84–90%, Table 2, entries 1–4). Nevertheless, this new annulation becomes less efficient for 3-en-1-ynamide **1f** bearing an oxazolidin-2-one to yield product **3f** in 64% yield (entry 5).


We altered the C(3)-substituents as in substrates **1g–1i**; their resulting products **3g–3h** (R¹ = isopropyl and cyclopropyl) were obtained in 74–79%, and **3i** (R¹ = Ph) with only 58% yield (entries 6–8). Notably, when a long *n*-butyl group was present as in species **1j** and **1k**, their corresponding reactions afforded compounds **3j/3j'** = 5/1 and **3k/3k'** = 11.1 : 1, respectively, in 55% and 68% yields (entries 9–10). For *E*-configured trisubstituted 3-en-1-yne **1l** (R¹ = Me, R² = Ph and R³ = H), 4*H*-azepine **3l**

and pyrrole **6l** were obtained in equal proportions (entry 11). When a cyclohexenyl group was present for alkene as in species **1m**, pyrrole product **6m** was dominant over azepine **3m** (entry 12). Accordingly, preferable 3-en-1-ynes comprise a small R² or R³ substituent whereas R¹ must be substituted. Herein, the structures of 4*H*-azepines **3b** and **3l**, and pyrrole species **6m** were confirmed with X-ray diffraction.¹¹

Isoxazoles of a wide scope are compatible with these [4+3]-annulations, as depicted in Table 3. The reaction of unsubstituted isoxazole **2b** with model 3-en-1-ynamide **1b** afforded the desired 4*H*-azepine **4a** in 84% yield, together with pyrrole **7a'** in only 8% yield (entry 1). Mono-substituted 3-methyl or 5-methyl isoxazoles **2c** and **2d** are also suitable for these annulations to afford compounds **4b** and **4c** in 75% and 87% yields, respectively (entries 2–3). We prepared additional 3,5-disubstituted isoxazoles **2e–2i** with R¹ = alkyl and phenyl, and R² = alkyl; their annulations proceed smoothly to produce desired **4d–4h** in 69–85% yields (entries 4–8). For di-substituted isoxazoles **2j** and **2k** bearing R² = Ph, 4*H*-azepines **4i** and **4j** were obtained in 61% and 71% yields respectively, together with their rearrangement products **5i** and **5j** in 15–30% yields (entries 9–10). Compounds **4a** and **5i** were characterized by X-ray diffraction.¹¹

Our convenient synthesis of 4*H*-azepines provides new synthetic utilities; several new functionalizations are depicted in Scheme 2. NaBH₄-reduction of species **3b** delivered an alcohol derivative **7a** in 84% yield. Selective hydrogenation of the same species afforded 2-aza-1,3-dien-5-one **7b** in 71% yield. A final treatment of 4*H*-azepine **3b** with NBS in acetone afforded compound **7c**, of which the molecular structure was determined by ¹H NOE spectra.

The Lewis-catalyzed rearrangement of 4*H*-azepines **3–4** to substituted pyridines **5** [eqn (3)] is unprecedented in 4*H*-azepine chemistry.¹⁰ We undertook such novel [4+2]-annulations



Scheme 2 New functionalization of 4*H*-azepines.

Table 4 [4+2]-Annulations between 3-en-1-ynamides and isoxazoles

Entry	(R ¹ , R ² , EWG)	1	(R ³ , R ⁴)	2	Time [h]	Yield [%]	5
(1)	Me, Me, Ts	1b	Me, Me	2a	19	73 (35) ^c	5a (X-ray)
(2)	<i>n</i> -Bu, Me, Ts	1k	Me, Me	2a	33	64	5b
(3)	<i>c</i> -Pr, Me, Ts	1h	Me, Me	2a	20	56	5c
(4)	<i>i</i> -Pr, Me, Ts	1g	Me, Me	2a	15	51	5d
(5)	Me, <i>n</i> -Bu, Ms	1a	Me, Me	2a	28	63	5e
(6)	Me, Me, Ts	1b	<i>n</i> -Bu, <i>n</i> -Bu	2f	19	78	5f
(7)	Me, Me, Ts	1b	Et, Et	2e	16	69	5g
(8)	Me, Me, Ts	1b	<i>n</i> Bu, <i>c</i> -Pr	2h	20	75	5h
(9)	Me, Me, Ts	1b	Ph, Ph	2j	24	80	5i (X-ray)
(10)	Me, Me, Ts	1b	Me, Ph	2k	30	75	5j


^a [1] = 0.15 M. ^b Product yields are reported after separation from a silica column. ^c The value in parentheses is reported using a mixture of IPrAuCl/AgNTf₂ (10 mol%) and Zn(OTf)₂ (20 mol%) in hot DCE (70 °C, 48 h); **3b** was also isolated in 28% yield.





This journal is © The Royal Society of Chemistry 2018

- F. G. West, *Chem. Commun.*, 2009, 5676; (d) M. J. Riveira, L. A. Marsili and M. P. Mischne, *Org. Biomol. Chem.*, 2017, **15**, 9255; (e) N. Shimada, C. Stewart and M. A. Tius, *Tetrahedron*, 2011, **67**, 5851.
- 3 (a) V. A. Guner, K. N. Houk and I. W. Davis, *J. Org. Chem.*, 2004, **69**, 8024; (b) R. V. Essen, D. Frank, H. W. Sunnemann, D. Vidovic, J. Magull and A. D. Meijere, *Chem.-Eur. J.*, 2005, **11**, 6583; (c) N. A. Magomedov, P. L. Ruggiero and Y. Tang, *J. Am. Chem. Soc.*, 2004, **126**, 1624; (d) E. N. Marvell, G. Caple, B. Schatz and W. Pippin, *Tetrahedron*, 1973, **29**, 3781; (e) J. R. Otero, *J. Org. Chem.*, 1999, **64**, 6842; (f) P. E. Tessier, N. Nguyen, M. D. Clay and A. G. Fallis, *Org. Lett.*, 2005, **7**, 767; (g) C. L. Benson and F. G. West, *Org. Lett.*, 2007, **9**, 2545; (h) G. A. Barcan, A. Patel, K. N. Houk and O. Kwon, *Org. Lett.*, 2012, **14**, 5388.
- 4 (a) R. B. Bates, W. H. Delnes, D. A. McCombs and D. E. Potter, *J. Am. Chem. Soc.*, 1969, **91**, 4608; (b) K. Marx and W. Eberbach, *Chem.-Eur. J.*, 2000, **6**, 2063; (c) M. Reisser and G. Mass, *J. Org. Chem.*, 2004, **69**, 4913; (d) T. Hübner, *Dissertation*, University of Freiberg, 1987; (e) A. Arany, D. Bendell, P. W. Groudwater, I. Garnett and M. Nyerges, *J. Chem. Soc., Perkin Trans. 1*, 1999, 2605.
- 5 (a) N. C. Deno, C. U. Pittman and J. O. Turner, *J. Am. Chem. Soc.*, 1965, **87**, 2153; (b) O. N. Faza, C. S. López, R. Alvarez and A. R. de Lera, *Chem.-Eur. J.*, 2009, **15**, 1944.
- 6 D. Alickmann, R. Fröhlich, A. H. Maulitz and E. U. Wurthwein, *Eur. J. Org. Chem.*, 2002, 1523.
- 7 (a) A. H. Zhou, Q. He, C. Shu, Y. F. Yu, S. Liu, T. Zhao, W. Zhang, X. Lu and L. W. Ye, *Chem. Sci.*, 2015, **6**, 1265; (b) X. Y. Xiao, A. H. Zhou, C. Shu, F. Pan, T. Li and L. W. Ye, *Chem.-Asian J.*, 2015, **10**, 1854; (c) W. B. Shen, X. Y. Xiao, Q. Sun, B. Zhou, X. Q. Zhu, J. Z. Yan, X. Lu and L. W. Ye, *Angew. Chem., Int. Ed.*, 2017, **56**, 605; (d) L. Li, T. D. Tan, Y. Q. Zhang, X. Liu and L. W. Ye, *Org. Biomol. Chem.*, 2017, **15**, 8483.
- 8 (a) H. Jin, L. Huang, J. Xie, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, **55**, 794; (b) H. Jin, B. Tian, X. Song, J. Xie, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, **55**, 12688.
- 9 (a) Z. Zeng, H. Jin, J. Xie, B. Tian, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Org. Lett.*, 2017, **19**, 1020; (b) M. Chen, N. Sun, H. Chen and Y. Liu, *Chem. Commun.*, 2016, **52**, 6324; (c) W. Xu, G. Wang, N. Sun and Y. Liu, *Org. Lett.*, 2017, **19**, 3307.
- 10 For synthesis of *nH*-azepines (*n* = 2 or 4), see selected examples: (a) U. Gockel, U. Hartmannsgruber, A. Steigel and J. Sauer, *Tetrahedron Lett.*, 1980, **21**, 599; (b) I. R. Dunkin, A. E. Ayeb and M. A. Lynch, *J. Chem. Soc., Chem. Commun.*, 1994, 1695; (c) K. Satake, R. Okuda, M. Hashimoto, Y. Fujiwara, H. Okamoto, M. Kimura and S. Morosawa, *J. Chem. Soc. Perkin Trans 1*, 1994, 1753; (d) Y. Luo and J. Wu, *Chem. Commun.*, 2011, **47**, 11137.
- 11 Crystallographic data of compounds **3b**, **3l**, **4a**, **5a**, **5i**, and **6m** were deposited in Cambridge Crystallographic Data Center: **3b**: CCDC 1589549, **3l**: CCDC 1589562, **4a**: CCDC 1589561, **5a**: CCDC 1589558, **5i**: CCDC 1589559 and **6m**: CCDC 1589560.†
- 12 For the aza-Nazarov cyclizations, see: (a) D. A. Klumpp, Y. Zhang, M. J. O'Connor, P. M. Esteves and L. S. de Almeida, *Org. Lett.*, 2007, **9**, 3085; (b) Z. X. Ma, S. He, W. Song and R. P. Hsung, *Org. Lett.*, 2012, **14**, 5736; (c) R. L. Sahani and R. S. Liu, *Angew. Chem., Int. Ed.*, 2017, **56**, 12736; (d) S. K. Pawar, R. L. Sahani and R. S. Liu, *Chem.-Eur. J.*, 2015, **21**, 10843; (e) C. Shu, Y. H. Wang, C. H. Shen, P. P. Ruan, X. Lu and L. W. Ye, *Org. Lett.*, 2016, **18**, 3254.
- 13 As suggested by one reviewer, an alternative mechanism is also possible for the Zn(II)-catalyzed rearrangement; this process involves an isomerization of initial species **3** to an unconjugated iminoyl ketone **H**, followed by a 6π -cyclization to generate species **I**. A subsequent Zn(II)-catalyzed aromatization of species **I** is expected to yield the final product **5**. In this process, species **H** is relatively higher than **3** in energy, but its feasibility is not excluded.



- 14 (a) S. M. Wang and L. Zhang, *Org. Lett.*, 2006, **8**, 4585; (b) G. Li, G. Zhang and L. Zhang, *J. Am. Chem. Soc.*, 2008, **130**, 3704; (c) S. B. Wagh and R. S. Liu, *Chem. Commun.*, 2015, **51**, 15462; (d) R. Chaudhuri, A. Das, H. Y. Liao and R. S. Liu, *Chem. Commun.*, 2010, **46**, 4601.
- 15 For reviews of gold-catalyzed cycloaddition or annulation reactions of alkynes, see (a) D. J. Gorin, B. D. Sherry and F. D. Toste, *Chem. Rev.*, 2008, **108**, 3351; (b) A. S. K. Hashmi, *Chem. Rev.*, 2007, **107**, 3180; (c) F. López and J. L. Mascareñas, *Beilstein J. Org. Chem.*, 2011, **7**, 1075; (d) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria and A. Simonneau, *Chem. Rev.*, 2011, **111**, 1954; (e) D. Garayalde and C. Nevado, *ACS Catal.*, 2012, **2**, 1462; (f) M. E. Muratore, A. Homes, C. Obradors and A. M. Echavarren, *Chem.-Asian J.*, 2014, **9**, 3066.

