

## PAPER

 View Article Online  
 View Journal | View Issue
Cite this: *RSC Adv.*, 2018, 8, 5542
 Received 5th December 2017  
 Accepted 27th January 2018

DOI: 10.1039/c7ra13070h

rsc.li/rsc-advances

# Direct C–H functionalization of difluoroboron dipyrromethenes (BODIPYs) at $\beta$ -position by iodonium salts†


 Wenming Ren,<sup>‡a</sup> Huaijiang Xiang,<sup>‡b</sup> Chengyuan Peng,<sup>a</sup> Zulipali Musha,<sup>ac</sup>  
 Jingjing Chen,<sup>ac</sup> Xin Li,<sup>‡b</sup> Ruimin Huang<sup>\*ac</sup> and Youhong Hu<sup>‡ac</sup>

A copper-catalyzed direct C–H arylation or vinylation of BODIPYs at the  $\beta$ -position by iodonium salts has been developed, which provides facile access to a variety of mono-substituted BODIPY dyes. Interestingly,  $\beta$ -styryl BODIPY compound **9b** exhibits apparent cytotoxicity after laser irradiation, which has great potential for photodynamic therapy.

## Introduction

Difluoroboron dipyrromethenes (BODIPYs) and their derivatives are most widely used as small molecule organic fluorophores due to its excellent features, such as high fluorescence quantum yield, narrow emission bandwidth with high peak intensity, good biocompatibility and high photophysical stability.<sup>1</sup> Modification of the BODIPY framework can tune for their fluorescence profile and functionality for the specific application such as bio-imaging and photodynamic therapy (PDT).<sup>2</sup> Besides the meso-derivation for chemosensors, the extension of BODIPY core by the conjugated groups at the  $\alpha$ - or  $\beta$ -position of could lead the advantage of red-shifted emission band and large Stokes shift.<sup>1a,3</sup> Introducing diverse functional groups based on the BODIPY directly is an efficient way to avoid the use of unstable intermediate and multiple-steps processes from pyrroles. Traditionally, the installation of functional conjugated groups has been proceeded by the coupling reaction from halogenated BODIPY or the condensation from active carbon (methyl group at  $\alpha$ -position and formyl group at  $\beta$ -position).<sup>3,4</sup> Direct C–H functionalization of BODIPY chromophores has become an attractive alternative due to the economy and efficiency.<sup>5</sup> Several groups successfully developed the C–H  $\alpha$ -arylation of BODIPYs from the different aryl sources.<sup>6</sup> However, the conjugated  $\beta$ -derivation is rarely explored.

Burgess and coworkers reported a few examples of palladium-catalyzed C–H Heck-type alkenylation at the  $\beta$ -position in 5 days conversion.<sup>7</sup> You and coworkers reported decarboxylative C–H arylation at the  $\beta$ -position from specific *ortho*-substituted benzoic acids in the harsh conditions to obtain the mono- and disubstituted products.<sup>8</sup> Herein, we report a direct C–H arylation or vinylation method for BODIPY postmodification by iodonium salts at the  $\beta$ -position for the mono-substitution as the major product, which could further result the diversity at  $\alpha$ - and  $\beta$ -positions of the BODIPYs (Scheme 1).

Scheme 1 Direct arylation or vinylation of BODIPY at  $\beta$ -position.

<sup>a</sup>State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zuchongzhi Road, Shanghai 201203, China. E-mail: yphu@simm.ac.cn; rmhuang@simm.ac.cn

<sup>b</sup>College of Pharmaceutical Sciences, Zhejiang University, 866 Yuhangtang Road, Hangzhou 310058, China

<sup>c</sup>University of Chinese Academy of Sciences, 19 Yuquan Road, Beijing 110039, China

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c7ra13070h

‡ These authors contributed equally.






Table 2 Photophysical properties of BODIPY dyes in CH<sub>3</sub>CN at room temperature

| Dyes | $\lambda_{\text{abs}}^a$ (nm) | $\epsilon$ (M <sup>-1</sup> cm <sup>-1</sup> ) | $\lambda_{\text{ex}}^b$ (nm) | $\lambda_{\text{em}}^b$ (nm) | $\Phi_F^{a,c}$ | Stokes shift (nm) |
|------|-------------------------------|------------------------------------------------|------------------------------|------------------------------|----------------|-------------------|
| 1    | 492                           | 83 400                                         | 491                          | 504                          | 0.98           | 13                |
| 3a   | 503                           | 86 500                                         | 502                          | 529                          | 0.71           | 27                |
| 3b   | 504                           | 79 500                                         | 503                          | 535                          | 0.70           | 32                |
| 3c   | 504                           | 35 200                                         | 503                          | 530                          | 0.78           | 27                |
| 3e   | 504                           | 78 700                                         | 504                          | 551                          | 0.24           | 47                |
| 3f   | 501                           | 72 000                                         | 501                          | 523                          | 0.86           | 22                |
| 3g   | 504                           | 83 100                                         | 502                          | 527                          | 0.91           | 25                |
| 3h   | 502                           | 81 600                                         | 502                          | 527                          | 0.70           | 25                |
| 3i   | 503                           | 69 600                                         | 502                          | 527                          | 0.66           | 25                |
| 3j   | 502                           | 80 300                                         | 502                          | 530                          | 0.77           | 28                |
| 3k   | 502                           | 85 100                                         | 501                          | 525                          | 0.80           | 24                |
| 5    | 519                           | 65 400                                         | 517                          | 574                          | 0.41           | 57                |
| 7    | 509                           | 65 400                                         | 510                          | 534                          | 0.67           | 24                |

<sup>a</sup> Data were measured in a concentration of  $3.0 \times 10^{-6}$  M. <sup>b</sup> Data were measured in a concentration of  $5.0 \times 10^{-6}$  M. <sup>c</sup> All fluorescence quantum yields ( $\Phi$ ) were calculated using fluorescein in 0.1 N NaOH solution ( $\Phi = 0.91$ , excitation = 488 nm) as the standard except 4l using Rhodamine 6G in anhydrous ethanol ( $\Phi = 0.95$ , excitation = 530 nm) as the standard.




Scheme 3 Direct C–H arylation of BODIPY 1 with different diaryliodonium triflates. <sup>a</sup> Condition A: 0.8 equiv. of 2 was used; <sup>b</sup> condition B: 2.4 equiv. of 2 was used; <sup>c</sup> isolated yield.

the asymmetrical diarylated BODIPY 5 containing D- $\pi$ -A (donor- $\pi$ -acceptor) structure was obtained in 52% yield. In addition, *meso*-phenyl BODIPY 6 could react with diaryliodonium 2a (Scheme 4B) under the condition A to achieve product 7 in 46% yield.


The photophysical properties of 3a–3k, 5 and 7 were evaluated in different solvents system (Table 2, Fig. S1–5†). As shown in Table 2, 2-aryl BODIPYs 3a–3k exhibited the absorption maxima and excitation maxima among 501–504 nm no matter electron-withdrawing or electron-donating substituents. All 2-aryl BODIPYs showed high absorption coefficients and relatively high fluorescence quantum yields. Methoxyl substituent 3e showed a lower fluorescence quantum yield (~24%) correspondingly. On the other hand, 3e exhibited the longest emission maxima at 551 nm and the largest Stokes shift (47 nm) among 2-aryl BODIPYs. These data indicate that slight modification of BODIPY could change their photophysical property significantly. The asymmetrical diarylated BODIPY 5 containing D- $\pi$ -A has the desired properties with red fluorescence, high quantum yield and large Stokes shift.

Next, we explored the reaction of BODIPY 1 with the variety of styryliodonium triflates 8 to generate the different conjugated



Scheme 4 Extension for  $\beta$ -arylation of BODIPY.





Scheme 5 Scope of the direct C–H vinylation of BODIPY 1 with different vinyl iodonium triflates.<sup>a</sup>Isolated yield.

systems. Due to the higher reactivity of vinyl iodonium than aryl iodonium salts,<sup>12b</sup> the reaction temperature at 50 °C gave the similar efficiency at 70 °C. To our delight, the different 2-styryl BODIPYs **9a–9e** were delivered (Scheme 5). Additionally, alkyl vinyl iodonium triflates can also be successfully applied in this reaction, maintaining 51% yield of 2-vinyl BODIPY **9f**.

Photophysical properties of **9a–9f** were investigated as shown in Table 3. The 2-vinyl BODIPYs **9** exhibited emission maxima up to 598–620 nm with large Stokes shifts. Probably due to the geometry relaxation of fluorophores upon photoexcitation, the quantum yields of those dyes were as low as 0.78%. Dye with photoproperties such as high extinction coefficient, low quantum yield and red emission could be used as photosensitizer for photodynamic therapy.<sup>2a,2d</sup> The BODIPY **9b** was thus chosen to evaluate photocytotoxic activity *in vitro*. The half maximal inhibitory concentration ( $\text{IC}_{50}$ ) of compound **9b** with or without light irradiation was assessed in human breast cancer cell line MDA-MB-231 and epidermoid carcinoma cell line A-431. Nine different concentrations (ranging from 0.195 to 50  $\mu\text{M}$ ) of **9b** were applied to cells and 12 h exposure to mercury lamp decreased the cell viability significantly in presence of **9b** ( $P < 0.0001$ ) in both cells (Fig. 1). For example, light irradiation reduced the  $\text{IC}_{50}$  value from 43.9  $\mu\text{M}$  to 4.0  $\mu\text{M}$  (~88% decline)




Fig. 1 Light irradiation (IR)-induced cytotoxicity by compound **9b** in MDA-MB-231 and A-431 cells using SRB assay. \* $P < 0.0001$ .




Fig. 2 Overlay of fluorescent and differential interference contrast (DIC) images of MDA-MB-231 cells incubated with of compound **9b** or vehicle control. Blue circles, laser irradiated regions; grey boxes, regions very near the laser-irradiated areas. Scale bars, 50  $\mu\text{m}$ .

in MDA-MB-231 cells. Photophysical property and laser-induced cytotoxicity of compound **9b** were also characterized and visualized by a Leica confocal microscopy. As shown in Fig. 2 and S7,<sup>†</sup> high fluorescence signals were detected in cytoplasm within 5 min after **9b** (2.5  $\mu\text{M}$ ) feeding. Then the photothermal treatment was performed by continuous irradiation of the cells using a 552 nm laser for 10 min, leading to a ~200  $\mu\text{m}$  diameter irradiation spot. Cell blebbing, a characteristic feature of injured cells, was only observed in compound **9b**-treated cells but not in the vehicle-treated cells. Similar photothermal effects were exhibited in **9a**- or **9f**-treated cells (Fig. S7<sup>†</sup>). These results suggest that 2-vinyl BODIPYs could be excellent photosensitizers and have potential for photodynamic therapy of cancer.

Table 3 Photophysical properties of BODIPY dyes in  $\text{CH}_3\text{CN}$  at room temperature

| Dyes | $\lambda_{\text{abs}}^a$ (nm) | $\epsilon$ ( $\text{M}^{-1} \text{cm}^{-1}$ ) | $\lambda_{\text{ex}}^b$ (nm) | $\lambda_{\text{em}}^b$ (nm) | $\Phi_F^{a,c}$ | Stokes shift (nm) |
|------|-------------------------------|-----------------------------------------------|------------------------------|------------------------------|----------------|-------------------|
| 9a   | 523                           | 50 100                                        | 517                          | 608                          | 0.0080         | 91                |
| 9b   | 520                           | 38 100                                        | 519                          | 620                          | 0.0078         | 101               |
| 9c   | 522                           | 52 400                                        | 505                          | 600                          | 0.0091         | 95                |
| 9d   | 520                           | 70 600                                        | 518                          | 610                          | 0.011          | 92                |
| 9e   | 523                           | 47 700                                        | 521                          | 607                          | 0.010          | 86                |
| 9f   | 511                           | 43 100                                        | 535                          | 598                          | 0.063          | 63                |

<sup>a</sup> Data were measured in a concentration of  $3.0 \times 10^{-6} \text{ M}$ . <sup>b</sup> Data were measured in a concentration of  $5.0 \times 10^{-6} \text{ M}$ . <sup>c</sup> The fluorescence quantum yields ( $\Phi$ ) were calculated using rhodamine 6G in anhydrous ethanol ( $\Phi = 0.95$ , excitation = 530 nm).



## Conclusions

In conclusion, we have developed the Cu-catalyzed direct C–H arylation and vinylation of BODIPYs at the  $\beta$ -position. This methodology provides a convenient synthetic procedure to prepare diversified BODIPY derivatives. These reactions exhibited good selectivity for mono-substituted product to allow the different modification of BODIPY at 2,6-position. Through the evaluation of photophysical properties of those dyes, 2-styryl BODIPYs showed intrinsic large Stokes shift up to 101 nm and low quantum yield, suggesting the potential for photodynamic therapy of cancer. In this way, the novel functionality of fluorophores with asymmetrical substitution can be synthesized economically and efficiently.

## Experimental

### General

All reactions were carried out under an atmosphere of argon in oven-dried glassware with magnetic stirring. All commercially available reagents were used as received. Chromatographic purifications were performed by flash chromatography with silica gel (40–63  $\mu$ m) packed in glass columns. The eluting solvent for the purification of each compound was determined by thin-layer chromatography (TLC) on glass plates coated with silica gel 60 F254 and visualized by UV light (254 nm or 365 nm).  $^1\text{H}$  NMR spectra and proton-decoupled  $^{13}\text{C}$  NMR spectra were obtained on a 400 MHz or 500 MHz Bruker NMR spectrometer.  $^1\text{H}$  chemical shifts ( $\delta$ ) are reported in parts per million (ppm) relative to TMS (s,  $\delta$  0).  $^{13}\text{C}$  NMR chemical shifts are reported relative to  $\text{CDCl}_3$  (t,  $\delta$  77.4). High-resolution mass data were obtained on an Agilent 6224 accurate-mass TOF LC/MS (ESI). Absorption spectra were acquired using a Varian Cary 300 spectrophotometer. Fluorescence measurements were carried out on a Horiba FluoroMax-4 spectrometer. Quantum yields were determined in reference to either fluorescein or rhodamine 6G and corrected for solvent refractive index. The extinction coefficients were determined through Beer's law plots. All data were measured at room temperature. Human breast cancer cell line MDA-MB-231 and human epidermoid carcinoma cell line A-431 were obtained from the American Type Culture Collection (Manassas, VA, USA) and were cultured in DMEM (high glucose) and RPMI 1640 medium respectively, supplemented with 10% fetal bovine serum (Hyclone, Logan, UT). Cells were incubated at 37 °C in 5%  $\text{CO}_2$  in air.

### General procedure for the Cu(II)-catalyzed mono-arylation or vinylation of BODIPYs (reaction condition A)

Copper(II) trifluoromethanesulfonate (7.23 mg, 0.02 mmol) was adequately suspended in 2 mL mixture solution (dichloromethane/toluene = 3/1, dried by 4 Å molecular sieves) in 10 mL sealed tube, then **1** (62.91 mg, 0.24 mmol), appropriate diaryliodonium salt (0.2 mmol) and 2,6-di-*tert*-butylpyridine (54  $\mu$ L, 0.24 mmol) were sequentially added to the solution. Seal up the tube with argon atmosphere. The resulting solution was stirred at 50 °C (R = styryl) or 70 °C (R = aryl group) for 6 h. The solution was concentrated and purified by silica gel column chromatography (hexane: dichloromethane: acetic ether = 300 : 100 : 0 to

300 : 100 : 5) to afford crude product. Single and double substituted products were validated by 400 MHz or 500 MHz Bruker NMR spectrometer. The residue was purified by TLC silica gel plate (10  $\times$  200  $\times$  200 mm, 10–40  $\mu$ m, separating solvent and ratio: dichloromethane/toluene = 1/5) or reversed phase chromatography (separating solvent and ratio:  $\text{H}_2\text{O}/\text{MeOH}$  = 15/85 to 5/95) to afford pure products **3a–k**, **5**, **7** and **9a–g**. The yield data was calculated by diaryliodonium stoichiometric equivalence.

### General procedure for the Cu(II)-catalyzed diarylation of BODIPYs (reaction condition B)

Copper(II) trifluoromethanesulfonate (14.5 mg, 0.04 mmol) was adequately suspended in 3 mL mixture solution in 10 mL sealed tube, then **1** (52.4 mg, 0.20 mmol), appropriate diaryliodonium salt (0.48 mmol) and 2,6-di-*tert*-butylpyridine (109  $\mu$ L, 0.48 mmol) were sequentially added to the solution. Seal up the tube with argon atmosphere. The purification procedures were the same with reaction condition A to afford **4a–4k**. The yield data was calculated by BODIPY **1** stoichiometric equivalence.

### Cell viability assay for photocytotoxic effect

Cells were seeded in 96-well plates at a density of 1600–3000 cells per well in triplicate. Twenty-four hours later, the cells were incubated with fresh medium with compounds at different concentrations (ranging from 0.195 to 50  $\mu$ M). The cells were exposed under the mercury lamp (HXP R120W/45C; Osram, Germany) for 12 h, and then were kept under normal culture condition for another 72 h. At the endpoint, the cells were fixed with 10% pre-cooled trichloroacetic acid (Sigma, St Louis, MO, USA) over 4 h followed by staining with 4 mg  $\text{mL}^{-1}$  sulforhodamine B (SRB; Sigma) in 1% acetic acid for 15 min. SRB in the cells was dissolved in 10 mmol  $\text{L}^{-1}$  Tris-HCl and measured at 560 nm using a microplate reader (SpectraMAX190; Molecular Devices, Sunnyvale, CA, USA). The half maximal inhibitory concentration ( $\text{IC}_{50}$ ) was decided using (log(inhibitor) vs. response – variable slope (four parameters)) method in the GraphPad Prism Software. The data were presented as the mean  $\pm$  SD. Differences were considered statistically significant at  $P < 0.0001$  by 2 way ANOVA test.

### Visualization of laser-induced cytotoxicity

Cells were grown on the 8-well chambered coverglass (Thermo Fisher Scientific, Waltham, MA, USA) for 36 h. Immediately after adding the compound with indicated concentrations to cells, a small region of interest (ROI) was randomly chosen and irradiated using the laser line at 552 nm from a confocal microscopy (Leica TCS SP8 STED, Germany) over 10 min. During that period, the fluorescent and differential interference contrast (DIC) images were acquired at 0 min, 5 min and 10 min, respectively. The power of the laser was kept at 80% output to ensure the consistent irradiation between experiments.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2-phenyl-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f'*][1,3,2]diazaborinine(3a).** An orange solid (yield: 35.9 mg, 53%, single substitute: double substitute = 91 : 9),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.43 (t,  $J$  = 7.5 Hz, 2H), 7.34 (t,  $J$  = 7.5 Hz, 1H), 7.21 (d,  $J$  = 6.9 Hz, 2H), 6.08 (s, 1H), 2.65 (s, 3H),



2.54 (s, 3H), 2.47 (s, 3H), 2.44 (s, 3H), 2.32 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  153.8, 152.2, 141.6, 141.1, 137.0, 133.9, 133.5, 132.4, 131.9, 130.4, 128.4, 127.1, 121.4, 17.5, 16.8, 15.4, 14.5, 13.2. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{20}\text{H}_{22}^{11}\text{BF}_2\text{N}_2^+$ : 339.1839; found: 339.1841,  $\Delta = -0.66$  ppm.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2-(*p*-tolyl)-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2]diazaborinine (3b).** An orange solid (yield: 25.9 mg, 37%, single substitute: double substitute = 93 : 7),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.24 (d,  $J = 7.7$  Hz, 2H), 7.10 (d,  $J = 7.7$  Hz, 2H), 6.07 (s, 1H), 2.64 (s, 3H), 2.54 (s, 3H), 2.47 (s, 3H), 2.43 (s, 3H), 2.41 (s, 3H), 2.32 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  153.5, 152.4, 141.5, 140.8, 137.2, 136.8, 133.5, 132.3, 131.9, 130.8, 130.3, 129.1, 121.3, 21.3, 17.5, 16.8, 15.4, 14.5, 13.3. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{21}\text{H}_{24}^{11}\text{BF}_2\text{N}_2^+$ : 353.1995; found: 353.1998,  $\Delta = -0.78$  ppm.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2-(*m*-tolyl)-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2]diazaborinine (3c).** An orange solid (yield: 24.2 mg, 34%, single substitute: double substitute = 91 : 9),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.31 (t,  $J = 7.5$  Hz, 1H), 7.16 (d,  $J = 7.6$  Hz, 1H), 7.00 (d,  $J = 11.3$  Hz, 2H), 6.08 (s, 1H), 2.65 (s, 3H), 2.54 (s, 3H), 2.47 (s, 3H), 2.44 (s, 3H), 2.40 (s, 3H), 2.32 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  153.6, 152.4, 141.5, 140.8, 138.0, 137.1, 133.8, 132.3, 131.9, 131.0, 128.2, 127.9, 127.5, 121.3, 21.5, 17.4, 16.8, 15.4, 14.5, 13.2. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{21}\text{H}_{24}^{11}\text{BF}_2\text{N}_2^+$ : 353.1995; found: 353.1992,  $\Delta = 0.92$  ppm.

**5,5-Difluoro-2-(4-methoxyphenyl)-1,3,7,9,10-pentamethyl-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo [1,2-*c*:2',1'-*f*] [1,3,2] diazaborinine (3e).** An orange solid (yield: 36.9 mg, 50%, single substitute: double substitute = 86 : 14),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.16–7.10 (m, 2H), 6.99–6.95 (m, 2H), 6.07 (s, 1H), 3.86 (s, 3H), 2.64 (s, 3H), 2.54 (s, 3H), 2.47 (s, 3H), 2.43 (s, 3H), 2.31 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  158.7, 153.5, 152.6, 141.46, 140.8, 137.2, 133.3, 132.3, 131.9, 131.5, 126.0, 121.2, 113.9, 55.3, 17.4, 16.8, 15.4, 14.5, 13.2. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{21}\text{H}_{24}^{11}\text{BF}_2\text{N}_2\text{O}^+$ : 369.1944; found: 369.1941,  $\Delta = 0.95$  ppm.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2-(4-(trifluoromethyl)phenyl)-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo [1,2-*c*:2',1'-*f*] [1,3,2] diazaborinine (3f).** An orange solid (yield: 29.5 mg, 36%, single substitute: double substitute = 96 : 4),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.69 (d,  $J = 8.0$  Hz, 2H), 7.33 (d,  $J = 8.0$  Hz, 2H), 6.12 (s, 1H), 2.66 (s, 3H), 2.55 (s, 3H), 2.47 (s, 3H), 2.45 (s, 3H), 2.32 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  155.1, 150.8, 142.0, 141.9, 137.9, 136.5, 132.8, 131.6, 130.7, 129.1, 128.7, 125.3, 123.1, 122.0, 17.5, 16.8, 15.3, 14.5, 13.1. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{21}\text{H}_{21}^{11}\text{BF}_5\text{N}_2^+$ : 407.1712; found: 407.1720,  $\Delta = -1.97$  ppm.

**Methyl 4-(5,5-difluoro-1,3,7,9,10-pentamethyl-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2]diazaborin-2-yl)benzoate (3g).** An orange solid (yield: 44.4 mg, 56%, single substitute: double substitute = 96 : 4),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.10 (d,  $J = 8.1$  Hz, 2H), 7.29 (d,  $J = 8.1$  Hz, 2H), 6.11 (s, 1H), 3.95 (s, 3H), 2.65 (s, 3H), 2.55 (s, 3H), 2.48 (s, 3H), 2.45 (s, 3H), 2.33 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  167.0, 154.9, 151.0, 141.9, 139.0, 136.6, 132.8, 132.1, 131.8, 130.4, 129.6, 128.8, 121.9, 52.2, 17.5, 16.9, 15.3, 14.5, 13.2. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{22}\text{H}_{24}^{11}\text{BF}_2\text{N}_2\text{O}_2^+$ : 397.1893; found: 397.1882,  $\Delta = 2.91$  ppm.

**5,5-Difluoro-2-(4-fluorophenyl)-1,3,7,9,10-pentamethyl-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*] [1,3,2] diazaborinine (3h).** An

orange solid (yield: 29.3 mg, 41%, single substitute: double substitute = 95 : 5),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.19–7.09 (m, 4H), 6.09 (s, 1H), 2.64 (s, 3H), 2.54 (s, 3H), 2.45 (s, 3H), 2.44 (s, 3H), 2.30 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  163.0, 161.1, 154.3, 151.7, 141.7, 141.4, 136.9, 132.5, 132.3, 132.0, 130.0, 131.7, 129.8, 121.6, 115.5, 17.5, 16.8, 15.3, 14.5, 13.1. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{20}\text{H}_{22}^{11}\text{BF}_3\text{N}_2^+$ : 357.1744; found: 357.1752,  $\Delta = -2.01$  ppm.

**2-(4-Chlorophenyl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*] [1,3,2] diazaborinine (3i).** An orange solid (yield: 32.3 mg, 43%, single substitute: double substitute = 94 : 6),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.40 (d,  $J = 8.3$  Hz, 2H), 7.14 (d,  $J = 8.3$  Hz, 2H), 6.10 (s, 1H), 2.64 (s, 3H), 2.54 (s, 3H), 2.45 (s, 3H), 2.44 (s, 3H), 2.30 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  154.5, 151.4, 141.8, 141.6, 136.7, 133.1, 132.6, 132.4, 132.0, 131.7, 128.6, 121.7, 17.5, 16.8, 15.3, 14.5, 13.1. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{20}\text{H}_{21}^{11}\text{BClF}_2\text{N}_2^+$ : 373.1449; found: 373.1452,  $\Delta = -0.33$  ppm.

**2-(4-Bromophenyl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*] [1,3,2] diazaborinine (3j).** An orange solid (yield: 28.9 mg, 35%, single substitute: double substitute = 93 : 7),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.55 (d,  $J = 8.3$  Hz, 2H), 7.07 (d,  $J = 8.3$  Hz, 2H), 6.09 (s, 1H), 2.63 (s, 3H), 2.54 (s, 3H), 2.45 (s, 3H), 2.43 (s, 3H), 2.30 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  154.8, 151.5, 142.0, 141.9, 136.9, 133.1, 132.9, 132.3, 131.9, 131.8, 122.0, 121.5, 17.7, 17.1, 15.5, 14.7, 13.3. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{20}\text{H}_{21}^{11}\text{BClF}_2\text{N}_2^+$ : 373.1449; found: 373.1452,  $\Delta = -0.33$  ppm.

**2-(3-Bromophenyl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5*H*-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*] [1,3,2] diazaborinine (3k).** An orange solid (yield: 29.3 mg, 35%, single substitute: double substitute = 90 : 10),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.48 (d,  $J = 7.9$  Hz, 1H), 7.37 (s, 1H), 7.30 (t,  $J = 7.8$  Hz, 1H), 7.14 (d,  $J = 7.5$  Hz, 1H), 6.11 (s, 1H), 2.65 (s, 3H), 2.55 (s, 3H), 2.47 (s, 3H), 2.45 (s, 3H), 2.32 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  154.8, 151.2, 141.9, 141.8, 136.7, 136.2, 133.3, 132.7, 131.8, 131.6, 130.2, 129.9, 129.1, 122.4, 121.8, 17.5, 16.9, 15.3, 14.6, 13.1. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{20}\text{H}_{21}^{11}\text{BrF}_2\text{N}_2^+$ : 417.0944; found: 417.943,  $\Delta = -0.28$  ppm.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2,8-diphenyl-5*H*-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2]diazaborinine (4a).** An orange solid, (yield: 41.4 mg, 50%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.47 (t,  $J = 7.5$  Hz, 4H), 7.38 (t,  $J = 7.5$  Hz, 2H), 7.26 (d,  $J = 7.0$  Hz, 4H), 2.75 (s, 3H), 2.53 (s, 6H), 2.38 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  152.4, 141.8, 137.1, 133.7, 132.2, 130.4, 128.4, 127.1, 17.3, 15.5, 13.3. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{26}\text{H}_{26}^{11}\text{BF}_2\text{N}_2^+$ : 415.2152; found: 415.2153,  $\Delta = -0.4$  ppm.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2,8-di-*p*-tolyl-5*H*-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2]diazaborinine (4b).** An orange solid, (yield: 15.2 mg, 17%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.25 (d,  $J = 7.9$  Hz, 5H), 7.11 (d,  $J = 7.9$  Hz, 4H), 2.70 (s, 3H), 2.49 (s, 6H), 2.41 (s, 6H), 2.34 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  152.4, 141.5, 137.0, 136.8, 133.5, 132.2, 130.8, 130.3, 129.1, 77.3, 77.0, 76.8, 29.7, 21.3, 17.2, 15.5, 13.3. HRMS (ESI):  $m/z$   $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{28}\text{H}_{30}^{11}\text{BF}_2\text{N}_2^+$ : 443.2465; found: 443.2461,  $\Delta = 0.86$  ppm.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2,8-di-*m*-tolyl-5*H*-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2]diazaborinine (4c).** An orange



solid, (yield: 20.5 mg, 23%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.33 (t,  $J = 7.5$  Hz, 2H), 7.17 (d,  $J = 7.5$  Hz, 2H), 7.05–7.00 (m, 4H), 2.71 (s, 3H), 2.49 (s, 6H), 2.41 (s, 6H), 2.35 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  152.3, 141.5, 137.9, 137.0, 133.7, 132.2, 131.0, 128.2, 127.8, 127.4, 21.5, 17.2, 15.5, 13.3. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{28}\text{H}_{29}^{11}\text{BFN}_2^+$ : 423.2402; found: 423.2409,  $\Delta = -1.64$  ppm.

**5,5-Difluoro-2,8-bis(4-methoxyphenyl)-1,3,7,9,10-pentamethyl-5H-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (4e).** An orange solid, (yield: 17.7 mg, 19%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.14 (d,  $J = 8.7$  Hz, 4H), 6.98 (d,  $J = 8.7$  Hz, 4H), 3.86 (s, 6H), 2.70 (s, 3H), 2.48 (s, 6H), 2.33 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  158.7, 152.4, 141.4, 137.0, 133.2, 132.1, 131.5, 126.0, 113.9, 55.3, 17.2, 15.5, 13.3. HRMS (ESI):  $m/z$  [ $\text{M} + \text{H}$ ] $^+$  calcd for  $\text{C}_{28}\text{H}_{29}^{11}\text{BF}_2\text{N}_2\text{O}_2^+$ : 455.2301; found: 455.2307,  $\Delta = -1.46$  ppm.

**5,5-Difluoro-1,3,7,9,10-pentamethyl-2,8-bis(4-(trifluoromethyl)phenyl)-5H-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (4f).** An orange solid, (yield: 21.1 mg, 19%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.71 (d,  $J = 7.9$  Hz, 4H), 7.36 (d,  $J = 7.9$  Hz, 4H), 2.74 (s, 3H), 2.50 (s, 6H), 2.36 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  152.4, 142.6, 137.6, 137.5, 132.4, 130.7, 129.5, 129.3, 125.4, 123.1, 17.4, 15.5, 13.3. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{28}\text{H}_{23}^{11}\text{BF}_7\text{N}_2^+$ : 531.1837; found: 531.1832,  $\Delta = 0.98$  ppm.

**Dimethyl 4,4'-(5,5-difluoro-1,3,7,9,10-pentamethyl-5H-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine-2,8-diyl) dibenzoate (4g).** An orange solid, (yield: 29.8 mg, 28%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.12 (d,  $J = 7.9$  Hz, 4H), 7.32 (d,  $J = 7.9$  Hz, 4H), 3.96 (s, 6H), 2.74 (s, 3H), 2.51 (s, 6H), 2.37 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  166.9, 152.4, 142.4, 138.7, 137.4, 132.8, 132.5, 130.4, 129.7, 128.9, 52.2, 17.4, 15.6, 13.3. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{30}\text{H}_{29}^{11}\text{BFN}_2\text{O}_4^+$ : 511.2199; found: 511.2196,  $\Delta = 0.55$  ppm.

**5,5-Difluoro-2,8-bis(4-fluorophenyl)-1,3,7,9,10-pentamethyl-5H-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (4h).** An orange solid, (yield: 18.2 mg, 20%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.16 (m, 8H), 2.71 (s, 3H), 2.47 (s, 6H), 2.32 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  163.1, 161.1, 152.4, 142.0, 137.3, 132.7, 132.1, 129.7, 115.5, 115.4, 17.3, 15.5, 13.2. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{26}\text{H}_{24}^{11}\text{BF}_3\text{N}_2^+$ : 431.2972; found: 431.2970,  $\Delta = 0.46$  ppm.

**2,8-Bis(4-chlorophenyl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5H-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (4i).** An orange solid, (yield: 49.3 mg, 51%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.42 (d,  $J = 8.4$  Hz, 4H), 7.16 (d,  $J = 8.4$  Hz, 4H), 2.71 (s, 3H), 2.48 (s, 6H), 2.33 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  152.3, 142.1, 137.3, 133.3, 132.5, 132.2, 131.7, 128.7, 17.3, 15.5, 13.2. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{26}\text{H}_{23}^{11}\text{BCl}_2\text{FN}_2^+$ : 463.1310; found: 463.1300,  $\Delta = -2.09$  ppm.

**2,8-Bis(4-bromophenyl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5H-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (4j).** An orange solid, (yield: 27.2 mg, 24%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.57 (d,  $J = 8.4$  Hz, 4H), 7.10 (d,  $J = 8.4$  Hz, 4H), 2.71 (s, 3H), 2.48 (s, 6H), 2.33 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  152.3, 142.1, 137.2, 132.7, 132.5, 132.3, 132.0, 131.7, 121.4, 17.3, 15.5, 13.2. HRMS (ESI):  $m/z$  [ $\text{M} + \text{H}$ ] $^+$  calcd for  $\text{C}_{26}\text{H}_{23}^{11}\text{B}^{79}\text{Br}_2\text{F}_2\text{N}_2^+$ : 570.0284; found: 570.0285,  $\Delta = 0.26$  ppm.

**2,8-Bis(3-bromophenyl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5H-4 $\lambda$ ,5 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (4k).** An

orange solid, (yield: 22.3 mg, 20%)  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.54–7.50 (m,  $J = 7.8$  Hz, 2H), 7.41 (t, 2H), 7.34 (t,  $J = 7.8$  Hz, 2H), 7.19 (d,  $J = 7.8$  Hz, 2H), 2.74 (s, 3H), 2.51 (s, 6H), 2.37 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  152.4, 142.3, 137.4, 136.0, 133.2, 132.3, 130.3, 130.0, 129.1, 122.5, 17.3, 15.5, 13.2. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{26}\text{H}_{23}^{11}\text{B}^{79}\text{Br}_2\text{FN}_2^+$ : 551.0300; found: 551.0293,  $\Delta = -1.20$  ppm.

**Methyl-4-(5,5-difluoro-8-(4-methoxyphenyl)-1,3,7,9,10-pentamethyl-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine-2-yl)benzoate (5).** An orange solid (second step's yield: 52%).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.11 (d,  $J = 7.3$  Hz, 2H), 7.32 (d,  $J = 7.4$  Hz, 2H), 7.14 (d,  $J = 7.6$  Hz, 2H), 6.99 (d,  $J = 7.7$  Hz, 2H), 3.95 (s, 3H), 3.86 (s, 3H), 2.72 (s, 3H), 2.50 (s, 6H), 2.35 (s, 6H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  166.6, 158.4, 153.6, 150.5, 141.4, 138.6, 137.5, 135.9, 133.5, 132.2, 131.7, 131.6, 131.0, 130.0, 129.2, 128.3, 125.3, 113.5, 54.9, 51.8, 16.9, 15.2, 15.0, 13.0, 12.8. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{29}\text{H}_{29}^{11}\text{BFN}_2\text{O}_3^+$ : 483.2250; found: 483.2258,  $\Delta = 1.66$  ppm.

**5,5-Difluoro-1,3,7,9-tetramethyl-2,10-diphenyl-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (7).** An orange solid (yield: 37.0 mg, 46%),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.55–7.48 (m, 3H), 7.40 (t,  $J = 7.4$  Hz, 2H), 7.37–7.29 (m, 3H), 7.19–7.15 (m, 2H), 6.03 (s, 1H), 2.61 (s, 3H), 2.55 (s, 3H), 1.41 (s, 3H), 1.32 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  155.6, 154.0, 143.2, 142.0, 139.2, 135.2, 133.7, 133.7, 131.7, 131.0, 130.2, 129.2, 129.0, 128.3, 128.0, 127.0, 121.4, 14.6, 14.4, 13.3, 12.7. HRMS (ESI):  $m/z$  [ $\text{M} + \text{H}$ ] $^+$  calcd for  $\text{C}_{25}\text{H}_{24}^{11}\text{BF}_2\text{N}_2^+$ : 401.2001; found: 401.2008,  $\Delta = 1.20$  ppm.

**(E)-5,5-Difluoro-1,3,7,9,10-pentamethyl-2-styryl-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (9a).** A brownish red solid (yield: 27.7 mg, 38%, single substitute: double substitute = 98 : 2).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.48 (d,  $J = 7.3$  Hz, 2H), 7.36 (t,  $J = 7.3$  Hz, 2H), 7.26 (t, 1H), 6.95 (d,  $J = 16.5$  Hz, 1H), 6.68 (d,  $J = 16.5$  Hz, 1H), 6.08 (s, 1H), 2.67 (s, 3H), 2.64 (s, 3H), 2.53 (s, 3H), 2.50 (s, 3H), 2.43 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  154.0, 152.9, 141.3, 141.2, 137.8, 136.6, 132.5, 132.0, 131.9, 128.7, 128.5, 127.5, 126.1, 121.5, 119.9, 77.3, 77.0, 76.8, 17.5, 17.0, 15.3, 14.5, 13.8. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{22}\text{H}_{23}^{11}\text{BFN}_2^+$ : 345.1933; found: 345.1935,  $\Delta = -0.73$  ppm.

**(E)-5,5-Difluoro-1,3,7,9,10-pentamethyl-2-(4-methylstyryl)-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (9b).** A brownish red solid (yield: 18.6 mg, 25%, single substitute: double substitute = 95 : 5).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.38 (d,  $J = 8.0$  Hz, 2H), 7.17 (d,  $J = 8.0$  Hz, 2H), 6.90 (d,  $J = 16.5$  Hz, 1H), 6.65 (d,  $J = 16.5$  Hz, 1H), 6.07 (s, 1H), 2.66 (s, 3H), 2.64 (s, 3H), 2.53 (s, 3H), 2.49 (s, 3H), 2.43 (s, 3H), 2.36 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  153.8, 153.1, 141.2, 141.0, 137.4, 136.5, 135.0, 132.5, 132.1, 129.4, 128.8, 126.1, 121.4, 118.9, 110.0, 21.2, 17.5, 16.9, 15.3, 14.5, 13.8. HRMS (ESI):  $m/z$  [ $\text{M} - \text{F}$ ] $^+$  calcd for  $\text{C}_{23}\text{H}_{25}^{11}\text{BFN}_2^+$ : 359.2089; found: 359.2098,  $\Delta = -2.39$  ppm.

**(E)-5,5-Difluoro-2-(4-fluorostyryl)-1,3,7,9,10-pentamethyl-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*][1,3,2] diazaborinine (9c).** A brownish red solid (yield: 16.6 mg, 22%, single substitute: double substitute = 96 : 4).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.47 (dd,  $J = 8.6$  Hz, 2H), 7.07 (t,  $J = 8.6$  Hz, 2H), 6.89 (d,  $J = 16.5$  Hz, 1H), 6.66 (d,  $J = 16.5$  Hz, 1H), 6.11 (s, 1H), 2.68 (s, 3H), 2.67 (s, 3H), 2.56 (s, 3H), 2.52 (s, 3H), 2.46 (s, 3H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  163.3, 161.3, 154.2, 152.7, 141.3, 136.5, 133.9, 132.6, 133.0, 130.6,



128.3, 127.6, 121.6, 119.7, 115.7, 115.5, 17.5, 17.0, 15.3, 14.5, 13.8. HRMS (ESI):  $m/z$   $[M - F]^+$  calcd for  $C_{22}H_{22}^{11}BF_2N_2^+$ : 363.1839; found: 363.1834,  $\Delta = 1.15$  ppm.

(*E*)-2-(4-Chlorostyryl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*] [1,3,2]diazaborinine (9d). A brownish red solid (yield: 53.4 mg, 67%, single substitute: double substitute = 96 : 4).  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.39 (d,  $J = 8.5$  Hz, 2H), 7.31 (d,  $J = 8.5$  Hz, 2H), 6.92 (d,  $J = 16.5$  Hz, 1H), 6.62 (d,  $J = 16.5$  Hz, 1H), 6.08 (s, 1H), 2.66 (s, 3H), 2.63 (s, 3H), 2.53 (s, 3H), 2.48 (s, 3H), 2.42 (s, 3H).  $^{13}C$  NMR (126 MHz,  $CDCl_3$ )  $\delta$  154.4, 152.6, 141.5, 141.4, 136.5, 136.3, 133.0, 132.7, 131.9, 130.3, 128.8, 128.1, 127.3, 121.7, 120.6, 17.5, 17.0, 15.2, 14.5, 13.8. HRMS (ESI):  $m/z$   $[M + H]^+$  calcd for  $C_{22}H_{22}^{11}BCl_2F_2N_2^+$ : 398.1527; found: 398.1522,  $\Delta = -1.42$  ppm.

(*E*)-2-(4-Bromostyryl)-5,5-difluoro-1,3,7,9,10-pentamethyl-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo [1,2-*c*:2',1'-*f*] [1,3,2]diazaborinine (9e). A brownish red solid (yield: 24.7 mg, 28%, single substitute: double substitute = 94 : 6).  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.49 (d,  $J = 8.4$  Hz, 2H), 7.36 (d,  $J = 8.5$  Hz, 2H), 6.97 (d,  $J = 16.5$  Hz, 1H), 6.62 (d,  $J = 16.5$  Hz, 1H), 6.11 (s, 1H), 2.68 (s, 3H), 2.66 (s, 3H), 2.56 (s, 3H), 2.51 (s, 3H), 2.45 (s, 3H).  $^{13}C$  NMR (126 MHz,  $CDCl_3$ )  $\delta$  154.4, 152.5, 141.5, 141.4, 136.7, 136.5, 132.7, 131.9, 131.8, 130.3, 128.0, 127.6, 121.7, 121.1, 120.7, 17.5, 17.0, 15.3, 14.5, 13.8. HRMS (ESI):  $m/z$   $[M - F]^+$  calcd for  $C_{22}H_{22}^{11}BBrFN_2^+$ : 423.1038; found: 423.1035,  $\Delta = -0.63$  ppm.

(*E*)-5,5-Difluoro-1,3,7,9,10-pentamethyl-2-(pent-1-en-1-yl)-5H-5 $\lambda$ ,6 $\lambda$ -dipyrrolo[1,2-*c*:2',1'-*f*] [1,3,2] diazaborinine (9f). A brownish red solid (yield: 33.6 mg, 51%, single substitute: double substitute = 89 : 11).  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  6.16 (d,  $J = 16.1$  Hz, 1H), 6.03 (s, 1H), 5.75 (dt,  $J = 16.1$ , 6.9 Hz, 1H), 2.59 (s, 3H), 2.56 (s, 3H), 2.51 (s, 3H), 2.40 (s, 3H), 2.38 (s, 3H), 2.21 (qd,  $J = 6.9$ , 2H), 1.50 (m,  $J = 7.4$  Hz, 2H), 0.97 (t,  $J = 7.4$  Hz, 3H).  $^{13}C$  NMR (126 MHz,  $CDCl_3$ )  $\delta$  153.2, 152.9, 141.0, 140.3, 136.5, 135.3, 132.1, 132.0, 129.5, 121.0, 120.6, 35.9, 22.7, 17.4, 16.8, 15.1, 14.4, 13.7, 13.6. HRMS (ESI):  $m/z$   $[M + H]^+$  calcd for  $C_{19}H_{26}^{11}BF_2N_2^+$ : 331.2152; found: 331.2159,  $\Delta = -0.73$  ppm.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This study was financially supported by the fund from the State Key Laboratory of Drug Research (grant no. SIMM1705KF06); Institutes for Drug Discovery and Development, Chinese Academy of Sciences (CASIMM0120163010), and One Hundred Talent Program of Chinese Academy of Sciences and the National Natural Science Foundation of China (81225022).

## References

- (a) A. Loudet and K. Burgess, *Chem. Rev.*, 2007, **107**, 4891–4932; (b) N. Boens, V. Leen and W. Dehaen, *Chem. Soc. Rev.*, 2012, **41**, 1130–1172.
- (a) A. Kamkaew, S. H. Lim, H. B. Lee, L. V. Kiew, L. Y. Chung and K. Burgess, *Chem. Soc. Rev.*, 2013, **42**, 77–88; (b) S. G. Awuah and Y. You, *RSC Adv.*, 2012, **2**, 11169–11183; (c) T. Kowada, H. Maeda and K. Kikuchi, *Chem. Soc. Rev.*, 2015, **44**, 4953–4972; (d) S. H. Lim, C. Thivierge, P. Nowak-Sliwinska, J. Han, H. van den Bergh, G. Wagnières, K. Burgess and H. B. Lee, *J. Med. Chem.*, 2010, **53**, 2865–2874.
- C. F. A. Gómez-Durán, I. Esnal, I. Valois-Escamilla, A. Urias-Benavides, J. Bañuelos, I. López Arbeloa, I. García-Moreno and E. Peña-Cabrera, *Chem.-Eur. J.*, 2016, **22**, 1048–1061.
- (a) X. Li, S. Huang and Y. Hu, *Org. Biomol. Chem.*, 2012, **10**, 2369–2372; (b) L. Jiao, C. Yu, J. Li, Z. Wang, M. Wu and E. Hao, *J. Org. Chem.*, 2009, **74**, 7525–7528; (c) N. Boens, B. Verbelen and W. Dehaen, *Eur. J. Org. Chem.*, 2015, **2015**, 6577–6595.
- H. M. L. Davies, J. Du Bois and J.-Q. Yu, *Chem. Soc. Rev.*, 2011, **40**, 1855–1856.
- (a) B. Verbelen, S. Boodts, J. Hofkens, N. Boens and W. Dehaen, *Angew. Chem., Int. Ed.*, 2015, **54**, 4612–4616; *Angew. Chem.*, 2015, **127**, 4695–4699; (b) B. Verbelen, V. Leen, L. Wang, N. Boens and W. Dehaen, *Chem. Commun.*, 2012, **48**, 9129–9131; (c) X. Zhou, Q. Wu, Y. Yu, C. Yu, E. Hao, Y. Wei, X. Mu and L. Jiao, *Org. Lett.*, 2016, **18**, 736–739; (d) J. L. Belmonte-Vázquez, R. Sola-Llano, J. Bañuelos, L. Betancourt-Mendiola, M. A. Vázquez-Guevara, I. López-Arbeloa and E. Peña-Cabrera, *Dyes Pigm.*, 2017, **147**, 246–259.
- C. Thivierge, R. Bandichhor and K. Burgess, *Org. Lett.*, 2007, **9**, 2135–2138.
- L. Luo, D. Wu, W. Li, S. Zhang, Y. Ma, S. Yan and J. You, *Org. Lett.*, 2014, **16**, 6080–6083.
- (a) R. Beaud, R. J. Phipps and M. J. Gaunt, *J. Am. Chem. Soc.*, 2016, **138**, 13183–13186; (b) L. Chan, A. McNally, Q. Y. Toh, A. Mendoza and M. J. Gaunt, *Chem. Sci.*, 2015, **6**, 1277–1281; (c) E. A. Merritt and B. Olofsson, *Angew. Chem., Int. Ed.*, 2009, **48**, 9052–9070; (d) V. V. Zhdankin and P. J. Stang, *Chem. Rev.*, 2008, **108**, 5299–5358; (e) Q. Y. Toh, A. McNally, S. Vera, N. Erdmann and M. J. Gaunt, *J. Am. Chem. Soc.*, 2013, **135**, 3772–3775; (f) X. Qian, J. Han and L. Wang, *Tetrahedron Lett.*, 2016, **57**, 607–610; (g) X. Wang, D. Huang, X. Wang, X. Zeng, X. Wang and Y. Hu, *Tetrahedron Lett.*, 2016, **57**, 4235–4238; (h) N. Gigant, L. Chausset-Boissarie, M.-C. Belhomme, T. Poisson, X. Pannecoucke and I. Gillaizeau, *Org. Lett.*, 2013, **15**, 278–281.
- N. Boens, B. Verbelen and W. Dehaen, *Eur. J. Org. Chem.*, 2015, **2015**, 6577–6595.
- R. J. Phipps, L. McMurray, S. Ritter, H. A. Duong and M. J. Gaunt, *J. Am. Chem. Soc.*, 2012, **134**, 10773–10776.
- (a) A. Bigot, A. E. Williamson and M. J. Gaunt, *J. Am. Chem. Soc.*, 2011, **133**, 13778–13781; (b) M. G. Suero, E. D. Bayle, B. S. L. Collins and M. J. Gaunt, *J. Am. Chem. Soc.*, 2013, **135**, 5332–5335.
- V. Lakshmi, T. Chatterjee and M. Ravikanth, *Eur. J. Org. Chem.*, 2014, **2014**, 2105–2110.

