

Cite this: *Chem. Sci.*, 2019, 10, 898

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 16th September 2018

Accepted 30th October 2018

DOI: 10.1039/c8sc04120b

rsc.li/chemical-science

X-ray Raman optical activity of chiral molecules†

Jérémy R. Rouxel,^{‡*a} Yu Zhang^{‡*b} and Shaul Mukamel^{*cd}

Resonant and off-resonant Raman Optical Activity signals in the X-ray regime (XROA) are predicted. XROA is a chiral-sensitive variant of the spontaneous Resonant Inelastic Scattering (RIXS) signal. Thanks to the highly localized nature of core excitations, these signals provide a direct probe of local chirality with high sensitivity to the molecular structure. We derive sum-over-states expressions for frequency domain XROA signals and apply them to tyrosine at the nitrogen and oxygen K-edges. Time-resolved extensions of ROA made possible by using additional pulses are briefly outlined.

1 Introduction

The recent development of intense X-ray light from synchrotron or X-ray free electron lasers (XFELs)^{1,2} makes it possible to extend well established infrared and optical nonlinear spectroscopic techniques^{3–6} to the X-ray regime. One notable advantage is that the light–matter interaction of X-ray light resonant with core excitations in molecules is highly spatially localized and element specific. Signals specifically targeting molecular chirality thus offer important structural information.⁷ Chirality is crucial for the biological activity of biomolecules and has been extensively studied using visible and IR techniques such as circular dichroism (CD), optical rotation⁸ and Raman Optical Activity (ROA).^{9,10} Chiral signals are intrinsically weak and considerable effort has been made to increase their magnitude^{11,12} which is vital, *e.g.* for sensor applications.^{13,14}

ROA employs circularly polarized light to measure the vibrational or electronic optical activity and is routinely used in the infrared¹⁵ and the visible¹⁶ regimes to obtain information on the absolute configuration of small chiral molecules and on the secondary and tertiary structure of proteins. This is particularly important for drug design since many important drugs are chiral. The X-ray extension of ROA should also provide favorable scaling since it increases as ω^5 compared to the conventional Raman background that scales as ω^4 .¹⁷ Expressions for XROA

signals are derived in Section 2 and simulation results of the amino acid tyrosine are presented and analyzed in Section 3. Our findings are summarized in Section 4.

2 The X-ray Raman optical activity signal

ROA has its origin in the pioneering work of Barron^{16,18,19} who demonstrated that the spontaneous Raman scattering intensity induced by circularly polarized light from chiral enantiomers is different. The ROA signal is usually measured by taking the difference between spontaneous Raman spectra induced by left and right polarized light at different scattering angles. The detector then measures all polarizations of the scattered light. This configuration is shown in Fig. 1. ROA signals can also be acquired in other ways by detecting, for example, the difference between the left and right polarized components of the scattered light when the incident light is linearly polarized or unpolarized.¹⁶ Mixed schemes in which both the polarization of the incoming light and the scattered light are varied have been used as well. By varying the polarization and detecting the signal at




Fig. 1 The ROA measurement scheme. Left and right circularly polarized X-ray pulses propagating along the *z* direction generate two Raman signals which are recorded as a function of the scattering angle. Their small difference constitutes the ROA signal, eqn (9). The detector polarization basis by *X* and *Y* provides an extra degree of freedom for the measurement.

^aEcole Polytechnique Fédérale de Lausanne, Laboratoire de Spectroscopie Ultrarapide, Faculté des Sciences de Base, ISIC-BSP, CH-1015 Lausanne, Switzerland. E-mail: jrrouxel@uci.edu

^bPULSE Institute, SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA. E-mail: yzhang15@slac.stanford.edu

^cDepartment of Chemistry, University of California, Irvine, CA 92697, USA. E-mail: smukamel@uci.edu

^dDepartment of Physics and Astronomy, University of California, Irvine, CA 92697, USA

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8sc04120b

‡ J. R. Rouxel and Y. Zhang have equal contributions.

In this paper, we study the difference in the Raman signal induced by the incoming left and right polarizations. The radiation/matter coupling is given by the multipolar Hamiltonian²² truncated at the electric quadrupole

Fig. 2(a), (b) and (c) represent the non-chiral electric dipole contribution, the magnetic dipole and the electric quadrupole, respectively. We shall denote the detected spontaneous Raman intensity by $S_{\text{RAM}}(\omega_{\text{X}}, \omega_{\text{s}}, \theta, \mathbf{e}_{\text{L/R}}, \mathbf{e}_{\text{s}})$, where ω_{X} is the incident pulse frequency, ω_{s} is the detected frequency, θ is the angle between the incident pulse and the detected light, and $\mathbf{e}_{\text{L/R}}$ and \mathbf{e}_{s} are the incident and diffracted pulse polarizations.

The XROA signal is given by the difference between left and right incoming polarizations:

Without loss of generality, in an isotropic sample, we assume that the incident pulse propagates along the z axis (this implies that $\mathbf{k}_x = \mathbf{e}_z$) and that the scattered Raman signal is measured in the xz plane (see Fig. 1).

The spontaneous Raman signal is defined as the change of photon number emitted in the detected direction


We shall neglect the magnetic field contribution because it is orders of magnitude smaller.²³ The time derivative can be calculated by using the expectation value of the Heisenberg equation of motion of the number operator.

The commutator can be calculated in a straightforward manner using the definition of the \mathbf{E}_s and \mathbf{B}_s operators

where \mathbf{k}_s is the wavevector of the emitted photon, Ω is the quantization volume and we have the boson commutation

Upon evaluating the commutator, the Raman signal (eqn (3)) becomes

where \Re and \Im denote the real and imaginary parts respectively. To obtain the ROA signal, we expand the exponent to second order in the incident field. Three terms then contribute to the signals. The leading term is given by the interaction with the electric dipoles only (diagram (a)) and does not contribute to the



ROA signal since it vanishes upon rotational averaging. Diagram (b) contains one magnetic dipole and diagram (c) has one electric quadrupole interaction.

pulse tuned at the N ($\omega_X \sim 389\text{--}397$ eV) and O ($\omega_X \sim 519\text{--}526$ eV) K-edges. The incoming X-rays are thus resonant with the excited transitions. The signals on D-tyrosine are identical but with an

$$S_{\text{RAM}}(\omega_X, \omega_s, \theta, \mathbf{e}_{\text{L/R}}, \mathbf{e}_s) = \left(\frac{\hbar\omega_s}{2\epsilon_0\Omega} \right)^2 \Re \int dt dt_3 dt_2 dt_1 d\mathbf{r}_4 d\mathbf{r}_3 d\mathbf{r}_2 d\mathbf{r}_1 \times e^{i\mathbf{k}_X \cdot (\mathbf{r}_1 - \mathbf{r}_3)} e^{-i\mathbf{k}_s \cdot (\mathbf{r}_2 - \mathbf{r}_4)} e^{i\omega_X(t-t_3)} e^{-i\omega_X(t-t_3-t_2-t_1)} e^{-i\omega_s t} e^{i\omega_s(t-t_3-t_2)} \\ \times [\mathbf{E}_X^*(\mathbf{r}_3, t-t_3) \otimes \mathbf{e}_s^* \otimes \mathbf{e}_s \otimes \mathbf{E}_X(\mathbf{r}_1, t-t_3-t_2-t_1) \langle \boldsymbol{\mu}(t_3) \boldsymbol{\mu}(t_2) \boldsymbol{\mu}(t_1) \boldsymbol{\mu}(0) \rangle \\ + \mathbf{E}_X^*(\mathbf{r}_3, t-t_3) \otimes \mathbf{e}_s^* \otimes \mathbf{e}_s \otimes \mathbf{B}_X(\mathbf{r}_1, t-t_3-t_2-t_1) \langle \boldsymbol{\mu}(t_3) \boldsymbol{\mu}(t_2) \boldsymbol{\mu}(t_1) \mathbf{m}(0) \rangle + \mathbf{E}_X^*(\mathbf{r}_3, t-t_3) \otimes \mathbf{e}_s^* \otimes \mathbf{e}_s \nabla \mathbf{E}_X(\mathbf{r}_1, t-t_3-t_2-t_1) \\ \times \langle \boldsymbol{\mu}(t_3) \boldsymbol{\mu}(t_2) \boldsymbol{\mu}(t_1) \mathbf{q}(0) \rangle + \text{permutations of magnetic and quadrupole interactions}] \quad (9)$$

where the $\mathbf{e}_{\text{L/R}}$ dependence of the signal is contained in the vector components of \mathbf{E}_X and \mathbf{B}_X . The three terms in brackets correspond to diagrams (a), (b) and (c), respectively in Fig. 2 and all operators evolve with the free molecular Hamiltonian in the interaction picture. Eqn (9) allows us to calculate time-resolved signals induced by broadband pulses. In that case, the spatial and temporal integrals can be then expressed as a Fourier transform. In this paper, we focus on ROA signals in the frequency domain measured with monochromatic light excitation. The total XROA signal is given by the sum of the magnetic and the quadrupole contributions:

$$S_{\text{XROA}}(\omega_X, \omega_s, \theta, \mathbf{e}_s) = S_{\text{XROA}}^{\text{mag}}(\omega_X, \omega_s, \theta, \mathbf{e}_s) + S_{\text{XROA}}^{\text{quad}}(\omega_X, \omega_s, \theta, \mathbf{e}_s) \quad (10)$$

Sum-over-states expressions for eqn (9) are given in Appendix B.

3 XROA signals of tyrosine

We have calculated the XROA signal, eqn (9), for a randomly oriented ensemble of L-tyrosine (Fig. 3(a)) with an X-ray incident

opposite sign and thus are not displayed. We focus on the zwitterionic structure (see Fig. 3(a)) of the molecule in water since this is its dominant form at physiological pH. The geometry was optimized at the B3LYP^{24,25}/aug-cc-pVTZ^{26,27} level of theory using the Gaussian program package.²⁸ The water solvent effect was considered with the PCM model.²⁹⁻³¹ All quantities needed for XROA signal calculations were calculated using the CD spectrum³²⁻³⁴ and the restricted excitation window time-dependent density functional (REW-TDDFT)³⁵⁻⁴⁰ implementations in the quantum chemistry package NWChem⁴¹ under the Tamm-Dancoff approximation (TDA).⁴²⁻⁴⁴ The computational protocol for the relevant transition moments (electric and magnetic dipoles, electric quadrupoles) has been presented in ref. 40. The B3LYP hybrid functional and the cc-pVTZ basis set²⁶ were used for all NWChem calculations. The water solvent effect was included using the COSMO model.⁴⁵ It is a common practice to apply constant shifts to the calculated core excitation energies from TDDFT to match the experimental results. Since in this study only the relative positions of spectroscopic features matter, we retain all original core excitation energies from the TDDFT calculations. We adopted a tight SCF convergence criterion




Fig. 3 Calculated spectra of L-tyrosine in its zwitterionic form. (a) UV-vis spectrum. (b) N K-edge X-ray absorption spectrum. (c) O K-edge X-ray absorption spectrum. The 3D molecular model is also shown and the x, y, z axes are marked in red, green and blue, respectively. The oxygen atom in the phenol group is labelled O₁ and the other two in the carbonyl group are labelled O_{2,3}. The separate contributions of these two types of O atoms to the total signal (in light blue) are plotted in red and black, respectively. The signal contributions are normalized to different scales.



(energy change of SCF cycles $<10^{-8}$ a.u.) and an ultrafine grid in all NWChem calculations.

It is well known that TDDFT-TDA in general produces excitation energies of identical or even better quality of full TDDFT, but possibly worse oscillator strengths because of violation of the sum rule.^{46,47} However, previous TDDFT-TDA studies on core spectroscopy, which is the major topic of this manuscript, showed that TDDFT-TDA and full TDDFT gave almost indistinguishable results.^{39,48} TDDFT-TDA is computationally much cheaper than full TDDFT, which facilitates its use in large systems. Moreover, it handles triplet instabilities much better than full TDDFT.⁴⁹ In addition, it also gives better nonadiabatic couplings by reducing the local-density-approximation (LDA)

error.⁵⁰ With all the considerations above, we use TDDFT-TDA in this study.

Under the multipolar expansion, only the first non-vanishing multipole is origin invariant. Thus, in the present work, the electric quadrupoles and the magnetic dipoles are origin dependent.⁵¹ Indeed, the spectroscopic observables calculated from exact wave functions must be independent of the origin.⁵² However, numerical truncation of the basis set can induce origin-dependence of the spectroscopic observables. A possible solution is to use Gauge Invariant Atomic Orbitals (GIAOs) or London orbitals.^{34,53–55} Alternatively, as shown by Reiher *et al.*,⁵² by evaluating multipoles in the velocity gauge, the corresponding spectroscopic tensors are origin-independent even with a truncated basis. The behavior of the spectroscopic





Fig. 4 2D XROA signals (eqn (9)) displayed vs. the incoming frequency ω_X and the Raman frequency shift $\Delta\omega = \omega_X - \omega_s$ at the N K-edge for various scattering angles. The left and right columns represent X and Y polarizations of the detected photons in the detector frame.



The XROA signal at the N K-edge *vs.* the incoming plane wave frequency and of the Raman shift $\omega_X - \omega_s$ is displayed in Fig. 4 for various scattering angles. The corresponding 2D Raman signal at the scattering angle of $\pi/2$ is given in the ESI.† The ratio of the chiral contribution to the nonchiral one is 1.7% at the main peak of the spectrum (392.2 eV excitation and 8.4 eV Raman shift). The former provides spectroscopic information on the core-excited manifold while the latter covers the valence

Fig. 6 depicts the total XROA signals calculated at various K-edges, nitrogen and two groups of oxygen atoms for both X- and Y-polarization detection at a scattering angle of $\pi/2$. The X- and Y-polarization detected signals show similar patterns. As can be seen in Section B-eqn (16), the ROA signal depends on multiple transitions and their orientations, hence there is no simple relationship between a single transition momentum and the signal. However, as a general rule, strong ROA signals require that the relevant core and valence excitations involve the same particle orbitals (since the corresponding transition moment is a single-electron operator), and the involved particle or hole orbital should be close to the chiral centers if they exist. We can take the strong N 1s ROA features as an example.






Fig. 6 2D XROA signals (eqn (9)) displayed vs. the incoming frequency ω_X and the frequency shift $\Delta\omega = \omega_X - \omega_s$ at various edges. O_1 refers to the oxygen atom in the OH group near the benzene ring while $O_{2,3}$ are the two equivalent oxygen atoms at the carbonyl group. These two atoms are spectroscopically equivalent since the hydrogen atom can easily migrate between them. The scattering angle is $\pi/2$.

Important molecular orbitals (MOs) (S1) and an analysis on relevant excitations (Table S1†) are provided in the ESI.† The strong feature around the point (390.6, 6.0) eV involves the weak N 1s core excitation at 390.6 eV and the strong valence excitation at 6.0 eV. The two excitations couple through the LUMO, which is distributed on the benzene ring (see S1). Since the LUMO (L) is not close to the N atom, the corresponding X-ray absorption is weak (Fig. 3). However, the relevant valence hole orbital HOMO-3 (H-3) lies on the N atom as well as the chiral center, and the benzene ring, so the corresponding valence excitation and the chiral ROA signal is strong. Another strong ROA feature around the point (393.4, 8.4) eV can be explained similarly. The core excitation


around 393.4 eV couples with the valence excitation around 8.4 eV through the L + 7 particle orbital. This orbital is delocalized over all atoms, and the corresponding valence hole orbital H-1 carries the information regarding the chiral center.

The O 1s ROA signals are more complicated since there are two inequivalent O atoms in the molecule. The total O 1s ROA signals and the contributions from the O_1 and $O_{2,3}$ atoms are displayed in Fig. 6. As discussed in our previous work on X-ray CD,⁵⁶ the chiral X-ray signal intensity depends on the distance between the X-ray chromophore and the molecular chiral center. This propensity is also clearly observed here by comparing the XROA signals of O_1 and $O_{2,3}$. One can see from



Fig. 7 depicts the magnetic and quadrupole contributions to the XROA signal at the N K-edge ($\omega_x = 393.25$ eV, $\theta = \pi/2$), eqn (16), for x and y polarization of the signal. The total signal measured at the N K-edge is given by the sum of the magnetic and the electric contributions. For x -polarized detection, the total signal (blue curve) is a mixture of magnetic dipole and electric quadrupole contributions of similar weight. The y -polarized signal under the same conditions is dominated by the magnetic interaction. The polarization direction of detection along which the quadrupole contribution is important is determined by the geometric and electronic structure of the system, but the fact that this contribution varies significantly for different polarizations of detection is general. The sensitivity of XROA signals to the electric quadrupole interaction opens a new window to molecular chirality compared to XCD signals which do not depend on the electric quadrupole because of cancellation caused by rotational averaging.

The signal in eqn (9) was calculated for a monochromatic excitation. It will be interesting to extend this technique to the time-domain by using ultrashort X-ray pulses. Time-resolved extension can be achieved by adding an actinic pulse with a controlled delay in an optical pump/X-ray probe scheme. This can be done by starting with eqn (9) and expanding the exponent to higher orders in the incoming pulses in order to trigger some chiral dynamics. XROA can then be used as a novel detection mode on top of a nonlinear process that would be sensitive to local change of the chiral structure in the vicinity of an X-ray chromophore. Time-domain XROA should be a powerful spectroscopy tool to track chiral molecules in chemical reactions.



Appendix A rotationally averaged tensors

A n -th rank tensor T is rotationally averaged using the averaging tensor $I^{(n)}$.⁶⁰ The correlation tensor involving electric and magnetic dipoles are only fourth rank tensors for ROA and the ones involving electric quadrupole are of rank 5.

$$I^{(4)} = \frac{1}{30} \begin{pmatrix} \delta_{i_1 i_2} \delta_{i_1 i_2} \\ \delta_{i_1 i_3} \delta_{i_2 i_4} \\ \delta_{i_1 i_4} \delta_{i_2 i_3} \end{pmatrix}^T \begin{pmatrix} 4 & -1 & -1 \\ -1 & 4 & -1 \\ -1 & -1 & 4 \end{pmatrix} \begin{pmatrix} \delta_{\lambda_1 \lambda_2} \delta_{\lambda_1 \lambda_2} \\ \delta_{\lambda_1 \lambda_3} \delta_{\lambda_2 \lambda_4} \\ \delta_{\lambda_1 \lambda_4} \delta_{\lambda_2 \lambda_3} \end{pmatrix} \quad (11)$$

$$I^{(5)} = \frac{1}{30} \begin{pmatrix} \varepsilon_{i_1 i_2 i_3} \delta_{i_4 i_5} \\ \varepsilon_{i_1 i_2 i_4} \delta_{i_3 i_5} \\ \varepsilon_{i_1 i_2 i_5} \delta_{i_3 i_4} \\ \varepsilon_{i_1 i_3 i_4} \delta_{i_2 i_5} \\ \varepsilon_{i_1 i_3 i_5} \delta_{i_2 i_4} \\ \varepsilon_{i_1 i_4 i_5} \delta_{i_2 i_3} \end{pmatrix}^T \begin{pmatrix} 3 & -1 & -1 & 1 & 1 & 0 \\ -1 & 3 & -1 & -1 & 0 & 1 \\ -1 & -1 & 3 & 0 & -1 & -1 \\ 1 & -1 & 0 & 3 & -1 & 1 \\ 1 & 0 & -1 & -1 & 3 & -1 \\ 0 & 1 & -1 & 1 & -1 & 3 \end{pmatrix} \begin{pmatrix} \varepsilon_{\lambda_1 \lambda_2 \lambda_3} \delta_{\lambda_4 \lambda_5} \\ \varepsilon_{\lambda_1 \lambda_2 \lambda_4} \delta_{\lambda_3 \lambda_5} \\ \varepsilon_{\lambda_1 \lambda_2 \lambda_5} \delta_{\lambda_3 \lambda_4} \\ \varepsilon_{\lambda_1 \lambda_3 \lambda_4} \delta_{\lambda_2 \lambda_5} \\ \varepsilon_{\lambda_1 \lambda_3 \lambda_5} \delta_{\lambda_2 \lambda_4} \\ \varepsilon_{\lambda_1 \lambda_4 \lambda_5} \delta_{\lambda_2 \lambda_3} \end{pmatrix} \quad (12)$$

where δ and ε are the Kronecker and Levi-Civita symbols respectively.

The $I^{(4)}$ tensor is used to rotationally average the magnetic correlation functions (e.g. $\langle \mu\mu\mu\mu \rangle$) and $I^{(5)}$ for the electric quadrupole ones (e.g. $\langle \mu\mu\mu q \rangle$). For example, we have

$$\langle \mu\mu\mu\mu \rangle_{ijkl} = (I^{(4)})_{ijkl}^{i'j'k'l'} \langle \mu\mu\mu\mu \rangle_{i'j'k'l'} \quad (13)$$

where we used Einstein summation convention for the Cartesian indices i, j, k , and l , and $\langle \dots \rangle_{\Omega}$ stands for rotational averaging.

Appendix B sum-over-states expressions of the XROA signals

Here, we provide the sum-over-states expressions of the three diagrams shown in Fig. 2. Assuming the incoming pulses are plane waves, the electric contribution (Fig. 2(a)) is given by

$$S_{\text{ROA}}^{\text{elec}}(\omega_X, \omega_s, \theta, \mathbf{e}_s) = C \Re \sum_{cc'v} c I_{cg}(\omega_X) I_{vg}^*(\omega_X - \omega_s) I_{cg}^*(\omega_X) \times \left\langle \mu_{gc'}^i \mu_{c'v}^{j\dagger} \mu_{vc}^k \mu_{cg}^{l\dagger} \right\rangle_{\Omega} (\bar{e}_L^i e_L^j \bar{e}_s^k e_L^l - \bar{e}_R^i e_R^j \bar{e}_s^k e_R^l) \quad (14)$$

where

$$C = \frac{2}{\hbar^4} |\mathcal{E}_X|^2 4\pi^2 \left(\frac{\hbar\omega_s}{2\varepsilon_0} \right)^2 \frac{N}{c} \quad (15)$$

$$I_{ab}(\omega) = \frac{1}{\omega - \omega_{ab} + i\Gamma_{ab}} \quad (16)$$

The magnetic contribution given in Fig. 2(b) is summed over the permutations of the magnetic interaction

$$S_{\text{ROA}}^{\text{mag}}(\omega_X, \omega_s, \theta, \mathbf{e}_s) = C \Re \sum_{cc'v} I_{cg}(\omega_X) I_{vg}^*(\omega_X - \omega_s) I_{cg}^*(\omega_X) \times \left(\left\langle m_{gc'}^i \mu_{c'v}^{j\dagger} \mu_{vc}^k \mu_{cg}^{l\dagger} \right\rangle_{\Omega} (\bar{b}_L^i e_L^j \bar{e}_s^k e_L^l - \bar{b}_R^i e_R^j \bar{e}_s^k e_R^l) \right. \\ + \left\langle \mu_{gc'}^i m_{c'v}^{j\dagger} \mu_{vc}^k \mu_{cg}^{l\dagger} \right\rangle_{\Omega} (\bar{e}_L^i b_L^j \bar{e}_s^k e_L^l - \bar{e}_R^i b_R^j \bar{e}_s^k e_R^l) \\ + \left\langle \mu_{gc'}^i \mu_{c'v}^{j\dagger} m_{vc}^k \mu_{cg}^{l\dagger} \right\rangle_{\Omega} (\bar{e}_L^i e_L^j \bar{b}_s^k e_L^l - \bar{e}_R^i e_R^j \bar{b}_s^k e_R^l) \\ \left. + \left\langle \mu_{gc'}^i \mu_{c'v}^{j\dagger} \mu_{vc}^k m_{cg}^{l\dagger} \right\rangle_{\Omega} (\bar{e}_L^i e_L^j \bar{e}_s^k b_L^l - \bar{e}_R^i e_R^j \bar{e}_s^k b_R^l) \right) \quad (17)$$

Finally, the quadrupole contribution is given in the diagram in Fig. 2(c)

$$S_{\text{ROA}}^{\text{quad}}(\omega_X, \omega_s, \theta, \mathbf{e}_s) = C \Re \sum_{cc'v} I_{cg}(\omega_X) I_{vg}^*(\omega_X - \omega_s) I_{cg}^*(\omega_X) \times \left(\left\langle q_{gc'}^{ij} \mu_{c'v}^{k\dagger} \mu_{vc}^l \mu_{cg}^{m\dagger} \right\rangle_{\Omega} (-i) k^j (\bar{e}_L^i e_L^j \bar{e}_s^k e_L^l - \bar{e}_R^i e_R^j \bar{e}_s^k e_R^l) \right. \\ + \left\langle \mu_{gc'}^i q_{c'v}^{jk\dagger} \mu_{vc}^l \mu_{cg}^{m\dagger} \right\rangle_{\Omega} i k_s^k (\bar{e}_L^i e_L^j \bar{e}_s^k e_L^l - \bar{e}_R^i e_R^j \bar{e}_s^k e_R^l) \\ + \left\langle \mu_{gc'}^i \mu_{c'v}^{j\dagger} q_{vc}^{kl} \mu_{cg}^{m\dagger} \right\rangle_{\Omega} (-i) k_s^l (\bar{e}_L^i e_L^j \bar{e}_s^k e_L^l - \bar{e}_R^i e_R^j \bar{e}_s^k e_R^l) \\ \left. + \left\langle \mu_{gc'}^i \mu_{c'v}^{j\dagger} \mu_{vc}^k q_{cg}^{lm\dagger} \right\rangle_{\Omega} i k^m (\bar{e}_L^i e_L^j \bar{e}_s^k e_L^l - \bar{e}_R^i e_R^j \bar{e}_s^k e_R^l) \right) \quad (18)$$

Note that the variable k and k_s are the wavevectors of the incoming and outgoing photons respectively while the superscripts k refer to the dummy summation index over cartesian components. The four-point correlation function of matter can be expanded in eigenstates. Chiral signals are measured in ensembles of randomly oriented chiral molecules. Thus, the four-point matter correlation function, eqn (17) and (18), must be rotationally averaged (see Appendix A). We use the following identities to simplify the polarization vector differences:

$$\bar{e}_L^i e_L^j - \bar{e}_R^i e_R^j = -i \varepsilon^{ilm} \hat{k}^m \quad (19)$$

$$\bar{b}_L^i e_L^j - \bar{b}_R^i e_R^j = i (\delta^{il} - \hat{k}^i \hat{k}^l) \quad (20)$$

Appendix C origin dependence upon coordinate change

The transition matrix elements can be computed either in the length or in the velocity gauge, denoted as r and p respectively. Their transformations under a coordinate translation \mathbf{a} from the origin \mathbf{O} are given by

$$\mu_{ij}^r(\mathbf{O} + \mathbf{a}) = \mu_{ij}^r(\mathbf{O}) \quad (21)$$

$$m_{ij}(\mathbf{O} + \mathbf{a}) = m_{ij}^r(\mathbf{O}) + \left(\frac{1}{2} \mathbf{a} \times \mathbf{p} \right)_{ij} \quad (22)$$



$$q_{ij}^r(\mathbf{O} + \mathbf{a}) = q_{ij}^r(\mathbf{O}) - \frac{3}{2}(\mathbf{a} \otimes \boldsymbol{\mu}_{ij} + \boldsymbol{\mu}_{ij} \otimes \mathbf{a}) + \mathbf{a} \cdot \boldsymbol{\mu}_{ij} \mathbb{1} \quad (23)$$

$$\boldsymbol{\mu}_{ij}^p(\mathbf{O} + \mathbf{a}) = \boldsymbol{\mu}_{ij}^p(\mathbf{O}) \quad (24)$$

$$q_{ij}^p(\mathbf{O} + \mathbf{a}) = q_{ij}^p(\mathbf{O}) - (\mathbf{a} \otimes \boldsymbol{\mu}_{ij} + \boldsymbol{\mu}_{ij} \otimes \mathbf{a}) \quad (25)$$

The magnetic dipole remains unchanged in the length and velocity gauge.⁵¹ The ROA spectroscopic signal is origin independent with truncated basis sets only in the velocity gauge. Thus the velocity gauge has been used to compute the signals.⁵²

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The support of the Chemical Sciences, Geosciences, and Biosciences division, Office of Basic Energy Sciences, Office of Science, U.S. Department of Energy through Award No. DE-FG02-04ER15571 and of the National Science Foundation (Grant No CHE-1663822) is gratefully acknowledged. J. R was supported by the DOE grant. Y. Z. would like to thank Prof. Jochen Autschbach for helpful discussions.

References

- 1 Y. Ding, Z. Huang, D. Ratner, P. Bucksbaum and H. Merdji, Generation of attosecond X-ray pulses with a multicycle two-color enhanced self-amplified spontaneous emission scheme, *Phys. Rev. Spec. Top.-Accel. Beams*, 2009, **12**, 060703.
- 2 M. Chini, K. Zhao and Z. Chang, The generation, characterization and applications of broadband isolated attosecond pulses, *Nat. Photonics*, 2014, **8**, 178.
- 3 J. D. Biggs, Y. Zhang, D. Healion and S. Mukamel, Two-dimensional stimulated resonance Raman spectroscopy of molecules with broadband X-ray pulses, *J. Chem. Phys.*, 2012, **136**(17), 174117.
- 4 Y. Zhang, J. D. Biggs and S. Mukamel, Understanding excitation energy transfer in metalloporphyrin heterodimers with different linkers, bonding structures, and geometries through stimulated X-ray Raman spectroscopy, *J. Mod. Opt.*, 2014, **61**, 558–567.
- 5 S.-K. Son, H. N. Chapman and R. Santra, Determination of multiwavelength anomalous diffraction coefficients at high x-ray intensity, *J. Phys. B: At., Mol. Opt. Phys.*, 2013, **46**, 164015.
- 6 K. Tamasaku, E. Shigemasa, Y. Inubushi, T. Katayama, K. Sawada, H. Yumoto, H. Ohashi, H. Mimura, M. Yabashi, K. Yamauchi, *et al.*, X-ray two-photon absorption competing against single and sequential multiphoton processes, *Nat. Photonics*, 2014, **8**, 313.
- 7 N. Berova, K. Nakanashi and R. W. Woody, in *Circular Dichroism, Principles and Applications*, Wiley, 2000.
- 8 A. Lázár, J. G. Ángyán, M. Hollósi, P. Huszthy and P. R. Surján, Theoretical CD spectrum calculations of the crown-ether aralkyl-ammonium salt complex, *Chirality*, 2002, **14**, 377–385.
- 9 L. Ashton, P. Pudney, E. Blanch and G. Yakubov, Understanding glycoprotein behaviours using Raman and Raman optical activity spectroscopies: Characterising the entanglement induced conformational changes in oligosaccharide chains of mucin, *Adv. Colloid Interface Sci.*, 2013, **199**, 66–77.
- 10 L. D. Barron, The development of biomolecular Raman optical activity spectroscopy, *Biomed. Spectrosc. Imaging*, 2015, **4**, 223–253.
- 11 Y. Tang and A. E. Cohen, Optical chirality and its interaction with matter, *Phys. Rev. Lett.*, 2010, **104**, 163901.
- 12 J. R. Rouxel, V. Y. Chernyak and S. Mukamel, Non-local real-space analysis of chiral optical signals, *Chem. Sci.*, 2016, **7**, 6824–6831.
- 13 H. Zhang and A. Govorov, Giant circular dichroism of a molecule in a region of strong plasmon resonances between two neighboring gold nanocrystals, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2013, **87**, 075410.
- 14 M. Schäferling, D. Dregely, M. Hentschel and H. Giessen, Tailoring enhanced optical chirality: design principles for chiral plasmonic nanostructures, *Phys. Rev. X*, 2012, **2**, 031010.
- 15 L. A. Nafie, Infrared and Raman vibrational optical activity: theoretical and experimental aspects, *Annu. Rev. Phys. Chem.*, 1997, **48**, 357–386.
- 16 L. Barron, in *Molecular light scattering and optical activity*, Cambridge, 2004.
- 17 J. Kapitán, L. D. Barron and L. Hecht, A novel Raman optical activity instrument operating in the deep ultraviolet spectral region, *J. Raman Spectrosc.*, 2015, **46**, 392–399.
- 18 L. Barron and A. Buckingham, Rayleigh and Raman scattering from optically active molecules, *Mol. Phys.*, 1971, **20**, 1111–1119.
- 19 L. Barron, M. Bogaard and A. Buckingham, Raman scattering of circularly polarized light by optically active molecules, *J. Am. Chem. Soc.*, 1973, **95**, 603–605.
- 20 T. Dartigalongue and F. Hache, Time-resolved circular dichroism in carbonmonoxy-myoglobin: The central role of the proximal histidine, *Chirality*, 2006, **18**, 273–278.
- 21 F. Hache, M.-C. Schanne-Klein, H. Mesnil, M. Alexandre, G. Lemerrier and C. Andraud, Nonlinear optical activity in chiral molecules: surface second harmonic generation and nonlinear circular dichroism, *C. R. Phys.*, 2002, **3**, 429–437.
- 22 A. Salam, *Molecular quantum electrodynamics: long-range intermolecular interactions*, John Wiley & Sons, 2010.
- 23 D. Abramavicius, B. Palmieri, D. V. Voronine, F. Sanda and S. Mukamel, Coherent multidimensional optical spectroscopy of excitons in molecular aggregates; quasiparticle *versus* supermolecule perspectives, *Chem. Rev.*, 2009, **109**, 2350–2408.



- 24 A. D. Becke, Density-functional thermochemistry. III. The role of exact exchange, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- 25 P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, *Ab Initio* Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields, *J. Phys. Chem.*, 1994, **98**, 11623–11627.
- 26 T. H. Dunning, Gaussian Basis Sets for Use in Correlated Molecular Calculations. I. The Atoms Boron through Neon and Hydrogen, *J. Chem. Phys.*, 1989, **90**, 1007–1023.
- 27 R. A. Kendall, T. H. Dunning Jr and R. J. Harrison, Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions, *J. Chem. Phys.*, 1992, **96**(9), 6796–6806.
- 28 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, and D. J. Fox, *Gaussian 09, Revision C.01*, Gaussian, Inc., Wallingford CT, 2009.
- 29 S. Miertus, E. Scrocco and J. Tomasi, Electrostatic Interaction of a Solute with a Continuum. A Direct Utilization of *ab initio* Molecular Potentials for the Prediction of Solvent Effects, *Chem. Phys.*, 1981, **55**(1), 117–129.
- 30 S. Miertus and J. Tomasi, Approximate Evaluations of the Electrostatic Free Energy and Internal Energy Changes in Solution Processes, *Chem. Phys.*, 1982, **65**, 239.
- 31 J. L. Pascual-Ahuir, E. Silla and I. Tuñón, GEPOL: An improved description of molecular-surfaces. 3. A new algorithm for the computation of a solvent-excluding surface, *J. Comput. Chem.*, 1994, **15**(10), 1127–1138.
- 32 J. Autschbach, T. Ziegler, S. J. A. van Gisbergen and E. J. Baerends, Chiroptical Properties from Time-dependent Density Functional Theory. I. Circular Dichroism Spectra of Organic Molecules, *J. Chem. Phys.*, 2002, **116**, 6930–6940.
- 33 J. Autschbach and T. Ziegler, Calculating Molecular Electric and Magnetic Properties from Time-dependent Density Functional Response Theory, *J. Chem. Phys.*, 2002, **116**, 891–896.
- 34 J. Autschbach, Time-Dependent Density Functional Theory for Calculating Origin-Independent Optical Rotation and Rotatory Strength Tensors, *ChemPhysChem*, 2011, **12**, 3224–3235.
- 35 M. Stener, G. Fronzoni and M. de Simone, Time Dependent Density Functional Theory of Core Electrons Excitations, *Chem. Phys. Lett.*, 2003, **373**, 115–123.
- 36 N. A. Besley and A. Noble, Time-Dependent Density Functional Theory Study of the X-ray Absorption Spectroscopy of Acetylene, Ethylene, and Benzene on Si(100), *J. Phys. Chem. C*, 2007, **111**, 3333–3340.
- 37 S. DeBeer-George, T. Petrenko and F. Neese, Time-dependent Density Functional Calculations of Ligand K-edge X-ray Absorption Spectra, *Inorg. Chim. Acta*, 2008, **361**, 965–972.
- 38 W. Liang, S. A. Fischer, M. J. Frisch and X. Li, Energy-Specific Linear Response TDHF/TDDFT for Calculating High-Energy Excited States, *J. Chem. Theory Comput.*, 2011, **7**, 3540–3547.
- 39 K. Lopata, B. E. V. Kuiken, M. Khalil and N. Govind, Linear-Response and Real-Time Time-Dependent Density Functional Theory Studies of Core-Level Near-Edge X-Ray Absorption, *J. Chem. Theory Comput.*, 2012, **8**, 3284–3292.
- 40 Y. Zhang, J. D. Biggs, D. Healion, N. Govind and S. Mukamel, Core and Valence Excitations in Resonant X-ray Spectroscopy Using Restricted Excitation Window Time-dependent Density Functional Theory, *J. Chem. Phys.*, 2012, **137**, 194306.
- 41 M. Valiev, E. Bylaska, N. Govind, K. Kowalski, T. Straatsma, H. V. Dam, D. Wang, J. Nieplocha, E. Apra, T. Windus, *et al.*, NWChem: A Comprehensive and Scalable Open-source Solution for Large Scale Molecular Simulations, *Comput. Phys. Commun.*, 2010, **181**, 1477–1489.
- 42 I. Tamm, Relativistic interaction of elementary particles, *J. Phys.*, 1945, **9**, 449.
- 43 S. M. Dancoff, Non-Adiabatic Meson Theory of Nuclear Forces, *Phys. Rev.*, 1950, **78**, 382–385.
- 44 S. Hirata and M. Head-Gordon, Time-dependent Density Functional Theory within the Tamm–Dancoff Approximation, *Chem. Phys. Lett.*, 1999, **314**, 291–299.
- 45 A. Klamt and G. Schüürmann, COSMO: a new approach to dielectric screening in solvents with explicit expressions for the screening energy and its gradient, *J. Chem. Soc., Perkin Trans. 2*, 1993, 799.
- 46 M. E. Casida and M. Huix-Rotllant, Progress in time-dependent density-functional theory, *Annu. Rev. Phys. Chem.*, 2012, **63**, 287–323.
- 47 A. Chantzis, A. D. Laurent, C. Adamo and D. Jacquemin, Is the Tamm–Dancoff approximation reliable for the calculation of absorption and fluorescence band shapes?, *J. Chem. Theory Comput.*, 2013, **9**, 4517–4525.
- 48 S. DeBeer George and F. Neese, Calibration of scalar relativistic density functional theory for the calculation of sulfur K-edge X-ray absorption spectra, *Inorg. Chem.*, 2010, **49**, 1849–1853.
- 49 M. J. Peach, M. J. Williamson and D. J. Tozer, Influence of triplet instabilities in TDDFT, *J. Chem. Theory Comput.*, 2011, **7**, 3578–3585.
- 50 C. Hu, O. Sugino and K. Watanabe, Performance of Tamm–Dancoff approximation on nonadiabatic couplings by time-dependent density functional theory, *J. Chem. Phys.*, 2014, **140**, 054106.



- 51 L. K. Sørensen, R. Lindh and M. Lundberg, What do we approximate and what are the consequences in perturbation theory?, arXiv preprint arXiv:1608.02399 2016.
- 52 S. Lubert and M. Reiher, Raman optical activity spectra of chiral transition metal complexes, *Chem. Phys.*, 2008, **346**, 212–223.
- 53 F. London, Théorie quantique des courants interatomiques dans les combinaisons aromatiques, *J. Phys. Radium*, 1937, **8**, 397.
- 54 T. Helgaker, K. Ruud, K. L. Bak, P. Jørgensen and J. Olsen, Vibrational Raman optical activity calculations using London atomic orbitals, *Faraday Discuss.*, 1994, **99**, 165–180.
- 55 M. Krykunov and J. Autschbach, Calculation of origin-independent optical rotation tensor components in approximate time-dependent density functional theory, *J. Chem. Phys.*, 2006, **125**(3), 034102.
- 56 Y. Zhang, J. R. Rouxel, J. Autschbach, N. Govind and S. Mukamel, X-ray circular dichroism signals: a unique probe of local molecular chirality, *Chem. Sci.*, 2017, **8**, 5969–5978.
- 57 J. Goulon, A. Rogalev, F. Wilhelm, N. Jaouen, C. Goulon-Ginet and C. Brouder, Optical activity probed with x-rays, *J. Phys.: Condens. Matter*, 2003, **15**, S633.
- 58 L. Alagna, T. Prosperi, S. Turchini, J. Goulon, A. Rogalev, C. Goulon-Ginet, C. R. Natoli, R. D. Peacock and B. Stewart, X-ray natural circular dichroism, *Phys. Rev. Lett.*, 1998, **80**, 4799.
- 59 B. Stewart, R. D. Peacock, L. Alagna, T. Prosperi, S. Turchini, J. Goulon, A. Rogalev and C. Goulon-Ginet, Circular Dichroism at the Edge: Large X-ray Natural CD in the 1s \rightarrow 3d Pre-Edge Feature of 2 [Co (en) 3Cl₃].NaCl·6H₂O, *J. Am. Chem. Soc.*, 1999, **121**, 10233–10234.
- 60 D. P. Craig and T. Thirunamachandran, *Molecular quantum electrodynamics: an introduction to radiation-molecule interactions*, Courier Corporation, 1984.

