



Cite this: *J. Anal. At. Spectrom.*, 2021, **36**, 954

Determination of the naturally occurring vanadium-complex amavadin in *Amanita muscaria* with HPLC-ICPMS†

Simone Braeuer, ‡* Martin Walenta, Lorenz Steiner and Walter Goessler

Amanita muscaria, also known as the fly agaric mushroom, can accumulate vanadium (V), with up to several hundred mg V kg⁻¹ dry mass. It is long known that V is present in *A. muscaria* as a complex called amavadin, but methods for the investigation of the distribution and biosynthesis of amavadin in mushrooms are missing. Here, we describe the development of the first sensitive method for the determination of amavadin and other V-containing compounds in environmental samples by employing high performance liquid chromatography (HPLC) and inductively coupled plasma mass spectrometry (ICPMS). A strong anion-exchange column serves as the stationary phase, and the mobile phase consists of an aqueous ammonium citrate buffer and ethylenediaminetetraacetate (EDTA). The concentration and pH of the mobile phase as well as the column temperature were evaluated to optimize the separation. With the final method, amavadin is eluted in less than 17 minutes, and its limit of detection is 0.05 μg V L⁻¹. Moreover, the compound's two isomers are separated from each other and can be quantified independently. The method was applied to extracts of fruit-body samples of *A. muscaria*. The extraction efficiency was 74 ± 12%, and amavadin accounted for 75–96% of the extracted V. In addition, significant concentrations of other V species could be detected, which have never been described before. Our results demonstrate that V speciation in mushrooms is more complex than assumed until now and that more in-depth investigations on this matter are needed. The developed method enables the investigation of organic and inorganic V species in the environment, even at low concentrations.

Received 19th December 2020
 Accepted 18th March 2021

DOI: 10.1039/d0ja00518e

rsc.li/jaas

Institute of Chemistry, University of Graz, Universitaetsplatz 1, 8010 Graz, Austria.
E-mail: simone.braeuer@uni-graz.at; Tel: +43 316 380 5318

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ja00518e

‡ Current address: Atomic & Mass Spectrometry Research Unit, Department of Chemistry, Ghent University, Krijgslaan 281-S12, 9000 Ghent, Belgium.

1 Introduction

1.1 Vanadium

Vanadium (V) is widely distributed in nature, being the 22nd most abundant element in the earth's crust.¹ The essentiality of V has been proven for some bacteria, terrestrial fungi, lichens



Simone Braeuer obtained her PhD in 2018 from the University of Graz, Austria, in the research group of Prof. Walter Goessler. She stayed there for one more year and then started a Postdoctoral fellowship at Ghent University, Belgium, in the Atomic & Mass Spectrometry research group of Prof. Frank Vanhaecke. Her work in Graz focused on the development and application of ICPMS and HPLC-ICPMS methods for the determination of trace element species in the environmental context. For her PhD, her main focus was on arsenic in mushrooms. Since mushrooms also contain other elements, sometimes in surprising concentrations and species, she soon became interested in some of those as well. The fascination with ICPMS and mushrooms grew so big that she is still investigating mushrooms in her current Postdoctoral project, but now also via multi-collector ICPMS and laser ablation ICPMS. She thinks that multidisciplinary and international collaborations are both fascinating and essential for environmental research. She also believes that high-quality, state-of-the-art analytical chemistry is of key importance – for environmental as well as for many other (interdisciplinary) studies.



and marine macro-algae,² but for other living organisms, the role of V is still unclear.³ Because of the similarity of vanadate and phosphate, it has been suggested that V could play a role in the regulation of the phosphate metabolism. A variety of V compounds are under investigation as potential anti-diabetic drugs⁴ and for several other medical applications.⁵ On the other hand, V can also be toxic to living organisms.^{6,7} It is usually thought that vanadate, the inorganic pentavalent form of V, is more toxic than V in the oxidation state +4 and that organic V species are less toxic than inorganic ones.⁸ However, the adverse health effects of V exposure are usually only observed in laboratory animals and people working in the V-processing industry,³ with a few exceptions.⁹

Ascidians (sea squirts) and some fan worms can accumulate incredible amounts of V. The highest concentrations, up to 350 mM (more than 17 000 mg V kg⁻¹), have been reported in *Ascidia gemmata*,¹⁰ where specialized cells store the element in its otherwise rather unstable trivalent form.¹¹ Until now, the actual function of V in ascidians remains elusive.^{10,12}

1.2 Amanita muscaria

Another living organism that accumulates V is the macrofungus *Amanita muscaria* (L.) Lam, also prominently known as fly agaric, along with a few closely related species. While most other mushrooms typically contain less than 0.5 mg V kg⁻¹ dry mass (dm),^{13–15} *A. muscaria* can easily take up more than 100 mg V kg⁻¹ dm.^{13,16–18}

The highest concentrations, namely up to 1000 mg V kg⁻¹ dm, have been found in the “bulb”, which is the lowest part of the stipe.^{17,19,20} This part of the fruit body is prone to contamination by soil, and no measures to minimize such contamination were described in the respective studies. However, the V concentrations in European top soils are reportedly between 1.28 and 537 mg kg⁻¹, with a median of only 60 mg kg⁻¹.⁹ This makes it unlikely that 1000 mg V kg⁻¹ in a mushroom sample are solely to be attributed to remaining soil particles.

Concerning the other parts of the fruit body, the V concentrations in the stipe, cap skin and cap flesh were at least two times lower than in the corresponding bulb sample, while they were again a little higher in the gills (but still lower than in the bulb; the concentrations were only shown in a rough graph, so that no exact numbers can be cited).¹⁹ In the spores, only 1–5 mg V kg⁻¹ dm were detected.¹⁹ Many other macrofungal species have been investigated for their V concentrations, including

several members of the genus *Amanita*, but only *A. regalis* and *A. velatipes* accumulated as much V as *A. muscaria*.^{17,21}

1.3 Amavadin

In 1972, Bayer and Kneifel found that V is present in *A. muscaria* as a negatively charged compound (pH 1–10), which they named “Amavadin”.²² They reported a tentative sum formula for the compound, and electron paramagnetic resonance (EPR) spectroscopy indicated that the oxidation state of V was +4. One year later, they published a first structure of the complex.²³ Further structural investigations by EPR, infrared (IR), and other techniques followed, which have been reviewed a few times over the decades.^{24–26} EPR spectra suggested that the different parts of the fruit body contained the same compound.^{20,27} The first synthesis of amavadin was reported in 1986,²⁸ and in the following years, a new structure of the complex was proposed²⁹ and confirmed by crystallography, X-ray absorption spectroscopy and nuclear magnetic resonance (NMR) spectroscopy.^{34,35}

Amavadin, as shown in Fig. 1, contains a rather unusual non-oxo V(IV) ion (the vanadyl ion VO²⁺ is more common) and can exist in two isomers, namely Δ-amavadin and Λ-amavadin. The two tetradentate ligands can coordinate to the V center in two distinct orientations, as shown in Fig. 1. It was reported that synthetic and also natural amavadin were present in solution in an almost equal mixture of the Δ- and Λ-isomers, although it was noted that isomerization could theoretically occur after extracting or dissolving the compound.^{30,31} The first enantioselective synthesis of amavadin was carried out by Hubregtse *et al.*, who found that the ratio of the diastereoisomers Λ : Δ was first 2.27, but later decreases to 0.8 at equilibrium.^{32,33}

Because of the reversible oxidation of V(IV) to V(V) in amavadin, several studies investigated its catalytic properties. It was found that a broad range of reactions can be catalyzed by the complex. This led to the synthesis and investigation of amavadin analogues with a different metal ion in the center or slightly modified ligands.³⁵

The biological role of amavadin is unknown. A few proposals have been made, but none have been proven so far. It has been speculated that amavadin was part of an ancient, primitive enzyme, which has been outdated through evolution.²⁶ Other suggestions are that it serves as a protective agent against external microbial pathogens or aids the self-regeneration of damaged fruit-body tissue by cross-linking thiol groups in proteins.^{32,34} The ability to mediate water oxidation gave the




Fig. 1 Structure of amavadin, schematic representation of its Δ- and Λ-helical forms, and structure of vanadyl acetate.



2 Experimental

2.1 Chemicals and instruments

Ultrapure water (18.2 M Ω cm) was used throughout the study. All chemicals were purchased from common chemical suppliers and used as received without further purification, if not described otherwise. A detailed description of all chemicals, including the purity and supplier, can be found in the ESI (Section 1 “Chemicals and instruments”†). The employed instruments are also listed in more detail in the ESI (Section 1†) and are only described in an abbreviated version here.

Air and moisture sensitive reactions were carried out following the standard Schlenk technique with Ar as an inert atmosphere. NMR spectra were recorded on a Bruker Advance III 300 MHz NMR spectrometer in deuterated solvents. IR spectra were recorded on a Bruker Alpha FTIR spectrometer.

Electrospray mass spectrometry (ESMS) was performed with a 6120 Quadrupole LC/MS (Agilent Technologies). A 1260 Infinity HPLC (Agilent Technologies) and a Nucleodur C18 Pyramid column (Macherey-Nagel) were used to separate the compounds for further analysis with ESMS. Total V concentrations were determined with a 7700x ICPMS (Agilent Technologies).

For speciation analysis, a 1260 Infinity HPLC (Agilent Technologies) was coupled to a 7700x ICPMS or a tandem ICPMS (ICPMS/MS, 8900, Agilent Technologies). A silica-based strong anion-exchange column (Zorbax SAX, 4.6 \times 150 mm, 5 μ m, Agilent Technologies) was used for the separation of the V species.

The water content of the mushroom samples was determined with a SMART Trac II Moisture & Fat Analyzer (CEM). An Ultraclave IV (MLS GmbH) was used for microwave assisted digestions. Extracts were centrifuged on a Rotina 420 R (Hettich Lab Technology). Nylon syringe filters (0.22 μ m, Simplepure) were used for filtering the extracts prior to analysis.

2.2 Synthesis

2.2.1 Dimethyl 2,2'-(hydroxyazanediy)l (2R,2'R)-dipropionate (1). In a flame dried, Ar flushed Schlenk flask, methyl-L-lactate (1.40 g, 13.4 mmol) was dissolved in dry methylene chloride (40 mL), and the solution was cooled to -60 $^{\circ}$ C. Tf₂O (1 M in dry methylene chloride, 14.7 mL) was added, followed by lutidine (3.16 g, 29.5 mmol) after 10 minutes which caused the solution to turn pale red immediately. After another 10 minutes, hydroxylamine solution (1 M in MeOH, 5.36 mL) was added, and the cooling bath was removed. The reaction mixture was slightly opaque, but after warming up to room temperature, all solids were dissolved. After stirring for 16 h, the mixture was quenched by the addition of water (20 mL), the phases were separated, and the aqueous phase was extracted with ethyl acetate (2 \times 20 mL). The combined organic phases were dried (MgSO₄) and the solvent was removed on a rotary evaporator. Purification by SiO₂ flash chromatography gave 664 mg (48%) of **1** as a pale-yellow resin. NMR spectra can be found in the ESI (Fig. S1 and S2†).

¹H NMR (300 MHz, CDCl₃) δ = 5.79 (s, 1H), 5.29 (s, 2H), 3.71 (q, J = 6.9, 2H), 3.70 (s, 6H), 1.33 (d, J = 6.9, 6H). ¹³C NMR (75 MHz, CDCl₃) δ = 173.2, 60.9, 53.4, 52.2, 12.6.

2.2.2 (2R,2'R)-2,2'-(Hydroxyazanediy)l dipropionic acid (2). In a round bottom flask, KOH (85% by weight, 310 mg, 6.5 mmol) was dissolved in water (20 mL). **1** (532 mg, 2.59 mmol) was added, and the mixture was stirred at room temperature for 1 hour. The product suspension was acidified to a pH of approx. 5 by the addition of 2 M HCl. Upon addition of Zn(OAc)₂ (1 M in H₂O, 3 mL), a colorless precipitate was formed. The suspension was transferred into test tubes, centrifuged for ten minutes, and the liquid was disposed of. The solid was washed twice with water and centrifuged again. The isolated colorless solid was dissolved in 5 M HCl (approx. 4 mL) and applied onto a column packed with Dowex® 50WX2 hydrogen form ion-exchange resin that was thoroughly rinsed with water prior to use, until the eluent had a steady pH of \sim 4. Fractions were collected and checked for product by spotting on a thin layer chromatographic plate, thorough drying and derivatization with KMnO₄ staining solution. The removal of water on a rotary evaporator gave 359 mg (86%) of **2** as a colorless solid. See ESI, Fig. S3 and S4† for NMR spectra and Fig. S5† for the IR spectrum.

¹H NMR (300 MHz, D₂O) δ = 4.02 (q, J = 7.0, 2H), 1.41 (d, J = 7.0, 6H). ¹³C NMR (75 MHz, D₂O) δ = 174.5, 62.7, 11.6. ESMS: m/z calc for [M + 1]⁺: 178, found: 178. IR: 1747 (ν_{CO}) cm⁻¹.

2.2.3 Vanadyl acetate, VO(OAc)₂. A suspension of V₂O₅ (1.00 g, 5.49 mmol) in acetic anhydride (50 mL) was brought to a reflux for 3 h, during which the suspension turned from orange to pale green. The hot product suspension was filtered, and the residue was washed with chloroform (3 \times 5 mL) and dried *in vacuo* to give 1.87 g (92%) of vanadyl acetate (Fig. 1) as a pale green solid. The IR spectrum is shown in ESI, Fig. S6.†

ESMS: m/z calculated for [M + 1]⁺: 186; m/z found: 158, corresponding to [VO(OOCH)₂ + H]⁺ – owing to the formic acid in the mobile phase. IR: 1508 (ν_{CO}), 894 (ν_{VO}) cm⁻¹.

2.2.4 Amavadin. A solution of **2** (219 mg, 1.24 mmol) in water (5 mL) was added to a suspension of vanadyl acetate (114 mg, 0.62 mmol) in water (3 mL). The mixture was stirred for 1 hour and the formed blue solution was filtered through a cotton plug, and the solvent was removed on a rotary evaporator. The crude product was dried *in vacuo* to give a 243 mg bright blue solid. It was a mixture of amavadin (74% w/w, 0.45 mmol, 73% yield) and vanadyl acetate. Because of the paramagnetic character of V(IV), no NMR of the product could be obtained. The IR spectrum is shown in ESI, Fig. S7.†

ESMS: m/z calc for [M + 3H]⁺: 402, found: 402. IR: 1561 (ν_{CO}) cm⁻¹.

2.3 Confirmation of the structure and concentration

2.3.1 Structural confirmation by ESMS. For HPLC-ESMS measurements, a reversed-phase Nucleodur C18 Pyramid column was used as the stationary phase. The mobile phase consisted of 0.1% formic acid in water (A) and methanol (B), with 70% A and 30% B. The ESMS was operated in positive ion mode. In scan mode, the m/z range 120–600 was scanned, and in SIM mode, m/z 158, 186 and 402 were monitored. More



details on the settings of the HPLC-ESMS measurements can be found in the ESI. Amavadin and vanadyl acetate were dissolved in ultrapure water (although the latter one was only poorly soluble⁶⁰) for the ESMS analyses.

2.3.2 Determination of the standard's concentration via ICPMS. Amavadin was dissolved in ultrapure water and then further diluted with 10% nitric acid. Vanadyl acetate was dissolved in 10% nitric acid and further diluted as well. The V concentration of both standards was determined with ICPMS in He collision gas mode. Details on the measurements are given in Table 1. An external calibration was used for quantification, with a V standard in 10% nitric acid, at concentrations from 0.1 to 10 $\mu\text{g L}^{-1}$. The Standard Reference Material (SRM) 1640a (trace elements in natural water, $n = 1$) was used to evaluate the measurement's accuracy (trueness: 93%).

Ge (m/z 74, 200 $\mu\text{g L}^{-1}$ in 10% nitric acid) was used as the internal standard. It was added online *via* a T-piece in front of the nebulizer to the calibration standards and the sample. Its tubing's diameter was 1/4 of the sample tubing.

2.4 HPLC-ICPMS method development

Because amavadin carries two negative charges, an anion-exchange column (Zorbax SAX) and an aqueous mobile phase with ammonium citrate (citric acid, pH adjustment with concentrated aqueous ammonia) were employed. Na_2EDTA (10

mM) was added to the mobile phase for complexing all present smaller V species and thus prohibiting unwanted precipitation or interconversion. Different column temperatures (5–45 °C), concentrations of citric acid (2–50 mM) and pH values (4.0–7.5) were tested to optimize the separation method. The evaluated parameters and also the final method are summarized in Table 1. A standard containing 10 $\mu\text{g V L}^{-1}$ of amavadin and vanadyl acetate was used for the experiments, dissolved and diluted in the initial mobile phase (Table 1, “starting values”).

The peaks were integrated at the baseline, with a vertical division at the valley between the two amavadin isomers.

The resolution R of the amavadin's two isomers was calculated according to the following formula, where t_{R1} and t_{R2} are the retention times at the peak maxima, and $w_{0.5 h1}$ and $w_{0.5 h2}$ are the peak widths at 50% peak height.

$$R = 1.18 \times \frac{(t_{R2} - t_{R1})}{(w_{0.5h1} + w_{0.5h2})}$$

The optimized method was validated by performing a calibration from 0.1 to 100 $\mu\text{g V L}^{-1}$ (calibration points: 0, 0.1, 0.5, 1, 5, 10, 50, and 100 $\mu\text{g V L}^{-1}$) of amavadin (sum of both isomers) and vanadyl acetate. The two standards were mixed together and diluted with the mobile phase (Table 1). The lowest standard was injected 5 times to determine the limit of detection (LOD) and limit of quantification (LOQ). The LOD was calculated as three times the standard deviation, and the LOQ ten times the standard deviation of the lowest standard. A standard solution with 10 $\mu\text{g V L}^{-1}$ of amavadin and vanadyl acetate was injected six times over the course of a 24 hour long measurement to evaluate the method's stability. The SRM 1640a ($n = 1$) was diluted 1 + 1 with the mobile phase and analyzed with the developed method to determine the method's accuracy.

2.5 Mushroom samples

2.5.1 Collection and cleaning. Three fruit bodies of *Amanita muscaria* (named A, B and C) were collected in Lachtal, Austria, in August 2020. The samples were kept at 4 °C overnight, and then brought to the lab and thoroughly cleaned with a small brush and moist paper towels. They were separated into their different parts, as shown in Fig. 2, namely the cap, stipe, gills, cap skin (hereafter referred to as “skin”, removed from the cap with a scalpel) and the outer part of the bulb (the lower part of the stipe), which was a mixture of bulb, mycelium and residual soil (hereafter referred to as “bulb”). In order to keep as much mycelium as possible, we refrained from removing the soil too thoroughly from the bulb samples. Each sample was homogenized with a small blender (M 20 Universal mill, IKA, Staufen, Germany). The homogenates were then stored at 4 °C and prepared for total element determination and speciation analysis within 24 hours of cleaning. The water content of the homogenized samples was determined with a SMART Trac II Moisture & Fat Analyzer.

2.5.2 Sample preparation. For the determination of the total V concentrations, around 2 g of cap, stipe and gills, and

Table 1 Details on the ICPMS settings, the optimized HPLC-ICPMS method and the parameters during method optimization

ICPMS measurements	
Instrument	Agilent 7700x
Collision gas mode	He (5 mL min ⁻¹)
Monitored m/z	51 (V), 74 (Ge, internal standard)
Integration time	0.1 s per m/z
Intensity of 1 $\mu\text{g V L}^{-1}$	40 kcps
Oxide ratio (m/z 156/140, ¹⁴⁰ Ce ¹⁶ O ⁺ / ¹⁴⁰ Ce ⁺)	0.5%
Doubly charged ratio (m/z 70/140; ¹⁴⁰ Ce ²⁺ / ¹⁴⁰ Ce ⁺)	1.3%
Signal stability (RSD)	2–3%
Optimized HPLC-ICPMS method	
Instruments	Agilent HPLC 1260 Infinity and ICPMS 7700x or ICPMS/MS 8900
Collision gas mode	He (5 mL min ⁻¹)
Monitored m/z	51 (V)
Integration time	0.3 s
Column	Zorbax SAX, 4.6 × 150 mm, 5 μm
Mobile phase	10 mM citric acid, 10 mM Na ₂ EDTA, pH adjusted to 5.0 with aqueous ammonia
Flow rate	1.0 mL min ⁻¹
Column temperature	13 °C
Injection volume	25 μL
Injection speed	100 $\mu\text{L min}^{-1}$
Evaluated parameters during HPLC-ICPMS method optimization (starting values in bold)	
Concentrations	2, 5, 10, 20 , 30, 40, 50 mM citric acid
pH values	4.0, 4.5, 5.0, 5.3 , 5.5, 6.0, 6.5, 7.0, 7.5
Temperatures	5, 10, 13, 15, 20, 25 , 30, 35, 40, 45 °C






Fig. 2 Different parts of the fruit body of *Amanita muscaria*. *The cap flesh is beneath the skin and not visible.

0.3–1 g of skin or bulb (less biomass was available) were weighed into quartz vessels, each sample in triplicates. 5 mL of nitric acid were added (2 mL in the case of skin samples), and the samples were digested in a microwave assisted autoclave, with a loading pressure of 40 bar Ar and a maximum temperature of 250 °C. The digests were diluted with ultrapure water to a final acidity of 10% (v/v) nitric acid. For quality control, the SRM 1573a (tomato leaves, $n = 4$) and blanks ($n = 6$) were digested as well.

For V speciation analysis, around 2 g of cap, stipe and gills and 1 g of bulb and skin were extracted with 20 mL (10 mL in the case of the bulb and skin) of mobile phase (Table 1). The parts of sample A were extracted in triplicates, the parts of samples B and C only one time each, and three blank samples were prepared. After the addition of the mobile phase, the samples were thoroughly mixed, put in an ultrasonic bath for 20 minutes, centrifuged for 10 minutes, and then filtered through 0.22 μm Nylon filters. The extracts were subjected to HPLC-ICPMS analysis without further dilution. For the determination of the extraction efficiency, the extracts were diluted 1 + 99 with 10% (v/v) nitric acid.

2.5.3 Determination of total V

The total V concentrations in the digests and in the diluted extracts were measured with ICPMS, as already described earlier for the amavadin standard (Section 2.3.2). In addition to the digested SRM 1573a, the SRM 1640a ($n = 1$) was diluted 1 + 9 (10% (v/v) nitric acid) and analyzed as well.

2.5.4 V speciation analysis. The pure extracts were transferred to polypropylene HPLC vials with polyethylene caps and a natural rubber septum, and investigated with the newly developed V speciation analysis method. One replicate of each A sample was monitored for 60 minutes to look for strongly retained compounds. An external calibration was prepared, including amavadin and vanadyl acetate at concentrations between 0.1 and 100 $\mu\text{g V L}^{-1}$ (plus a blank). In the case of amavadin, the calibration line was calculated for the sum of its two isomers. Since the response of ICPMS is compound-independent, this calibration was used for the quantification of the two individual isomers and also for unknown V species.

The identity and accuracy of amavadin and vanadyl acetate in the mushroom extracts were confirmed by spiking one sample (gills A) online *via* a HPLC autosampler with standard solutions. For the confirmation of amavadin, 2 μL of extract were spiked with 20 μL of 100 $\mu\text{g V L}^{-1}$ of amavadin (see ESI, Fig. S11[†]). In the case of vanadyl acetate, 20 μL of extract were spiked with 1 μL of a 5 $\mu\text{g V L}^{-1}$ solution of vanadyl acetate. The ratios were selected to achieve a doubling of the peaks in question.

3 Results and discussion


3.1 Synthesis

Amavadin was synthesized employing a slightly modified version of the procedure reported by Hubregtse *et al.*³² The ligand was prepared from lactic acid methyl ester, starting with the conversion of the hydroxyl moiety into the corresponding triflate with triflic anhydride in the presence of lutidine, and subsequent nucleophilic substitution with hydroxylamine to give **1** (see Scheme 1). The hydrolysis of the methyl esters in aqueous KOH and precipitation with zinc acetate gave the desired ligand as the corresponding zinc salt. Ion exchange chromatography yielded a free ligand (**2**), that reacted with vanadyl acetate to form amavadin. A sample of 243 mg of dark blue crystals was obtained this way. Later analyses with ICPMS, HPLC-ICPMS and co-chromatography with vanadyl acetate revealed that 26% (w/w) of the obtained solid still was vanadyl acetate, and 74% (w/w) was amavadin (see also Section 3.2). This means that amavadin was synthesized with a yield of 180 mg, or 73%.

Due to the d^1 electron configuration of the V(IV) center, amavadin and vanadyl acetate both are paramagnetic, and thus NMR could not be performed on the product. The oxidation of V(IV) to V(V) in DMSO was attempted, as reported in the literature.³¹ A sample of amavadin in DMSO- d_6 was exposed to the atmosphere for several weeks, which caused the solution to turn from blue to red; however still no NMR spectrum was obtained. It is unclear whether amavadin could not be oxidized completely or the presence of paramagnetic vanadyl acetate made the sample immeasurable by standard NMR techniques.

The successful synthesis of amavadin was proven by ESMS and IR-spectroscopy. The IR spectrum (ESI, Fig. S7[†]) shows distinct peaks at wavenumbers reported in the literature.³² The mass spectrum of amavadin is shown in ESI, Fig. S8 and S9.[†]





Scheme 1 Synthesis of amavadin starting from (L)-lactic acid methyl ester.

The molecule peak for $[M + 3H]^+$ can be seen at m/z 402, besides lower peaks at m/z 401 and 403, and also sodium adducts at m/z 423, 424 and 425 are visible.

3.2 Method development

Because of the negative charges of amavadin, even at very low pH values,²² anion-exchange chromatography was selected as the best option for its chromatographic separation. Citric acid at pH 5.3 (adjusted with ammonia) in ultrapure water was selected as the starting point for the mobile phase. At this pH, citric acid has a dianionic character (pK_a values: 3.13, 4.76 and 6.40⁶¹), like amavadin, which should increase the elution strength of the buffer.

Amavadin was at first dissolved in ultrapure water. During method development, it was found that there was another V-containing compound in the amavadin standard. The first logical candidate was vanadyl acetate, the precursor that was used in the synthesis of amavadin. Thus, vanadyl acetate was dissolved in the initial mobile phase (Table 1, “starting values”) and analyzed along with amavadin. The first results from ESMS and HPLC-ICPMS showed that the impurity was indeed vanadyl acetate, which was subsequently included in the calibration standards. Quantification *via* ICPMS and HPLC-ICPMS (in the latter case with vanadyl acetate as the quantification standard and SRM 1640a as the reference material) showed that 43% of the total V of the “amavadin standard” was present in the form of vanadyl acetate (meaning that 26% of the solid material is vanadyl acetate).

Due to the poor solubility of vanadyl acetate in pure water, the amavadin standard was henceforth also dissolved in the mobile phase, to ensure that the vanadyl acetate moiety in the amavadin standard was quantitatively dissolved. In order to have similar concentrations of the two V species, additional vanadyl acetate was added to the amavadin standard. Over the course of method development, the relative peak areas of both species remained constant, which indicates that no interconversion of the two species was taking place.

Na_2EDTA was included in the mobile phase to ensure a quantitative elution of vanadyl acetate. This can also be beneficial for other small V species, which were not a target of the developed method, but can nevertheless occur in environmental samples. Vanadyl acetate (that is to say, very likely vanadyl complexed by EDTA) was eluting significantly earlier than amavadin as a clean, symmetric peak at all settings of the method development and will not be further discussed in this context.

Finally, 20 mM citric acid at pH 5.3 (adjusted with ammonia) with 10 mM Na_2EDTA was selected as the starting point (see also Table 1). The column oven was set at 25 °C, and the flow rate was 1 mL min^{-1} . With these first settings, a double peak appeared in the chromatogram at 7.1 and 7.5 minutes, corresponding to the two isomers of amavadin. To achieve a good resolution of these two peaks, the buffer concentration and pH value, as well as the column temperature, were optimized. The evaluated conditions are summarized in Table 1, and the resulting chromatograms are shown in Fig. 3.

The concentration of citric acid was varied between 2 and 50 mM. As can be seen from the resolution values depicted in Fig. 4A and the chromatograms in Fig. 3A, lowering the concentration of the buffer improved the separation of the two isomers. However, this also increased the retention time significantly, up to 20 and 21 minutes at 2 mM citric acid. Based on these findings, we found that a concentration of 10 mM provides a good balance between the retention time (second isomer: 8.9 min at the peak maximum, resulting in a total chromatogram time of 10 min) and resolution ($R = 0.56$) of the amavadin peaks.

As the next step, the column temperature was varied between 5 and 45 °C. With colder settings, the two peaks were better separated (Fig. 3 and 4B) but also slightly broadened. There was not much difference between 5 and 10 °C, and so 10 °C was selected as the best option ($R = 0.80$). However, it was later observed that under the given laboratory conditions longer experiments at 10 °C led to condensation in the column oven, which eventually resulted in a leak error in the HPLC. To overcome this problem, the temperature was slightly increased to 13 °C. This enabled a stable measurement for more than 24 hours, with still good resolution.

Finally, the influence of the pH value on the separation was evaluated. As can be seen in Fig. 3C and 4C, the two isomers were not separated any more at pH values above 6.5. Lower pH values led to a higher resolution, but while the retention times were 13.4 and 14.4 minutes at pH 5.0 ($R = 0.90$), they increased dramatically to over 26 minutes at pH 4.5 and more than 60 minutes at pH 4.0. This is not surprising, because citric acid will only have one negative charge at those pH values, while amavadin will still have a charge of -2 .

In summary, as can also be seen in Table 1, the final method included a Zorbax SAX (4.6 × 150 mm 5 μm) silica-based anion-exchange column and a mobile phase of 10 mM citric acid, pH 5.0, and 10 mM Na_2EDTA . The pH was adjusted with ammonia. The flow rate was 1.0 mL min^{-1} and the column temperature






Fig. 3 HPLC-ICPMS chromatograms (V at m/z 51) of amavadin ($5 \mu\text{g V L}^{-1}$, later-eluting double peak, corresponding to the two isomers) and vanadyl acetate ($5 \mu\text{g V L}^{-1}$, first peak) during method optimization. (A) Varying concentration of citric acid, at 25 °C and pH 5.3. The lowest concentration, 2 mM, is shown in the inset due to the much longer retention times. (B) Varying temperature, at 10 mM citric acid and pH 5.3. (C) Varying pH, at 10 mM citric acid and 10 °C. The chromatograms at pH 4.0 and 4.5 are shown in the inset due to the much longer retention times. Offsets of the y-axis in A, B and C: +3 kcps from the previous line. Exception: +1.5 kcps for pH 4.5 in the inset of C.

13 °C. The injection volume was set to 25 μL , and the injection speed to 100 $\mu\text{L min}^{-1}$.

Exemplary chromatograms of 10 and 0.1 $\mu\text{g V L}^{-1}$ are drawn in Fig. 5. Note that the retention times are a bit longer than in

the chromatograms obtained during method development (Fig. 3). An already used column was employed during method development. To make sure that no previous use had led to a modification of the stationary phase and therefore influence






Fig. 4 Resolution of the two amavadin isomers at different chromatographic settings. (A) Varying concentration of citric acid, at 25 °C and pH 5.3. (B) Varying temperature, at 10 mM citric acid and pH 5.3. (C) Varying pH, at 10 mM citric acid and 10 °C.

the separation of amavadin, a virgin column was used for method validation and application to mushroom samples. As mentioned, this led to an increase in the retention times, but the resolution of the two isomers was not affected.

3.3 Method validation

Although the two isomers of amavadin were almost baseline-separated, they were considered as one compound for method validation and calibration. The values obtained for the individual isomers were used to calculate the ratio of the two isomers.

The method's figures of merit are listed in Table 2. They include the LOD, LOQ, correlation coefficient R , stability of a 10 $\mu\text{g V L}^{-1}$ standard during a 24 hour measurement and the average repeatability of three replicates of the five mushroom parts of sample A. The low LODs of 0.04 and 0.05 $\mu\text{g V L}^{-1}$ for vanadyl acetate and amavadin, respectively, allow the detection of traces of amavadin or similar species. With a dilution factor of 10 (due to the extraction process), the LODs for mushroom

samples are still below 1 $\mu\text{g V kg}^{-1}$ (namely 0.4 and 0.5 $\mu\text{g V kg}^{-1}$).

The method is very stable; the detected amavadin concentrations only vary by around 2.6% (2.8% for vanadyl acetate) in the re-measured standard. Also, for the mushroom samples, the repeatability is lower than 5%. The amavadin concentrations in the three replicates of sample A varied between 1.7 and 5.6% for the five different fruit-body parts, while the variations of the vanadyl acetate concentrations were $4.6 \pm 2.1\%$. The highest relative standard deviations (RSDs) were found in the skin sample (5.6% and 7.8% for amavadin and vanadyl acetate, respectively), which can be explained by the difficult homogenization of this part of the mushroom. The accuracy of the measured SRM, NIST 1640a, was excellent, namely 98.4% of the total V. The V speciation in this SRM is not defined, but V had the same retention time as vanadyl acetate (probably eluted as vanadyl EDTA). The method's accuracy was further confirmed by the recoveries after online spiking of an extract (gills A) with amavadin and vanadyl acetate, which were 90 and 105%, respectively. Another indirect proof of the method's accuracy is




Fig. 5 HPLC-ICPMS chromatograms of V (m/z 51) of the lowest (0.1 $\mu\text{g V L}^{-1}$) and a medium concentrated (10 $\mu\text{g V L}^{-1}$) standard of amavadin and vanadyl acetate, obtained with the optimized method parameters.



Table 2 Method figures of merit; calibration range: 0.1–100 $\mu\text{g V L}^{-1}$; stability = RSD of six times re-measured 10 $\mu\text{g V L}^{-1}$ standard; repeatability = mean \pm standard deviation of the RSD of the three replicates of each mushroom part of sample A. Recovery = results from online spiking an extract (gills A) with the standards

	LOD [$\mu\text{g V L}^{-1}$]	LOQ [$\mu\text{g V L}^{-1}$]	R	Stability [%]	Repeatability [%]	Recovery [%]
Vanadyl acetate	0.04	0.13	0.9999	2.8	4.6 \pm 2.1	90
Amavadin	0.05	0.17	0.9999	2.6	3.7 \pm 1.7	105

the quantitative column recovery of the mushroom extracts (see below, Section 3.4.3).

3.4 Mushroom samples

3.4.1 Water content. The water content varied between 88 and 94% for all samples, except the bulb samples, which contained 72, 86 and 84% water (samples A, B and C, respectively). Concentrations will be given on a wet mass basis, because the whole experiments were performed with fresh samples.

3.4.2 Total V and extractable V concentrations. In the digested SRM 1573a, we detected 90 \pm 1% ($n = 4$) of the certified V concentration, and for the SRM 1640a, the trueness was 95% ($n = 1$) when determining the V concentrations in the digested samples, and 98% ($n = 1$) when the total extractable V was analyzed.

Concerning the samples of *Amanita muscaria*, the highest V concentrations, namely 38–97 mg kg^{-1} , were found in the bulb samples. In the other samples, the V concentrations were between 6.4 and 33 mg kg^{-1} . Within each mushroom, the concentrations in the stipe and gills were very similar. Slightly lower concentrations were found in the skin, and the caps had the lowest V concentrations among all fruit-body parts. All total V concentrations, along with the extractable V concentrations and V species, are listed in Table 3.

Overall, 74 \pm 12% of the total V was extractable with the mobile phase. The repeatability of the samples of fruit-body A

was between 1 and 7% (three replicate extractions per sample). The bulb samples had the highest extraction efficiencies of 81–93%, while the lowest extraction efficiencies were found for the skin samples, with 50–63%. It will be interesting to see in future studies whether an extended extraction time or higher temperature can lead to an increased extraction efficiency in *Amanita muscaria* samples.

3.4.3 V species. The concentrations of the detected V species in the fruit-body samples are given in Table 3. The identity of amavadin was confirmed by spiking a diluted extract of gills A with a standard solution, containing 100 $\mu\text{g V L}^{-1}$ of amavadin and vanadyl acetate (see chromatograms in ESI, Fig. S11†). The data showed quantitative column recoveries (sum of all detected V species divided by the extractable V), with 98 \pm 9%. Amavadin accounted for 4.0–83 mg V kg^{-1} , which is 75–96% of the extractable V. It was not possible to determine which peak was the Λ - and which one the Δ -isomer, and thus the earlier eluting isomer will be referred to as isomer 1, and the later eluting one as isomer 2. The ratio of isomer 2 to isomer 1 was very constant, between 1.9 and 2.2, in all samples except the bulbs. There, the ratio was 2.4, 2.4 and 2.3 for samples A, B and C, respectively. Interestingly, the ratio was only 0.90 \pm 0.05 in the synthetic standard solutions. In other words, the two isomers were present at almost equal concentrations in the synthetic amavadin, while the concentrations of isomer 2 were more than twice as high as that of isomer 1 in *Amanita muscaria*

Table 3 Concentrations of total V, extractable V and V species in three fresh fruit bodies of *Amanita muscaria*, in mg V kg^{-1} , and extraction efficiencies in % of total V. Sample A was extracted three times, whereas B and C were only extracted once.

Sample	Total V	Extractable V	Extraction efficiency	Amavadin	Vanadyl acetate ^a	Unknown 1	Other species
Skin A	8.5 \pm 0.6	4.2 \pm 0.3	49 \pm 4	4.0 \pm 0.2	0.0040 \pm 0.0003	0.27 \pm 0.01	0.32 \pm 0.03
Cap A	6.4 \pm 0.3	5.1 \pm 0.3	80 \pm 5	4.8 \pm 0.2	Traces ^b	0.32 \pm 0.03	0.19 \pm 0.02
Gills A	13.5 \pm 0.3	9.3 \pm 0.1	69 \pm 1	8.1 \pm 0.1	0.0035 \pm 0.0001	0.50 \pm 0.03	0.20 \pm 0.01
Stipe A	16.4 \pm 2.2	11.0 \pm 0.8	67 \pm 5	10.3 \pm 0.6	0.037 \pm 0.001	0.48 \pm 0.04	0.28 \pm 0.09
Bulb A	38.1 \pm 0.4	31.0 \pm 1.0	81 \pm 3	26.2 \pm 0.8	0.60 \pm 0.01	1.21 \pm 0.03	0.59 \pm 0.13
Skin B	18.0 \pm 1.3	11	63	11	0.0088	0.63	0.26
Cap B	13.1 \pm 0.4	11	85	11	0.0030	0.63	0.17
Gills B	25.6 \pm 0.8	20	79	16	0.0048	0.92	0.27
Stipe B	25.3 \pm 0.4	18	70	17	0.075	0.71	0.31
Bulb B	58.1 \pm 1.6	51	88	38	1.8	1.6	1.1
Skin C	19.5 \pm 0.9	12	60	11	< 0.0004	0.92	0.35
Cap C	18.1 \pm 0.5	13	73	12	0.0033	0.66	0.34
Gills C	33.0 \pm 1.2	23	71	22	0.0086	1.1	0.58
Stipe C	27.5 \pm 0.7	23	84	21	0.068	0.82	0.51
Bulb C	97.4 \pm 0.7	90	93	83	1.4	2.9	1.2

^a Species detected at the same retention time as the vanadyl acetate standard (vanadyl EDTA). ^b Traces: between the LOD and LOQ (0.0004 and 0.0013 mg V kg^{-1}).



4 Conclusion and outlook

We synthesized amavadin and developed a sensitive anion-exchange HPLC-ICPMS method for the separation and detection of amavadin and its two diastereoisomers with a limit of quantification of $0.17 \mu\text{g V L}^{-1}$.

We applied the method on the extracts of fruit-body samples of *Amanita muscaria*. 75–96% of the extracted V was found in the form of amavadin. We also found other V species in the extracts, of which one was eluting with a similar double-peak profile to the two diastereoisomers of amavadin. Additionally, long retained V containing signals were detected in all the investigated parts of the mushroom (bulb, stipe, cap, skin, and gills).

Our results add valuable and unique information to our knowledge on amavadin in mushrooms. The new method allows vanadium speciation analysis at trace concentrations in mushrooms and other environmental samples.

Until now, it was assumed that only *Amanita muscaria* and its few closely related species contain amavadin, because of their extraordinary V concentrations. With the developed method, it can be explored whether traces of amavadin are present in other environmental samples as well. This will help in elucidating the biogeochemical pathway of V and its role in the environment.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors want to thank Evelyn Waldhauser, Antonia Toedling and Linda Maierhofer for their support in the lab. The authors acknowledge the financial support by the University of Graz.

References

- Agency for Toxic Substances and Disease Registry (ATSDR), *Toxicological Profile for Vanadium*, U.S. Department of Health and Human Services, Public Health Service, Atlanta, GA, 2012.
- D. Rehder, Vanadium in health issues, *ChemTexts*, 2018, **4**, 20.
- D. Rehder, The future of/for vanadium, *Dalton Trans.*, 2013, **42**, 11749–11761.
- S. Treviño and A. Diaz, Vanadium and insulin: Partners in metabolic regulation, *J. Inorg. Biochem.*, 2020, **208**, 111094.
- D. Rehder, The potentiality of vanadium in medicinal applications, *Inorg. Chim. Acta*, 2020, **504**, 119445.
- T. I. Fortoul, V. Rodríguez-Lara, A. González-Villalva, M. Rojas-Lemus, G. Cano-Gutiérrez, M. Ustarroz-Cano, L. Colín-Barenque, P. Bizarro-Nevarés, I. García-Pealez, L. F. Montaña, R. S. Jimenez-Martinez, N. Lopez-Valdez, M. L. Ruiz-Guerrero, N. A. Meléndez-García, F. A. García-Ibarra, V. Martínez-Baez, D. Z. Alfaro, A. Muñoz-Rivera-Cambas, L. S. López-Zepeda, E. M. Quezada-Maldonado and S. Cervantes-Yépez, Inhalation of vanadium pentoxide and its toxic effects in a mouse model, *Inorg. Chim. Acta*, 2014, **420**, 8–15.
- F. L. Assem and L. S. Levy, A review of current toxicological concerns on vanadium pentoxide and other vanadium compounds: Gaps in knowledge and directions for future research, *J. Toxicol. Environ. Health, Part B*, 2009, **12**, 289–306.
- K. Gruzewska, A. Michno, T. Pawelczyk and H. Bielarczyk, Essentiality and toxicity of vanadium supplements in health and pathology, *J. Physiol. Pharmacol.*, 2014, **65**, 603–611.
- J. P. Gustafsson, Vanadium geochemistry in the biogeosphere –speciation, solid-solution interactions, and ecotoxicity, *Appl. Geochem.*, 2019, **102**, 1–25.
- T. Ueki, N. Yamaguchi, Romaidi, Y. Isago and H. Tanahashi, Vanadium accumulation in ascidians: A system overview, *Coord. Chem. Rev.*, 2015, **301–302**, 300–308.
- J. Costa Pessoa, E. Garribba, M. F. A. Santos and T. Santos-Silva, Vanadium and proteins: Uptake, transport, structure, activity and function, *Coord. Chem. Rev.*, 2015, **301–302**, 49–86.
- D. Rehder, The role of vanadium in biology, *Metallomics*, 2015, **7**, 730–742.
- Z. Řanda and J. Kučera, Trace elements in higher fungi (mushrooms) determined by activation analysis, *J. Radioanal. Nucl. Chem.*, 2004, **259**, 99–107.
- P. Kalač, Trace element contents in European species of wild growing edible mushrooms: A review for the period 2000–2009, *Food Chem.*, 2010, **122**, 2–15.
- J. Vetter, Vanadium content of some common edible, wild mushroom species, *Acta Aliment.*, 1999, **28**, 39–48.
- J. Falandysz, T. Kunito, R. Kubota, K. Lipka, A. Mazur, J. J. Falandysz and S. Tanabe, Selected elements in fly agaric *Amanita muscaria*, *J. Environ. Sci. Health, Part A: Toxic/Hazard. Subst. Environ. Eng.*, 2007, **42**, 1615–1623.
- E. Koch, H. Kneifel and E. Bayer, Das Vorkommen von Amavadin in Pilzen der Gattung *Amanita* [in German], *Z. Naturforsch.*, 1987, **873–878**.
- J. Vetter, Mineral composition of basidiomes of *Amanita* species, *Mycol. Res.*, 2005, **109**, 746–750.
- H.-U. Meisch, J. A. Schmitt and W. Reinle, Schwermetalle in Höheren Pilzen, III * Vanadium und Molybdän/Heavy Metals in Higher Fungi, II I. Vanadium and Molybdenum, *Z. Naturforsch., C: J. Biosci.*, 1978, **33**, 1–6.
- R. D. Gillard and R. J. Lancashire, Electron spin resonance of vanadium in *Amanita muscaria*, *Phytochemistry*, 1984, **23**, 179–180.
- H.-U. Meisch, W. Reinle and J. A. Schmitt, High vanadium content in mushrooms is not restricted to the fly agaric (*Amanita muscaria*), *Sci. Nat.*, 1979, **66**, 620–621.
- E. Bayer and H. Kneifel, Isolation of Amavadin, a Vanadium Compound Occurring in *Amanita Muscaria*, *Z. Naturforsch., B: Anorg. Chem., Org. Chem., Biochem., Biophys., Biol.*, 1972, **27**, 207.
- H. Kneifel and E. Bayer, Determination of the Structure of the Vanadium Compound, Amavadin, from Fly Agaric, *Angew. Chem., Int. Ed.*, 1973, **12**, 508.



- 24 C. D. Garner, E. M. Armstrong, R. E. Berry, R. L. Beddoes, D. Collison, J. J. A. Cooney, S. N. Ertok and M. Helliwell, Investigations of Amavadin, *J. Inorg. Biochem.*, 2000, **80**, 17–20.
- 25 J. A. L. da Silva, F. da Silva, J. R. João and A. J. L. Pombeiro, Amavadin, a vanadium natural complex: Its role and applications, *Coord. Chem. Rev.*, 2013, **257**, 2388–2400.
- 26 F. J. J. R. da Silva, Vanadium in biology—the case of the *Amanita* toadstools, *Chem. Speciation Bioavailability*, 1989, **1**, 139–150.
- 27 P. Krauß, E. Bayer and H. Kneifei, Elektronenspinresonanz-Untersuchungen an Amavadin, einem vanadiumhaltigen Naturstoff/Electron Spin Resonance Investigations of Amavadin, a Natural Product Containing Vanadium, *Z. Naturforsch., B: Anorg. Chem., Org. Chem.*, 1984, **39**, 829–832.
- 28 H. Kneifel and E. Bayer, Stereochemistry and total synthesis of amavadin, the naturally occurring vanadium compound of *Amanita muscaria*, *J. Am. Chem. Soc.*, 1986, **108**, 3075–3077.
- 29 E. Bayer, E. Koch and G. Anderegg, Amavadin, an Example for Selective Binding of Vanadium in Nature: Studies of Its Complexation Chemistry and a New Structural Proposal, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 545–546.
- 30 E. M. Armstrong, R. L. Beddoes, L. J. Calviou, J. M. Charnock, D. Collison, N. Ertok, J. H. Naismith and C. D. Garner, The Chemical Nature of Amavadin, *J. Am. Chem. Soc.*, 1993, **115**, 807–808.
- 31 E. M. Armstrong, D. Collison, N. Ertok and C. D. Garner, NMR studies on natural and synthetic Amavadin, *Talanta*, 2000, **53**, 75–87.
- 32 T. Hubregtse, E. Neeleman, T. Maschmeyer, R. A. Sheldon, U. Hanefeld and I. W. C. E. Arends, The first enantioselective synthesis of the amavadin ligand and its complexation to vanadium, *J. Inorg. Biochem.*, 2005, **99**, 1264–1267.
- 33 T. Hubregtse, H. Kooijman, A. L. Spek, T. Maschmeyer, R. A. Sheldon, I. W. C. E. Arends and U. Hanefeld, Study on the isomerism in meso-amavadin and an amavadin analogue, *J. Inorg. Biochem.*, 2007, **101**, 900–908.
- 34 C. M. M. Matoso, A. J. L. Pombeiro, J. J. R. F. da Silva, M. F. C. G. da Silva, J. A. L. da Silva, J. L. Baptista-Ferreira and F. Pinho-Almeida, in *Vanadium Compounds*, ed. A. S. Tracey and D. C. Crans, American Chemical Society, Washington, DC, 1998, vol. 711, pp. 241–247.
- 35 M. Domarus, M. L. Kuznetsov, J. Marçalo, A. J. L. Pombeiro and J. A. L. da Silva, Amavadin and Homologues as Mediators of Water Oxidation, *Angew. Chem., Int. Ed.*, 2016, **55**, 1489–1492.
- 36 V. Ugone, D. Sanna, G. Sciortino, D. C. Crans and E. Garribba, ESI-MS Study of the Interaction of Potential Oxidovanadium(IV) Drugs and Amavadin with Model Proteins, *Inorg. Chem.*, 2020, **59**, 9739–9755.
- 37 J.-H. Huang, F. Huang, L. Evans and S. Glasauer, Vanadium: Global (bio)geochemistry, *Chem. Geol.*, 2015, **417**, 68–89.
- 38 K. Pyrzyńska and T. Wierbicki, Determination of vanadium species in environmental samples, *Talanta*, 2004, **64**, 823–829.
- 39 G. P. Dechaine and M. R. Gray, Chemistry and Association of Vanadium Compounds in Heavy Oil and Bitumen, and Implications for Their Selective Removal, *Energy Fuels*, 2010, **24**, 2795–2808.
- 40 T. Jakusch, J. Costa Pessoa and T. Kiss, The speciation of vanadium in human serum, *Coord. Chem. Rev.*, 2011, **255**, 2218–2226.
- 41 T. Iglesias-González, C. Sánchez-González, M. Montes-Bayón, J. Llopis-González and A. Sanz-Medel, Absorption, transport and insulin-mimetic properties of bis(maltolato)oxovanadium (IV) in streptozotocin-induced hyperglycemic rats by integrated mass spectrometric techniques, *Anal. Bioanal. Chem.*, 2012, **402**, 277–285.
- 42 K. De Cremer, M. van Hulle, C. Chéry, R. Cornelis, K. Strijckmans, R. Dams, N. Lameire and R. Vanholder, Fractionation of vanadium complexes in serum, packed cells and tissues of Wistar rats by means of gel filtration and anion-exchange chromatography, *J. Biol. Inorg. Chem.*, 2002, **7**, 884–890.
- 43 V. Nischwitz, J. T. Davies, D. Marshall, M. González, J. L. Gómez Ariza and H. Goenaga-Infante, Speciation studies of vanadium in human liver (HepG2) cells after *in vitro* exposure to bis(maltolato)oxovanadium(IV) using HPLC online with elemental and molecular mass spectrometry, *Metallomics*, 2013, **5**, 1685.
- 44 N. Kilbarda, S. E. Afton, J. M. Harrington, F. Yan and K. E. Levine, Rapid speciation and determination of vanadium compounds using ion-pair reversed-phase ultra-high-performance liquid chromatography inductively coupled plasma-sector field mass spectrometry, *J. Chromatogr. A*, 2013, **1304**, 121–126.
- 45 X. S. Li and X. C. Le, Speciation of vanadium in oils and coke and bacterial culture by high performance liquid chromatography inductively coupled plasma mass spectrometry, *Anal. Chim. Acta*, 2007, **602**, 17–22.
- 46 T. V. Komarova, O. N. Obrezkov and O. A. Shpigun, Ion chromatographic behaviour of anionic EDTA complexes of vanadium(IV) and vanadium(V), *Anal. Chim. Acta*, 1991, **254**, 61–63.
- 47 L. Minelli, E. Veschetti, S. Giammanco, G. Mancini and M. Ottaviani, Vanadium in Italian waters: monitoring and speciation of V(IV) and V(V), *Microchem. J.*, 2000, **67**, 83–90.
- 48 F. Aureli, S. Ciardullo, M. Pagano, A. Raggi and F. Cubadda, Speciation of vanadium(IV) and (V) in mineral water by anion exchange liquid chromatography-inductively coupled plasma mass spectrometry after EDTA complexation, *J. Anal. At. Spectrom.*, 2008, **23**, 1009–1016.
- 49 M. A. Larsson, M. D'Amato, F. Cubadda, A. Raggi, I. Öborn, D. B. Kleja and J. P. Gustafsson, Long-term fate and transformations of vanadium in a pine forest soil with added converter lime, *Geoderma*, 2015, **259–260**, 271–278.
- 50 M. Colina, P. H. E. Gardiner, Z. Rivas and F. Troncone, Determination of vanadium species in sediment, mussel and fish muscle tissue samples by liquid chromatography-inductively coupled plasma-mass spectrometry, *Anal. Chim. Acta*, 2005, **538**, 107–115.



- 51 Z. L. Chen and G. Owens, Trends in speciation analysis of vanadium in environmental samples and biological fluids—A review, *Anal. Chim. Acta*, 2008, **607**, 1–14.
- 52 Z. Chen and R. Naidu, On-column complexation and simultaneous separation of vanadium(IV) and vanadium(V) by capillary electrophoresis with direct UV detection, *Anal. Bioanal. Chem.*, 2002, **374**, 520–525.
- 53 K. Pyrzyńska and T. Wierzbicki, Solid-phase extraction for preconcentration and separation of vanadium species in natural waters, *Microchim. Acta*, 2004, **147**, 59–64.
- 54 J.-Y. Yang and Y. Tang, Accumulation and Biotransformation of Vanadium in *Opuntia microdasys*, *Bull. Environ. Contam. Toxicol.*, 2015, **94**, 448–452.
- 55 D. Wang and S. A. Sañudo-Wilhelmy, Development of an analytical protocol for the determination of V(IV) and V(V) in seawater: Application to coastal environments, *Mar. Chem.*, 2008, **112**, 72–80.
- 56 M. Žižić, T. Dučić, D. Grolimund, D. Bajuk-Bogdanović, M. Nikolic, M. Stanić, S. Križak and J. Zakrzewska, X-ray absorption near-edge structure micro-spectroscopy study of vanadium speciation in *Phycomyces blakesleeanus* mycelium, *Anal. Bioanal. Chem.*, 2015, **407**, 7487–7496.
- 57 P. Frank, K. Kustin, W. E. Robinson, L. Linebaugh and K. O. Hodgson, Nature and Ligation of Vanadium within Whole Blood Cells and Henze Solution from the Tunicate *Ascidia ceratodes*, As Investigated by Using X-ray Absorption Spectroscopy, *Inorg. Chem.*, 1995, **34**, 5942–5949.
- 58 M. Jensen-Fontaine, W. P. Norwood, M. Brown, D. G. Dixon and X. C. Le, Uptake and Speciation of Vanadium in the Benthic Invertebrate *Hyalella azteca*, *Environ. Sci. Technol.*, 2014, **48**, 731–738.
- 59 V. T. Kütter, M. Montes-Bayón, S. M. Sella, A. Sanz-Medel and E. V. Silva-Filho, Vanadium-binding protein in marine plankton from tropical south atlantic ocean, *J. Braz. Chem. Soc.*, 2014, **25**, 1116–1123.
- 60 D. C. Crans, J. J. Smee, E. Gaidamauskas and L. Yang, The Chemistry and Biochemistry of Vanadium and the Biological Activities Exerted by Vanadium Compounds, *Chem. Rev.*, 2004, **104**, 849–902.
- 61 A. M. N. Silva, X. Kong and R. C. Hider, Determination of the pK_a value of the hydroxyl group in the α -hydroxycarboxylates citrate, malate and lactate by ^{13}C NMR: implications for metal coordination in biological systems, *BioMetals*, 2009, **22**, 771–778.

