

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Supramolecular assemblies of semiconductor quantum dots and a bis(bipyridinium) derivative: luminescence quenching and aggregation phenomena

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Marek Oszajca,^{a,b} Christophe Lincheneau,^a Matteo Amelia,^a Massimo Baroncini,^a Serena Silvi,^a Konrad Szaciłowski,^{b,c} and Alberto Credi*^a

We have synthesized CdSe and CdSe-ZnS core-shell luminescent nanocrystal quantum dots and studied their interaction with a ditopic bis(bipyridinium) compound in solution. The latter strongly quenches the luminescence of the quantum dots by a static mechanism, indicating that the nanocrystal and molecular components undergo association in the ground state. Photoexcitation of these inorganic-organic hybrids causes an electron-transfer process from the conduction band of the nanocrystal to the LUMO of the molecule. The ability of the bipyridinium-type species to trigger association of the quantum dots is evidenced by spectrofluorimetric titrations and DLS measurements in solution, and confirmed by TEM experiments on surfaces. The quantum dot-molecule complexes can be disassembled in solution by addition of a calixarene host capable of encapsulating the bipyridinium units of the molecular connector. Our results demonstrate that supramolecular chemistry offers convenient ways to control the aggregation of semiconductor nanocrystals, a crucial task for the generation of nanostructured arrays with well defined properties.

Introduction

Quantum dots (QDs) are semiconductor nanocrystals with diameters ranging between 1 and 15 nm and are characterized by the quantum confinement of the charge carriers.¹ Quantum confinement confers peculiar optoelectronic properties² to QDs – in particular, a large molar absorption coefficient in the UV-visible and an intense size-tunable luminescence band with a narrow spectral profile – that render them valid substitutes for molecular luminophores³ for a variety of technological applications.⁴ Moreover such properties, together with a high thermal and photochemical stability, and versatile surface functionalization, make QDs attractive components for the construction of photoactive nanocrystal-molecule assemblies.^{5,6}

The vast majority of QD-molecule nano hybrids reported to date are based on the covalent attachment of the molecular units to the nanocrystal surface.⁶ Such a strategy requires the synthesis of derivatives in which the molecule of interest is connected to a ligand moiety for the QD surface, and the optimization of exchange procedures in order to replace the native ligands of the nanocrystal with the functional ones. Several recent reports, however, have shown^{7,8} that bare semiconductor nanocrystals and functional molecular

components can be self-assembled in an efficient manner taking advantage of supramolecular interactions in solution. This approach is advantageous because (i) no exchange of the capping ligands of the quantum dots is required, (ii) the molecular components do not need to have a surface docking unit, and (iii) the association process, relying on non-covalent interactions, is reversible under appropriate conditions.

In a previous investigation⁹ we showed that CdSe-ZnS core-shell QDs form remarkably stable supramolecular complexes with 1,1'-dialkyl-4,4'-bipyridinium salts in nonpolar solvents, most likely owing to a combination of van der Waals forces between the guest and the surface-bound molecular monolayer, and solvophobic effects experienced by the positively charged guest. These complexes may also be stabilized by electrostatic interactions between the bipyridinium units and the negatively charged ZnS surface.¹⁰ The formation of these complexes is signalled by the decrease of the luminescence intensity of the nanocrystals, which is quenched because of a photoinduced electron-transfer process from the QD conduction band to the LUMO of the nearby bipyridinium unit.^{9,11,12} The original luminescence intensity of the nanoparticles could be quantitatively restored upon addition of a calix[6]arene host able to compete with the QDs for the bipyridinium quenchers.⁹

in hot water and to the clear mixture a saturated aqueous solution of NH_4PF_6 was added dropwise. The white precipitate was filtered, washed with water and THF, and dried in vacuo overnight to yield the title compound in quantitative yield as a pale yellow powder. M.p. = $>180^\circ\text{C}$ (dec.); $^1\text{H-NMR}$ (400 MHz, acetone- d_6): δ = 9.48 (d, $J=7$ Hz, 4H), 9.43 (d, $J=7$ Hz, 4H), 8.30 (m, 8H), 7.80 (s, 4H), 6.23 (s, 4H), 4.95 (m, 4H), 2.09 (m, 4H), 1.25–1.52 (m, 28H), 0.89 (m, 6H); $^{13}\text{C-NMR}$ (101 MHz, acetone- d_6): δ = 155.37, 150.25, 144.40, 143.87, 133.32, 130.65, 128.54, 128.42, 65.01, 62.43, 33.12, 32.69, 30.82, 30.78, 30.57, 30.42, 27.43, 24.12, 15.62. MS (ESI): m/z 1133.9 $[\text{M-PF}_6]^+$.

CDSE NANOCRYSTALS. The synthesis of CdSe core QDs was achieved following the procedures developed by Peng and coworkers,²⁰ with minor modifications.²¹ In brief, the nucleation of the QDs was obtained by reacting the cadmium(II) precursor with elemental selenium coordinated by TBP at 270°C in a solvent with a high boiling point. ODE was used as the solvent, and TOPO and HDA were employed as the surface ligands. In the growth phase, the temperature was lowered to 250°C . In a typical synthetic batch, 0.20 mmol of CdO and 0.80 mmol of OA were heated at 170°C in a three-neck flask connected to a Schlenk line whilst stirring and under an argon flow until the reaction was completed (the solution turned from reddish to clear). After cooling the mixture to room temperature, 0.5 g of TOPO, 1.5 g of HDA and 2.0 g of ODE were added, and the mixture was heated to 270°C . At this temperature a swift injection of a room-temperature Se stock solution (2.0 mmol Se, 0.45 g TBP and 1.4 g of ODE, prepared in a glove box and transferred by syringe) was performed. The temperature was set at 250°C immediately after the injection in order to allow the nanocrystals to grow. After cooling to room temperature the QDs were purified by 3 extraction cycles with a 1:1 methanol/hexane mixture.

CDSE-ZNS CORE-SHELL NANOCRYSTALS. CdSe-ZnS nanocrystals were prepared by overcoating a CdSe core with a ZnS shell using the SILAR procedure.²⁰ In brief, ZnO (0.4 mmol) was dissolved in a mixture of ODA (1 mL) and ODE (9 mL) at 250°C , and S (0.4 mmol) was dissolved in ODE (10 mL) at 200°C . The precalculated amounts of Zn cations and S anions were heteroepitaxially deposited through an air-free standard syringe procedure onto the previously synthesized CdSe nanocrystals. Four layers of ZnS were deposited on top of the core QDs. As the shell thickness for a full monolayer of ZnS is 0.31 nm,²² the QD diameter increases of 2.5 nm after the deposition of the four monolayers.

Results and Discussion

CdSe Nanocrystals

The investigated CdSe core nanocrystals in CHCl_3 exhibit a broad UV-visible absorption and the characteristic low energy exciton peak ($\lambda_{\text{max}} = 560$ nm) arising from quantum confinement of the charge carriers.¹⁻³ The diameter of the nanocrystals, determined from the position of their low-energy

absorption peak according to a published procedure,²³ resulted to be 3.3 nm, while the molar absorption coefficient of the exciton peak is $\epsilon_{\text{max}} = 134000 \text{ M}^{-1} \text{ cm}^{-1}$. TEM Images revealed that the particles are spherical and well monodispersed, as expected from the applied synthetic methodology. The luminescence spectrum of the QDs in air equilibrated CHCl_3 at room temperature shows the intense and narrow luminescence band ($\lambda_{\text{max}} = 578$ nm, fwhm = 30 nm) typical of radiative exciton recombination. Such an emission exhibit a decay that could be satisfactorily fitted with a double exponential function, yielding an average lifetime of 33 ns.[†]

Because the bis(bipyridinium) compound $\mathbf{1}\cdot\text{4PF}_6$ is insoluble in CHCl_3 , its properties were examined in $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1 (v/v). The absorption spectrum of $\mathbf{1}^{4+}$ shows an intense band in the near UV ($\lambda_{\text{max}} = 255$ nm, $\epsilon_{\text{max}} = 51500 \text{ M}^{-1} \text{ cm}^{-1}$) and no luminescence, in line with the behaviour of 1,1'-dibenzyl-4,4'-bipyridinium.²⁴




Fig. 1 (a) Luminescence spectral changes observed upon titration of 1.5 μM CdSe QDs with $\mathbf{1}^{4+}$. (b) Filled circles, left scale: titration plot obtained from the QD luminescence intensity at 578 nm in the presence of increasing amounts of $\mathbf{1}^{4+}$. The dashed and dotted lines show the tangent to the curve at two different slopes and are a guide for the eye, not a fit. Empty triangles, right scale: titration plot obtained from the luminescence lifetime values corresponding to the CdSe emission. Conditions: air equilibrated $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1, room temperature, $\lambda_{\text{exc}} = 500$ nm.

A 1.5 μM solution of CdSe QDs was titrated with $\mathbf{1}^{4+}$ in $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1 (v/v) and the QD emission at 578 nm ($\lambda_{\text{exc}} = 500$ nm) was monitored as a function of the concentration of $\mathbf{1}^{4+}$. As shown in Figure 1a, the addition of $\mathbf{1}^{4+}$ causes a strong quenching of the QD luminescence. Such an emission decrease cannot be ascribed to dynamic quenching, because of the relatively short lifetime of the nanocrystal emission and the very small concentration of the bipyridinium quencher. Even assuming that dynamic quenching is diffusion controlled ($k_d = 1.2 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$),²⁵ the contribution of dynamic quenching to the emission decrease at the highest employed concentration of $\mathbf{1}^{4+}$ would be less than 0.2%. Moreover, the luminescence lifetime of the QDs is just slightly affected by the addition of the quencher (Figure 1b). All these findings suggest that the quenching is static, that is, the nanocrystals and $\mathbf{1}^{4+}$ are associated in the ground state, as previously observed for related systems.^{7,9}

No evidence for ground-state association, however, could be found in the absorption spectra, as the addition of $\mathbf{1}^{4+}$ causes no substantial changes in the absorption features of the QDs (Figure 2a). Nevertheless, a slight increase in absorbance at around 400 nm and between 500 and 700 nm is observed; the differential spectrum (Figure 2b) shows the appearance of new bands with $\lambda_{\text{max}} = 401$ and 612 nm, which can be unequivocally ascribed to the presence of bipyridinium radical cations.²⁴

As these spectral changes are not observed if the QD/ $\mathbf{1}^{4+}$ solution is kept in the dark, reduction of the dicationic bipyridinium units of $\mathbf{1}^{4+}$ to the corresponding radical cations has to be caused by the light excitation employed in the spectrofluorimetric experiments. Indeed, the photoinduced generation of bipyridinium radical cations was obtained upon visible irradiation of CdSe QD-bipyridinium solutions in the presence of a sacrificial reductant, confirming that the QD luminescence quenching can be ascribed to oxidative electron transfer.^{11,12} In the present case, it is likely that small amounts of adventitious species (e.g., water) scavenge the photogenerated hole left in the QD valence band, thereby preventing the back electron transfer from the reduced bipyridinium unit to the oxidized nanocrystal. In any instance, the absorption bands of the bipyridinium radical cation disappear after a few minutes in air equilibrated solution, due to slow reoxidation by dioxygen.

In addition to charge transfer, other possible quenching mechanisms in nanocrystal/chromophore complexes are energy transfer (FRET) and formation of non-radiative surface states arising from changes in the capping monolayer.²⁶ FRET Quenching can be ruled out in the present case because $\mathbf{1}^{4+}$ cannot act as an energy acceptor for the CdSe QDs. On the other hand, $\mathbf{1}^{4+}$ does not possess functional groups capable of binding to the CdSe surface, and control experiments show that the PF_6^- counterions have no effect on the QD emission. Moreover, the luminescence quenching occurs immediately after the addition of $\mathbf{1}^{4+}$ and is constant thereafter, whereas ligand exchange typically occurs on the time scale of minutes.²⁶ The absence of kinetic effects and the reversibility of the quenching observed upon addition of the competitive

calixarene host (see below) are consistent with the non-covalent nature of the nanocrystal-quencher association.




Fig. 2 (a) Absorption spectrum of a 1.5 μM solution of CdSe QDs before (full line) and after the spectrofluorimetric titration with 4.6 equivalents of $\mathbf{1}^{4+}$ as described in Fig. 1. (b) Difference absorption spectrum of the traces shown in (a), indicating the formation of bipyridinium radical cations. Conditions: air equilibrated $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1, room temperature, $\lambda_{\text{exc}} = 500$ nm.

Insightful information on the interaction between the QDs and $\mathbf{1}^{4+}$ can be gathered from a careful examination of the titration curve (Figure 1b). At least three different trends for the intensity decrease can be identified, depending on the $\mathbf{1}^{4+}/\text{QD}$ ratio. It can be noticed that the tangent to the initial part of the curve (dashed line in Figure 1b) intercepts the x -axis – corresponding to zero emission intensity – at about 0.25 equivalents of $\mathbf{1}^{4+}$. This observation suggests that at low $\mathbf{1}^{4+}/\text{QD}$ ratios one molecule of $\mathbf{1}^{4+}$ (i) can associate with up to four nanocrystals, and (ii) it is able to completely quench the luminescence of any of them.

Because of its size and shape in comparison with those of the quantum dots, it is difficult to imagine that $\mathbf{1}^{4+}$ can directly link together more than two nanocrystals. It may be hypothesized, however, that such QD pairs could exhibit a

stronger ability to bind other nanoparticles than individual QDs, because of the larger surface available for non-covalent interparticle interactions.¹⁴

On increasing the number of $\mathbf{1}^{4+}$ equivalents, non-emissive adducts in which one QD is complexed with two molecules of $\mathbf{1}^{4+}$ are also formed (dotted line in Figure 1b). Not surprisingly, complexes with higher $\mathbf{1}^{4+}$: QD stoichiometries are obtained as the titration goes on.




Fig. 3 (a) Luminescence spectral changes observed upon titration of 0.47 μM CdSe-ZnS QDs with $\mathbf{1}^{4+}$. (b) Titration plot obtained from the QD luminescence intensity at 608 nm in the presence of increasing amounts of $\mathbf{1}^{4+}$. The dashed line shows the tangent to the first part of the curve (it is a guide for the eye, not a fit). Conditions: air equilibrated $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1, room temperature, $\lambda_{\text{exc}} = 500$ nm.

CdSe-ZnS Core-shell nanocrystals

CdSe-ZnS core-shell QDs are more interesting for technological applications than bare CdSe nanocrystals, because they possess improved stability and photoluminescence efficiency.²⁷ Hence, we studied the effect of the bis(bipyridinium) species $\mathbf{1}^{4+}$ on QDs composed of a CdSe core ($d = 3.8$ nm) coated with a ZnS shell 1.2 nm thick. These

nanocrystal show the typical low-energy absorption ($\lambda_{\text{max}} = 579$ nm, $\epsilon_{\text{max}} = 200000 \text{ M}^{-1} \text{ cm}^{-1}$)²³ and emission ($\lambda_{\text{max}} = 608$ nm, $\text{fwhm} = 39$ nm) peaks of their CdSe core.

The addition of $\mathbf{1}^{4+}$ to a $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1 solution of the CdSe-ZnS QDs caused a quenching of the emission (Figure 3a), that is, a qualitatively similar result to that observed for the CdSe nanocrystals. The shape of the titration curve, however (Figure 3b), is different from that shown in Figure 1b. In fact, the emission decrease is much less steep than in the case of the CdSe QDs and a full luminescence quenching is observed after addition of a significantly larger number of $\mathbf{1}^{4+}$ equivalents. The initial part of the curve, whose tangent intercepts the x -axis at about 2 equivalents, indicates that at least two $\mathbf{1}^{4+}$ molecules are required to completely quench the luminescence of one QD.

This result contrasts with that obtained for the bare CdSe QDs and can be rationalized on the basis of the different size and structure of the CdSe-ZnS QDs. On the one hand, aggregates in which several QDs are bridged by one $\mathbf{1}^{4+}$ molecule are unlikely to form because of the larger size of the core-shell particles. On the other hand, the presence of the ZnS shell insulates the CdSe core from the external quencher, thus slowing down the photoinduced electron-transfer process at the basis of quenching. This observation is confirmed by the fact that the absorption bands of the bipyridinium radical cations, observed for the CdSe QDs as a consequence of photoexcitation in the spectrofluorimetric titration, were not detected during the same experiments in the case of the CdSe-ZnS QDs. Photoreduction of the bipyridinium units of $\mathbf{1}^{4+}$ was obtained, though, upon illumination of the CdSe-ZnS QDs with high-intensity visible light ($\lambda > 495$ nm) in the presence of 24 equivalents of $\mathbf{1}^{4+}$. From the intensity of the absorption band at 401 nm we estimated that, under these conditions, irradiation for 5 min in deoxygenated solution affords ca. 38% reduction of the bipyridinium units.[†]

QD Aggregation triggered by $\mathbf{1}^{4+}$

To investigate the ability of the bis(bipyridinium) compound $\mathbf{1}^{4+}$ to non-covalently link together the nanocrystals we performed dynamic light scattering (DLS) and transmission electron microscopy (TEM) experiments on CdSe QD solutions with and without the presence of $\mathbf{1}^{4+}$. The CdSe QDs described in the previous section could not be utilized in DLS experiments because they absorb and emit light at the wavelength (633 nm) used by the instrument to detect light scattering. Hence, the DLS and TEM measurements were performed on smaller CdSe QDs, exhibiting the first exciton absorption peak at $\lambda_{\text{max}} = 517$ nm, corresponding to a core diameter of 2.5 nm.²³ The hydrodynamic diameter of the nanocrystals (Figure 4a) resulted to be 4.8 ± 0.1 nm, a figure which is consistent with the core diameter estimated from spectroscopic data and the thickness of the capping ligand shell.²⁸ Indeed, the addition of 0.5 equivalents of $\mathbf{1}^{4+}$ to the QD solution caused a growth of the mean hydrodynamic diameter of the particles to 8.1 ± 0.7 nm as well as an increase in their polydispersity (Figure 4b). This observation is in full agreement

with the formation of aggregates consisting of 2 to 4 nanocrystals, as inferred from luminescence titration data.




Fig. 4 Differential (red bars) and cumulative (blue line) number distributions of particle size obtained from DLS measurements for the CdSe QDs (a) alone and (b) in the presence of 0.5 equivalents of 1^{4+} . Conditions: air equilibrated $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1, room temperature.




Fig. 5 TEM Images recorded on substrates obtained from deposition of dilute solutions of (a, b) CdSe QDs, (c, d) CdSe QDs and 1^{4+} in a 1:1 ratio, and (e, f) CdSe QDs and 1^{4+} in a 1:4 ratio. Arrows in (d) indicate small QD aggregates. Scale bar is 10 nm for (a, c, e) and 50 nm for (b, d, f).

Further support to the QD aggregation induced by 1^{4+} came from TEM experiments.[†] The images recorded on substrates obtained from deposition of a dilute solution of the CdSe QDs showed mainly the presence of individual particles in regions with both higher (Figure 5a) and lower (Figure 5b) populations of nanocrystals. When 1 equivalent of 1^{4+} was added to the QD solution prior to deposition, single nanocrystals could still be detected (Figure 5c) but small clusters were also found (Figure 5d). In the presence of 4 equivalents of 1^{4+} the QDs occurred only in the form of large aggregates and no isolated nanocrystals could be observed (Figure 5ef).

Luminescence recovery by supramolecular encapsulation of 1^{4+}

As pointed out in the introduction, we showed earlier⁹ that QD-bipyridinium adducts could be disrupted by complexing the bipyridinium species with the calix[6]arene derivative **2**, which can efficiently encapsulate bipyridinium-type guests in organic solution.^{18,29} Hence, we investigated whether the addition of **2** could have an effect on the interaction of 1^{4+} with the luminescent nanocrystals.

First of all we performed spectrophotometric titrations to determine the nature and stability of the complex between the bis(bipyridinium) guest and the calixarene host. Upon addition of **2** to 1^{4+} in $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1 an absorption increase in the 450-550 nm region, typical of the inclusion of a bipyridinium unit within the cavity of **2**,²⁹ was observed (Figure 6a). The corresponding titration curve obtained from the absorbance values at 460 nm (Figure 6b) could be satisfactorily fitted with a binding model implying the formation of two adducts (eqs. 1 and 2);³⁰ namely, a 1:1 complex in which one molecule of **2** encircles a bipyridinium unit of 1^{4+} ($K_{1:1} = 1.3 \times 10^5 \text{ M}^{-1}$) and a 2:1 complex in which two calixarene molecules surround both bipyridinium units of one 1^{4+} species ($K_{2:1} = 2.5 \times 10^3 \text{ M}^{-1}$).

The value of $K_{1:1}$ is in line with those found in previous investigations for the 1:1 complex between **2** and 1,1'-dialkyl-4,4'-bipyridinium guests. On the other hand, the significantly lower value of $K_{2:1}$ indicates that the presence of a first calixarene unit around 1^{4+} hampers the association of a second molecule of **2**. This finding is not unexpected if one considers that the distance between the two bipyridinium units of 1^{4+} is short in relation to the length of **2**, resulting in a mutual steric hindrance of the two calixarene hosts in the 2:1 complex.

Variable amounts of calixarene **2** were added to assemblies of 1^{4+} and either CdSe ($\lambda_{\text{max,abs}} = 560 \text{ nm}$, $d = 3.3 \text{ nm}$) or CdSe-ZnS QDs in $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1 and spectrofluorimetric measurements were carried out on the resulting solutions. Upon addition of 5 equivalents of **2** (with respect to 1^{4+}) to a solution containing CdSe QDs ($1.5 \mu\text{M}$) and 1^{4+} in a 1:4.6 ratio, only less than 1% of the initial emission intensity was restored. Similarly, the addition of 38 equivalents of **2** (with respect to 1^{4+}) to a solution containing CdSe-ZnS QDs ($0.47 \mu\text{M}$) and 1^{4+} in a 1:20 ratio caused <1% recovery of the luminescence. Notably, the emission of CdSe-ZnS QDs ($0.43 \mu\text{M}$) and 1,1'-dibenzyl-4,4'-bipyridinium in a 1:19 ratio was fully restored after the addition of 20 equivalents of **2** (with respect to the bipyridinium guest).⁹ The comparison of these results suggests

that the binding of the bis(bipyridinium) 1^{4+} species to the nanocrystals is stronger than that of a structurally related compound bearing a single bipyridinium unit. The larger size of 1^{4+} , its twice as large electric charge, and its lower affinity for chloroform may be among the reasons for such a behaviour. It may also be considered that complexation of 1^{4+} by **2** could be hampered by aggregation of the nanocrystals triggered by the former.






Fig. 6 (a) Absorption spectral changes observed upon titration of 25 μM 1^{4+} with **2**. (b) Titration plot obtained from the absorption values at 460 nm. The grey curve is the data fitting according to the 1:1 and 2:1 binding model described in the text (eqs. 1 and 2). Conditions: air equilibrated $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1, room temperature.



Nevertheless, the titration of a 1.5 μM 1:1 mixture of CdSe QDs and 1^{4+} (in which ca. 87% of the nanocrystals luminescence is quenched, see Figure 1b) with **2** shows a recovery of up to 30% of the original emission intensity of the

QDs after the addition of a large excess (215 equivalents) of the calixarene host (Figure 7). Conversely, the addition of the same excess of **2** to the QDs (in the absence of 1^{4+}) caused a 8% emission decrease. These observations indicate that complexation of 1^{4+} by **2** causes the disassembly of the non-luminescent adducts consisting of the QDs and the bis(bipyridinium) species.




Fig. 7 Luminescence spectral changes at 578 nm observed upon titration of a 1:1 mixture of CdSe QDs and 1^{4+} (1.5 μM) with the calixarene host **2**. Conditions: air equilibrated $\text{CHCl}_3/\text{CH}_3\text{CN}$ 9:1, room temperature, $\lambda_{\text{exc}} = 500$ nm.

Conclusions

We have investigated the interaction of CdSe and CdSe-ZnS luminescent nanocrystals with the ditopic bis(bipyridinium) species 1^{4+} in organic solution. Our results show that the nanocrystal and molecular components undergo association in the ground state, yielding complexes in which the luminescence of the quantum dots is strongly quenched. Interestingly, 1^{4+} can associate and quench as many as four CdSe nanocrystals, possibly owing to cooperative interparticle binding phenomena. The ability of 1^{4+} to promote aggregation of the quantum dots is confirmed by DLS and TEM experiments.

Visible light irradiation of the adducts causes the formation of bipyridinium radical cations, confirming that the luminescence quenching mechanism is an electron-transfer process from the excited electron in the conduction band of the nanocrystal to the LUMO of the bipyridinium quencher. The presence of a ZnS shell around the CdSe core does not prevent the electron-transfer quenching process but it does reduce its efficiency.

Calix[6]arene **2**, which is able to encapsulate both bipyridinium units of 1^{4+} by virtue of supramolecular interactions, was used to promote the decomplexation of 1^{4+} from the quantum dots. Indeed, the addition of an excess of **2** to a solution containing the nanocrystal/ 1^{4+} complexes causes a recovery of the luminescence of the quantum dots, indicating the partial disassembly of the complexes.

These results demonstrate that supramolecular interactions are a convenient way to control the aggregation of semiconductor nanocrystals. The combination of molecular recognition and self-assembly concepts with luminescent quantum dots is a promising route for the generation of nanostructured arrays with controlled morphologies and properties. Such a task is important for improving the performances of quantum dot-based optoelectronic devices.

Acknowledgements

This work was supported by the European Union (EU) (Hysens Project, grant no. 263091, ATOMIN project, grant no. POIG.02.01.00-12-023/08), the Ministero dell'Università e della Ricerca (PRIN 2010CX2TLM) and the University of Bologna. M. O. thanks the National Science Centre (grant no. UMO 2011/01/N/ST5/02550) and the Foundation for Polish Science for a MPD programme fellowship co-financed by the EU European Regional Development Fund. K. S. acknowledges financial support from the National Science Centre (grant no. UMO-2011/03/B/ST5/01495).

Notes and references

^a Photochemical Nanosciences Laboratory, Dipartimento di Chimica "G. Ciamician", Università di Bologna, Via Selmi 2, 40126 Bologna, Italy. Fax: +39 051 2099456; Tel: +39 051 2099540; E-mail: alberto.credi@unibo.it

^b Faculty of Chemistry, Jagiellonian University, Ingardena 3, 30-060 Krakow, Poland.

^c Faculty of Non-Ferrous Metals, AGH University of Science and Technology, al. Mickiewicza 30, 30-060 Kraków, Poland.


† Electronic Supplementary Information (ESI) available: UV-Visible absorption spectra, luminescence spectra and lifetimes, and TEM images. See DOI: 10.1039/b000000x/

1. *Semiconductor nanocrystal quantum dots*, Ed. A. L. Rogach, Springer-Verlag, Wien, Austria, 2008.
2. A. M. Smith and S. Nie, *Acc. Chem. Res.*, 2010, **43**, 190.
3. U. Resch-Genger, M. Grabolle, S. Cavaliere-Jaricot, R. Nitschke and T. Nann, *Nat. Methods*, 2008, **5**, 763.
4. *Quantum dot sensors: Technology and commercial applications*, Eds. J. F. Callan and F. M. Raymo, Pan Stanford Publishing, Singapore, 2013.
5. I. Yildiz, E. Deniz and F. M. Raymo, *Chem. Soc. Rev.*, 2009, **38**, 1859; I. L. Medintz and H. Mattoussi, *Phys. Chem. Chem. Phys.*, 2009, **11**, 17; N. Hildebrandt, *ACS Nano*, 2011, **5**, 5286; T. L. Doane and C. Burda, *Chem. Soc. Rev.*, 2012, **41**, 2885; R. Freeman and I. Willner, *Chem. Soc. Rev.*, 2012, **41**, 4067; R. Freeman, J. Girsh and I. Willner, *ACS Appl. Mater. Interf.*, 2013, **5**, 2815; T. Avellini, C. Lincheneau, F. Vera, S. Silvi and A. Credi, *Coord. Chem. Rev.*, 2014, **263-264**, 151.
6. See, e.g.: Y. Chen, R. Thakar and P. T. Snee, *J. Am. Chem. Soc.* 2008, **130**, 3744; I. L. Medintz, M. H. Stewart, S. A. Trammell, K. Susumu, J. B. Delehanty, B. C. Mei, J. S. Melinger, J. B. Blanco-Canosa, P. E. Dawson and H. Mattoussi, *Nat. Mater.*, 2010, **9**, 676; Z. Erno, I. Yildiz, B.

- Gorodetsky, F. M. Raymo and N. R. Branda, *Photochem. Photobiol. Sci.*, 2010, **9**, 249; M. Amelia, A. Lavie-Cambot, N. D. McClenaghan and A. Credi, *Chem. Commun.*, 2011, **47**, 325; W. R. Algar, D. Wegner, A. L. Huston, J. B. Blanco-Canosa, M. H. Stewart, A. Armstrong, P. E. Dawson, N. Hildebrandt and I. L. Medintz, *J. Am. Chem. Soc.*, 2012, **134**, 1876; S. Impellizzeri, B. McCaughan, J. F. Callan, and F. M. Raymo, *J. Am. Chem. Soc.*, 2012, **134**, 2276; R. Freeman, T. FINDER, L. Bahshi, R. Gill and I. Willner, *Adv. Mater.*, 2012, **24**, 6416; J. Aguilera-Sigalat, V. F. Pais, A. Domenech-Carbo, U. Pischel, R. E. Galian and J. Perez-Prieto, *J. Phys. Chem. C*, 2013, **117**, 7365.
7. M. Amelia, M. Font and A. Credi, *Dalton Trans.*, 2011, **40**, 12083; M. Amelia, A. Credi, *Inorg. Chim. Acta*, 2012, **381**, 247.
8. A. Boulesbaa, A. Issac, D. Stockwell, Z. Huang, J. Huang, J. Guo and T. Lian, *J. Am. Chem. Soc.*, 2007, **129**, 15132; S.-C. Cui, T. Tachikawa, M. Fujitsuka and T. Majima, *J. Phys. Chem. C*, 2011, **115**, 1824; J. Aguilera-Sigalat, J. M. Casas-Solvas, M. C. Morant-Minana, A. Vargas-Berenguel, R. E. Galian and J. Perez-Prieto, *Chem. Commun.*, 2012, **48**, 2573; A. Iagatti, R. Flamini, M. Nocchetti and L. Latterini, *J. Phys. Chem. C*, 2013, **117**, 23996.
9. B. Gadenne, I. Yildiz, M. Amelia, F. Ciesa, A. Secchi, A. Arduini, A. Credi and F.M. Raymo, *J. Mater. Chem.*, 2008, **17**, 2022.
10. J. Bebie, M. A. A. Schoonen, M. Fuhrmann, D. R. Strongin, *Geochim. Cosmochim. Acta* 1998, **62**, 633; M. Tiemann, F. Marlow, F. Brieler, M. Lindén, *J. Phys. Chem. B*, 2006, **110**, 23142; M. Wang, Q. Zhang, W. Hao, Z.-X. Sun, *Chem. Cent. J.*, 2011, **5**, 73; J. Saikia, B. Sahab, G. Das, *J. Coll. Interf. Sci.*, 2014, **416**, 235.
11. I. Yildiz and F. M. Raymo, *J. Mater. Chem.*, 2006, **16**, 1118; I. Yildiz, M. Tomasulo and F. M. Raymo, *Proc. Natl. Acad. Sci. USA*, 2006, **103**, 11457.
12. S. Logunov, T. Green, S. Marguet and M. A. El-Sayed, *J. Phys. Chem. A*, 1998, **102**, 5652; C. Burda, T. C. Green, S. Link and M. A. El-Sayed, *J. Phys. Chem. B*, 1999, **103**, 1783.
13. M. Grzelczak, J. Vermant, E. M. Furst and L. M. Liz-Marzán, *ACS Nano*, 2010, **4**, 3591; R. Schreiber, J. Do, E.-M. Roller, T. Zhang V. J. Schüller, P. C. Nickels, J. Feldmann and T. Liedl, *Nat. Nanotech.*, 2014, **9**, 74.
14. T. Wang, D. LaMontagne, J. Lynch, J. Zhuang and Y. C. Cao, *Chem. Soc. Rev.*, 2013, **42**, 2804.
15. H. Zhang, E. W. Edwards, D. Y. Wang and H. Möhwald, *Phys. Chem. Chem. Phys.*, 2006, **8**, 3288; Y. Gao and Z. Tang, *Small*, 2011, **7**, 2133; C. Zhang, R. J. Macfarlane, K. L. Young, C. H. J. Choi, L. Hao, E. Auyeung, G. Liu, X. Zhou and C. A. Mirkin, *Nat. Mater.*, 2013, **12**, 741.
16. D. Vanmaekelbergh and P. Liljeroth, *Chem. Soc. Rev.*, 2005, **34**, 299; D. V. Talapin, J. S. Lee, M. V. Kovalenko and E. V. Shevchenko, *Chem. Rev.*, 2010, **110**, 389.
17. R. J. Alvarado, J. Mukherjee, E. J. Pacsial, D. Alexander, F. M. Raymo, *J. Phys. Chem. B*, 2005, **109**, 6164.
18. A. Credi, S. Dumas, S. Silvi, M. Venturi, A. Arduini, A. Pochini and A. Secchi, *J. Org. Chem.*, 2004, **69**, 5881.

19. A. Credi and L. Prodi, *J. Mol. Struct.*, 2014, in press; DOI: 10.1016/j.molstruc.2014.03.028.
20. Z. A. Peng and X. Peng, *J. Am. Chem. Soc.*, 2001, **123**, 183; J. Li, Y. A. Wang, W. Guo, J. C. Keay, T. D. Mishima, M. B. Johnson and X. Peng, *J. Am. Chem. Soc.*, 2003, **125**, 12567.
21. Z. A. Peng and X. Peng, *J. Am. Chem. Soc.*, 2002, **124**, 3343; C. R. Bullen, P. Mulvaney, *Nano Lett.*, 2004, **4**, 2303.
22. D. Chen, F. Zhao, H. Qi, M. Rutherford and X. Peng, *Chem. Mater.*, 2010, **22**, 1437.
23. W. W. Yu, L. Qu, W. Guo and X. Peng, *Chem. Mater.* 2003, **15**, 2854.
24. R. Ballardini, A. Credi, M. T. Gandolfi, C. Giansante, G. Marconi, S. Silvi and M. Venturi, *Inorg. Chim. Acta*, 2007, **360**, 1072.
25. M. Montalti, A. Credi, L. Prodi and M.T. Gandolfi, *Handbook of photochemistry, 3rd Ed.*, 2006, CRC Press, Boca Raton.
26. T. Blaudeck, E. I. Zenkevich, M. Abdel-Mottaleb, K. Szwaykowska, D. Kowanko, F. Cichos and C. von Borczyskowski, *ChemPhysChem*, 2012, **13**, 959.
27. M. A. Hines and P. Guyot-Sionnest, *J. Phys. Chem.*, 1996, **100**, 468.
28. J. M. Tsay, S. Doose and S. Weiss, *J. Am. Chem. Soc.*, 2006, **128**, 1639.
29. A. Arduini, R. Bussolati, A. Credi, A. Secchi, S. Silvi, M. Semeraro and M. Venturi, *J. Am. Chem. Soc.*, 2013, **135**, 9924; M. Semeraro, A. Secchi, S. Silvi, M. Venturi, A. Arduini and A. Credi, *Inorg. Chim. Acta*, 2014, **417**, 258.
30. R. A. Binstead, SPECFIT; Spectrum Software Associates: Chapel Hill, NC, 1996.

TABLE OF CONTENTS

Supramolecular assemblies of semiconductor quantum dots and a bis(bipyridinium) derivative: luminescence quenching and aggregation phenomena

Non-emissive aggregates of CdSe and CdSe-ZnS nanocrystals are formed in solution in the presence of a bis(bipyridinium) species, and disrupted upon addition of a macrocyclic host for the bipyridinium units.