


 Cite this: *RSC Adv.*, 2022, 12, 13014

Noncovalently bound excited-state dimers: a perspective on current time-dependent density functional theory approaches applied to aromatic excimer models†

Amy C. Hancock and Lars Goerigk *

Excimers are supramolecular systems whose binding strength is influenced by many factors that are ongoing challenges for computational methods, such as charge transfer, exciton coupling, and London dispersion interactions. Treating the various intricacies of excimer binding at an adequate level is expected to be particularly challenging for Time-Dependent Density Functional Theory (TD-DFT) methods. In addition to well-known limitations for some TD-DFT methods in the description of charge transfer or exciton coupling, the inherent London dispersion problem from ground-state DFT translates to TD-DFT. While techniques to appropriately treat dispersion in DFT are well-developed for electronic ground states, these dispersion corrections remain largely untested for excited states. Herein, we aim to shed light on current TD-DFT methods, including some of the newest developments. The binding of four model excimers is studied across nine density functionals with and without the application of additive dispersion corrections against a wave function reference of SCS-CC2/CBS(3,4) quality, which approximates select CCSDR(3)/CBS data adequately. To our knowledge, this is the first study that presents single-reference wave function dissociation curves at the complete basis set level for the assessed model systems. It is also the first time range-separated double-hybrid density functionals are applied to excimers. In fact, those functionals turn out to be the most promising for the description of excimer binding followed by global double hybrids. Range-separated and global hybrids—particularly with large fractions of Fock exchange—are outperformed by double hybrids and yield worse dissociation energies and inter-molecular equilibrium distances. The deviation between each assessed functional and reference increases with system size, most likely due to missing dispersion interactions. Additive dispersion corrections of the DFT-D3(BJ) and DFT-D4 types reduce the average errors for TD-DFT methods but do so inconsistently and therefore do not offer a black-box solution in their ground-state parametrised form. The lack of appropriate description of dispersion effects for TD-DFT methods is likely hindering the practical application of the herein identified more efficient methods. Dispersion corrections parametrised for excited states appear to be an important next step to improve the applicability of TD-DFT methods and we hope that our work assists with the future development of such corrections.

 Received 16th March 2022
 Accepted 12th April 2022

DOI: 10.1039/d2ra01703b

rsc.li/rsc-advances

1 Introduction

Exciplexes are short-lived, heterodimeric “excited complexes” that are stable in the electronic excited state while dissociative in the electronic ground state.^{1–3} Their formation can be considered as the association between an excited monomer M^* and a second monomer N in the ground state, as depicted by the following reaction scheme:



where the asterisk denotes an electronically excited species. Ideally, the monomers are polar or polarisable species able to facilitate attractive charge-transfer (CT) interactions in an excited state.¹ In the homodimeric case ($M = N$), exciplexes are called “excimers”, which are the focus of this work.

The excited-state properties of the dimeric species are unique from those of either monomer.⁴ The stabilising effect in exciplexes and excimers can be rationalised with molecular orbital (MO) theory and a collision model of the reaction in eqn (1). When the two monomers collide, the predominant interactions occur between the highest occupied (HOMO) and lowest unoccupied (LUMO) molecular orbitals of each species to form

School of Chemistry, The University of Melbourne, Parkville, Australia. E-mail: lars.goerigk@unimelb.edu.au; Tel: +61-3-8344-6784

† Electronic supplementary information (ESI) available. See <https://doi.org/10.1039/d2ra01703b>





Fig. 1 Simplified molecular orbital diagram describing the formation of an excimer from molecular orbitals of the monomers as described in eqn (1). "H" and "L" refer to the highest occupied and lowest unoccupied molecular orbitals, respectively; based on similar examples in ref. 1 and 5–7.

a new set of orbitals as shown in the simplified diagram in Fig. 1.^{1,5–7} In the case of two ground-state molecules, the resulting constructive and destructive-interference contributions cancel, resulting in minimal to no net stabilisation. In an excimer or exciplex, the electronically excited species causes two interacting orbitals to be singly occupied resulting in stabilisation through formation of the complex.⁵ This electronic stabilisation effect is short-lived, as relaxation to the ground state occurs quickly, causing repulsive intermolecular forces to oppose the relatively weak attractive forces. Energy decomposition analysis of the excimer/exciplex stabilisation energy revealed the following contributing components: electrostatics, Pauli repulsion, CT, exciton coupling and London dispersion.^{8,9}

Aromatic excimers were discovered experimentally in the fluorescence spectrum of pyrene in cyclohexane solution by a broad, structureless emission band that occurred at a lower energy than the associated monomer emission.¹⁰ Since then, excimers have also been revealed to be crucial species in contemporary applications of technological and biological relevance. Contrary to their notoriety as an undesired energy trap for singlet-fission,¹¹ some studies have shown potential advantages in organic electronics, such as excimer states mediating intramolecular electron transfer¹² and charge separation¹³ or broadening the emission for white organic light-emitting diodes;¹⁴ see ref. 15 for a review. Additionally, excimers have applications as chemosensors,^{16,17} molecular rulers,^{18,19} and industrial-scale lasers.²⁰ Excimers also occur in biological systems, playing a role in the photo-damage of DNA as they can occur between nucleobases such as adenine²¹ and guanine.²²

Computational chemistry techniques, particularly Kohn–Sham Density Functional Theory^{23,24} (DFT) approaches, also called density functional approximations (DFAs), are frequently used in applications of technological relevance, including the description of exciplexes and excimers. In order to have predictive character, DFAs must be robust; *i.e.*, they must yield results with equal accuracy or error margins across a variety of different chemical problems. Comprehensive benchmark studies have identified such generally applicable DFT approaches for ground-state problems including for thermochemistry, kinetics, and geometries, to name a few examples.^{25–31} In all studies, it was evident that the accurate treatment of noncovalent interactions (NCIs), in both inter- and intramolecular cases, is crucial to achieve the desired accuracy and robustness of a method. In the context of DFT, it is particularly important to properly address the correct treatment of London dispersion effects. It has been known since the mid-1990s that conventional DFAs do not capture those effects correctly.^{32–35} This sparked the development of various dispersion-corrected DFT techniques, some of which later turned out to be less effective than initially claimed,^{36–38} and others having now become the recommended default for ground-state DFT applications.^{39,40} We refer interested readers to recently published reviews^{41,42} directed at users that are unfamiliar with the field of dispersion-corrected DFT applications for ground-state problems as well as reviews that recommend the current best practice in the field,^{43–45} which includes dispersion-corrected double-hybrid^{46,47} DFAs (DHDFAs) when feasible.

The binding of excimers and exciplexes inherently relies on the system's excited-state properties and the dominating types of NCIs, which introduces additional complexity compared to the ground-state case. Linear-response time-dependent DFT within the adiabatic approximation^{48–50} (TD-DFT) has become the method of choice to treat excited-state problems. Recent advances^{51–53} and detailed benchmark studies^{54–64} on single-molecule cases have shed light on the quality of TD-DFAs for the calculation of excitation energies. As recently summarised for readers unfamiliar with the field, lower rungs on the Jacob's Ladder of DFT⁶⁵ should be avoided due to the emergence of artificial ghost states and large red shifts in excitation energies.⁶⁶ Global-hybrid DFAs can be suitable for local valence excitations, subject to having the right amount of nonlocal Fock exchange (FE)—about 40% (ref. 57 and 66)—but fail for CT and other long-range transitions.^{52,57,62,67–71} While the range-separation (RS) technique^{72–75} applied to the exchange part of DFAs solves the long-range problem, local valence excitations tend to be blue-shifted.^{52,58,62,66,76} Time-dependent DHDFAs⁷⁷ depend additionally on a nonlocal, perturbative electron-correlation component⁷⁸ that compensates for many of the systematic errors of hybrids.^{47,51,54,55,57,58,66,79} In fact, our group recently developed RS-DHDFAs^{52,53} that belong to some of the currently most robust TD-DFT methods for a variety of different local and long-range valence excitations in organic molecules.^{52,53,62–64} A slightly different group of RS-DHDFAs have also proven to be very promising when used within the Tamm–Dancoff Approximation⁸⁰ (TDA-DFT).^{81–83}



Compared to single-molecule cases, there are fewer systematic studies that explore NCIs in excited states of molecular complexes. However, it is expected that TD-DFT methods inherit the problems of the underlying ground-state DFA, such as the inability to properly describe London dispersion interactions. Additionally, the TD-DFT formalism itself carries problems that may further complicate the treatment of excited-state NCIs. In 2008, Huenerbein and Grimme presented a study of excimers and one exciplex across a limited number of TD-DFTs.⁵ The study was unique in the sense that a London dispersion correction was rigorously applied to an excited-state study, namely the DFT-D2 (ref. 84) correction developed in 2006. Although the dispersion coefficients used in DFT-D2 were based on ground-state polarisabilities, the combination of DFT-D2 with high-FE global-hybrid DFAs, showed a qualitatively better agreement with experimental data compared to dispersion-uncorrected TD-DFT, both for the stability of the dimers and the inter-monomer equilibrium distance. The authors' argument was that a ground-state based correction should be seen as a lower bound for the dispersion energy in an excited-state system, as C_6 coefficients are different, and most likely larger, for more polarisable excited states. Some of the subsequent TD-DFT based studies on excimer and exciplex systems either do not explore the dispersion problem of DFT^{85,86} or erroneously justified the use of Minnesota functionals⁸⁷ for consideration of dispersion effects,^{22,88} despite ground-state based studies that disprove claims they are able to treat such interactions.^{25,27,37,38,89} Some other studies recognised the dispersion problem, but for a lack of a better choice, followed Huenerbein and Grimme's recommendation to use a global-hybrid combined with DFT-D2,^{6,90–95} while fewer applied the newer, ground-state based DFT-D3 method in its zero-damping⁹⁶ [DFT-D3(0)] or in its Becke–Johnson-damping^{97–99} [DFT-D3(BJ)]⁴⁰ form.^{100–106} To our knowledge, only three studies have considered adjusting dispersion corrections for excited states. In the first, Ikabata and Nakai¹⁰⁷ calculated excited-state dispersion coefficients within the local-response dispersion^{108,109} (LRD) model to explore the interaction energies of exciton-localised molecular complexes from the S66 (ref. 110) benchmark set as well as three molecular excimers. Then, Briggs and Besley¹¹¹ empirically varied the dispersion coefficients and van der Waals radii in DFT-D2 for small model systems of ethene–argon and formaldehyde–methane complexes. And finally, Johnson and co-workers calculated dispersion coefficients within the exchange-hole dipole moment^{97–99,112} (XDM) model for excited states of conjugated hydrocarbons, pull–pull chromophores, and CT complexes.¹¹³ While these studies considered suitable model systems, only a limited selection of DFAs were applied. These three studies, among others,^{114,115} would allow the conclusion that state-specific dispersion corrections for excited states should be developed, but further characterisation of current methods is necessary to provide guidance towards such future developments. The present paper intends to provide such characterisation.

Most of the aforementioned exciplex studies used older global-hybrid DFAs, while some^{85,86,90,91,102,105,106,116} used range-separated hybrids. Of the two studies that investigated

DHDFAs^{94,102} both utilised TDA-DFT, instead of the full TD-DFT scheme, but only Krueger and Blanquart¹⁰² considered a dispersion correction. Interestingly, most of the methods previously used to study exciplex systems are not those recommended by some of the latest single-molecule excitation studies,^{52,53,57,62,64,66} which by itself is reason enough to investigate the latter in the exciplex/excimer context. The majority of excimer and exciplex studies investigated these methods *via* applications,^{22,86,88,91–94,100,101,104,116} or carried out benchmark analyses against reference values that were not well defined and mostly based on experiment.^{5,90} Both benchmarks and application-based excimer studies focussed on comparison of methods by the calculation of the dissociation energy, as this explores the binding of the complex, while fewer studies have explored other spectrometric parameters such as the fluorescence, absorption and repulsion energies.^{7,85,86,104,107,116–121} Potential energy curves of the lowest excited state reported relative to the asymptote of the dissociation curve of the ground state⁵ are standard practice to derive these spectroscopic parameters. Ground-state studies commonly explore NCIs by focusing entirely on interaction energies that can be calculated, *e.g.*, as the difference between the dimer and individual monomer energies or alternatively as the difference between the total energies of the dimer and a system in which the two monomers have been dissociated to a large distance from one another. By this definition, a negative interaction energy indicates a stable complex. The dissociation energy of an excimer is inherently an interaction energy and provides a convenient metric to discuss excited-state binding and NCIs in one go. However, the computational studies of excimers and exciplexes do not report excited-state interaction energy curves, other than a handful of exceptions we are aware of,^{7,8,85,102} we advocate to adopt them more frequently, as this allows direct comparison with what has become the standard in ground-state treatments of NCIs. Note that according to that definition of an interaction energy, the dissociation energy D_e is simply its absolute value.

There have been many rigorously conducted NCI studies for ground-state systems that are usually based on a high-quality reference and compared different computational approaches on an equal footing; the benefits of using high-level computational data as opposed to experimental references has been well-established and explained elsewhere.^{27,43,51,52,66} To the best of our knowledge, similar studies are missing to systematically study NCIs in excited states. In this work, we intend to initiate the first step towards more systematic studies of excited-state NCIs by restudying excimer model systems under incorporation of the latest state-of-the-art TD-DFT methods. The four model systems discussed herein are the benzene, naphthalene, anthracene and pyrene dimers (see Fig. 2) which have been studied in combination before by two prior studies.^{7,90} We discuss the change of the interaction energy for each dimer system upon dissociation, *i.e.* we discuss equilibrium and non-equilibrium geometries. In each case, we discuss the first excited state, which is the excimer state. In the following section, Section 2, we briefly outline general computational details before analysing various wave-function theory (WFT) levels based on truncated basis sets and at the complete-basis-





Fig. 2 Excimer benchmark cases used in this study consisting of the following monomers: benzene (a), naphthalene (b), anthracene (c) and pyrene (d).

set (CBS) limit in Section 3. The purpose of this analysis is to identify a suitable level of theory that can serve as a reliable, yet fast benchmark for the subsequent discussion of TD-DFT methods in various subsections of Section 4. The latter is split into different aspects: the impact of non-local FE, the behaviour of dispersion-uncorrected TD-DFT methods, the impact of using ground-state based corrections of the DFT-D3(BJ)^{40,96} and DFT-D4 (ref. 122 and 123) type, and an analysis of method performance averaged over the four systems. In a nutshell, our study is sufficiently comprehensive to fill an important gap in our knowledge of current TD-DFT methods and additive dispersion corrections for the description of excimers.

2 General computational details

TURBOMOLE 7.3 (ref. 50 and 124–126) was used for geometry optimisations of the lowest-lying singlet excited states of each dimer at the spin-component-scaled^{127,128} approximate coupled cluster singles doubles (SCS-CC2 (ref. 129)) level with the def2-TZVP¹³⁰ triple- ζ (TZ) atomic-orbital (AO) basis set and a geometry convergence criterion of $10^{-7} E_h$. Use of the RICC2 module in TURBOMOLE limits the symmetry consideration to Abelian point groups such that all excimers are calculated with D_{2h} symmetry. The excited-state optimised structures were used to generate non-relaxed dissociation curves as follows: the internal coordinates of each monomer were frozen, then total energies of the lowest-lying singlet excited states were calculated across a range of inter-monomer separations up to 16 Å, and then each energy was taken relative to 16 Å, which represents the asymptote for the given state. The geometry optimised excimer structures were not comprised of perfectly planar monomers, which has been previously noted during both TD-DFT and CC2 (ref. 131) optimisations of these excimers⁹⁰ and exciplexes comprised of the same monomers.⁸⁵ Therefore, we defined the inter-monomer separation for our dissociation curves as the distance between two central carbon atoms, opposite each other on each monomer; this parameter is visually defined for each excimer structure in Fig. S1 (ESI[†]). D_e , is defined as the difference between the total energy of the first excited state at the 16 Å point and the total energy of the same state at the minimum point. The effect of geometric relaxation falls outside the scope of this study. However, frozen-monomer structures at the SOS-CIS(D_0)¹³² level have been shown to give good estimates of the equilibrium inter-monomer distance, with excimer dissociation energies that are generally overestimated by *ca.* 2 kcal mol⁻¹ in the case of the benzene excimer.¹³³ As this study uses the same

structures throughout, internal comparison of the dissociation energies is reasonably justified.

Herein, the figures display interaction energy curves such that a negative interaction energy indicates a stable dimer. In the context of this paper, D_e corresponds to the dissociation energy of an excimer and therefore the terms interaction and dissociation energy are used interchangeably. Equilibrium inter-monomer distances r_e , corresponding to the distance at which the minimum energy arises, are also discussed in parts of our discussion.

TURBOMOLE was also used for all SCS-CC2 and CC2 single-point calculations along the dissociation curves. The frozen-core and resolution of the identity¹³⁴ (RI) approximations were employed with appropriate auxiliary basis sets.¹³⁵ Coupled cluster singles doubles with perturbative triples excitation correction¹³⁶ [CCSDR(3)] calculations for selected points along the dissociation curves were carried out with Dalton2016,¹³⁷ also utilising the frozen-core approximation. The WFT calculations were used to identify suitable reference data for the subsequent assessment of TD-DFT methods.

Single-point, linear-response TD-DFT calculations within the adiabatic approximation were conducted for all dissociation curves for a series of DFAs as listed in Table 1. For the study of the influence of FE we apply a series of PBE¹³⁸- and BLYP^{139–141}-based methods with varying amounts of non-local exchange. However, as lower-rung DFAs have shown to be unreliable for excited-state problems,^{55,66} we limit our final benchmarking study to global hybrids, range-separated hybrids, global double hybrids and range-separated double hybrids; the exact functional types are detailed in Table 1. Those DFAs have been chosen either based on popularity or known accuracy for excited states of organic molecules; see ref. 66 for recommendations and insights. Note that time-dependent double-hybrid calculations require a conventional TD-DFT step for its hybrid portion followed by a configuration interaction singles with perturbative doubles⁷⁸ [CIS(D)] correction as initially suggested by Grimme and Neese;⁷⁷ see ref. 66 for a comprehensive, free-access review on this methodology and some of its latest advances.

Self-consistent-field (SCF) steps in this work were carried out with energy convergence criteria of $10^{-7} E_h$ (TURBOMOLE) and $10^{-8} E_h$ (ORCA). The frozen-core approximation was applied across all TURBOMOLE calculations, and in ORCA^{147,148} calculations it was used along with the RI approximation for the perturbative parts of the double hybrids. Large quadrature-grid options “7” (TURBOMOLE) and “grid6 finalgrid7” (ORCA) were



Table 1 List of TD-DFT methods applied

Program	Method	Comments ^a
TURBOMOLE 7.3	BLYP, ^{139–141} PBE ¹³⁸ B3LYP, ^{142,143} PBE20 BHLYP, ¹⁴⁴ PBE50 PBE38, ⁹⁶ BLYP38	General Gradient Approximation (GGA); no FE Global hybrids; 20% FE Global hybrids; 50% FE Global hybrids; 37.5% ($\frac{3}{8}$) FE
ORCA V4.1.0	PBE75, BLYP75 CAM-B3LYP ⁷⁴ ω B97X ¹⁴⁵ B2PLYP ⁴⁶ B2GP-PLYP ¹⁴⁶	Global hybrids; 75% FE RS hybrid, but without 100% FE in the long range RS hybrid with 100% FE in the long range Global double hybrid; 53% FE; 27% CIS(D) correlation Global double hybrid; 65% FE; 36% CIS(D) correlation
Local version of ORCA4 ^b	ω B2PLYP, ⁵² ω B2GP-PLYP ⁵²	RS versions of B2(GP-)PLYP with 100% FE in the long range

^a GGAs: rung 2; global and RS hybrids: rung 4; global and RS double hybrids: rung 5. FE: Fock exchange. ^b Available in ORCA 4.2 and ORCA5.

used to ensure smooth dissociation curves. Dispersion corrections of the type DFT-D3(BJ)^{40,96} and DFT-D4 (ref. 122 and 123) were carried out with the DFTD3 and DFTD4 standalone programs.^{149,150} Damping parameters for the various functionals were taken from the respective reference that published them first. For DFT-D3(BJ) these are ref. 40 for B3LYP, ref. 25 for PBE38, BHLYP, and CAM-B3LYP, ref. 151 for B2(GP-)PLYP, and ref. 152 for ω B2(GP-)PLYP. All DFT-D4 parameters were taken from ref. 123 except for PBE38 and the two RS-DHDFAs which were taken from ref. 153 and 152, respectively.

A series of different basis sets—ranging from double- ζ (DZ) to quadruple- ζ (QZ)—were employed throughout this work, with some of them used for extrapolations to the CBS limit. More information is provided in the following sections.

3 Establishing a suitable reference

Prior to analysing the performance of TD-DFT methods, a sufficiently reliable reference method that is still feasible for larger systems must be identified. An appropriate reference method should minimise the computational cost while maintaining reasonable comparability to a higher level of theory. Coupled-cluster WFT methods are considered the gold standard of chemical accuracy.^{154,155} For ground states, coupled-cluster singles-and-doubles with perturbative triples, CCSD(T),¹⁵⁶ at the CBS limit is the ideal that many aim to achieve in contemporary benchmarking.^{27,157,158} It constitutes a very accurate approximation to the true interaction energy of noncovalently bound complexes in their electronic ground state.^{27,158–160} One excited-state equivalent to ground-state treatments with CCSD(T) is linear-response CCSD enhanced with a different type of perturbative triples correction, such as the aforementioned CCSDR(3).¹³⁶ It delivers excitation energies that are similar to the more costly linear-response approximate coupled cluster singles doubles triples (CC3);¹⁶¹ for examples of CCSDR(3) and CC3 being established as benchmarks and comparisons between both, see ref. 51, 52 and 162–165. While the ground-state gold standard has, in recent years, become increasingly feasible for large systems of up to several hundred atoms,^{166,167} the excited-state equivalent remains prohibitively expensive.¹⁵⁵

For most of our systems, CCSDR(3) is therefore not achievable for computational reasons, and in the following we identify if instead CC2 or its spin-component-scaled¹²⁸ variant, SCS-CC2, can be used as a low-cost alternative.

Establishing a reference method also requires the careful choice of an appropriate basis set. Large basis sets quickly become computationally prohibitive, especially for large systems, while small basis sets are rife with errors due to being incomplete. Studies on obtaining vertical excitation energies, including studies on DHDFAs, have shown that large TZ basis sets are often sufficient to obtain nearly converged results, with little change when using QZ basis sets.^{51,66} That being said, as interaction energies in the excited state have not been studied frequently, little is known about the effects of truncated basis sets on interaction energies in excited states.

The basis set superposition error (BSSE), for example, plagues the treatment of ground-state NCIs with small AO basis sets in both WFT and DFT methods. This well-known error is caused by the limited number of AOs available on a molecular fragment ‘borrowing’ the basis functions from other fragments, which artificially stabilises the multi-fragment system—such as a dimer—relative to its separate fragments.¹⁶⁸ Small basis sets also suffer from basis set incompleteness error (BSIE) which can artificially destabilise complexes through a failure to correctly describe intermolecular electron density.¹⁶⁸ In the case of interaction energies, BSIE is not consistent with varied intermonomer separation and will therefore not be cancelled by similar error of infinitely separated monomers in the calculation of interaction energies.¹⁶⁹ If use of a larger basis set is not suitable, additive corrections have been developed for ground states to minimise the BSSE.^{170–172} BSIE corrections^{173,174} have also been developed although they are generally scarce and more computationally demanding. To our knowledge BSSE corrections have not yet been developed for excited states, and basis set convergence studies for such systems are sparse.^{121,169} Some previous exciplex studies^{7,8,88,90,102,104,120,121,169} opted to utilise the counterpoise correction¹⁷⁵ to account for BSSE in exciplex binding. Our own basis set study below utilises BSSE-uncorrected interaction energies in interest of time and avoiding corrections that have not been thoroughly assessed



specifically for excited states. In that way we are also able to provide a picture of current methodology that can be easily applied by method users. Moreover, we strive to use large basis sets for CBS extrapolations, which further reduces the impact of BSSE. That being said, it is worthwhile to explore the influence of counterpoise or similar corrections in later studies. A BSIE correction for excited states was recently developed by Loos and co-workers that was shown to recover chemically accurate vertical excitation energies of small organic molecules with augmented double- ζ basis sets, with the exception of diffuse excited states.¹⁷⁶ This suggests that work to establish basis set error corrections for excited states is underway, however, current methods do not seem to be robust enough yet to provide reliable corrections for our present study.

Note that we explore the CBS limit with extrapolation techniques that were originally developed for ground-state problems and have to our knowledge never been assessed for NCI energies in excited states.

3.1 Which CC2 variant is more appropriate?

For this section, non-relaxed dissociation curves for the first excited state of the benzene dimer were generated with CC2 and SCS-CC2 and various AO basis sets. While CC2 and SCS-CC2 both yield reliable excitation energies, albeit still above the related chemical-accuracy threshold of 0.1 eV,^{66,177} benchmark studies suggest that the latter does not show consistent improvement to vertical excitation energies in general.^{177,178} However, enhanced excited-state geometries and vibrational frequencies could lead to more accurate 0–0 transition energies for $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ excitations.^{179–181} Therefore, any potential improvements for SCS-CC2 seem to be problem specific, hence we need to carry out a comparison with CC2.

To decide between CC2 and SCS-CC2, we conduct a study with DZ-, TZ-, and QZ-quality basis sets both with and without additional diffuse functions. This comparison involves both Ahlrichs [def2- n ZVP(D)]^{130,182} and Dunning [(aug)-cc-pVnZ]^{183,184} basis sets; where n corresponds to 'S/D', 'T', 'Q', respectively. All dissociation curves are shown in Fig. 3 including selected points for CCSDR(3)/def2-TZVP near the minimum to allow for an initial evaluation. Note that due to its computational cost, generating complete dissociation curves with CCSDR(3) was technically not feasible. Numerical values for interaction energies and intermolecular distances at the respective curve minima are shown in Table S1.†

It is obvious that all CC2 minima—regardless of the basis set—are deeper than the SCS-CC2 ones, meaning that CC2 consistently gives larger dissociation energies (Fig. 3). CC2 results indicate systematic overstabilisation, exhibiting dissociation energies that are almost twice as large as CCSDR(3). For instance, with the def2-TZVP basis set, the CCSDR(3) interaction energy at an intermolecular distance r of 3.00 Å (a point in proximity of the expected minimum for this level of theory) is $-11.96 \text{ kcal mol}^{-1}$ (Table S2†) and the well depths of SCS-CC2 and CC2 are $-13.18 \text{ kcal mol}^{-1}$ ($r_e = 2.99 \text{ Å}$) and $-20.19 \text{ kcal mol}^{-1}$ ($r_e = 2.90 \text{ Å}$), respectively. While the differences between CC2 and SCS-CC2 are quite striking, the general




Fig. 3 Benzene excimer dissociation curves for various CC2 and SCS-CC2 based levels of theory compared to selected CCSDR(3)/def2-TZVP points.

trends show parallels to ground-state studies, where it has been established that conventional MP2 overestimates interaction energies in dispersion-driven complexes, while SCS-MP2 provides a more balanced description.^{28,128} CC2 has also been shown to overestimate the dispersion contribution in the same excimer complexes as studied here⁹⁰ and it appears, by its closer proximity to CCSDR(3), that SCS-CC2 reduces this overestimation. Despite SCS-CC2 being in closer proximity to CCSDR(3), both CC2 variants across all basis sets explored overstabilise the excimer relative to CCSDR(3)/def2-TZVP. When inspecting Fig. 3, we observe that the well depth increases for both CC2 variants with increasing cardinal number. For instance, we observe a change in interaction energy from $-12.60 \text{ kcal mol}^{-1}$ (def2-SVP) to $-13.29 \text{ kcal mol}^{-1}$ (def2-QZVP) for SCS-CC2, and from -18.22 (def2-SVP) to $-20.64 \text{ kcal mol}^{-1}$ (def2-QZVP) for CC2 (Table S1†).

Basis sets including diffuse functions—aug-cc-pVTZ and def2-TZVPD—produce minima that are considerably lower than basis sets without diffuse functions, meaning that the absolute interaction energies are larger for both CC2 variants in those cases. As CC2 more greatly overestimates the interaction energy minima, additional diffuse functions give well depths furthest from the CCSDR(3) reference. Interestingly, def2-TZVPD produces a deeper minimum than aug-cc-pVTZ. For SCS-CC2, def2-TZVPD and aug-cc-pVTZ differ by $0.58 \text{ kcal mol}^{-1}$, while for CC2 they differ by $0.38 \text{ kcal mol}^{-1}$. Without diffuse functions, Dunning and Ahlrichs basis sets are closer in energy; for instance, def2-TZVP and cc-pVTZ minima differ by $0.19 \text{ kcal mol}^{-1}$ and $0.24 \text{ kcal mol}^{-1}$ for SCS-CC2 and CC2 respectively (Table S1†). The choice to include diffuse functions is highly system and method dependent as in some cases their addition can increase the BSSE of the system.¹⁸⁵ In the case of



exciplex binding, diffuse functions giving a worse result may suggest that the BSSE and BSIE do not decrease at the same rate.¹⁶⁹ The impact of diffuse functions is of interest for future studies. Herein, we choose to discard them for pragmatic reasons, as the CCSDR(3) data could not be obtained with diffuse functions. However, this decision will not influence our subsequent TD-DFT benchmark study, as long as the same type of basis set is used therein. Our final findings and conclusions are therefore unlikely to be affected by this decision.

Based on the herein discussed findings, we rule out using the CC2 method in the remainder of the study due to its greater tendency to overestimate the well depths relative to CCSDR(3). Interaction-energy curves for Ahlrichs and Dunning basis sets without diffuse functions, show reasonable agreement for SCS-CC2, with the def2-TZVP well being only by 0.19 kcal mol⁻¹ deeper than for cc-pVTZ. Given that Ahlrichs basis sets are computationally more efficient due to relying on fewer primitive Gaussian-type orbitals, they are our preference in this study, particularly when considering the larger dimers. In order to further clarify the best choice of basis set for SCS-CC2 as a wave function reference, CBS values are generated for CCSDR(3) and SCS-CC2 in the following section.

3.2 Complete basis set extrapolations

The total energy calculated by a given method is expected to converge to a finite value with an increase in AO basis set size. This also leads to converging interaction energies, as indicated by the series of interaction energies ranging from DZ to QZ basis sets discussed in the previous section, with changes between TZ and QZ being smaller than between DZ and TZ. CBS extrapolations take advantage of this convergence behaviour to estimate the fully converged energy for a given family of basis sets. While the practice of CBS extrapolation is well established for ground-state studies, CBS extrapolations are scarcely conducted for excited-state problems due to a lack of established extrapolation methods. Prior to the development of established excited-state extrapolation methods, application of ground-state extrapolations should at least improve the ground state at the base of excited-state energies. This improvement of the ground-state energy is expected to extrapolate the convergence behaviour enough to give us further insight into the best comparison of SCS-CC2 to CCSDR(3). We therefore chose to employ the same extrapolation formulae and the same extrapolation exponents as in ground-state studies. Further studies to comprehensively investigate basis set convergence behaviour of NCIs in excited states, such as that conducted by Krueger and Blanquart for exciplex systems,¹⁶⁹ are strongly encouraged but such characterisation falls outside the scope of this study.

In ground-state extrapolations the HF and correlation energies are extrapolated separately due to their different convergence behaviours. The equivalent to ground-state HF energy in excited-state coupled-cluster treatments is the CCS total energy as it does not include electron correlation. We conducted linear-response calculations, so CCS total energies (E_{CCS}) were obtained by adding the CCS excitation energy to the HF ground-state total energy. Such excited-state total energies for two

truncated basis sets were then used to obtain the resulting CBS-limit energy with the familiar formula:¹⁸⁶

$$E_{CCS}(CBS) = \frac{E_{CCS}(X)\exp(-\alpha\sqrt{Y}) - E_{CCS}(Y)\exp(-\alpha\sqrt{X})}{\exp(-\alpha\sqrt{Y}) - \exp(-\alpha\sqrt{X})}, \quad (2)$$

where X and Y are the successive cardinal numbers of the two basis sets used, and α is a constant specific to the family of basis set used. We adopted the value of α for ground-state Ahlrichs-basis set extrapolations, namely $\alpha = 10.39$ for DZ–TZ and $\alpha = 7.88$ for TZ–QZ extrapolations.¹⁸⁷

The electron-correlation contribution E_C was obtained from the differences between CCS and SCS-CC2 or CCSDR(3) excitation energies. Those contributions were extrapolated to the CBS limit with the familiar formula for the correlation energy (E_C):¹⁸⁸

$$E_C(CBS) = \frac{X^\beta E_C(X) - Y^\beta E_C(Y)}{X^\beta - Y^\beta}, \quad (3)$$

where the basis-set specific constant β has values of 2.40 (DZ–TZ) or 2.97 (TZ–QZ), respectively.¹⁸⁷

The resulting energies $E_{CCS}(CBS)$ and $E_C(CBS)$ were added together to obtain the excited-state total energies. These were then used to calculate the interaction energies at the CBS level. Ideally, the basis sets used in such extrapolations should be as large as technically possible. For CCSDR(3), only DZ and TZ calculations were feasible for the benzene dimer, while TZ and QZ calculations were possible for SCS-CC2. The resulting CCSDR(3)/CBS(2,3) interaction energies for the same four selected intermolecular distances discussed for CCSDR(3)/def2-TZVP in the previous section are shown in Fig. 4 (numerical values in Table S2†) alongside the SCS-CC2 curves with truncated Ahlrichs basis sets and CBS(3,4). SCS-CC2/CBS interaction energies are slightly more negative than CCSDR(3)/CBS for three




Fig. 4 Benzene excimer interaction energies around the minimum energy well for SCS-CC2 and CCSDR(3), including CBS-extrapolated results.



of the four points with the differences ranging from 0.34 kcal mol⁻¹ ($r = 2.90$ Å) to 0.1 kcal mol⁻¹ ($r = 3.00$ Å). Both levels of theories agree for $r = 3.05$ Å with $\Delta E = -13.42$ kcal mol⁻¹. The agreement between both levels is therefore better near the minimum region of CCSDR(3)/CBS, which lies close to 3 Å (see Fig. 4).

Fig. 4 also shows parts of the dissociation curves of the three truncated basis set levels for SCS-CC2. We can clearly rule out the DZ level from further consideration. The TZ and QZ curves are relatively close to one another with the difference between both their minima being only 0.11 kcal mol⁻¹, suggesting the result is close to convergence (Table S1†). The SCS-CC2/def2-TZVP minimum lies 0.34 kcal mol⁻¹ above the CCSDR(3)/CBS interaction energy at $r = 3.00$ Å. In contrast, the SCS-CC2/def2-QZVP minimum is only 0.23 kcal mol⁻¹ higher. Considering that the CCSDR(3)/CBS interaction energy at $r = 3.00$ Å is -13.52 kcal mol⁻¹, the absolute percentage errors for the SCS-CC2/TZ, QZ and CBS(3,4) minima are only 2.5, 1.7 and 0.96%, respectively; see Table 2 for the SCS-CC2/CBS(3,4) minimum.

When choosing a reliable benchmark level of theory for the subsequent TD-DFT study, the accuracy of a method must be considered alongside the time and resources needed to treat the system sizes involved in this study as well as the large number of points needed to generate smooth dissociation curves. As SCS-CC2/def2-QZVP calculations are feasible across all excimer systems, obtaining SCS-CC2/CBS(3,4) curves offers the potential for a reference at a standard comparable with ground-state studies. Ideally, to establish the accuracy of using this level of theory, we would want to obtain CCSDR(3)/def2-TZVP data for the larger excimer systems to offer a comparison. However, at this stage, our computational resources prevent this proposed extension. Given the low percentage errors for the benzene test case for truncated basis sets, discussed above, we obtain SCS-CC2 values for all excimers with TZ, QZ and CBS(3,4). Table 2 lists the values of D_e and r_e from minima across these basis set treatments, while the corresponding dissociation curves are given as Fig. S2–S5.† Without higher-level data for all systems we cannot make a definitive statement as to the accuracy of these results. That being said, prior to extrapolation r_e appears already close to convergence across all systems and D_e differs within 0.11 to 1.52 kcal mol⁻¹.

Table 2 SCS-CC2 values for truncated basis sets and at the CBS limit for all excimer structures. The CBS values serve as reference values for the subsequent TD-DFT benchmark study

Basis	Benzene	Naphthalene	Anthracene	Pyrene
Dissociation energy D_e (kcal mol⁻¹)				
def2-TZVP	13.18	29.18	29.86	35.64
def2-QZVP	13.29	28.50	28.84	34.12
CBS(3,4)	13.65	28.32	28.40	36.33
Equilibrium distance r_e (Å)				
def2-TZVP	2.99	3.06	3.20	3.20
def2-QZVP	2.99	3.05	3.19	3.20
CBS(3,4)	2.97	3.04	3.18	3.19

Our main incentive for the TD-DFT benchmark is to assess the performance of modern TD-DFT methods, with and without applied dispersion corrections, based on their resulting dissociation energy curves. For this purpose, we choose SCS-CC2/CBS(3,4) as the reference level for the subsequent study. We would like to point out that the curves discussed herein are an improvement on what has been acceptable in the field as binding energies with basis sets of QZ quality or higher are rare for aromatic exciplexes. Nevertheless, we recommend a future study dedicated solely to (single-reference) WFT methods that addresses CBS extrapolations and BSSE corrections more in detail. As such studies are beyond the scope of this work, we continue with the discussion of the TD-DFT results.

4 Benchmarking TD-DFT methods

Having established SCS-CC2/CBS(3,4) as a reference method for the four excimer models, it is now possible to move on to benchmarking various DFAs. Of the four excimers (Fig. 2), benzene is the smallest and has therefore received the most attention in previous computational studies.^{5,90,116,120,121,133,189} While the other excimers have received some attention, they are typically studied individually^{92,93,104,118,119} rather than comparatively.^{7,8,90} The binding of an excimer involves a change in the geometry upon excitation with the eclipsed dimer, also called the “perfect sandwich” structure,^{3,190} widely accepted as the most stable conformation of an excimer.^{7,8,88,133,190} The excitation responsible for excimer formation causes a displaced ground-state dimer to move into this eclipsed form, which is also associated with a reduction in distance between the monomers.^{3,5} The most energetically stable intermolecular separation as determined experimentally for aromatic excimers is reported in the range of 3.0–3.6 Å.⁴ Our SCS-CC2/CBS(3,4) r_e values, reported in Table 2, fall within 0.03 Å of this range which is reasonable given high-accuracy theoretical calculations tend to predict shorter r_e .⁷

The close and parallel stacking of the eclipsed formation facilitates excimer binding interactions: electrostatics, Pauli-repulsion, CT, exciton coupling and London dispersion.⁸ For example, parallel stacking and short intermolecular distance increases orbital overlap, promoting exciton delocalisation.¹⁰² The interplay of these various interactions may present a challenge for DFT methods to properly describe the dissociation energies and inter-molecular distances. The main focus of this section is to investigate DFAs with this in mind. Before we discuss each model system individually, we study the impact of the amount of FE on the TD-DFT dissociation energies. Each model system is then first discussed without the addition of dispersion corrections before the impact of said corrections is studied separately. This section ends with an overarching discussion across all four dimers by means of statistical analysis.

4.1 Dependence on Fock exchange

In this section, we analyse the influence of FE on the description of excimer binding by global hybrids. This analysis was



performed by varying percentages of FE, between 0 and 75%, for two underlying exchange–correlation (XC) approximations, PBE and BLYP. The resulting functionals of varied FE are detailed in Table 1. PBE and BLYP were chosen as they are the XC approximations behind many popular (double-)hybrids, and as using two different approximations may help us to better indicate the individual influence of FE or the underlying XC expression. The functionals are analysed through their ability to recover the minimum of the dissociation curve of each excimer which can be separated into the equilibrium distance (r_e) between the two monomers and the stability of the dimers represented by the dissociation energy (D_e). The ability of each method to describe these parameters is assessed by quantitative comparison to those calculated with our SCS-CC2/CBS(3,4) reference by use of signed percentage error:

$$\% \text{ ERROR}_i = \frac{\text{METHOD}_i - \text{REFERENCE}_i}{\text{REFERENCE}_i} \times 100\% \quad (4)$$

where i represents either D_e or r_e . In this definition of percentage error, positive values represent overestimation while negative ones represent underestimation of the quantity. Fig. 5 details the resulting signed percentage errors for D_e (top panel) and r_e (bottom panel). The numerical values and corresponding dissociation energy curves are given in Section SI.4.†

As Fig. 5 shows, the percentage errors in D_e and r_e are considerably large across all systems regardless of FE percentage. Almost all functionals underestimate D_e , with a magnitude that tends to increase with amount of FE and

system size; averaging across structures these errors range from about -83 to -38% . The size-dependence of the error can be partially explained with the fact that dispersion effects are expected to increase with system size¹⁹¹ and that dispersion corrections have not been applied at this stage. While percentage errors in r_e are predominantly positive, negative errors seem to be associated more with higher amounts of FE, with average magnitudes ranging from 4 to 17%. Recovery of each minimum characteristic does not meet conveniently at the same functionals, *i.e.* a good description of r_e is not accompanied by small errors in D_e across all the functionals. Despite the overall poor performance from global hybrids regardless of FE percentage we can still gain an insight into the impact FE has for the complicated effects of excimer binding.

Functionals with large amounts of FE tend to worsen the description of excimer binding across the four tested systems. Functionals with 20% FE yield the smallest errors in D_e for each excimer, but the errors do increase with system size, yielding errors that range from -70.8 to 3.4%. Functionals containing 75% FE (-97.5 to -41.7% error range in D_e) are comparable to those with 0% (-97.4 to -22.7% error range in D_e) which are well-known to produce large errors for excited states and offer no exception here. GGAs are therefore not explored in the benchmarking study of the following section. The poor performance associated with 37.5–75% FE, however, is somewhat surprising given that global hybrids with large amounts of FE generally describe CT and other long-range excitation effects better than smaller amounts of FE.⁶⁰ High percentages of FE combined with PBE exchange and correlation appear to improve the description of r_e in larger excimers. However this improved description of r_e cannot overcome the poor description of D_e by these functionals.

The PBE-based functionals generally yield larger dissociation energies than BLYP-based functionals and are closer to the reference despite still underestimating the excimer stability; for example, PBE20 yields errors in D_e ranging between -62.1 and 3.4% while B3LYP errors range between -70.8 and -21.7% . A notable difference is seen for BLYP and BLYP75; they are unable to bind the anthracene and pyrene excimers. Interaction energy curves of CCS, which can be thought of as the HF equivalent for electronic excited states, were plotted to offer a comparison to a method with no electron-correlation but 100% FE (see Fig. S7†). The CCS interaction energy curves possess only very shallow minima for the anthracene ($D_e = 0.31 \text{ kcal mol}^{-1}$) and pyrene ($D_e = 0.70 \text{ kcal mol}^{-1}$) excimers indicating that electron correlation plays a significant role in the stabilisation of those systems. PBE-based functionals can be thought to better describe the electron correlation contributions to the excimer states than BLYP based functionals. The difference between the two underlying XC functionals for the excimer state therefore parallels that of ground states, where PBE is more attractive than BLYP in the treatment of NCIs.^{42,84,192}

No simple trend in performance with increasing FE is observed for the description of both D_e and r_e . Despite the generally improved performance of PBE based functionals, DFAs based on both XC expressions greatly underestimate the dissociation energy and overestimate r_e with error trends that



Fig. 5 Percentage error in dissociation energy (D_e , top) and equilibrium inter-monomer separation (r_e , bottom) of each excimer with respect to SCS-CC2/CBS(3,4) for varied Fock exchange with PBE and BLYP exchange–correlation functionals. BLYP and BLYP75 results are not shown for the anthracene and pyrene excimers as they gave repulsive dissociation curves.



Table 3 Dissociation energies (D_e in kcal mol⁻¹) and equilibrium inter-monomer distances (r_e in Å) for each excimer across the nine uncorrected functionals. SCS-CC2/CBS(3,4) numbers are shown as a reference

Method	Benzene		Naphthalene		Anthracene		Pyrene	
	D_e	r_e	D_e	r_e	D_e	r_e	D_e	r_e
Reference	13.65	2.97	28.32	3.04	28.40	3.18	36.33	3.19
B3LYP	10.68	3.15	14.69	3.37	9.18	3.61	10.60	3.67
PBE38	5.69	3.17	15.30	3.03	9.04	3.26	10.24	3.30
BHLYP	8.10	3.13	11.79	3.29	6.88	3.51	8.22	3.54
CAM-B3LYP	8.58	3.09	13.32	3.24	8.18	3.45	9.71	3.50
ω B97X ^a	11.99	3.08	18.95	3.24	16.01	3.36	17.97	3.37
B2PLYP	11.36	3.05	20.58	3.19	17.23	3.36	20.51	3.38
B2GP-PLYP	11.99	3.03	22.43	3.14	19.82	3.31	23.53	3.32
ω B2PLYP	13.07	2.99	22.72	3.09	20.10	3.25	23.11	3.27
ω B2GP-PLYP	13.90	2.98	24.67	3.07	22.70	3.23	26.29	3.25

^a Approximated and based on observed minima; see explanation in text and problematic curves in Fig. S6.

increase with FE and system size indicating the requirement of more sophisticated functionals for an accurate description of excimer binding. The ability of a global hybrid TD-DFA to describe each quantity seems to require a trade-off in accuracy to the other. As both D_e and r_e must be correctly described in order to predict excimer binding, global-hybrid TD-DFA do not offer reliable results. Methods from the higher rungs of Jacob's Ladder and additive dispersion corrections have been effective in addressing some of short-comings associated with global hybrids for ground states, so their performance will be addressed in the following sections.

4.2 Individual discussion of excimer models

In this section, the interaction energy curves for each excimer across a range of TD-DFA are analysed relative to the reference of SCS-CC2/CBS(3,4). The chosen range of TD-DFA is based on either popularity, established accuracy for single-molecule excitations, or novelty; they are: B3LYP, PBE38, BHLYP (global hybrids), CAM-B3LYP, ω B97X (RS hybrids), B2PLYP, B2GP-PLYP (global DHDFAs), ω B2PLYP and ω B2GP-PLYP (RS-DHDFAs). These functionals occupy the top two rungs of Jacob's Ladder

with varying exchange–correlation components and FE percentages, with and without RS, as detailed in Table 1. For all the dissociation curves discussed, the values associated with the minima, *i.e.* D_e and r_e , are listed in Table 3. Dissociation curves are shown in Fig. 6–9. ω B97X interaction energy curves are not shown here, but instead in Fig. S6† due to observed problems with getting smooth curves that could not be fixed with any of the usual convergence techniques. ω B97X will however be discussed as part of the overall statistical analysis (Section 4.4) with the values based on the actually observed minima under the assumption that they are good approximations to the true minima. Hereafter, deviations from the reference and associated percentage errors will be given as absolute values. Indication to under- or overestimation will be discussed qualitatively.

4.2.1 The benzene excimer. For the benzene excimer, excited-state interaction energies calculated across varied inter-monomer distances at the SCS-CC2/CBS(3,4) level of theory (Fig. 6) yield a dissociation energy of 13.65 kcal mol⁻¹ at 2.97 Å (r_e). The ability to describe the shape and depth of the potential energy well differs from DFA to DFA. The assessed global hybrids (left panel in Fig. 6) consistently underbind the benzene



Fig. 6 Dissociation energy curves of the lowest-lying singlet excited state of the fully-stacked benzene dimer.





Fig. 7 Dissociation energy curves of the lowest-lying singlet excited state of the fully-stacked naphthalene dimer.

excimer, underestimating D_e by 2.97–7.96 kcal mol⁻¹ (22–58% error) and overestimating r_e by 0.16–0.20 Å (5–7% error), corresponding to the largest errors for this system. As discussed in the FE study, global hybrids are not capable of capturing the binding interactions of the benzene excimer. Poor performance of global hybrids is consistent with their inability to accurately describe the CT, exciton coupling and dispersion interactions inherent to excimer binding.^{8,42,79,193} The long-range correction in CAM-B3LYP reduces the overestimation in r_e compared to its uncorrected counterpart B3LYP with an improvement from a deviation of 0.18 to 0.12 Å (central panel in Fig. 6); however it also worsens the description of D_e (2.97 kcal mol⁻¹ underestimation for B3LYP vs. 5.08 kcal mol⁻¹ for CAM-B3LYP). Double hybrids yield an improved description of interaction energies along the dissociation curve, with B2GP-PLYP being closer to the reference than B2PLYP (central panel in Fig. 6), both of which underestimate D_e by 1.66 and 2.29 kcal mol⁻¹, respectively (12 and 17% error).

Double hybrids provide a more balanced description of the excimer binding than global and RS hybrids. This improvement is consistent with benchmarking trends of double-hybrid robustness for excitation energies and absorption spectra due to their perturbative correction.^{47,66,77} Improved performance by DHDFAs additionally parallels the finding that only double hybrids were able to properly describe an exciton-coupled ECD spectrum,⁷⁹ for which WFT methods had to be applied earlier.¹⁹⁴ However, as recently re-emphasised, global double hybrids still fail to correctly describe CT excitations.⁶³ The combination of range-separation and perturbative correction improves the description of the benzene excimer dissociation energy curve, giving the curves closest to the reference at all intermolecular distances (right panel in Fig. 6). The assessed RS-DHDFAs, ω B2PLYP and ω B2GP-PLYP, improve upon their uncorrected counterparts, with the former underestimating D_e by 0.58 kcal mol⁻¹ (4% error) and the latter slightly overestimating it by 0.25 kcal mol⁻¹ (2% error). Close comparison of RS-DHDFAs with the reference interaction energy curves of the benzene excimer display potential to corroborate claims of their robustness for local-valence and long-range excitations.^{52,53,62,64}

In summary, it is observed that, for the benzene excimer, the description of binding is improved by climbing Jacob's Ladder with rung-five functionals showing considerable improvement over those belonging to rung four (see Table 1 for the Jacob's Ladder classification of each assessed functional).

4.2.2 The naphthalene excimer. The naphthalene excimer is more than twice as stable as the benzene excimer, yielding a SCS-CC2/CBS(3,4) dissociation energy of 28.32 kcal mol⁻¹ at 3.04 Å (Fig. 7). When assessing DFA dissociation energies, one has to consider the well-documented issue of conventional TD-DFT methods often struggling to correctly order the first two excited states of the naphthalene monomer.¹⁹⁵ For instance, TD-B3LYP and TD-BHLYP incorrectly predict the ¹L_a state of the monomer to be lower than the ¹L_b state.^{52,58,195} In the fully-stacked dimer, the first excimer state has, in fact, ¹L_a character, but at the dissociation limit this turns out to be the second excited state.^{7,118} Indeed, we observe this for our reference method and most other methods, but want to point out that the aforementioned problem for the naphthalene monomer is also observed here, which can lead to the incorrect calculation of dissociation energies for B3LYP and BHLYP if this problem is not spotted. This is particularly a problem when a calculation is carried out without any symmetry, but less so if the programs used distinguish between excited-state symmetries, as the symmetries of the first two excited states differ.

The chosen exchange–correlation functionals uniformly predict a weaker binding with absolute errors of D_e ranging between 3.65 and 16.53 kcal mol⁻¹ (13–58% error). r_e is largely overestimated with absolute deviations ranging from 0.01 to 0.33 Å (0.3–11% error), the exception being PBE38, which underestimates the distance by only 0.01 Å. The fourth-rung functionals behave differently between naphthalene and benzene excimers whereas fifth-rung functionals exhibit similar performance trends. For the naphthalene excimer PBE38 offers an r_e closer to the reference than the BLYP based functionals, however underestimation of D_e is on par with other DFAs belonging to this rung (46% error), consistent with results from our previous FE study in Section 4.1. B3LYP binds the excimer with a D_e comparable to those of other global hybrids (48%






Fig. 8 Dissociation energy curves of the lowest-lying singlet excited state of the fully-stacked anthracene dimer.




Fig. 9 Dissociation energy curves of the lowest-lying singlet excited state of the fully-stacked pyrene dimer.

error) while its long-range corrected counterpart, CAM-B3LYP, again improves the description of r_e (reduced deviation from 0.33 to 0.20 Å) but yields D_e akin to the global hybrids (53% error). While the relative trends between rung-four functionals differ between benzene and naphthalene excimers, the finding remains that global hybrids provide the worst results.

Double hybrids improve the description of excimer binding considerably with absolute errors in the order of 21–27% and 3–5% for D_e and r_e , respectively. Further improvement on conventional DHDFAs results from the inclusion of range-separation with absolute errors in the order of 13–20% and 1–2% for D_e and r_e , respectively. Despite the closer resemblance of RS-DHDFAs curves with the reference curve (right panel in Fig. 7) the binding strength is insufficiently described. The best method, ω B2GP-PLYP underestimates the dissociation energy by 3.65 kcal mol⁻¹, which is a larger error than for the benzene dimer in Section 4.2.1. Most likely, this error can be attributed to missing dispersion, as its importance increases with the number of electrons.¹⁹¹ This will be further discussed in Section 4.4.

4.2.3 The anthracene excimer. The anthracene excimer is slightly more stable than the naphthalene excimer, with an SCS-CC2/CBS(3,4) dissociation energy of 28.40 kcal mol⁻¹ at a larger equilibrium monomer separation of 3.18 Å (Fig. 8). The herein

tested TD-DFAs uniformly underbind the anthracene excimer with absolute errors in D_e between 5.71 and 21.52 kcal mol⁻¹ and overestimate r_e by between 0.05 and 0.43 Å. Trends in functional performance are similar to those of the naphthalene excimer, although with larger errors in D_e . Each functional predicts a smaller excimer dissociation energy for the anthracene excimer than it does for the naphthalene excimer, so comparison with the larger reference value for the anthracene excimer results in larger errors for D_e (20–76%).

It was previously noted that the D_e of the anthracene excimer being larger than that of the naphthalene excimer may be due to a larger dispersion contribution to the excimer binding.⁷ Given that the TD-DFT methods discussed in this section are unable to describe dispersion, this further supports the study of dispersion-corrected functionals for excimer binding in Section 4.3.

4.2.4 The pyrene excimer. The pyrene excimer (Fig. 9) is more stable than the anthracene excimer, with an SCS-CC2/CBS(3,4) dissociation energy of 36.33 kcal mol⁻¹ at 3.19 Å. Similarly to the naphthalene dimer, we also observed the wrong order of states at the dissociation limit for some functionals, however, difficulties with the analysis can be avoided when using symmetry in the calculation. Each functional predicts similar dissociation energies as for the anthracene case (within



1.20–3.71 kcal mol⁻¹), which relative to the larger reference D_e of the pyrene excimer yields larger errors (28–77%). The trends observed for the naphthalene and anthracene excimers hold true for the pyrene excimer, with an improved description of excimer stabilisation moving up Jacob's Ladder.

The analysis of r_e is similar between pyrene and anthracene, showing errors for the same methods within 0.05 Å. Range-separation improves the equilibrium distance, with RS-DHDFAs predicting r_e within an absolute error of 0.08 Å (3% error).

To summarise Section 4.2, the relative performance of the tested functionals follows a general trend: global hybrids and CAM-B3LYP perform the worst in terms of D_e and r_e , with a slight improvement in r_e through range-separation; double hybrids improve upon the description of both well characteristics while range-separated double hybrids show the best description of both characteristics across the tested functionals. We saw an increase in the error of D_e with system size which may be due to the lack of properly treating dispersion interactions. In the following section we explore the application of ground-state optimised dispersion corrections to assess their potential in accounting for the missing dispersion in excimer binding.

4.3 Impact of dispersion corrections

So far we have only discussed dispersion-uncorrected results. As dispersion-uncorrected TD-DFAs are able to partially predict excimer binding it is clear that interactions beyond dispersion are important, which aligns with the findings from an energy decomposition analysis reported in 2018.⁸ In our introduction we have pointed out how some studies make use of dispersion-uncorrected methods for the treatment of excimers, but our results clearly show that dispersion effects should not be neglected. Herein, we demonstrate the impact of ground-state dispersion corrections for the two extreme cases in our study, namely the benzene dimer as the smallest and the pyrene dimer as the largest system.

Ground-state dispersion corrections are often applied to excited-state studies justified by seemingly good outcomes from a few studies.^{5,114,115} However, it is reasonable to expect dispersion interactions to change upon electronic excitation¹¹³ rendering any benefit application-dependent. Herein, we test these justifications for two schemes of additive DFT-D type dispersion corrections as applied to the excimer state without any state-specific adjustment. The DFT-D3(BJ) and DFT-D4—hereafter dubbed “D3(BJ)” and “D4”—dispersion energies were calculated for all points along the dissociation curves and added to the respective total excited-state energies. The analysis of these dispersion-corrected dissociation energy curves focuses on four functionals B3LYP, CAM-B3LYP and B2PLYP and ω B2PLYP, *i.e.* two global functionals and their range-separated counterparts (minima given in Table 4). A more generalised picture across all eight dispersion-corrected functionals will be provided in the overall statistical analysis in Section 4.4.

4.3.1 Binding region of potential energy well. For the benzene excimer, both dispersion corrections overcorrect the

Table 4 Dissociation energies (D_e in kcal mol⁻¹) and equilibrium inter-monomer distances (r_e in Å) for the benzene and pyrene excimers for selected dispersion-corrected and uncorrected functionals. SCS-CG2/CBS(3,4) numbers are shown as a reference

Functional	Uncorrected		DFT-D3(BJ)		DFT-D4	
	D_e	r_e	D_e	r_e	D_e	r_e
Benzene excimer						
Reference	13.65	2.97	—	—	—	—
B3LYP	10.68	3.15	21.24	3.03	21.47	3.03
CAM-B3LYP	8.58	3.09	15.24	3.04	15.77	3.03
B2PLYP	11.36	3.05	16.94	3.00	16.96	3.01
ω B2PLYP	13.07	2.99	13.20	2.99	13.19	2.99
Pyrene excimer						
Reference	36.33	3.19	—	—	—	—
B3LYP	10.60	3.67	33.41	3.31	34.37	3.31
CAM-B3LYP	9.71	3.50	27.39	3.33	28.41	3.32
B2PLYP	20.51	3.38	34.28	3.26	34.60	3.27
ω B2PLYP	23.11	3.27	23.63	3.27	23.66	3.27

binding predicted by each TD-DFA except ω B2PLYP (Fig. 10). Whether D4 or D3(BJ) is closer to the reference depends on whether the dispersion-corrected method under- or overbinds the excimer; if the former is the case, the D4 version is closer to the reference, but in the latter case the D3(BJ) variant is closer. Across the four functionals shown here, absolute deviations in D_e range from 0.46 to 7.82 kcal mol⁻¹ for D4 and from 0.45 to 7.59 kcal mol⁻¹ for D3(BJ). The corrections offer an improved but overestimated description of the optimal inter-monomer separation for B3LYP, CAM-B3LYP, and B2PLYP with errors between 0.03 and 0.07 Å for D3(BJ) and between 0.04 and 0.06 Å for D4. However, the improvement to r_e is drastically overshadowed by the greatly overstabilised dissociation energy. Dispersion-corrected and -uncorrected ω B2PLYP give the same overestimation of r_e by 0.02 Å. As the smallest of the model systems, benzene excimer binding contains a smaller dispersion contribution than the larger models. With less dispersion to account for, more complex functionals are able to reasonably account for the other contributions to the dissociation energy without dispersion correction. The ground-state optimised corrections, thus, have a tendency to overestimate the dispersion contribution to the benzene excimer dissociation energy.

The pyrene excimer is larger and more strongly bound than the benzene excimer, which uncorrected TD-DFAs fail to predict with errors ranging from 13.21 to 26.62 kcal mol⁻¹ (Fig. 11). For this larger system, dispersion-corrected TD-DFAs offer some improvement to the description of the pyrene excimer binding with D4 corrections predicting stronger binding than D3(BJ) for all functionals, and therefore yields smaller errors: D4 and D3(BJ) underestimate D_e with errors ranging from 1.74 to 12.67 kcal mol⁻¹ and from 2.06 to 12.70 kcal mol⁻¹, respectively.

Both dispersion corrections considerably improve the B3LYP dissociation energy to within 2.91 kcal mol⁻¹ accompanied by an improved r_e with an overestimation of 0.12 Å. Dispersion-corrected CAM-B3LYP does not show the same improvement






Fig. 10 Dissociation energy curves of the lowest-lying singlet excited state of the fully-stacked benzene dimer for selected functionals with and without dispersion corrections.

to the dissociation energy as its global counterpart, yielding an underestimation of up to $8.93 \text{ kcal mol}^{-1}$ for D_e and an overestimation of r_e by 0.14 and 0.13 Å for D3(BJ) and D4, respectively. B2PLYP-D3(BJ) and -D4 give dissociation energies with errors comparable to dispersion-corrected B3LYP (underbound by 2.05 and 1.73 kcal mol^{-1}), but with a smaller geometry error (overestimation of 0.07 and 0.08 Å). Dispersion corrected ω B2PLYP offers only a small improvement to the uncorrected functional such that ω B2PLYP-D3(BJ) and ω B2PLYP-D4 underbind the pyrene excimer by 12.70 and 12.67 kcal mol^{-1} , respectively, and overestimate r_e by 0.08 Å. A negligible impact of D3(BJ) and D4 on both ω B2PLYP and ω B2GP-PLYP has also been noticed in ground-state benchmarking on the complete GMTKN55 (ref. 27) database and its NCI category.¹⁵²

For the larger system size of the pyrene excimer, the ground-state dispersion corrections improve TD-DFA dissociation energies and equilibrium distances. This is likely why others^{5,93,117,119} have recommended the use of ground-state optimised dispersion corrections for the calculation of this excimer's dissociation energy. While DFT-D type dispersion corrections offer some improvement to the description of excimer binding, their performance is far from a black-box solution for excited states. Inconsistent improvement by ground-state

dispersion corrections for excited states reinforces the need for state-specific dispersion corrections for reliable and predictive TD-DFT methods.

4.3.2 Unphysical repulsion in the mid-range. Beyond the binding region, in the medium-to-long range inter-monomer separation, TD-DFA interaction energy curves falsely predict a repulsive region. For the benzene excimer, all functionals as well as SCS-CC2/def2-TZVP exhibit that positive region falling between 4.3 Å and the asymptote. An extrapolation of SCS-CC2 to the CBS limit [CBS(3,4)] mostly corrects the unphysical repulsion of SCS-CC2/def2-TZVP, reducing the repulsion to less than $0.01 \text{ kcal mol}^{-1}$, which is way within the expected numerical noise for that method and a negligible value (see Fig. S13†). Larger excimers exhibit a similar repulsive region for global hybrids and range-separated hybrids (see Fig. S14–S17† for further details). Herein, we focus on the benzene excimer by observing B3LYP and B2PLYP with and without dispersion corrections in the range between 4.00 and 12.00 Å (Fig. 12). The extent of the repulsion for dispersion-uncorrected B3LYP is over twice that of B2PLYP, with maxima of 0.35 and 0.12 kcal mol^{-1} , respectively. The D3(BJ) correction reduces the repulsion yielding maxima of up to 0.02 kcal mol^{-1} , which brings the two functionals into close comparison with each other (maxima




Fig. 11 Dissociation energy curves of the lowest-lying singlet excited state of the fully-stacked pyrene dimer for selected functionals with and without dispersion corrections.






Fig. 12 Unphysical repulsive region for the benzene excimer state dissociation curve and its correction by ground-state optimised dispersion corrections.

within 0.001 kcal mol⁻¹ of each other). The D4 curves are also comparable in this region, although they exhibit larger maxima than D3(BJ) curves: 0.07 and 0.06 kcal mol⁻¹ for B3LYP-D4 and B2PLYP-D4, respectively. The positive behaviour that remains after the applied corrections would not be expected with corrections appropriately parametrised for excited states. Interestingly, in a recent study of exciplex interaction energy curves, a similar unphysical repulsive hump exhibited by ω B97 also saw correction by ω B97X-D3 (ref. 196) with DFT-D3(0), something that has not been noticed by the authors.¹⁰² As excimer binding is comprised of more than just dispersion effects,⁸ dispersion-uncorrected functionals are able to predict some attraction in the binding region, with more complex functionals performing reasonably in binding the benzene excimer. In the medium-range, however, the dispersion-uncorrected functionals predict an unphysical repulsive

region. Medium-range inter-monomer separations appear to be governed by dispersion, and can therefore only be accounted for by inclusion of the dispersion energy missing from the description by dispersion-uncorrected TD-DFA.

With this closer look at ground-state dispersion corrections for the binding of benzene and pyrene excimers, it is clear that accounting for dispersion is important for accurate calculation of dissociation curves in both the binding region and beyond. While ground-state optimised dispersion corrections offer some improvement to excimer binding, a state-specific reparametrisation for excited states would be necessary for robust and reliable TD-DFA results. An overarching discussion of the performance of dispersion corrections for all benchmarked DFAs is presented in the following section.

4.4 Averaged functional performance

Following our previous analysis of the interaction energy curves, we continue with an overall discussion of each functional's performance averaged across all systems both with and without dispersion corrections. Functional performance is almost universally assessed by mean absolute deviations (MADs) as the metric for benchmarking of quantum-chemical methods.¹⁹³ Here, the MAD for each characteristic averaged for each method is calculated by following the general form:

$$\text{MAD}_i = \frac{\sum_i^N |\text{METHOD}_i - \text{REFERENCE}_i|}{N} \quad (5)$$

where METHOD_i and REFERENCE_i are the values of the property for the i th excimer, either D_e or r_e , and N the number of systems. An analysis of MADs is simply presented as a convenient metric to summarise the performance of a method across our four model structures, and we acknowledge a sample size of four does not offer MADs with statistical significance comparable to more comprehensive excited-state benchmarks.^{51,55,57,59,163-165,197} MADs for the nine functionals, with and without D3(BJ) and D4 dispersion corrections, compared to the SCS-CC2/CBS(3,4) reference are presented in Fig. 13, with corresponding numerical values reported in Table S9.†




Fig. 13 Mean absolute deviations for D_e (left) and r_e (right). * We assessed the pure ω B97X. Its various dispersion-corrected variants all depend on slightly different underlying XC expressions,^{31,145,196,198-200} which is why dispersion-corrected results are not provided.



For the description of D_e and r_e , dispersion-uncorrected functionals display a descending trend for the MADs while ascending Jacob's Ladder (Fig. 13). It should be noted that, for dispersion-uncorrected functionals, these deviations overwhelmingly correspond to underestimations of D_e and overestimations of r_e , whereas dispersion-corrected functionals vary in this regard (see ESI†). The largest errors in D_e correspond to global hybrids and CAM-B3LYP yielding MADs in the order of 15.38 to 17.93 kcal mol⁻¹. Similarly, the largest MADs in r_e range between 0.23 and 0.36 Å and are attributed to CAM-B3LYP, B3LYP and B3LYP, *i.e.* the global and RS hybrids with Becke88 exchange and LYP correlation. PBE38 gives a much smaller MAD of 0.10 Å, comparable to higher-rung DHDFAs.

For global hybrids, an increase in FE is associated with an increase in MAD for D_e , corroborating the results of the FE study (Section 4.1). This increase in deviation supports the tendency of global hybrids with large amounts of FE to cause issues for describing excitations in TD-DFT.^{67–70} For example, global hybrids with large components of FE tend to produce blue-shifted excitation energies in single molecules.^{55,57,66} Despite the established better description of CT excitations with range-separated hybrids, CAM-B3LYP performs worse than B3LYP. On the other hand, ω B97X, despite the aforementioned problems with obtaining smooth dissociation curves, proves to be the best tested dispersion-uncorrected hybrid in this study with MADs of 10.45 kcal mol⁻¹ and 0.17 Å.

We have already mentioned problems in state order for the naphthalene and pyrene dimers in Sections 4.2.2 and 4.2.4, but still need to elaborate more in detail on the fact that the dimers consist of polycyclic aromatic hydrocarbons (PAHs). Single-molecule studies of PAHs have established that the first two π - π^* excitations in PAHs—called L_a and L_b according to Platt²⁰¹—are poorly described by many TD-DFT methods, including global hybrids.^{195,202} RS hybrids can improve the description of L_a but blueshift L_b excitations.^{76,203} Double hybrids, particularly the latest range-separated ones, have to date yielded the most accurate and balanced description of both states in PAHs.^{52,53,58} This, in addition to the aforementioned better description of exciton coupling,⁷⁹ most likely explains why DHDFAs are the best-performing methods in our study. The global DHDFAs B2PLYP and B2GP-PLYP have MADs of 9.26 and 7.23 kcal mol⁻¹ for D_e and of 0.15 Å and 0.11 Å in r_e . The two dispersion-uncorrected RS-DHDFAs show even better MADs, most likely due to the better description of CT effects, but their MADs in D_e are still well above the accepted chemical-accuracy threshold of 0.1 kcal mol⁻¹ for NCIs with values of 6.92 and 4.91 kcal mol⁻¹ for ω B2PLYP and ω B2GP-PLYP. Their values for r_e are 0.05 and 0.04 Å, respectively.

In conclusion, all tested dispersion-uncorrected functionals fail to recover excimer binding of SCS-CC2/CBS(3,4) quality.

The application of D3(BJ) and D4 dispersion corrections shows significantly reduced MADs across all tested functionals, excluding ω B2(GP)-PLYP, with larger reductions observed for D_e than for r_e . However, the improvement to excimer binding by the tested dispersion corrections shown through this reduction in MADs is not predictable. While dispersion-uncorrected functionals yield errors due to underestimation of D_e and

overestimation of r_e , dispersion corrections under or over-correct depending on the system and functional. D4 and D3(BJ) yield similar D_e MADs in the order of 1.96–6.58 kcal mol⁻¹ and 1.48–6.59 kcal mol⁻¹, respectively. For the same underlying functional, the two iterations of DFT-D yield comparable MADs where differences between D4 and D3(BJ) MADs do not exceed 0.03 Å or 1.16 kcal mol⁻¹, with the exception of B2GP-PLYP where D4 gives considerably higher D_e deviations than D3(BJ).

Dispersion-corrected global double hybrids give the best description of excimer binding with greatly improved r_e (ranging from 0.05 to 0.07 Å) and the smallest deviations in D_e despite their tendency to overbind the excimer systems: their MADs for D_e range from 1.48 kcal mol⁻¹ [B2GP-PLYP-D3(BJ)] to 3.77 kcal mol⁻¹ [B2GP-PLYP-D4]. Overestimation of exciplex binding from dispersion-corrected double hybrids was also noted in the study by Krueger and Blanquart where B2PLYP-D3(BJ) significantly overbound the exciplexes analysed which the authors acknowledged was largely due to the inclusion of a dispersion correction.¹⁰² Their tendency to overestimate D_e suggests that excited-state parametrised dispersion corrections would be necessary for a reliable description by TD-DFT methods.

Dispersion-corrected RS-DHDFAs only slightly improve on dispersion-uncorrected results, with MADs for D_e that range from 4.90 [ω B2GP-PLYP-D3(BJ)/D4] to 6.59 kcal mol⁻¹ [ω B2PLYP-D3(BJ)]. MADs in r_e are unchanged from dispersion-uncorrected RS-DHDFAs (0.04–0.05 Å) remaining the best performing across all functionals dispersion-corrected or otherwise. As the first application of ground-state parametrised dispersion corrections to RS-DHDFAs for excited-state problems, these functionals display potential for TD-DFT dissociation energies that approach SCS-CC2/CBS(3,4) quality. However, their tendency to underestimate D_e reinforces the necessity of excited-state parametrised dispersion corrections for a reliable description by TD-DFT methods.

From the discussion of dispersion-uncorrected functionals, double hybrids and range-separated double hybrids performed well for the more complicated excited-state interactions comprising excimer binding. As with range-separated double hybrids, the ground-state parametrised dispersion corrections do not reliably account for excited-state dispersion interactions, but without accounting for this missing dispersion TD-DFT cannot provide reliable results for excited states. While with current dispersion corrections both types of DHDFAs seem to be adequate, it is safe to assume that the better description of CT with RS-DHDFAs means that they will prevail once paired with state-specific corrections.

Global hybrids and CAM-B3LYP show a significant error reduction upon dispersion correction yielding MADs for D_e in the order of 3.20–5.12 kcal mol⁻¹, comparable to the MADs of higher-rung dispersion-corrected functionals, alongside an improved but still overestimated r_e with MADs ranging from 0.05 to 0.12 Å. The dispersion corrections offer considerable improvement to excimer binding for these less sophisticated functionals which gave the largest MADs when dispersion-uncorrected. However, even with dispersion-corrected, global hybrids and CAM-B3LYP are still largely outperformed by



DHDFAs. PBE38 seems to be able to compete with the best methods, but we would like to reiterate that it does not describe CT⁵⁷ nor do related PBE-based hybrids describe exciton coupling correctly,⁷⁹ which indicates that its good performance is most likely influenced by fortuitous error cancellation.

5 Summary and conclusion

The binding of four different aromatic excimer models was analysed by means of dissociation curves. To our knowledge, this is the first study to provide single-reference wave function curves at the complete basis set (CBS) limit for aromatic excimer systems. More specifically, linear SCS-CC2/CBS(3,4) was established as a reliable reference that allowed us to shed light onto various TD-DFT methods including a detailed analysis of excimers with double-hybrid density functionals and the first application of range-separated double hybrids to such systems. Our main goal was to address the impact of Fock exchange, range separation, the perturbative nonlocal correction used in double hybrids, and London dispersion corrections. We analysed two quantities that characterise the minima along the dissociation curves, namely dissociation energies D_e and intermonomer equilibrium distances r_e .

Overall, it turned out to be challenging to obtain a good description of both D_e and r_e with hybrid functionals, whereas double hybrids provided a more robust picture. High admixtures of Fock exchange in global hybrids usually led to smaller dissociation energies or even repulsive curves. Dispersion-uncorrected global hybrids with 50% or less Fock exchange, range-separated hybrids, as well as global and range-separated double hybrids, all gave curves with distinct minima, with the latter functional type giving the best curves, most likely due to a better description of charge-transfer and exciton coupling. That being said, all dispersion-uncorrected TD-DFT methods produced large errors that increased considerably with system size. For the smallest system, the benzene excimer, dispersion-uncorrected methods see the best results from range-separated double hybrids, approaching the accuracy of SCS-CC2/CBS(3,4) quality. However, for larger systems there was greater disparity between even the most accurate TD-DFT methods and the reference. D_e values were usually underestimated and r_e values overestimated, which points to missing dispersion interactions as the likely reason.

The application of ground-state parametrised dispersion corrections generally reduced the errors of dispersion-uncorrected functionals but did not reach chemical accuracy most likely due to not having been designed to describe excited states. We have noticed overstabilisation of some systems for some functionals, which occurred more often for the smaller systems. For some methods and systems, improvements of the minimum-energy regions were observed, but dissociation energies were still underestimated. To our knowledge, we were also the first to point out that most TD-DFT methods were unphysically repulsive in the mid range, something that could be reduced by applying dispersion corrections, which in turn indicated that dispersion was the most dominant contribution in that range.

Our study has shown that some of the latest and most modern TD-DFT methods, namely range-separated double hybrids, belong to the most robust and accurate when treating excited states, which parallels single-molecule studies.^{52,53,62–64,66} However, we have also shown that there is a need for the development of state-specific London dispersion corrections to achieve a reliable and robust TD-DFT description of excimers and related systems for all tested methods, including range-separated double hybrids.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

The authors are grateful to Dr Rika Kobayashi from the National Computational Infrastructure (NCI) in Canberra for her technical assistance with the Dalton program. A. C. H. acknowledges a Melbourne Research Scholarship from The University of Melbourne. L. G. is grateful for generous allocations of computational resources from the NCI within the National Computational Merit Allocation Scheme (project fk5), and Research Platform Services (ResPlat) at The University of Melbourne (project punim0094).

References

- 1 N. J. Turro, *Modern Molecular Photochemistry*, University Science Books, Sausalito, 1991.
- 2 J. B. Birks, *Rep. Prog. Phys.*, 1975, **38**, 903–974.
- 3 H. Saigusa and E. C. Lim, *Acc. Chem. Res.*, 1996, **29**, 171–178.
- 4 T. Förster, *Angew. Chem., Int. Ed. Engl.*, 1969, **8**, 333–343.
- 5 R. Huenerbein and S. Grimme, *Chem. Phys.*, 2008, **343**, 362–371.
- 6 M. C. Heissenbüttel, P. Marauhn, T. Deilmann, P. Krüger and M. Rohlfing, *Phys. Rev. B*, 2019, **99**, 035425.
- 7 S. Shirai, S. Iwata, T. Tani and S. Inagaki, *J. Phys. Chem. A*, 2011, **115**, 7687–7699.
- 8 Q. Ge and M. Head-Gordon, *J. Chem. Theory Comput.*, 2018, **14**, 5156–5168.
- 9 Q. Ge, Y. Mao and M. Head-Gordon, *J. Chem. Phys.*, 2018, **148**, 64105.
- 10 T. Förster and K. Kasper, *Z. Elektrochem.*, 1955, **59**, 976–980.
- 11 C. B. Dover, J. K. Gallaher, L. Frazer, P. C. Tapping, A. J. Petty, M. J. Crossley, J. E. Anthony, T. W. Kee and T. W. Schmidt, *Nat. Chem.*, 2018, **10**, 305–310.
- 12 J. Sung, A. Nowak-Król, F. Schlosser, B. Fimmel, W. Kim, D. Kim and F. Würthner, *J. Am. Chem. Soc.*, 2016, **138**, 9029–9032.
- 13 T. Ye, R. Singh, H. J. Butt, G. Floudas and P. E. Keivanidis, *ACS Appl. Mater. Interfaces*, 2013, **5**, 11844–11857.
- 14 D. Thirion, M. Romain, J. Rault-Berthelot and C. Poriol, *J. Mater. Chem.*, 2012, **22**, 7149–7157.
- 15 J. Vollbrecht, *New J. Chem.*, 2018, **42**, 11249–11254.



- 16 P. Thirupathi, J.-Y. Park, L. N. Neupane, M. Y. L. N. Kishore and K.-H. Lee, *ACS Appl. Mater. Interfaces*, 2015, **7**, 14243–14253.
- 17 K.-S. Focsaneanu and J. C. Scaiano, *Photochem. Photobiol. Sci.*, 2005, **4**, 817–821.
- 18 M. E. Østergaard and P. J. Hrdlicka, *Chem. Soc. Rev.*, 2011, **40**, 5771–5788.
- 19 S. Karuppappan and J.-C. Chambron, *Chem.-Asian J.*, 2011, **6**, 964–984.
- 20 D. Basting and G. Marowsky, *Excimer Laser Technology*, Springer, 2005.
- 21 L. M. Ibele, P. A. Sánchez-Murcia, S. Mai, J. J. Nogueira and L. González, *J. Phys. Chem. Lett.*, 2020, **11**, 7483–7488.
- 22 R. Improta, *Chem.-Eur. J.*, 2014, **20**, 8106–8115.
- 23 P. Hohenberg and W. Kohn, *Phys. Rev.*, 1964, **136**, B864.
- 24 W. Kohn and L. J. Sham, *Phys. Rev.*, 1965, **140**, A1133.
- 25 L. Goerigk and S. Grimme, *Phys. Chem. Chem. Phys.*, 2011, **13**, 6670–6688.
- 26 N. Mardirossian and M. Head-Gordon, *Mol. Phys.*, 2017, **115**, 2315–2372.
- 27 L. Goerigk, A. Hansen, C. Bauer, S. Ehrlich, A. Najibi and S. Grimme, *Phys. Chem. Chem. Phys.*, 2017, **19**, 32184–32215.
- 28 N. Mehta, M. Casanova-Páez and L. Goerigk, *Phys. Chem. Chem. Phys.*, 2018, **20**, 23175–23194.
- 29 A. Najibi and L. Goerigk, *J. Chem. Theory Comput.*, 2018, **14**, 5725–5738.
- 30 G. Santra, N. Sylvetsky and J. M. L. Martin, *J. Phys. Chem. A*, 2019, **123**, 5129–5143.
- 31 A. Najibi and L. Goerigk, *J. Comput. Chem.*, 2020, **41**, 2562–2572.
- 32 S. Kristyán and P. Pulay, *Chem. Phys. Lett.*, 1994, **229**, 175–180.
- 33 J. Pérez-Jordá and A. D. Becke, *Chem. Phys. Lett.*, 1995, **233**, 134–137.
- 34 P. Hobza, J. Šponer and T. Reschel, *J. Comput. Chem.*, 1995, **16**, 1315–1325.
- 35 J. Šponer, J. Leszczynski and P. Hobza, *J. Comput. Chem.*, 1996, **17**, 841–850.
- 36 L. Goerigk, *J. Chem. Theory Comput.*, 2014, **10**, 968–980.
- 37 L. Goerigk, *J. Phys. Chem. Lett.*, 2015, **6**, 3891–3896.
- 38 N. Mardirossian and M. Head-Gordon, *J. Chem. Theory Comput.*, 2016, **12**, 4303–4325.
- 39 O. A. Vydrov and T. Van Voorhis, *J. Chem. Phys.*, 2010, **133**, 244103.
- 40 S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.*, 2011, **32**, 1456–1465.
- 41 J. F. Dobson, *Int. J. Quantum Chem.*, 2014, **114**, 1157–1161.
- 42 S. Grimme, A. Hansen, J. G. Brandenburg and C. Bannwarth, *Chem. Rev.*, 2016, **116**, 5105–5154.
- 43 L. Goerigk and N. Mehta, *Aust. J. Chem.*, 2019, **72**, 563–573.
- 44 P. Morgante and R. Peverati, *Int. J. Quantum Chem.*, 2020, **120**, e26332.
- 45 J. M. L. Martin and G. Santra, *Isr. J. Chem.*, 2020, **60**, 787–804.
- 46 S. Grimme, *J. Chem. Phys.*, 2006, **124**, 034108.
- 47 L. Goerigk and S. Grimme, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2014, **4**, 576–600.
- 48 E. K. U. Gross and W. Kohn, *Adv. Quantum Chem.*, 1990, **21**, 255–291.
- 49 M. E. Casida, in *Time-Dependent Density Functional Response Theory for Molecules*, World Scientific, Singapore, 1995, pp. 155–192.
- 50 R. Bauernschmitt and R. Ahlrichs, *Chem. Phys. Lett.*, 1996, **256**, 454–464.
- 51 T. Schwabe and L. Goerigk, *J. Chem. Theory Comput.*, 2017, **13**, 4307–4323.
- 52 M. Casanova-Páez, M. B. Dardis and L. Goerigk, *J. Chem. Theory Comput.*, 2019, **15**, 4735–4744.
- 53 M. Casanova-Páez and L. Goerigk, *J. Chem. Theory Comput.*, 2021, **17**, 5165–5186.
- 54 L. Goerigk and S. Grimme, *J. Phys. Chem. A*, 2009, **113**, 767–776.
- 55 L. Goerigk, J. Moellmann and S. Grimme, *Phys. Chem. Chem. Phys.*, 2009, **11**, 4611–4620.
- 56 D. Jacquemin, V. Wathelet, E. A. Perpète and C. Adamo, *J. Chem. Theory Comput.*, 2009, **5**, 2420–2435.
- 57 L. Goerigk and S. Grimme, *J. Chem. Phys.*, 2010, **132**, 184103.
- 58 L. Goerigk and S. Grimme, *J. Chem. Theory Comput.*, 2011, **7**, 3272–3277.
- 59 S. S. Leang, F. Zahariev and M. S. Gordon, *J. Chem. Phys.*, 2012, **136**, 104101.
- 60 A. D. Laurent and D. Jacquemin, *Int. J. Quantum Chem.*, 2013, **113**, 2019–2039.
- 61 E. Brémond, M. Savarese, C. Adamo and D. Jacquemin, *J. Chem. Theory Comput.*, 2018, **14**, 3715–3727.
- 62 M. Casanova-Páez and L. Goerigk, *J. Chem. Phys.*, 2020, **153**, 064106.
- 63 M. Casanova-Páez and L. Goerigk, *J. Comput. Chem.*, 2021, **74**, 3–15.
- 64 J. Van Dijk, M. Casanova-Páez and L. Goerigk, *ChemRxiv*, 2022, DOI: [10.26434/chemrxiv-2022-xztxl-v2](https://doi.org/10.26434/chemrxiv-2022-xztxl-v2), preprint.
- 65 J. P. Perdew and K. Schmidt, *AIP Conf. Proc.*, 2001, 1–20.
- 66 L. Goerigk and M. Casanova-Paéz, *Aust. J. Chem.*, 2021, **74**, 3–15.
- 67 D. J. Tozer, R. D. Amos, N. C. Handy, B. O. Roos and L. Serrano-Andres, *Mol. Phys.*, 1999, **97**, 859–868.
- 68 D. J. Tozer, *J. Chem. Phys.*, 2003, **119**, 12697–12699.
- 69 A. Dreuw, J. L. Weisman and M. Head-Gordon, *J. Chem. Phys.*, 2003, **119**, 2943–2946.
- 70 A. Dreuw and M. Head-Gordon, *J. Am. Chem. Soc.*, 2004, **126**, 4007–4016.
- 71 N. T. Maitra, *J. Phys.: Condens. Matter*, 2017, **29**, 423001.
- 72 T. Leininger, H. Stoll, H.-J. Werner and A. Savin, *Chem. Phys. Lett.*, 1997, **275**, 151–160.
- 73 H. Iikura, T. Tsuneda, T. Yanai and K. Hirao, *J. Chem. Phys.*, 2001, **115**, 3540–3544.
- 74 T. Yanai, D. P. Tew and N. C. Handy, *Chem. Phys. Lett.*, 2004, **393**, 51–57.
- 75 R. Baer and D. Neuhauser, *Phys. Rev. Lett.*, 2005, **94**, 043002.
- 76 R. M. Richard and J. M. Herbert, *J. Chem. Theory Comput.*, 2011, **7**, 1296–1306.



- 77 S. Grimme and F. Neese, *J. Chem. Phys.*, 2007, **127**, 154116.
- 78 M. Head-Gordon, R. J. Rico, M. Oumi and T. J. Lee, *Chem. Phys. Lett.*, 1994, **219**, 21–29.
- 79 L. Goerigk, H. Kruse and S. Grimme, in *Theoretical Electronic Circular Dichroism Spectroscopy of Large Organic and Supramolecular Systems*, Wiley-Blackwell, 2012, ch. 22, pp. 643–673.
- 80 S. Hirata and M. Head-Gordon, *Chem. Phys. Lett.*, 1999, **314**, 291–299.
- 81 D. Mester and M. Kállay, *J. Chem. Theory Comput.*, 2021, **17**, 927–942.
- 82 D. Mester and M. Kállay, *J. Chem. Theory Comput.*, 2021, **17**, 4211–4224.
- 83 D. Mester and M. Kállay, *J. Chem. Theory Comput.*, 2022, **18**, 865–882.
- 84 S. Grimme, *J. Comput. Chem.*, 2006, **27**, 1787–1799.
- 85 R. A. Krueger and G. Blanquart, *Phys. Chem. Chem. Phys.*, 2019, **21**, 10325–10335.
- 86 G. F. Velardez, H. T. Lemke, D. W. Breiby, M. M. Nielsen, K. B. Møller and N. E. Henriksen, *J. Phys. Chem. A*, 2008, **112**, 8179–8187.
- 87 R. Peverati and D. G. Truhlar, *Philos. Trans. R. Soc., A*, 2014, **372**, 20120476.
- 88 D. Casanova, *Int. J. Quantum Chem.*, 2015, **115**, 442–452.
- 89 L. Goerigk, H. Kruse and S. Grimme, *ChemPhysChem*, 2011, **12**, 3421–3433.
- 90 M. Kolaski, C. R. Arunkumar and K. S. Kim, *J. Chem. Theory Comput.*, 2013, **9**, 847–856.
- 91 M. C. Capello, F. J. Hernández, M. Broquier, C. Dedonder-Lardeux, C. Jouvét and G. A. Pino, *Phys. Chem. Chem. Phys.*, 2016, **18**, 31260–31267.
- 92 N. O. Dubinets, A. A. Safonov and A. A. Bagaturyants, *J. Phys. Chem. A*, 2016, **120**, 2779–2782.
- 93 J. Hoche, H. C. Schmitt, A. Humeniuk, I. Fischer, R. Mitrić and M. I. Röhr, *Phys. Chem. Chem. Phys.*, 2017, **19**, 25002–25015.
- 94 A. A. Safonov, A. A. Bagaturyants and V. A. Sazhnikov, *J. Mol. Model.*, 2017, **23**, 164.
- 95 D. Kokkin, M. Ivanov, J. Loman, J.-Z. Cai, B. Uhler, N. Reilly, R. Rathore and S. A. Reid, *J. Chem. Phys.*, 2018, **149**, 134314.
- 96 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 97 A. D. Becke and E. R. Johnson, *J. Chem. Phys.*, 2005, **122**, 154104.
- 98 E. R. Johnson and A. D. Becke, *J. Chem. Phys.*, 2005, **123**, 024101.
- 99 E. R. Johnson and A. D. Becke, *J. Chem. Phys.*, 2006, **124**, 174104.
- 100 N. Elfers, I. Lyskov, J. D. Spiegel and C. M. Marian, *J. Phys. Chem. C*, 2016, **120**, 13901–13910.
- 101 I. Anger, E. Rykova and A. Bagaturyants, *ChemistrySelect*, 2017, **2**, 9495–9500.
- 102 R. A. Krueger and G. Blanquart, *J. Phys. Chem. A*, 2019, **123**, 1796–1806.
- 103 C. Ye, V. Gray, K. Kushwaha, S. Kumar Singh, P. Erhart and K. Börjesson, *Phys. Chem. Chem. Phys.*, 2020, **22**, 1715–1720.
- 104 Y. Gao, H. Liu, S. Zhang, Q. Gu, Y. Shen, Y. Ge and B. Yang, *Phys. Chem. Chem. Phys.*, 2018, **20**, 12129–12137.
- 105 M. T. do Casal and T. M. Cardozo, *Theor. Chem. Acc.*, 2020, **139**, 144.
- 106 A. Diaz-Andres and D. Casanova, *J. Phys. Chem. Lett.*, 2021, **12**, 7400–7408.
- 107 Y. Ikabata and H. Nakai, *J. Chem. Phys.*, 2012, **137**, 124106.
- 108 T. Sato and H. Nakai, *J. Chem. Phys.*, 2009, **131**, 5648.
- 109 T. Sato and H. Nakai, *J. Chem. Phys.*, 2010, **133**, 194101.
- 110 J. Řezáč, K. E. Riley and P. Hobza, *J. Chem. Theory Comput.*, 2011, **7**, 2427–2438.
- 111 E. A. Briggs and N. A. Besley, *Phys. Chem. Chem. Phys.*, 2014, **16**, 14455–14462.
- 112 A. D. Becke and E. R. Johnson, *J. Chem. Phys.*, 2007, **127**, 154108.
- 113 X. Feng, A. Otero-de-la Roza and E. R. Johnson, *Can. J. Chem.*, 2018, **96**, 730–737.
- 114 V. Barone, M. Biczysko and M. Pavone, *Chem. Phys.*, 2008, **346**, 247–256.
- 115 A. Fabrizio and C. Corminboeuf, *J. Phys. Chem. Lett.*, 2018, **9**, 464–470.
- 116 J. C. Amicangelo, *J. Phys. Chem. A*, 2005, **109**, 9174–9182.
- 117 B. Shi, D. Nachtigallova, A. J. Aquino, F. B. Machado and H. Lischka, *J. Phys. Chem. Lett.*, 2019, **10**, 5592–5597.
- 118 S. Shirai, Y. Kurashige and T. Yanai, *J. Chem. Theory Comput.*, 2016, **12**, 2366–2372.
- 119 S. Reiter, M. K. Roos and R. Vivie-Riedle, *ChemPhotoChem*, 2019, **3**, 881–888.
- 120 T. Rocha-Rinza and O. Christiansen, *Chem. Phys. Lett.*, 2009, **482**, 44–49.
- 121 T. Rocha-Rinza, L. De Vico, V. Veryazov and B. O. Roos, *Chem. Phys. Lett.*, 2006, **426**, 268–272.
- 122 E. Caldeweyher, C. Bannwarth and S. Grimme, *J. Chem. Phys.*, 2017, **147**, 034112.
- 123 E. Caldeweyher, S. Ehlert, A. Hansen, H. Neugebauer, S. Spicher, C. Bannwarth and S. Grimme, *J. Chem. Phys.*, 2019, **150**, 154112.
- 124 F. Furche, R. Ahlrichs, C. Hättig, W. Klopper, M. Sierka and F. Weigend, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2014, **4**, 91–100.
- 125 M. Häser and R. Ahlrichs, *J. Comput. Chem.*, 1989, **10**, 104–111.
- 126 C. Hättig and F. Weigend, *J. Chem. Phys.*, 2000, **113**, 5154–5161.
- 127 S. Grimme, *J. Chem. Phys.*, 2003, **118**, 9095–9102.
- 128 S. Grimme, L. Goerigk and R. F. Fink, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2012, **2**, 886–906.
- 129 A. Hellweg, S. A. Grün and C. Hättig, *Phys. Chem. Chem. Phys.*, 2008, **10**, 4119–4127.
- 130 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 131 O. Christiansen, H. Koch and P. Jørgensen, *Chem. Phys. Lett.*, 1995, **243**, 409–418.
- 132 D. Casanova, Y. M. Rhee and M. Head-Gordon, *J. Chem. Phys.*, 2008, **128**, 164106.
- 133 D. Kim, *Bull. Korean Chem. Soc.*, 2014, **35**, 2738–2742.



- 134 O. Vahtras, J. Almlöf and M. Feyereisen, *Chem. Phys. Lett.*, 1993, **213**, 514–518.
- 135 F. Weigend, A. Köhn and C. Hättig, *J. Chem. Phys.*, 2002, **116**, 3175–3183.
- 136 O. Christiansen, H. Koch and P. Jørgensen, *J. Chem. Phys.*, 1996, **105**, 1451–1459.
- 137 K. Aidas, C. Angeli, K. L. Bak, V. Bakken, R. Bast, L. Boman, O. Christiansen, R. Cimiraglia, S. Coriani, P. Dahle, E. K. Dalskov, U. Ekström, T. Enevoldsen, J. J. Eriksen, P. Ettenhuber, B. Fernández, L. Ferrighi, H. Fliegl, L. Frediani, K. Hald, A. Halkier, C. Hättig, H. Heiberg, T. Helgaker, A. C. Hennum, H. Hettema, E. Hjertenæs, S. Høst, I. M. Høyvik, M. F. Iozzi, B. Jansík, H. J. A. Jensen, D. Jonsson, P. Jørgensen, J. Kauczor, S. Kirpekar, T. Kjærgaard, W. Klopper, S. Knecht, R. Kobayashi, H. Koch, J. Kongsted, A. Krapp, K. Kristensen, A. Ligabue, O. B. Lutnæs, J. I. Melo, K. V. Mikkelsen, R. H. Myhre, C. Neiss, C. B. Nielsen, P. Norman, J. Olsen, J. M. H. Olsen, A. Osted, M. J. Packer, F. Pawłowski, T. B. Pedersen, P. F. Provasi, S. Reine, Z. Rinkevicius, T. A. Ruden, K. Ruud, V. V. Rybkin, P. Salek, C. C. Samson, A. S. de Merás, T. Saue, S. P. Sauer, B. Schimmelpfennig, K. Snegov, A. H. Steindal, K. O. Sylvester-Hvid, P. R. Taylor, A. M. Teale, E. I. Tellgren, D. P. Tew, A. J. Thorvaldsen, L. Thøgersen, O. Vahtras, M. A. Watson, D. J. Wilson, M. Ziolkowski and H. Ågren, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2014, **4**, 269–284.
- 138 J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865–3868.
- 139 A. D. Becke, *Phys. Rev. A: At., Mol., Opt. Phys.*, 1988, **38**, 3098–3100.
- 140 C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1988, **37**, 785–789.
- 141 B. Miehlich, A. Savin, H. Stoll and H. Preuss, *Chem. Phys. Lett.*, 1989, **157**, 200–206.
- 142 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- 143 P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, *J. Phys. Chem.*, 1994, **98**, 11623–11627.
- 144 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 1372–1377.
- 145 J.-D. Chai and M. Head-Gordon, *J. Chem. Phys.*, 2008, **128**, 084106.
- 146 A. Karton, A. Tarnopolsky, J.-F. Lamère, G. C. Schatz and J. M. L. Martin, *J. Phys. Chem. A*, 2008, **112**, 12868–12886.
- 147 F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2012, **2**, 73–78.
- 148 F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2018, **8**, e1327.
- 149 S. Grimme, *DFT-D3 V3.1*, University of Bonn, 2015.
- 150 E. Caldeweyher, S. Ehlert and S. Grimme, *DFT-D4 Version 2.0*, Mulliken Center for Theoretical Chemistry, University of Bonn, Germany, 2019.
- 151 L. Goerigk and S. Grimme, *J. Chem. Theory Comput.*, 2011, **7**, 291–309.
- 152 A. Najibi, M. Casanova-Páez and L. Goerigk, *J. Phys. Chem. A*, 2021, **125**, 4026–4035.
- 153 G. Santra and J. M. Martin, *J. Chem. Theory Comput.*, 2021, **17**, 1368–1379.
- 154 R. J. Bartlett and M. Musiał, *Rev. Mod. Phys.*, 2007, **79**, 291–352.
- 155 R. Izsák, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2020, **10**, e1445.
- 156 K. Raghavachari, G. W. Trucks, J. A. Pople and M. Head-Gordon, *Chem. Phys. Lett.*, 1989, **157**, 479–483.
- 157 A. Karton, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2016, **6**, 292–310.
- 158 P. Jurečka, J. Šponer, J. Černý and P. Hobza, *Phys. Chem. Chem. Phys.*, 2006, **8**, 1985–1993.
- 159 J. Řezáč and P. Hobza, *J. Chem. Theory Comput.*, 2013, **9**, 2151–2155.
- 160 N. Mehta, T. Fellowes, J. M. White and L. Goerigk, *J. Chem. Theory Comput.*, 2021, **17**, 2783–2806.
- 161 H. Koch, O. Christiansen, P. Jørgensen, A. M. Sanchez De Merás and T. Helgaker, *J. Chem. Phys.*, 1997, **106**, 1808–1818.
- 162 S. P. Sauer, M. Schreiber, M. R. Silva-Junior and W. Thiel, *J. Chem. Theory Comput.*, 2009, **5**, 555–564.
- 163 P. F. Loos, A. Scemama, A. Blondel, Y. Garniron, M. Caffarel and D. Jacquemin, *J. Chem. Theory Comput.*, 2018, **14**, 4360–4379.
- 164 P.-F. Loos, A. Scemama, M. Boggio-Pasqua and D. Jacquemin, *J. Chem. Theory Comput.*, 2020, **16**, 3720–3736.
- 165 M. Vénil, A. Scemama, M. Caffarel, F. Lipparini, M. Boggio-Pasqua, D. Jacquemin and P. F. Loos, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2021, **11**, e1517.
- 166 C. Riplinger, B. Sandhoefer, A. Hansen and F. Neese, *J. Chem. Phys.*, 2013, **139**, 134101.
- 167 C. Riplinger, P. Pinski, U. Becker, E. F. Valeev and F. Neese, *J. Chem. Phys.*, 2016, **144**, 024109.
- 168 F. Jensen, *Introduction to Computational Chemistry*, John Wiley & Sons, Chichester, 3rd edn, 2017.
- 169 R. A. Krueger and G. Blanquart, *Int. J. Quantum Chem.*, 2019, **119**, e25819.
- 170 A. Galano and J. R. Alvarez-Idaboy, *J. Comput. Chem.*, 2006, **27**, 1203–1210.
- 171 F. Jensen, *J. Chem. Theory Comput.*, 2010, **6**, 100–106.
- 172 H. Kruse and S. Grimme, *J. Chem. Phys.*, 2012, **136**, 04B613.
- 173 A. Otero-De-La-Roza and G. A. DiLabio, *J. Chem. Theory Comput.*, 2017, **13**, 3505–3524.
- 174 P. F. Loos, B. Pradines, A. Scemama, E. Giner and J. Toulouse, *J. Chem. Theory Comput.*, 2020, **16**, 1018–1028.
- 175 S. F. Boys and F. Bernardi, *Mol. Phys.*, 1970, **19**, 553–566.
- 176 E. Giner, A. Scemama, J. Toulouse and P.-F. Loos, *J. Chem. Phys.*, 2019, **151**, 144118.
- 177 L. Goerigk and S. Grimme, *J. Chem. Theory Comput.*, 2010, **6**, 107–126.
- 178 A. Tajti and P. G. Szalay, *J. Chem. Theory Comput.*, 2019, **15**, 5523–5531.
- 179 N. O. C. Winter, N. K. Graf, S. Leutwyler and C. Hättig, *Phys. Chem. Chem. Phys.*, 2013, **15**, 6623–6630.



- 180 C. Schon, W. Roth, I. Fischer, J. Pfister, C. Kaiser, R. F. Fink and B. Engels, *Phys. Chem. Chem. Phys.*, 2010, **12**, 9339–9346.
- 181 J. Pfister, C. Schon, W. Roth, C. Kaiser, C. Lambert, K. Gruss, H. Braunschweig, I. Fischer, R. F. Fink and B. Engels, *J. Phys. Chem. A*, 2011, **115**, 3583–3591.
- 182 D. Rappoport and F. Furche, *J. Chem. Phys.*, 2010, **133**, 1–11.
- 183 T. H. Dunning, *J. Chem. Phys.*, 1989, **90**, 1007–1023.
- 184 R. A. Kendall, T. H. Dunning and R. J. Harrison, *J. Chem. Phys.*, 1992, **96**, 6796–6806.
- 185 E. Papajak, J. Zheng, X. Xu, H. R. Leverentz and D. G. Truhlar, *J. Chem. Theory Comput.*, 2011, **7**, 3027–3034.
- 186 A. Karton and J. M. L. Martin, *Theor. Chem. Acc.*, 2006, **115**, 330–333.
- 187 F. Neese and E. F. Valeev, *J. Chem. Theory Comput.*, 2011, **7**, 33–43.
- 188 A. Halkier, T. Helgaker, P. Jørgensen, W. Klopper, H. Koch, J. Olsen and A. K. Wilson, *Chem. Phys. Lett.*, 1998, **286**, 243–252.
- 189 D. Kim, *Bull. Korean Chem. Soc.*, 2017, **38**, 763–771.
- 190 B. S. Sudhindra and A. K. Chandra, *Curr. Sci.*, 1977, **46**, 406–409.
- 191 A. Stone, *The Theory of Intermolecular Forces*, Oxford University Press, 2nd edn, 2013.
- 192 S. Grimme, *J. Comput. Chem.*, 2004, **25**, 1463–1473.
- 193 B. Ruscic, *Int. J. Quantum Chem.*, 2014, **114**, 1097–1101.
- 194 L. Goerigk and S. Grimme, *ChemPhysChem*, 2008, **9**, 2467–2470.
- 195 S. Grimme and M. Parac, *ChemPhysChem*, 2003, **4**, 292–295.
- 196 Y.-S. Lin, G.-D. Li, S.-P. Mao and J.-D. Chai, *J. Chem. Theory Comput.*, 2013, **9**, 263–272.
- 197 P. F. Loos, F. Lipparini, M. Boggio-Pasqua, A. Scemama and D. Jacquemin, *J. Chem. Theory Comput.*, 2020, **16**, 1711–1741.
- 198 J.-D. Chai and M. Head-Gordon, *Phys. Chem. Chem. Phys.*, 2008, **10**, 6615–6620.
- 199 N. Mardirossian and M. Head-Gordon, *Phys. Chem. Chem. Phys.*, 2014, **16**, 9904–9924.
- 200 A. Najibi and L. Goerigk, *J. Chem. Theory Comput.*, 2018, **14**, 5725–5738.
- 201 J. R. Platt, *J. Chem. Phys.*, 1949, **17**, 484–496.
- 202 M. Parac and S. Grimme, *Chem. Phys.*, 2003, **292**, 11–21.
- 203 K. Lopata, R. Reslan, M. Kowalska, D. Neuhauser, N. Govind and K. Kowalski, *J. Chem. Theory Comput.*, 2011, **7**, 3686–3693.

