ChemComm



COMMUNICATION

View Article Online



Cite this: Chem. Commun., 2019, 55 11123

Received 25th June 2019, Accepted 16th August 2019

DOI: 10.1039/c9cc04795f


rsc.li/chemcomm

Pd-Catalyzed decarboxylative cross-coupling reactions of epoxides with α,β -unsaturated carboxylic acids†

Xiao-Yu Lu, *\oldsymbol{0}** Jin-Song Li, *\oldsymbol{a} Shi-Qun Wang, *\oldsymbol{a} Yu-Jing Zhu, *\oldsymbol{a} Yue-Ming Li, *\oldsymbol{a} Lu-Yu Yan,^a Jia-Mei Li,^a Jin-Yu Wang, D^a Hai-Pin Zhou^a and Xiu-Tao Ge^a

A Pd-catalyzed decarboxylative cross-coupling of α,β-unsaturated carboxylic acids with cyclic and acyclic epoxides has been developed. Both β-monosubstituted and β-disubstituted unsaturated carboxylic acids, as well as conjugated diene unsaturated carboxylic acids are suitable reaction substrates. Substituted homoallylic alcohols were obtained in moderate to good yields. The product was obtained as a mixture of diastereomers favoring the anti diastereomer of the cyclic epoxides. This work provides a method for the modification of complex organic molecules containing α,β-unsaturated carboxylic acids.

Transition metal-catalyzed cross-couplings are among the most vital C-C bond-forming reactions in modern organic synthesis.¹ Alcohols are valuable compounds that play vital roles in organic synthesis. One synthesis method for alcohols is through ringopening reactions of epoxides where C-C bonds are constructed.² In the past few decades, various epoxide ring-opening reactions have been realized, 3 including transition-metal-catalyzed Negishitype⁴ and Suzuki-type⁵ ring-opening/cross-coupling reactions. In addition, reductive ring-opening/coupling reactions of epoxides have been reported.⁶ For example, reductive coupling of alkynes and epoxides does not break the C-C bond to form an alcohol as implied in Scheme 1a but rather forms the same types of homoallylic alcohols made in this publication. 6c Furthermore, Heck-type reaction⁷ and C-H activation reaction⁸ of epoxides have also been reported. For example, palladium-catalyzed Hecktype reaction of epoxides is shown in Scheme 1b.7a Carboxylic acids are one of the most important classes of organic structures, and are found in many active molecules.9 Recently, decarboxylative cross-coupling has emerged as a powerful tool for the construction of C-C bonds. 10 We hypothesized that decarboxylative crosscouplings could be used with epoxide ring-opening reactions to synthesize alcohols. To the best of our knowledge, there have been



Scheme 1 Decarboxylative cross-coupling reactions of epoxides with α,β -unsaturated carboxylic acids.

no reports on decarboxylative cross-coupling of carboxylic acids with epoxides.11

Herein, we report the first example of Pd-catalyzed decarboxylative cross-coupling reactions of α,β -unsaturated carboxylic acids with epoxides. The product was obtained as a mixture of diastereomers favoring the anti diastereomer of the cyclic epoxides (Scheme 1c). In addition to β-monosubstituted unsaturated carboxylic acids, β-disubstituted unsaturated carboxylic acids and conjugated diene unsaturated carboxylic acids also give products with moderate to good reaction yields. This methodology provides access to a diverse array of synthetically valuable substituted homoallylic alcohols.¹² It also provides a method for the modification of complex organic molecules containing α,β -unsaturated carboxylic acids.

We began our study by selecting cinnamic acid (1a) and cyclohexene oxide (2a) as model reaction substrates (Table 1). When using Pd(PPh₃)₄ as a catalyst, dppf as a ligand, Cy₂NMe as a base and PhCF₃ as a solvent, the desired product was observed in a low yield (entry 1). First, we assessed the reaction by modifying the ligand and solvent in the presence of Pd(PPh₃)₄ as the catalyst. Using dioxane instead of PhCF₃ as the solvent resulted in almost no increase in reaction yield (entry 2). Switching the ligand to Xantphos decreased the reaction yield (entries 3 and 4).

^a School of Materials and Chemical Engineering, ChuZhou University, Chu Zhou, 239000, China. E-mail: xiaoyulu@mail.ustc.edu.cn

^b School of Chemistry and Chemical Engineering, AnHui University, He Fei, 230601,

[†] Electronic supplementary information (ESI) available. See DOI: 10.1039/c9cc04795f

Table 1 Optimization of the reaction conditions

	COOH +	Pd(PPh ₃) ₄ , 5 mol% Ligand, 6 mol% Nal, 30 mol% Solvent	HO,,	HO
1a	2a	100 °C or 90 °C	3a (trans)	4a (cis)

Entry	Catalyst	Ligand	Base	Solvent	Yield% 3a:4a (d.r.)
1 ^a	Pd(PPh ₃) ₄	dppf	Cy ₂ NMe	PhCF ₃	28(4.5:1)
2^a	$Pd(PPh_3)_4$	dppf	Cy ₂ NMe	Dioxane	31(3.5:1)
3^a	$Pd(PPh_3)_4$	Xantphos	Cy ₂ NMe	$PhCF_3$	10(4.2:1)
4^a	$Pd(PPh_3)_4$	Xantphos	Cy ₂ NMe	Dioxane	15(3.2:1)
5^b	$Pd(PPh_3)_4$	dppf	Cy ₂ NMe	$PhCF_3$	50(4:1)
6^b	$Pd(PPh_3)_4$	dppe	Cy ₂ NMe	PhCF ₃	10(4.2:1)
7^b	$Pd(PPh_3)_4$	dppp	Cy ₂ NMe	PhCF ₃	2
8^b	$Pd(PPh_3)_4$	PCy_3	Cy ₂ NMe	$PhCF_3$	Trace
9^b	$Pd(PPh_3)_4$	dppf	K_3PO_4	$PhCF_3$	8(4:1)
10^b	$Pd(PPh_3)_4$	dppf	Cs_2CO_3	$PhCF_3$	5(4:1)
11^b	Pd(PPh ₃) ₄	dppf	(iPr) ₂ NEt	$PhCF_3$	10(4.1:1)
12^c	Pd(PPh ₃) ₄	dppf	Cy ₂ NMe	$PhCF_3$	62(5:1)
13 ^c	$Pd(PPh_3)_4$	dppf	Cy ₂ NMe	Toluene	46(5.5:1)
14^c	$Pd(PPh_3)_4$	dppf	Cy ₂ NMe	DMA	21(4.2:1)
15^c	$Pd(PPh_3)_4$	dppf	Cy ₂ NMe	PhCF ₃ /DMA	28(4.3:1)
16^d	$Pd(PPh_3)_4$	dppf	Cy ₂ NMe	PhCF ₃ /Diox	81(3.9:1)
17	_	dppf	Cy ₂ NMe	PhCF ₃ /Diox	0
18	$Pd(PPh_3)_4$		Cy ₂ NMe	PhCF ₃ /Diox	(3.1:1)
19^e	$Pd(PPh_3)_4$	dppf	Cy ₂ NMe	PhCF ₃ /Diox	Trace

^a Reaction conditions: **1a** (0.6 mmol), **2a** (0.3 mmol), base (1.5 equiv.) in 2 mL solvent at 100 °C for 24 h. ^b **1a** (0.3 mmol), **2a** (0.6 mmol), base (1.5 equiv.) in 2 mL solvent at 100 °C for 24 h. ^c **1a** (0.3 mmol), **2a** (0.6 mmol) in 2 mL solvent (10:1) at 90 °C for 24 h. ^d 1.5 mL solvent (5:1), **1a** (0.3 mmol), **2a** (2.5 equiv.) and NaI/Et₃N·HCl at 90 °C for 30 h. ^e No NaI. The yield was determined by GC using benzophenone as internal standard. Diox = dioxane.

We then adjusted the substrate proportions under the initial reaction conditions (entry 5) and the reaction yield was significantly improved. Using the new substrate proportions, we screened a series of ligands, such as dppe, dppp and PCy₃, but they gave reaction yields of less than 10% (entries 6-8). These results indicate that dppf is a relatively good ligand for the targeted reaction. Under the condition of dppf as a ligand, we examined other bases, but saw no improvement in the reaction yield (entries 9-11). Lowering the reaction temperature has certain benefits for improving the reaction yield (entry 12). Finally, we obtained the optimal reaction conditions by using a mixed solvent system of PhCF3 and dioxane (81% GC yield, d.r. = 3.9:1, entry 16). We confirmed the presence of two isomers by NMR (see ESI†). Reactions performed without Pd(PPh₃)₄ did not result in product formation (entry 17). On removal of dppf, the yield will be significantly reduced (entry 18). A control experiment indicated that the reaction completely failed without the addition of NaI (entry 19). Using other catalysts, such as Pd(OAc)2, significantly reduced the reaction yield (see ESI†).

We next examined the substrate scope using the optimized conditions described above (Table 2). Many of the substituted unsaturated carboxylic acids coupled with cyclic epoxides in moderate to good yields and preferential anti-selectivity on the cyclic ring. Both electron-rich and electron-poor unsaturated carboxylic acids afforded the products. The functional groups ether (3b, 3c), halogen (3e), trifluoromethoxy (3j), amino (3h)

Table 2 Scope of β-monosubstituted unsaturated carboxylic acids

and trifluoromethyl (3i) were well tolerated and afforded good yields. In particular, the ester (3f), alcohol (3m), ketal (3g) and amide (3p) substitutions were compatible with the reaction conditions. Moreover, carboxylic acids bearing *ortho*-chlorine (3k), bicyclic naphthyl (3r), thiophene (3s) and furan (3t) groups also participated in the reaction.

Conjugated dienes are found in many complex active molecules and pharmaceuticals and play vital roles in organic synthesis and materials science. We found that conjugated diene unsaturated carboxylic acids furnished the corresponding products in good yields (3aa and 3ab). In particular, sorbic acid showed satisfactory reactivity towards product formation (Scheme 2), which is notable because it is difficult to synthesize diene substituted homoallylic alcohols by other methods. ¹⁴

COOH

ChemComm Communication

Scheme 2 Decarboxylation of sorbic acids.

In addition to β-monosubstituted unsaturated carboxylic acids, β-disubstituted unsaturated carboxylic acids also give the products. Many β-disubstituted unsaturated carboxylic acids coupled with epoxides in moderate yields (Table 3). Some substituent groups, such as methyl (3bb) and methoxy (3bc), were tolerated and afforded moderate yields. Thiophene (3bd) and naphthalene (3be) can also be present in the reaction.

For monosubstituted and 1,1-disubstituted acyclic epoxides the products formed in these reactions are the ester that would result from direct nucleophilic attack of the carboxylic acid on the epoxide (see ESI†). So we tested 1,2-disubstituted acyclic epoxide as the reaction substrate. The trans-2,3-epoxybutane couples with cinnamic acid to form the product in 63% yield (Scheme 3a). The treatment of cis 3cb with cinnamic acid afforded the product in 41% yield (Scheme 3b). These results indicated that the yield of acyclic epoxide was worse than cyclic epoxide. Unfortunately, for styrene epoxides, such as transstilbene oxide, we almost got 1,2-diphenylethan-1-one (see ESI†).

An unsymmetrical epoxy (Scheme 4) substrate was investigated and two regioselective products, tertiary alcohol (3da) and secondary alcohol (3db) were obtained. The regioselectivity at the less-substituted positions was superior (3da:3db=2.5:1). Therefore, the selectivity of the asymmetric epoxy mainly depends on steric hindrance (the configuration of the isomers was assigned by ¹H NMR and H-H COSY spectra).

We next demonstrated the decarboxylative cross-coupling reactions of epoxides for late-stage modification of biologically

Table 3 Scope of β-disubstituted unsaturated carboxylic acids

Scheme 3 Examples of acyclic epoxides.

$$\begin{array}{c} Pd(PPh_3)_4,\ 8\ mol\%\\ dppf,\ 10\ mol\%\\ \hline Nal,\ Et_8NHCl\\ PhCF_3/dioxane,\ 2\ mL\\ Cy_2NMe,\ 90\ ^\circ C,\ 36\ h\\ \hline cinnamic\ acid \\ \end{array}$$

Scheme 4 Example of asymmetric epoxide

Scheme 5 The modification of complex molecules.

active molecules containing α,β -unsaturated carboxylic acids. The treatment of retinoic acid with cyclohexene oxide afforded the product (3ea) in good yield (Scheme 5a). Modification of acitretin with cyclohexene oxide resulted in the formation of 3eb in high yield (Scheme 5b). The treatment of abscisic acid with cyclohexene oxide obtained the product in 38% yield (Scheme 5c). The treatment of estrone derivative with cyclohexene oxide afforded the product (3ed) in 72% yield (Scheme 5d). These results fully demonstrate the value of this decarboxylative reaction in the modification of complex active molecules.

The mechanism of the reaction was investigated through several experiments. Reactions using 0.5 equiv. of radical trapping agent 2,2,6,6-tetramethylpiperidineoxy (TEMPO) were largely inhibited. Then, we performed a reaction of an in situ formed palladium(0) complex of dppf with iodohydrin. It formed cyclohexanone and cyclohexanol (ESI†). Another reaction of the palladium(0) complex with cyclohexene oxide gave similar results in the presence of 1 equiv. of NaI and Et₃N·HCl (ESI†). On removal of NaI, no production of cyclohexanone and cyclohexanol was observed. We observed the product with 16% yield when using iodohydrin instead of the epoxide and in the absence of sodium iodide (see ESI†). These results are similar to previous reports.^{7a} The cyclooctadiene monoxide couples with cinnamic acid to form primarily [3.3.0]-bicyclooctanols (ESI†).6b Moreover, (Z)-cinnamic acid also afforded the same trans-alkene product, and the stereoconvergent reaction was considered to be carried out by a radical pathway (ESI†).15

On the basis of the above results and previous reports, 11,16 a possible reaction mechanism is depicted in ESI.† Initially, the Communication ChemComm

β-iodohydrin, which is *in situ* produced from the epoxide and NaI, reacts with (dppf)palladium(0) species to give a β-hydroxyalkyl radical (I) and an L_n PdI complex. Addition of the hydroxyalkyl radical (I) at the α-position of the C—C double bond in cinnamic acid would give the benzylic radical II. Next, the radical II combines with (L_n) PdI to form alkylpalladium(II) species III. Finally, decarboxylation of the Pd(II) species occurs to generate the product while regenerating the Pd(0) species.

In summary, we have developed the first Pd-catalyzed decarboxylative cross-coupling of α,β -unsaturated carboxylic acids with epoxides. The products were obtained as a mixture of diastereomers favoring the anti diastereomer of the cyclic epoxides. Reactions with β -monosubstituted and β -disubstituted unsaturated carboxylic acids proceed with moderate to good yields and conjugated diene unsaturated carboxylic acids showed moderate reactivity. This methodology also provides access to a diverse array of substituted homoallylic alcohols, which are valuable structural fragments in organic synthesis. It also provides a route for the modification of complex organic molecules containing α,β -unsaturated carboxylic acids.

This work was supported by 2017qd11, 16030801108 and KJ2019A0635.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 (a) F. Diederich and P. J. Stang, Metal-Catalyzed Cross-Coupling Reactions, Wiley-VCH, Weinheim, 1998; (b) R. Jana, T. P. Pathak and M. S. Sigman, Chem. Rev., 2011, 111, 1417; (c) S. Z. Tasker, E. A. Standley and T. F. Jamison, Nature, 2014, 509, 299; (d) A. Suzuki, Angew. Chem., Int. Ed., 2011, 50, 6722; (e) N. Kambe, T. Iwasaki and J. Terao, Chem. Soc. Rev., 2011, 40, 4937.
- 2 (a) C.-Y. Huang and A. G. Doyle, Chem. Rev., 2014, 114, 8153;(b) J. He, J. Ling and P. Chiu, Chem. Rev., 2014, 114, 8037.
- 3 (a) P. Crotti and M. Pineschi, Aziridines and Epoxides in Organic Synthesis, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 271–313; (b) V. V. Fokin and P. Wu, Aziridines and Epoxides in Organic Synthesis, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 443–477; (c) P. A. S. Lowden, Aziridines and Epoxides in Organic Synthesis, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 399–442; (d) H. Ohno, Aziridines and Epoxides in Organic Synthesis, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 37–71; (e) B. Olofsson and P. Somfai, Aziridines and Epoxides in Organic Synthesis, Wiley-VCH Verlag GmbH & Co. KGaA, 2006, pp. 315–347.
- 4 (a) C. Y. Huang and A. G. Doyle, J. Am. Chem. Soc., 2012, 134, 9541;
 (b) K. L. Jensen, E. A. Standley and T. F. Jamison, J. Am. Chem. Soc., 2014, 136, 11145;
 (c) D. K. Nielsen, C. Y. Huang and A. G. Doyle, J. Am. Chem. Soc., 2013, 135, 13605.
- 5 (a) M. L. Duda and F. E. Michael, J. Am. Chem. Soc., 2013, 135, 18347;
 (b) X. Y. Lu, C. T. Yang, J. H. Liu, Z. Q. Zhang, X. Lu, X. Lou, B. Xiao and Y. Fu, Chem. Commun., 2015, 51, 2388; (c) Y. Takeda, Y. Ikeda,

- A. Kuroda, S. Tanaka and S. Minakata, *J. Am. Chem. Soc.*, 2014, 136, 8544; (d) E.-A. M. A. Ahmed, X. Lu, T.-J. Gong, Z.-Q. Zhang, B. Xiao and Y. Fu, *Chem. Commun.*, 2017, 53, 909; (e) A. Ebrahim-Alkhalil, Z.-Q. Zhang, T.-J. Gong, W. Su, X.-Y. Lu, B. Xiao and Y. Fu, *Chem. Commun.*, 2016, 52, 4891; (f) X.-Y. Lu, J.-S. Li, J.-Y. Wang, S.-Q. Wang, Y.-M. Li, Y.-J. Zhu, R. Zhou and W.-J. Ma, *RSC Adv.*, 2018, 8, 41561.
- 6 (a) Y. Zhao and D. J. Weix, J. Am. Chem. Soc., 2013, 136, 48;
 (b) Y. Zhao and D. J. Weix, J. Am. Chem. Soc., 2015, 137, 3237;
 (c) C. Molinaro and T. F. Jamison, J. Am. Chem. Soc., 2003, 125, 8076;
 (d) M. G. Beaver and T. F. Jamison, Org. Lett., 2011, 13, 4140.
- 7 (a) S. Teng, M. E. Tessensohn, R. D. Webster and J. S. Zhou, ACS Catal., 2018, 8, 7439; (b) Y. Ikeda, H. Yorimitsu, H. Shinokubo and K. Oshima, Adv. Synth. Catal., 2004, 346, 1631.
- G. Cheng, T. J. Li and J. Q. Yu, J. Am. Chem. Soc., 2015, 137, 10950.
 A. Tortajada, F. Julia-Hernandez, M. Borjesson, T. Moragas and R. Martin, Angew. Chem., Int. Ed., 2018, 57, 15948–15982.
- (a) S. Bloom, C. Liu, D. K. Kölmel, J. X. Qiao, Y. Zhang, M. A. Poss, W. R. Ewing and D. W. C. MacMillan, Nat. Chem., 2017, 10, 205; (b) J. T. Edwards, R. R. Merchant, K. S. McClymont, K. W. Knouse, T. Qin, L. R. Malins, B. Vokits, S. A. Shaw, D. H. Bao, F. L. Wei, T. Zhou, M. D. Eastgate and P. S. Baran, Nature, 2017, 545, 213; (c) C. P. Johnston, R. T. Smith, S. Allmendinger and D. W. C. MacMillan, Nature, 2016, 536, 322; (d) C. Li, J. Wang, L. M. Barton, S. Yu, M. Tian, D. S. Peters, M. Kumar, A. W. Yu, K. A. Johnson, A. K. Chatterjee, M. Yan and P. S. Baran, Science, 2017, 356, eaam7355; (e) T. Qin, J. Cornella, C. Li, L. R. Malins, J. T. Edwards, S. Kawamura, B. D. Maxwell, M. D. Eastgate and P. S. Baran, Science, 2016, 352, 801; (f) M.-C. Fu, R. Shang, B. Zhao, B. Wang and Y. Fu, Science, 2019, 363, 1429.
- 11 (a) S. Han, X. Ren, Q. Wu, A. Liang, J. Li, D. Zou, Y. Wu and Y. Wu, Adv. Synth. Catal., 2018, 360, 2308; (b) R. Kancherla, K. Muralirajan, B. Maity, C. Zhu, P. E. Krach, L. Cavallo and M. Rueping, Angew. Chem., Int. Ed., 2019, 58, 3412; (c) C. Wang, Y. Lei, M. Guo, Q. Shang, H. Liu, Z. Xu and R. Wang, Org. Lett., 2017, 19, 6412; (d) G. Prina Cerai and B. Morandi, Chem. Commun., 2016, 52, 9769; (e) K. Xu, Z. Tan, H. Zhang, J. Liu, S. Zhang and Z. Wang, Chem. Commun., 2017, 53, 10719; (f) J. J. Zhang, J. C. Yang, L. N. Guo and X. H. Duan, Chem. Eur. J., 2017, 23, 10259.
- (a) L. F. Tietze, T. Kinzel and C. C. Brazel, Acc. Chem. Res., 2009,
 367; (b) Y. Chen, D. C. Blakemore, P. Pasau and S. V. Ley, Org. Lett., 2018,
 (c) T. Takeda, S. Amarume, I. Sekioka and A. Tsubouchi, Org. Lett., 2015,
 17, 1150; (d) C. Wang and H. Yamamoto, J. Am. Chem. Soc., 2014,
 136, 1222.
- (a) M. A. Bohn, A. Schmidt, G. Hilt, M. Dindaroglu and H. G. Schmalz, Angew. Chem., Int. Ed., 2011, 50, 9689; (b) L. Liao, R. Jana, K. B. Urkalan and M. S. Sigman, J. Am. Chem. Soc., 2011, 133, 5784; (c) N. J. McAlpine, L. Wang and B. P. Carrow, J. Am. Chem. Soc., 2018, 140, 13634; (d) E. McNeill and T. Ritter, Acc. Chem. Res., 2015, 48, 2330; (e) V. T. Nguyen, H. T. Dang, H. H. Pham, V. D. Nguyen, C. Flores-Hansen, H. D. Arman and O. V. Larionov, J. Am. Chem. Soc., 2018, 140, 8434; (f) A. Tortajada, R. Ninokata and R. Martin, J. Am. Chem. Soc., 2018, 140, 2050.
- 14 (a) P. A. Wender, D. A. Holt and S. M. Sieburth, J. Am. Chem. Soc., 1983, 105, 3348–3350; (b) J. Montgomery and M. Song, Org. Lett., 2002, 4, 4009.
- 15 G. Li, T. Wang, F. Fei, Y.-M. Su, Y. Li, Q. Lan and X.-S. Wang, *Angew. Chem., Int. Ed.*, 2016, 55, 3491.
- 16 (a) J. N. Jaworski, S. D. McCann, I. A. Guzei and S. S. Stahl, Angew. Chem., Int. Ed., 2017, 56, 3605; (b) J. N. Jaworski, C. V. Kozack, S. J. Tereniak, S. M. M. Knapp, C. R. Landis, J. T. Miller and S. S. Stahl, J. Am. Chem. Soc., 2019, 141, 10462; (c) D. Wang, A. B. Weinstein, P. B. White and S. S. Stahl, Chem. Rev., 2018, 118, 2636.