

# Dalton Transactions

An international journal of inorganic chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. A. Nasrullah, T. Taeufer, J. Pospech, E. Barath and C. Hering-Junghans, *Dalton Trans.*, 2025, DOI: 10.1039/D5DT01495F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## ARTICLE

## FLP-type Alkenylation of a Phosphafluorene


Ayu Afifah Nasrullah,<sup>a,b</sup> Tobias Täufer,<sup>a</sup> Jola Pospech,<sup>a</sup> Eszter Baráth\*,<sup>a</sup> and Christian Hering-Junghans\*,<sup>a</sup>Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Phosphafluorenes, in which the chemically vulnerable C-9 position of the fluorene core has been replaced with phosphorus, demonstrate enhanced stability compared to conventional fluorene-based materials and belong to the general class of dibenzophospholes (DBPs). Phosphafluorene formation is commonly observed in the reductive functionalization of terphenyl-based dihalophosphanes. In here, we outline a high-yielding route towards phosphafluorene **1**, starting from readily available <sup>Dipp</sup>TerPCL<sub>2</sub> (<sup>Dipp</sup>Ter = 2,6-Dipp-C<sub>6</sub>H<sub>3</sub>; Dipp = 2,6-*i*Pr<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>). Different functionalization strategies of **1** are outlined, including a complex with AuCl, its BH<sub>3</sub> adduct, and its corresponding sulphide. Interestingly, facile alkenylation of **1** was achieved in the presence of B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and various phenylacetylene derivatives. This FLP-type alkenylation is *E*-selective giving zwitterionic phosphonium borates.

## Introduction

Phosphanylidene phosphoranes of the general form RP(PR')<sub>3</sub> are also referred to as Phospha-Wittig reagents.<sup>1</sup> Specifically, aryl-substituted variants are commonly used in phosphaalkene synthesis in the so-called phospha-Wittig reaction. The first aryl-substituted phosphanylidene phosphoranes, ArP(PMe<sub>3</sub>), were synthesized by Protasiewicz and co-workers in 1998 through the combination of Ar-PCl<sub>2</sub> with an excess of PMe<sub>3</sub> and zinc powder.<sup>2</sup> To date a limited number of ArP(PMe<sub>3</sub>) (Ar = 2,4,6-*t*Bu<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>, Mes\*,<sup>2</sup> 2,6-(2,6-Cl<sub>2</sub>C<sub>6</sub>H<sub>2</sub>)<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>, <sup>Cl</sup>Ter,<sup>3</sup> 2,6-(2,4,6-Me<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>)<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>, <sup>Mes</sup>Ter,<sup>2, 4</sup> 2,6-(2,4,6-*i*Pr<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>)<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>, <sup>Tipp</sup>Ter,<sup>5</sup> 2,6-(2,6-*i*Pr<sub>2</sub>-C<sub>6</sub>H<sub>2</sub>)<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>, <sup>Dipp</sup>Ter,<sup>6</sup> 1,1,3,3,5,5,7,7-octaethyl-1,2,3,5,6,7-hexahydro-s-indacen-4-yl, EIND<sup>7</sup>) have been reported and their utility in Wittig-type reactions has been explored. Moreover, it was shown that Phospha-Wittig reagents offer a pathway for the generation of arylphosphinidenes, Ar-P, under photochemical conditions. For example, when <sup>Tipp</sup>TerP(PMe<sub>3</sub>) was irradiated at 365 nm the formation of a phosphafluorene was described, through insertion of the phosphinidene into one of the C<sub>Ar</sub>-C<sub>iPr</sub> bonds of the Tipp groups.<sup>5</sup> Our group observed <sup>Dipp</sup>Ter-P generation and deactivation to give phosphafluorene **1** via the thermal decomposition of the isolated phosphagallene, <sup>Dipp</sup>TerPGaCp\* at 50 °C for 2 h (Scheme 1, a), or through the irradiation (396 nm) of <sup>Dipp</sup>TerP(PMe<sub>3</sub>) in toluene-*d*<sub>8</sub> at room temperature (Scheme 1, b).<sup>8</sup>



**Scheme 1.** (a) Thermal phosphinidene release from a phosphagallene, (b) Photochemical PMe<sub>3</sub> release from <sup>Dipp</sup>TerP(PMe<sub>3</sub>); (c,d) Lewis acid induced phosphafluorene formation in terphenyl-substituted halophosphanes; (e,f) Reductive approaches towards phosphafluorenes from dichlorophosphanes (e, f); and summary of this work.

These findings complement earlier work on phosphafluorene formation via the reduction of <sup>Tipp</sup>TerPCL<sub>2</sub> with magnesium (Scheme 1, f).<sup>9</sup> It was later shown that the reducing agent plays an important role in the outcome of the reaction.<sup>10</sup> Related cyclizations of terphenyl-based dichlorophosphanes were also

<sup>a</sup> Leibniz Institut für Katalyse e.V. (LIKAT), A.-Einstein-Str. 29a, 18059 Rostock (Germany). Email: [eszter.barath@catalysis.de](mailto:eszter.barath@catalysis.de), [Christian.hering-junghans@catalysis.de](mailto:Christian.hering-junghans@catalysis.de)

<sup>b</sup> Pusat Persediaan Sains dan Teknologi, Universiti Malaysia Sabah, Jln UMS, 88400, Kota Kinabalu, Sabah (Malaysia).

Supplementary Information available: Synthesis of compounds, NMR spectra and crystallographic details. CCDC 2466772-2466775, 2482916. See DOI: 10.1039/x0xx00000x




## ARTICLE

## Dalton Transactions

described by Marshall<sup>11</sup> in the reduction with potassium (Scheme 1, e), and by Wehmschulte and co-workers in  $\text{AlCl}_3$ -mediated Friedel-Crafts type reactions (Scheme 1, c).<sup>12</sup> Similar intramolecular cyclizations have also been described for terphenyl-substituted phosphonium cations, allowing the isolation of the 9-phospha fluorenium ions (Scheme 1, d).<sup>13</sup> Phosphafluorenes, in which the chemically vulnerable C-9 position of the fluorene is replaced with phosphorus, belong to the broader class of dibenzophospholes (DBPs),<sup>14</sup> which are a family of compounds that have garnered significant interest due to their unique electronic properties and potential applications in materials science and catalysis. DBPs are important building blocks for organic  $\pi$ -conjugated compounds, which have enormous potential as optoelectronic materials for example in light-emitting diodes (OLED), organic field-effect transistors, nonlinear optical devices, and organic solar cells.<sup>15–18</sup> In recent years, monodentate non-symmetrical dibenzophospholes have emerged as ligands in rhodium-catalyzed hydroformylations.<sup>19</sup> Phosphafluorenes, demonstrate enhanced stability compared to conventional fluorene-based materials. They are usually blue-light-emitting materials,<sup>15</sup> and the photophysical properties can be precisely tuned through functionalization, especially at the phosphorus atom. Examples of functionalization strategies on the P center of phosphafluorenes include adduct formation with  $\text{BH}_3$ ,<sup>20</sup> oxidation,<sup>21</sup> sulfurization,<sup>22, 23</sup> complexation with tungsten, gold,<sup>22, 24</sup> iron or ruthenium,<sup>12</sup> and related transformation. DBPs can also be viewed as sterically encumbered phosphines, rendering them suitable candidates for frustrated Lewis pair (FLP) applications when paired with sterically hindered Lewis acids. The functionalization of alkynes using FLPs has been intensively studied.<sup>25</sup> The outcome of the reactions greatly depends on the basicity of the phosphine used. In the case of P/B-FLPs two outcomes are possible: (a) phosphonium alkynylborates, through deprotonation of the alkyne, or (b) addition to the alkynes resulting in zwitterionic alkenyl phosphonium borates, when less basic phosphines were employed.<sup>26–28</sup> In this contribution, we report on different functionalization products of phosphafluorene **1** using  $\text{BH}_3$ ,  $\text{S}_8$  or  $\text{AuCl}$ . Moreover, the reactivity of **1** towards terminal alkynes (4-R- $\text{C}_6\text{H}_4\text{-C}\equiv\text{C-H}$ ) in the presence of  $\text{B}(\text{C}_6\text{F}_5)_3$  was investigated, proving an FLP-type alkenylation strategy for DBPs (Scheme 1, bottom).


## Results and Discussion

Rather than starting from  $\text{DippTerP}=\text{GaCp}^*$  or  $\text{DippTerP}(\text{PMe}_3)$ , phosphafluorene **1** is conveniently prepared through reduction of  $\text{DippTerP}(\text{PCl}_2)$  with two equivalents of activated magnesium ( $\text{Mg}^*$ , also see Scheme 1, f).<sup>9</sup> After work-up, **1** was afforded as light orange, fluffy solid in 77 % yield (Scheme 2, top). The formation of **1** was confirmed by  $^{31}\text{P}$  NMR spectroscopy in  $\text{C}_6\text{D}_6$  ( $\delta(^{31}\text{P}\{^1\text{H}\}) = -1.1$  ppm) and a complex  $^1\text{H}$  NMR spectrum in line with  $\text{C}_1$  symmetry in solution, shows a characteristic upfield-shifted pseudo-triplet ( $\delta(^1\text{H}) = 0.26$  ppm) for one of Me-groups of the P-*i*Pr moiety.



**Scheme 2.** Synthesis of phosphafluorene **1** (top). Functionalization of the phosphorus center in **1**: (i)  $\text{AuCl-SMe}_2$ , toluene, 16 h, r.t.; (ii)  $\text{H}_3\text{B-SMe}_2$ , *n*-hexane, 16 h, r.t.; (iii)  $1/8 \text{ S}_8$ ,  $\text{CH}_2\text{Cl}_2$ , 16 h, r.t.

Next, chemical modifications at the P-center of **1**, particularly through coordination to transition metals, sulfurization, and reactions with Lewis acids were targeted (Scheme 2, bottom). The combination of **1** with  $(\text{Me}_2\text{S})\text{AuCl}$  in toluene afforded complex **2** (Scheme 2, i), which was recrystallized from a 1:5  $\text{CH}_2\text{Cl}_2$ :*n*-hexane solution at  $-30^\circ\text{C}$  giving colourless single crystals suitable for single crystal X-ray diffraction (SCXRD) analysis and **2** was isolated in 50 % yield. The  $^{31}\text{P}\{^1\text{H}\}$  NMR shift in  $\text{C}_6\text{D}_6$  at 32.4 ppm indicated the formation of complex **2** and a tetracoordinate P atom. Upon coordination of  $\text{AuCl}$ , the  $^3J_{\text{P-H}}$  coupling constant of the shielded Me-group ( $\delta(^1\text{H}) = 0.04$  ppm) increases and results in a clear doublet of doublet splitting of the signal. Phosphafluorene  $\text{BH}_3$  adduct **3** (Scheme 2, ii) was obtained when stirring **1** and  $\text{H}_3\text{B-SMe}_2$  in *n*-hexane. After crystallization from *n*-heptane at  $-30^\circ\text{C}$  as colourless single crystalline material in 65% yield. The  $^{31}\text{P}\{^1\text{H}\}$  and  $^{11}\text{B}\{^1\text{H}\}$  NMR spectra of **3** in  $\text{C}_6\text{D}_6$  displayed characteristic shifts at 37.3 ppm and  $-38.4$  ppm, respectively, indicating tetracoordinate P and B atoms. Additionally, the  $\text{BH}_3$  moiety is detected in the IR spectrum, with B-H stretching modes centred at  $2345 \text{ cm}^{-1}$ .<sup>29</sup>



**Figure 1.** Molecular structures of **2** (left), **3** (middle) and **4** (right). Ellipsoids drawn at 50 % with C-H atoms omitted, Dipp- and one of the *i*Pr-groups rendered as wire-frame for clarity. Selected bond lengths [Å] and angles [°] of **2**: C1-P1 1.845(2), C4-P1 1.798(2), C13-P1 1.805(2), P1-Au1 2.2247(6), Cl1-Au1 2.2885(6); C4-P1-C13 91.83(10), C4-P1-C1 106.46(10), C13-P1-C1 105.84(10), C4-P1-Au1 114.61(7), C13-P1-Au1 121.62(7), C1-P1-Au1 113.71(8); C5-C4-P1-C13 179.9(2). **3**: P1-C4 1.7991(13), P1-C13 1.8102(13), P1-C1 1.8435(14), P1-B1 1.9269(16); C4-P1-C13 91.52(6), C4-P1-C1 108.29(6), C13-P1-C1 107.20(6), C4-P1-B1 111.38(7), C13-P1-B1 121.08(6), C1-P1-B1 114.62(7); C13-P1-C4-C9  $-178.84(12)$ ; **4**: P1-C1 1.809(1), P1-C20 1.797(1), P1-C28




1.866(1), P1–S1 1.9462(5); C20–P1–C1 91.78(6), C20–P1–C28 106.39(6), C1–P1–C28 104.83(6), C28–P1–S1 112.84(5); C20–P1–C1–C2 175.58(12).

In the  $^1\text{H}$  NMR spectrum, the expected quartet resonance for  $\text{BH}_3$  is significantly broadened and overlaps with the signals of the *i*Pr-Me groups and could therefore not be unambiguously assigned. Similarly to **2**, the shielded Me-groups of the *i*Pr-groups on the P atom changes into a doublet of doublets.

The reaction of **1** with elemental sulfur cleanly furnished thiophosphafluorene **4** as a white solid in 61 % isolated yield. The  $^{31}\text{P}$  NMR signal of **4** ( $\delta(^{31}\text{P}\{^1\text{H}\}) = 53.4$  ppm;  $\Delta_1(\delta(^{31}\text{P})) = +54.5$  ppm) is notably deshielded compared to **2** and **3**, consistent with P=S bond formation. A similar deshielding is observed in related dibenzophosphapentaphenes upon sulfurization (cf.  $\Delta(\delta(^{31}\text{P})) = +48.9$  ppm).<sup>22</sup> Single-crystal X-ray diffraction experiments (SC-XRD, Figure 1) revealed that **2**, **4** and **3** crystallize in the monoclinic space groups  $P2_1/c$  and  $P2_1/n$ , respectively. **2-4** show a four-coordinate P atom ( $\Sigma(\angle\text{P}) = 304.1(1)$  (**2**), 307.01(6) (**3**), 303.0(6)° (**4**)). The P–C<sub>Ar</sub> distances (**2**: 1.798(2), 1.805(2); **3**: 1.799(1), 1.810(1); **4**: 1.797(1), 1.809(1) Å) are shorter compared to the related free phosphafluorene derived from  $\text{Ti}^{\text{IPr}}\text{P}(\text{Cl})_2$  (cf. 1.805(11), 1.817(4) Å).<sup>9</sup> Similarly, the P–C<sub>iPr</sub> (**2**: 1.845(2); **3**: 1.844(1); **4**: 1.866(1) Å) bonds in **2** and **3** are contracted compared to the related phosphafluorene. The P1–Au1 bond distance (2.2247(6) Å) in **2** aligns with reported gold(I) P-heterocycle complexes,<sup>24, 30</sup> the P–B distance in  $\text{BH}_3$ -adduct **3** (1.9269(16) Å) is in the range of reported dibenzophosphole  $\text{BH}_3$  adducts,<sup>31</sup> and the P–S distance aligns with that in dibenzophosphapentaphene sulfides.<sup>22</sup>

In an effort to identify alternative Lewis acid to  $\text{H}_3\text{B-SMe}_3$ , we explored bulkier derivatives such as  $\text{B}(\text{C}_6\text{F}_5)_3$ . When **1** and  $\text{B}(\text{C}_6\text{F}_5)_3$  were combined in a 1:1 ratio in  $\text{C}_6\text{D}_6$ , no discernible reaction was observed according to  $^{31}\text{P}$  NMR spectroscopy. This lack of reactivity suggests frustrated Lewis pair (FLP) behaviour arising from steric congestion between the Lewis acid and base. Upon addition of one equivalent phenylacetylene to the reaction mixture in  $\text{C}_6\text{D}_6$ , the color of the solution changes from light orange to dark red. NMR data of the reaction mixture showed complete conversion to a new species. Layering of the reaction mixture in  $\text{C}_6\text{D}_6$  with *n*-hexane at ambient temperature and standing overnight afforded a white powder of  $[\text{P}]\text{C}(\text{Ph})=\text{C}(\text{H})\text{B}(\text{C}_6\text{F}_5)_3$  (**5**;  $[\text{P}] = 1-(2,6\text{-diisopropylphenyl})-5,9\text{-diisopropyl-9-phosphafluorene}$ ) and was isolated in 61% yield. Compound **5** features a  $^1\text{H}$  NMR resonance at 7.84 ppm ( $d, ^3J_{\text{P-H}} = 38.0$  Hz) assigned to the alkenyl C=CH moiety (cf.  $\text{Ph}_3\text{P}(\text{PhC}=\text{CH})\text{B}(\text{C}_6\text{F}_5)_3$ :  $\delta(^1\text{H}) = 8.32$  ppm,  $^3J_{\text{P-H}} = 36.0$  Hz).<sup>26</sup> The  $^{31}\text{P}$  NMR signal at 36.1 ppm is in line with a phosphonium center, whereas the  $^{19}\text{F}$  NMR spectrum showed three sets of signals for the *o*-, *p*-, and *m*-fluorine atoms of the  $\text{C}_6\text{F}_5$  groups located at  $\delta(^{19}\text{F})$  –130.5, –161.7, and –166 ppm, respectively, in line with reported FLP-type activations of phenylacetylene and free rotation about the B–C axis.<sup>26</sup> In the  $^{11}\text{B}$  NMR spectrum, a signal at –15.4 ppm ( $d, ^3J_{\text{P-H}} = 18.0$  Hz) is in line with a four-coordinate boron center and a phosphonium borate zwitterionic species. Collectively these data, suggested that **5** is exclusively the 1,2-addition product (Scheme 3, bottom). This

assumption was confirmed by X-ray crystallography, showing that the phosphine and borane added to the alkyne in (E)-fashion with the B adding to the CH terminus of the alkyne (Figure 2, left).<sup>27</sup> The solid-state structure reveals P–C<sub>Ar</sub> (1.792(2), 1.801(2) Å) and P–C<sub>iPr</sub> (1.828(2) Å) distances similar to those found in **2** and **3** (*vide supra*). The P–C<sub>C</sub> distance (1.828(2) Å) is in the range of a contracted P–C single bond ( $\Sigma r_{\text{cov}}(\text{P-C}) = 1.86$  Å)<sup>32</sup> and minimally longer than in  $\text{Ph}_3\text{P}(\text{PhC}=\text{CH})\text{B}(\text{C}_6\text{F}_5)_3$  (cf. 1.806(1) Å).<sup>26</sup> The C31–C32 distance (1.344(2), cf.  $\Sigma r_{\text{cov}}(\text{C}=\text{C}) = 1.34$  Å)<sup>32</sup> clearly indicates activation of the alkyne unit, with the P and B atoms being in *trans*-orientation with respect to the C=C bond. Notably, the C=CH proton is oriented towards the phosphafluorene moiety.



**Scheme 3.** Reactivity of **1** towards  $\text{B}(\text{C}_6\text{F}_5)_3$  in the absence (top) and presence of phenylacetylene derivatives, giving alkenyl phosphonium borates **5-9**.

Motivated by these findings, we next tested a series of *para*-substituted phenylacetylene ( $4\text{-R-C}_6\text{H}_4\text{-C}\equiv\text{CH}$  where  $\text{R} = \text{CH}_3$  (**6**), *t*Bu (**7**), OMe (**8**),  $\text{CF}_3$  (**9**)) to investigate possible substituent effects on this FLP reactivity. Reaction of these alkynes with the **1**/ $\text{B}(\text{C}_6\text{F}_5)_3$  FLP system proceeded analogously to **5**, yielding diverse zwitterionic alkenyl phosphonium borate products (Scheme 3), which underlines the generality of this functionalization strategy.<sup>27</sup> This FLP-type alkenylation of a P-heterocycle is also related to the chemistry of phosphinine-borane adducts towards phenylacetylene.<sup>33</sup> The  $^{31}\text{P}$  NMR shifts for **6-9** are clustered near 36.0 ppm, clearly showing minimal influence of the *para*-substituent. All compounds (**6-9**) displayed  $^{19}\text{F}$  NMR signals for the  $\text{B}(\text{C}_6\text{F}_5)_3$  moiety in the range of –130.5 (*ortho*-F), –161.7 (*para*-F), and –166.0 (*meta*-F), mirroring **5**, and thereby indicating a similar  $\text{B}(\text{C}_6\text{F}_5)_3$  moiety in all alkenylation products **5-9**. Notably, **9** showed an additional singlet at –63.3 ppm in the  $^{19}\text{F}$  NMR spectrum assignable to the






## ARTICLE

## Dalton Transactions

CF<sub>3</sub> group. The <sup>11</sup>B NMR spectra of **6–9** uniformly exhibited resonances near –15.0 ppm (d, <sup>3</sup>J<sub>P–B</sub> = 18 Hz) as expected for borate-type activation products. These observations suggest minimal electronic influence from the *para*-substituents of phenylacetylenes (EDG/EWG) on the regio- and chemoselectivity of the alkenylation. In all derivatives the C=CH resonance is deshielded at ca. 8 ppm and a <sup>3</sup>J<sub>P–H</sub> coupling constant of ca. 38 Hz.




**Figure 2.** Molecular structures of **5** and **9**. Ellipsoids drawn at 50 % with C-H-atoms omitted (except those at C32 (**5**) and C29(**9**)), Dipp-, C<sub>6</sub>F<sub>5</sub>-, CF<sub>3</sub>- and one of the *i*Pr-groups rendered as wire-frame for clarity. Selected bond lengths [Å] and angles [°] of **5**: P1–C4 1.7915(18), P1–C13 1.8014(16), P1–C1 1.8282(18), P1–C31 1.8282(17), B1–C32 1.648(2), C31–C32 1.344(2); C4–P1–C13 93.34(8), C4–P1–C1 110.49(8), C13–P1–C1 113.17(8), C4–P1–C31 107.19(8), C13–P1–C31 112.12(7), C1–P1–C31 117.64(8), C32–C31–P1 112.79(12), C31–C32–B1 138.96(15); C4–P1–C31–C32 65.08(14). **9**: C1–P1 1.8007(14), C12–P1 1.7807(14), C28–P1 1.7969(13), C55–P1 1.8326(14), C28–C29 1.3433(19), C29–B1 1.642(2), C12–P1–C28 108.96(6), C12–P1–C1 93.52(6), C28–P1–C1 122.55(6), C12–P1–C55 108.64(7), C28–P1–C55 113.19(6), C1–P1–C55 107.71(6), C29–C28–P1 116.86(10), C28–C29–B1 136.79(13); C29–C28–P1–C12 137.25(11).

These results underscore the robustness of FLP-mediated alkyne activation, where steric and electronic perturbations at the *para*-position are effectively reduced by the conjugated aromatic system, preserving the core electronic structure of the adduct.


Single crystals of **9** suitable for SC-XRD were obtained from layering the C<sub>6</sub>D<sub>6</sub> reaction mixture with *n*-heptane at room temperature. As previously observed in **5**, the alkyne activation is (*E*)-selective, with boron attached to the CH terminus (Figure 2, right). All key structural parameters are similar to those found in **5**. However, the orientation of the C=CH proton is different, which is not oriented towards the P-*i*Pr group.

Lastly, the absorption and emission profiles of **1** and its derivatives in MeCN solution were investigated. Phosphafluorene **1** shows absorption maxima in the UVB region at 303 and 312 nm and shows weak blue fluorescence with a broad emission maximum at 369 nm. As a representative for P-functionalized species **2–4**, phosphafluorene **4** was studied and shows two bathochromically shifted absorptions maxima at 313 and 323 nm, while the emission maximum is hypsochromically shifted to 361 nm (Figure 3). A second emission maximum is evident at 453 nm. Notably, the emission band is rather broad and tails into the visible region, in line with blue fluorescence in MeCN solution. Interestingly, the absorption spectra of alkenylated species **5–9** are nearly identical with absorption maxima at ca. 330 nm (Figure 4), indicating minimal influence of the *para*-substituent in the employed phenylacetylene

derivatives. The emission spectrum of **5** features a maximum at 381 nm with a shoulder at 400 nm (Figure S66). Broad emission maxima between 380–390 nm are detected for **6**, **7** and **9**. Compound **8** displays distinct photophysical characteristics, featuring a minor absorption shoulder at 385 nm, while the emission maximum is markedly red-shifted to 472 nm. In the absence of lifetime data, we tentatively attribute this pronounced red-shift to phosphorescence rather than fluorescence or a chemical transformation in the excited state. Generally, P functionalization in **4** results in a red-shift of the absorption maximum, which is even more pronounced for the alkenylated species **5–9**. This agrees with previous studies that have clearly shown that P-functionalization will result in bathochromically shifted absorption bands.<sup>15, 22</sup> The absorption and emission spectra of **4** and **8** show no mirror relation, indicating significant structural rearrangements in the excited state. Consistent with previous reports, the Stokes shift in phosphafluorene luminescence is rather large.



**Figure 3.** Normalized absorption- (compact) and emission (dashed) spectra of **1** and **4**.



**Figure 4.** Normalized absorption- (compact) and emission (dashed) spectra of **1** and **5–9**.



## Conclusions

We outline a facile synthetic route towards phosphafluorene **1** starting from readily accessible  $\text{DippTerPCl}_2$  in a reductive approach. Quarternization of **1** with  $\text{AuCl}$ ,  $\text{BH}_3$ , and sulphur afforded P-functionalized species **2-4**, respectively. When **1** was combined with  $\text{B}(\text{C}_6\text{F}_5)_3$  no reaction was observed. When adding phenylacetylene to the mixture, clean FLP-type alkenylation of the P in **1** was noted giving (*E*)-configured alkenyl phosphonium borate **5**. This novel alkenylation strategy for DBPs was extended and facile conversion with *para*-substituted phenylacetylenes was achieved giving species **6-9**. To the best of our knowledge this is the first report on FLP-type functionalization of DBPs and the generality of the concept with respect to the DBP-precursors is currently investigated. It needs to be stated that all compounds in this paper are P-chiral and paths to render these transformations enantioselective will be studied in the future. Furthermore, the photophysical properties of the alkenylation products will be studied in more detail.

## Author contributions

A.A.N. carried out the experimental work, wrote the supporting information and drafted the manuscript. C.H.-J. was responsible for the conceptualization, supervision of the experimental investigations, solved and refined the SCXRD structures. T.T. and J.P. carried out the photophysical characterization. J.P., C.H.-J. and E.B. finalized the manuscript. All authors agreed to the submitted content.

## Conflicts of interest

There are no conflicts to declare.

## Data availability

The data supporting this article have been included as part of the Supplementary Information. This includes: Synthesis and characterization of compounds, NMR spectra, additional spectroscopic data and crystallographic details.

Crystallographic data **2** (2466772), **3** (2466773), **4** (2482916), **5** (2466774) and **9** (2466775) have been deposited at the CCDC and can be obtained from <https://www.ccdc.cam.ac.uk>.

## Acknowledgements

A.A.N. thanks the Universiti Malaysia Sabah for funding. We thank our technical and analytical staff for assistance, especially Dr. Anke Spannenberg (LIKAT) and Dr. Jonas Surkau (University of Rostock) for their support regarding X-ray analysis.

## Notes and references

View Article Online  
DOI: 10.1039/D5DT01495F

- J. D. Protasiewicz, *Eur. J. Inorg. Chem.*, 2012, **2012**, 4539-4549.
- S. Shah and J. D. Protasiewicz, *Chem. Commun.*, 1998, 1585-1586.
- R. C. Smith, T. Ren and J. D. Protasiewicz, *Eur. J. Inorg. Chem.*, 2002, **2002**, 2779-2783.
- S. Shah, G. P. A. Yap and J. D. Protasiewicz, *J. Organomet. Chem.*, 2000, **608**, 12-20.
- S. Shah, M. C. Simpson, R. C. Smith and J. D. Protasiewicz, *J. Am. Chem. Soc.*, 2001, **123**, 6925-6926.
- P. Gupta, J.-E. Siewert, T. Wellnitz, M. Fischer, W. Baumann, T. Beweries and C. Hering-Junghans, *Dalton Trans.*, 2021, **50**, 1838-1844.
- K. Takeuchi, H.-o. Taguchi, I. Tanigawa, S. Tsujimoto, T. Matsuo, H. Tanaka, K. Yoshizawa and F. Ozawa, *Angew. Chem. Int. Ed.*, 2016, **55**, 15347-15350.
- T. Taeufer, F. Dankert, D. Michalik, J. Pospech, J. Bresien and C. Hering-Junghans, *Chem. Sci.*, 2023, **14**, 3018-3023.
- B. Twamley, C. D. Soffield, M. M. Olmstead and P. P. Power, *J. Am. Chem. Soc.*, 1999, **121**, 3357-3367.
- R. C. Smith, S. Shah and J. D. Protasiewicz, *J. Organomet. Chem.*, 2002, **646**, 255-261.
- A. S. Ionkin and W. J. Marshall, *Heteroatom Chem.*, 2003, **14**, 360-364.
- A. A. Diaz, J. D. Young, M. A. Khan and R. J. Wehmschulte, *Inorg. Chem.*, 2006, **45**, 5568-5575.
- M. Olaru, D. Duvinage, Y. Naß, L. A. Malaspina, S. Mebs and J. Beckmann, *Angew. Chem. Int. Ed.*, 2020, **59**, 14414-14417.
- P. Hibner-Kulicka, J. A. Joule, J. Skalik and P. Bałczewski, *RSC Advances*, 2017, **7**, 9194-9236.
- J. Yin, R.-F. Chen, S.-L. Zhang, Q.-D. Ling and W. Huang, *J. Phys. Chem. A*, 2010, **114**, 3655-3667.
- H.-C. Su, O. Fadhel, C.-J. Yang, T.-Y. Cho, C. Fave, M. Hissler, C.-C. Wu and R. Réau, *J. Am. Chem. Soc.*, 2006, **128**, 983-995.
- C.-H. Lin, C.-W. Hsu, J.-L. Liao, Y.-M. Cheng, Y. Chi, T.-Y. Lin, M.-W. Chung, P.-T. Chou, G.-H. Lee, C.-H. Chang, C.-Y. Shih and C.-L. Ho, *J. Mat. Chem.*, 2012, **22**, 10684-10694.
- X. Deng, S. Liu, Y. Sun, D. Zhong, D. Jia, X. Yang, B. Su, Y. Sun, G. Zhou, B. Jiao and Z. Wu, *Dyes and Pigments*, 2023, **209**, 110885.
- A. Oukhrib, L. Bonnafoux, A. Panossian, S. Waifang, D. H. Nguyen, M. Urrutigoity, F. Colobert, M. Gouygou and F. R. Leroux, *Tetrahedron*, 2014, **70**, 1431-1436.
- K. V. Rajendran and D. G. Gilheany, *Chemical Commun.*, 2012, **48**, 817-819.
- H. A. van Kalker, S. H. A. M. Leenders, C. R. A. Hommersom, F. P. J. T. Rutjes and F. L. van Delft, *Chem. Eur. J.*, 2011, **17**, 11290-11295.
- P.-A. Bouit, A. Escande, R. Szűcs, D. Szieberth, C. Lescop, L. Nyulászi, M. Hissler and R. Réau, *J. Am. Chem. Soc.*, 2012, **134**, 6524-6527.
- J. A. Macor, J. L. Brown, J. N. Cross, S. R. Daly, A. J. Gaunt, G. S. Girolami, M. T. Janicke, S. A. Kozimor, M. P. Neu, A. C. Olson, S. D. Reilly and B. L. Scott, *Dalton Trans.*, 2015, **44**, 18923-18936.
- R. Inaba, K. Oka, T. Iwami, Y. Miyake, K. Tajima, H. Imoto and K. Naka, *Inorg. Chem.*, 2022, **61**, 7318-7326.
- J. Guo, M. Yan and D. W. Stephan, *Org. Chem. Front.*, 2024, **11**, 2375-2396.
- M. A. Dureen and D. W. Stephan, *J. Am. Chem. Soc.*, 2009, **131**, 8396-8397.
- M. A. Dureen, C. C. Brown and D. W. Stephan, *Organometallics*, 2010, **29**, 6594-6607.
- C. Jiang, O. Blacque and H. Berke, *Organometallics*, 2010, **29**, 125-133.



## ARTICLE

## Dalton Transactions

- 29 C. A. Jaska, H. Dorn, A. J. Lough and I. Manners, *Chem. Eur. J.*, 2003, **9**, 271-281.
- 30 S. Attar, W. H. Bearden, N. W. Alcock, E. C. Alyea and J. H. Nelson, *Inorg. Chem.*, 1990, **29**, 425-433.
- 31 V. Diemer, A. Berthelot, J. Bayardon, S. Jugé, F. R. Leroux and F. Colobert, *J. Org. Chem.*, 2012, **77**, 6117-6127.
- 32 P. Pyykkö and M. Atsumi, *Chem. Eur. J.*, 2009, **15**, 12770-12779.
- 33 J. Lin, F. Wossidlo., N. T. Coles, M. Weber, S. Steinhauer, T. Böttcher and C. Müller, *Chem. Eur. J.*, 2022, **28**, e202104135.

View Article Online  
DOI: 10.1039/D5DT01495F



**Data Availability Statement**View Article Online  
DOI: 10.1039/D5DT01495F

The data supporting this article have been included as part of the Supplementary Information. This includes: Synthesis and characterization of compounds, NMR spectra, additional spectroscopic data and crystallographic details.

Crystallographic data 2 (2466772), 3 (2466773), 4 (2482916), 5 (2466774) and 9 (2466775) have been deposited at the CCDC and can be obtained from <https://www.ccdc.cam.ac.uk>.

