




Cite this: *RSC Adv.*, 2018, 8, 41422

Effects of methyl jasmonate and melatonin treatments on the sensory quality and bioactive compounds of harvested broccoli

Feng Luo, Jia-Hui Cai, Xuan Zhang, Dong-Bing Tao, Xin Zhou, Qian Zhou, Ying-Bo Zhao, Bao-Dong Wei, Shun-Chang Cheng and Shu-Juan Ji *

Harvested broccoli is prone to decline in quality with regard to its appearance and nutrition. In this study, freshly harvested broccoli was treated with methyl jasmonate (MeJA) and melatonin (MT) and stored at 20 °C and the changes in sensory qualities and bioactive compounds were analyzed. The control samples began yellowing on day 2, whereas MeJA and MT treatments delayed the yellowing by 2 and 4 days, respectively. Upon yellowing, sweetness and bitterness of control samples increased sharply, accompanied by the accumulation of bioactive compounds, except for sulforaphane; however, no significant change in volatile components was detected. When the samples started losing their green color, MeJA alleviated the bitterness while increasing the sweetness and sulforaphane content. The bitterness, astringency, umami level, and the content of sulfurous volatiles improved significantly in the MT-treated samples. Moreover, these samples showed high antioxidant activity; the protective effect on V_C and carotenoids was extremely significant.

Received 26th September 2018

Accepted 29th November 2018

DOI: 10.1039/c8ra07982j

rsc.li/rsc-advances

1 Introduction

Vegetables are important components of diet, supplying a multitude of nutrients required for health. Broccoli (*Brassica oleracea* L. var. *italica*) has been the focus of much research¹ because of its potential to provide health-promoting phytochemicals in the diet. Broccoli heads are fresh and green when they are harvested, but are prone to deterioration, which results in the yellowing of the florets and a rapid decrease in their nutritional content upon storage. Moreover, on deterioration, the flavor and aroma of broccoli can become unacceptable to customers. To improve the post-harvest quality of broccoli, treatment with a variety of exogenous compounds, such as 1-methylcyclopropene, sugars, ethanol, salicylic acid, and *N*-benzylaminopurine, has been investigated.^{2,3}

Methyl jasmonate (MeJA) is a plant hormone and functions as a signaling molecule.⁴ It is also an important fragrance component of jasmine flowers. Exogenous application of MeJA can promote effective accumulation of nutrients in fruits and vegetables by induction of hormone signal transduction in cells.⁵ Kang *et al.*⁶ reported that MeJA treatment could significantly increase glucosinolate biosynthesis in broccoli florets. Recently, researchers found that *N*-acetyl-5-methoxytryptamine (melatonin, MT) was effective in removing free radicals from cells.⁷ Several metabolites formed by the interaction of MT with

free radicals were also found to be highly effective scavengers of the free radicals.⁸ Besides, MT also stimulated the activity of protective antioxidant enzymes (glutathione reductase, glutathione peroxidase, and superoxide dismutase). Under extreme stress conditions, MT could effectively restore the germination of cucumber seeds that has been exposed to chilling injury.⁹ Moreover, melatonin treatment effectively increased the tolerance of cabbage seeds to toxic copper ion stress, increased the herbicidal ability of rice and alleviated the germination of *Pennisetum alopecuroides* L. Spreng seeds under NaCl stress.^{9,10} However, the use of MT as an effective treatment against oxidation has rarely been reported in studies on the postharvest quality of broccoli.

Sensory quality, which includes color, taste, aroma, and tissue morphology, is the most direct index used to measure the quality of fruits and vegetables. Aroma is determined by a complex mixture of volatile components emitted by a product, which are perceived by the taste receptor; it might be decided by the off-odor produced by deterioration of the product quality. Usually, aroma is analyzed by gas chromatography-mass spectrometry (GC-MS). The taste, which could be sour, salty, sweet, bitter, astringent, or umami, is perceived by the tongue in mammals. Electronic tongues, developed recently, can simulate the taste receptors of mammals, and can reflect the taste of products and help in determining the effects of different treatments on the taste of products.¹¹

Broccoli heads are made up of dense buds. The newly harvested broccoli is green, with the fragrance of fresh vegetables and the unique flavor of cruciferous vegetables. At room

Department of Food Science, Shenyang Agricultural University, No. 120 Dongling Road, Shenyang 110866, PR China. E-mail: jsjsyau@yeah.net; jsjsyau@sina.com; 403880675@qq.com; Fax: +86 (0) 24 88498337; Tel: +86 (0) 24 88498337



temperature, the color of broccoli gradually turns yellow during the shelf life and its aroma and taste also change significantly. In severe cases, it produces off-odor, which seriously affects its commodity value. There have only been few reports on the flavor and volatile components of broccoli. Lv *et al.*¹² surmised that Se treatment can have a positive effect in maintaining the quality of broccoli and in enhancing its sensory quality through the release of volatile compounds.

The beneficial roles of broccoli in the prevention of cancer, cardiovascular disease, tumor, and senescence are well-recognized, and are associated with the health-promoting effects of the bioactive compounds present in it.¹³ The main bioactive compounds of broccoli are flavonoids, vitamin C, carotenoids, and sulforaphane.¹⁴ The nutrient content and antioxidant activity were extremely important during post-harvest broccoli, which was susceptible to yellowing after harvest, and has a negative impact on its nutritional and commercial value.¹⁵ UV, steam and ethanol treatments effectively inhibited the reduction of antioxidant activity in post-harvest broccoli.^{16–18} Glucosinolates are unique secondary metabolites of cruciferous plants, which can produce a series of important biologically active substances upon hydrolysis by myrosinase, of which, sulforaphane is an effective anticancer compound.¹⁹ Among the vegetables of the Brassica family, broccoli is the ideal material for obtaining sulforaphane and, therefore, its consumption is increasing.

The causes and mechanisms of yellowing of broccoli have been investigated in a number of studies; however, research on the changes in sensory quality and bioactive compounds of broccoli that accompany the yellowing process remains scant. In the present study, we determined the changes in the sensory quality and bioactive compounds accompanying the yellowing of untreated broccoli during the shelf life. We also investigated the effects of MeJA and MT treatment on the above-mentioned parameters of broccoli during its shelf life after harvest.

2 Materials and methods

2.1 Plant material and treatments

'Naihan-Youxiu' broccoli heads were harvested at a commercial field in Shenyang, Liaoning Province, China. The standard for broccoli harvest was tight florets with buds of uniform size and color. Two or three leaves were left intact at the base of the ballflower. The harvested products were placed in plastic boxes, with head pointing upwards and no stacking, and were immediately transported to the laboratory where they were selected based on similar maturity and size and absence of physical damage and pests. The samples were divided into three groups (control, MeJA, and MT samples). The samples in the three groups were treated as follows: 10 freshly broccoli heads were used for the analysis on the day of harvest; 60 broccoli heads were sprayed evenly with distilled water; 600 μL MeJA (Sigma-Aldrich, USA) standard (95%) was dissolved in distilled water with the addition of 1% ethanol solution. Then, the MeJA solution ($0.5 \mu\text{mol L}^{-1}$) was used for spraying on the 60 broccoli heads; 60 samples were sprayed with the aqueous solution of melatonin (MT) (Sigma-Aldrich, USA) ($100 \mu\text{mol L}^{-1}$). All the

broccoli heads were dried uniformly after the treatments, and were subsequently used for the storage experiments.

The samples were packed in polyethylene bags (thickness: 0.03 mm), with ten heads per bag, and the bags were sealed by folding. Finally, the samples were stored at $20 \text{ }^\circ\text{C}$ under 78–80% relative humidity. The test cycle was divided into two phases. In the first stage (day 0–4), the samples were tested once daily, whereas during the second stage (day 4–8), the samples were tested at intervals of two days. Besides, the samples of all treatments were taken on sampling time until the broccoli completely yellowed. All the measurements were performed restoring the samples to the room temperature. 3 replicates of 6 broccoli heads were taken for each sampling date and treatment, and the whole experiment was repeated three times.

2.2 Analysis of the volatile compounds

The methods described by Lv *et al.*¹⁴ were used for the extraction and determination of the broccoli volatiles. A sample of broccoli florets (3.0 g) was placed in a vial with 20 mL headspace and 2.5 g NaCl and 10 μL 2-nonanone internal standard (0.25 mg mL^{-1}) were added; the vial was sealed with a Teflon septum aluminum cap and the contents were mixed. For manual solid-phase microextraction (SPME) analysis, the samples were equilibrated at $50 \text{ }^\circ\text{C}$ for 40 min and then exposed to a fiber coated with 65 μm of polydimethylsiloxane (Supelco Co., Bellefonte, PDMS) at $50 \text{ }^\circ\text{C}$ for 40 min. Subsequently, the volatiles were desorbed over 180 s at $250 \text{ }^\circ\text{C}$ into the splitless injection port. The volatiles were analyzed with a GC-MS apparatus (7890A-5975C, Agilent Technologies, USA) equipped with HP-INNOWAX ($3 \times 10^4 \text{ mm length} \times 0.25 \text{ mm i.d.} \times 2.5 \times 10^{-4} \text{ mm film thickness}$) fused silica capillary column. The heating program was as follows: the injector was maintained at $250 \text{ }^\circ\text{C}$; the initial oven temperature was $40 \text{ }^\circ\text{C}$, which was held for 3 min; the temperature then increased to $120 \text{ }^\circ\text{C}$ at $5 \text{ }^\circ\text{C min}^{-1}$, and was subsequently raised to $200 \text{ }^\circ\text{C}$ at $10 \text{ }^\circ\text{C min}^{-1}$ and held for 5 min. The MS conditions were as follows: the temperature of the electron ionization source was maintained at $200 \text{ }^\circ\text{C}$ and the mass scanning was done in the 35 to 500 m/z range. The flow rate of helium was 1.0 mL min^{-1} . The NIST 11 spectra library was used to identify the volatile compounds. By comparing the linear retention indices and the EI mass spectra with those of the reference compounds, the identities of most of the compounds were confirmed. The compounds were quantified using the internal standard method, where the concentrations of various volatile components were normalized to that of 2-nonanone.

2.3 Electronic tongue measurement

The gustatory sensation during the storage process of broccoli in the different treatment groups was determined by using the electronic tongue (SA402B, INSENT, Japan). The first step was the self-examination and preparation of the electronic tongue system until the sensors (ZZ, BA, BB, CA, GA, HA, and JB) were rinsed with distilled water for several minutes to obtain constant readings using water and a reference electrode (Ag/AgCl 3 M KCl) (Metrohm AG), immersed in the liquid sample.



The tissue of broccoli florets (50 g) was ground and filtered through six layers of gauze; three biological replications were performed. Finally, 100 mL filtrate was injected into the measuring cup of the automatic sampler to start the experiment. The test parameters were set three times, and the sourness, saltiness, sweetness, bitterness, astringency, and umami tastes were tested.

2.4 Determination of the content of bioactive compounds

2.4.1 Determination of flavonoid content. Flavonoids were extracted from the broccoli. The 2.0 g tissue (florets) was triturated with 70% ethanol and placed in a water bath at 85 °C for 1.5 h.²⁰ And then the total flavonoid content was determined by the spectrophotometric method. 2 mL extract was mixed with 0.6 mL of 5% NaNO₂ solution and incubated for 5 min avoid light. And then the mixture was added with 0.6 mL 10% Al(NO₃)₃ for incubating 6 min in the dark. Add 4 mL of 4% NaOH solution to the mixture, dilute to 10 mL with deionized water, and place in the dark for 20 min. Finally, the absorbance was measured by using spectrophotometer at 510 nm.

The standard solution of quercetin in different concentrations is obtained by repeating the same procedure. The quercetin standard was obtained from DingGuo Changsheng Biotechnol CO. LTD (Beijing). The flavonoid content was calculated using the following calibration curve: $Y = -0.0134X + 12.188$, $R = 0.9968$, where X is the flavonoid content in mg g⁻¹ and Y is the absorbance. The total flavonoid content was expressed as mg of quercetin equivalent/g of fresh weight (FW). Three biological replicates were used for each treatment, and the experiments were performed three times.

2.4.2 Determination of V_C content. The content of V_C was determined as described in a previous report²¹ and analyzed by high performance liquid chromatography-mass spectrometer (HPLC-MS). The ice-cold extraction solution consisted of 2 mmol L⁻¹ disodium-dihydrogen-EDTA and 6% meta-phosphoric acid. 5 g broccoli buds tissue was triturated with 5 mL ice-cold extraction solution, and the additional 5 mL solution was added in the extract. And then, the extract was homogenized for 2 min in the absence of light. The homogenate was filtered twice by filter paper, and the residue was washed with 5 mL ice-cold extraction solution. Then, the total homogenate was filtered again. Finally, determination solution consisted of 100 μL filtrate and 400 μL 0.87% ortho-phosphoric acid solution containing disodium-dihydrogen-EDTA (1 mmol L⁻¹) and Trizma Base (100 mmol L⁻¹), which was used for the determination of V_C content.

The HPLC-MS analysis was carried out using the LCMS-8050 spectrometer (Shimadzu, Kyoto, Japan). The flow rate was set at 1.0 mL min⁻¹, and the injection volume was 10 μL. Separation was performed on a 2 μm Slum-pack GIS C18 column (75 mm × 2.1 mm; Shimadzu, Kyoto, Japan) at 25 °C. The mobile-phase solvent system used was acetonitrile (2%): NaH₂PO₄ (50 mmol L⁻¹): dodecyltrimethyl ammonium chloride (2.5 mmol L⁻¹): disodiumdihydrogen-EDTA (1.25 mmol L⁻¹) (39/13/13/13, by volume), and isocratic flow terminated each run. V_C standards were obtained from DingGuo Changsheng Biotechnol CO.LTD

(Beijing, China). The quantification of V_C was carried out by the standard calibration method and the results were expressed as mg of ascorbic acid/100 g of fresh weight (FW). The peak areas for V_C were evaluated in triplicate for each treatment.

2.4.3 Extraction of carotenoids and HPLC-MS analysis. Carotenoids were extracted from broccoli florets as described previously and were analyzed by HPLC.²² The broccoli bud tissue was triturated with methanol/chloroform/water (1 : 2 : 1, v/v/v), and the extract was centrifuged at 8000 × g for 15 min. The aqueous phases were extracted with chloroform several times until the residue was colorless. The residue was dried under a gentle stream of nitrogen. The reaction was terminated by addition of 6% (w/v) KOH (in methanol) and incubation in a water bath at 60 °C for 30 min in the absence of light. The dried residues were re-suspended with the nitrogen stream and applied to the HPLC-MS.

The HPLC-MS analysis was carried out using the liquid chromatograph mass spectrometer (LCMS-8050, Shimadzu, Kyoto, Japan). The injection volume was 10 μL, and flow rate was set to 0.2 mL min⁻¹. Separation was performed on a 2 μm × 75 mm × 2.1 mm column (Slum-pack GIS C18; Shimadzu, Kyoto, Japan) and operated at 40 °C. The mobile-phase solvent system used was acetonitrile: methanol (0.1 M ammonium formate): dichloromethane: water (65/20/6/9, by volume), and isocratic flow terminated each run. After equilibrating the column, the carotenoids contained (zeaxanthin, β-cryptoxanthin and β-carotene) in the samples were separated and quantified using the standard calibration curve. The standards were obtained from DingGuo Changsheng Biotechnol CO. LTD. (Beijing, China). Three biological replicates were used for each treatment, and the experiments were performed three times. The sum of zeaxanthin, β-cryptoxanthin and β-carotene content was taken as the carotenoids content in broccoli.

2.4.4 Determination of sulforaphane content. Sulforaphane was extracted according to the method described in a previous report.²³ The floret tissue (0.2 g) was milled in liquid nitrogen using a pre-cooled mortar and mixed with 2 mL of 0.5 U mL⁻¹ myrosinase solution. Acetonitrile (1 mL) was added to the liquid for dissolving the residue and the homogenate was passed through a PTFE (0.45 μm) membrane syringe filter before injecting into the HPLC. The samples were separated at 30 °C on a Kromasil C18 column (250 × 4.6 mm) using acetonitrile and water at a flow rate of 1.0 mL min⁻¹. Sulforaphane (Sigma) was used as an external standard for HPLC analysis. The absorbance was measured at 254 nm.

2.4.5 Measurement of antioxidant activity. The antioxidant activity was measured in terms of DPPH radical scavenging activity, which was determined using the modified described by Hatano *et al.*²⁴ after some modification. The DPPH standard was dissolved in anhydrous ethanol. The initial concentration of DPPH was 10 × 10⁻⁵ mol L⁻¹, which was diluted into solution of five different concentrations. A calibration curve was drawn by measuring the absorbance of the standard solutions at 765 nm using a UV spectrophotometer. The DPPH standard (0.2 g) was dissolved in anhydrous ethanol and a 2 × 10⁻⁴ mol L⁻¹ DPPH diluent was prepared. Tissue from broccoli buds (5.0 g) was milled in 50% ethanol solution (50 mL) and the extract was



centrifuged at $8000 \times g$ for 10 min. The following reaction fluid was added to the extract: A (2 mL DPPH and 2 mL anhydrous alcohol), A₀ (anhydrous alcohol), B (2 mL DPPH and 2 mL sample solution), and B₀ (2 mL anhydrous alcohol and 2 mL sample solution); the absorbance was measured at 765 nm. Finally, the absorbance value was converted to the molar concentration of DPPH according to the standard curve. The results were expressed as milligram of gallic acid equivalent per gram fresh weight. The formula used for the calculation of DPPH radical scavenging activity was as follows:

$$C_1 = (1 - C_{\text{DPPH}}/C_0) \times 100\%$$

C_1 (%) is the DPPH radical scavenging active, C_{DPPH} (mol L^{-1}) is the absorbance of the sample, and C_0 (mol L^{-1}) is the initial concentration of DPPH.

2.5 Data analysis

The results of electronic tongue measurements were evaluated by multivariate pattern recognition algorithms with the help of Statistica v. 9.0 software (StatSoft, Inc., Tulsa, OK). Data are presented as means \pm standard deviation of three replicates and were analyzed using SPSS v.20 software (SPSS Inc., Chicago, IL, USA). Differences among the groups were evaluated by one-way analysis of variance, and the relationship between the indices was analyzed by Pearson's correlation. The figures were prepared using Prism 5.0 software (Graph Pad Inc., La Jolla, CA, USA).

3 Results

3.1 Effect of different treatments on the color of broccoli

The color change is an important sensory index for broccoli that intuitively reflects the effect of post-harvest handling. As shown in Fig. 1, broccoli heads began to turn yellow on the second day of harvest, when stored at 20 °C without any treatment. In contrast, the color of broccoli heads treated with MeJA and MT barely showed any change. On the fourth day of the shelf life, the control samples were so severely yellowed that they had no commercial value, whereas the MeJA-treated samples showed slight signs of yellowing and the color of broccoli heads treated with MT remained bright green. Compared to that in the control samples, in the MeJA and MT treated samples, the yellowing process of broccoli heads was effectively delayed by 2 and 4 days, respectively. Based on subsequent studies, it was clear that the control, MeJA, and MT samples turned yellow on day 2, 4, and 6, respectively.

3.2 Effects of different treatments on the volatile components

As shown in Table 1, only three kinds of sulfurous volatile compounds were detected in fresh samples, namely hexanal, (*E*)-2-hexenal, and low levels of volatile alcohols and esters. Compared to the fresh samples, the sulfides were unchanged when the control and MeJA samples turned yellow, whereas those in the MT-treated product increased from three to five. The




Fig. 1 Color changes in broccoli on day 2, 4, 6, and 8 in the different treatment groups, viz. control, 3 methyl jasmonate (MeJA), and melatonin (MT).

content of sulfides in the control was 13.2% lower than that in the fresh samples, whereas that in the MeJA and MT-treated samples was significantly decreased and increased ($P < 0.05$) by 72.9% and 40.9%, respectively, compared to their content in the control samples. Moreover, no significant difference in the composition of sulfide compounds was found between the control and fresh samples. In the MeJA-treated samples, dimethyl disulfide and dimethyl trisulfide were the main components whose contents were changed with respect to those in the control samples, whereas methyl disulfide was a unique sulfide in the MT-treated samples when broccoli began to turn yellow.

In addition to sulfides, nine kinds of esters were detected in broccoli, including ethenyl hexanoate, (*Z*)-3-hexen-1-ol acetate, ethyl acetate, (*Z*)-3-hexen-1-ol propanoate, *cis*-3-hexenyl iso-butylate, isothiocyanates, butyl isothiocyanate, *N*-valeric acid *cis*-3-hexenyl ester, and 2-methylbutyl isothiocyanate. Six (control), five (MeJA), and three (MT) different kinds of volatile esters were detected on the day of turning yellow, while the day of harvesting was five. Unlike for the fruity plant, the abundance of esters in broccoli was not high, although broccoli had a rich variety of esters. When broccoli buds turned yellow from green, there was almost no difference in the kind and content of volatile esters ($P < 0.05$) in the control samples. However, compared to the control samples, the MT treatment caused a 12.8% drop in the content of volatile esters, whereas no significant change ($P < 0.05$) was observed in the MeJA-treated samples. *Cis*-3-hexenyl iso-butylate only existed in the fresh samples, whereas 2-methylbutyl isothiocyanate was detected in



Table 1 Changes of the volatile compounds in control, methyl jasmonate (MeJA), and melatonin (MT) samples on the day of harvesting and turning yellow^a

Volatile compounds ($\mu\text{g kg}^{-1}$)	RT (min)	RI		Fresh samples	Control	MeJA	MT
		LRI	RE				
Methanethiol	1.44	—	444	0.28 \pm 0.2c	0.12 \pm 0.1d	0.55 \pm 0.07b	1.52 \pm 0.05a
Dimethyl disulfide	1.65	—	508	2.20 \pm 0.6a	1.66 \pm 0.4a	1.21 \pm 0.09c	7.66 \pm 0.4a
Trichloromethane	2.45	—	601	ND	ND	0.18 \pm 0.04a	0.09 \pm 0.03b
1-Penten-3-ol	2.76	—	677	ND	0.04 \pm 0.01 ns	0.06 \pm 0.02 ns	ND
2-Ethyl-furan	3.28	698	690	0.05 \pm 0.02 ns	0.02 \pm 0.01 ns	ND	ND
3-Pentanol	3.56	702	688	ND	ND	ND	0.05 \pm 0.02 ns
3-Methyl-1-butanol	3.77	739	739	ND	ND	0.15 \pm 0.03 ns	0.10 \pm 0.05 ns
2-Methyl-butanoic acid	4.21	762	756	0.23 \pm 0.01a	0.11 \pm 0.02b	0.28 \pm 0.04a	ND
Hydrogen sulfide	4.42	783	771	ND	ND	0.09 \pm 0.01c	0.22 \pm 0.01a
Hexanal	5.66	784	788	0.45 \pm 0.01a	0.28 \pm 0.09b	0.51 \pm 0.2a	ND
Dimethyl sulfide	6.26	814	805	0.38 \pm 0.04a	0.22 \pm 0.01b	0.32 \pm 0.3a	0.42 \pm 0.2a
(E)-2-Hexenal	6.59	841	853	3.26 \pm 0.2b	3.08 \pm 0.3b	5.08 \pm 0.6a	1.37 \pm 0.4c
2-Hexenal	6.81	859	860	1.2 \pm 0.3b	0.8 \pm 0.4b	1.2 \pm 0.2b	3.3 \pm 0.16a
Cis-3-Hexen-1-ol	7.32	862	869	0.40 \pm 0.01a	0.22 \pm 0.1c	0.21 \pm 0.06c	0.22 \pm 0.07c
(Z)-2-Hexen-1-ol	7.54	874	868	ND	1.89 \pm 0.24a	ND	0.45 \pm 0.11c
1-Hexanol	7.82	876	860	0.31 \pm 0.07b	0.25 \pm 0.11b	ND	ND
Methoxy-phenyl-oxime	8.44	881	864	2.04 \pm 0.5	1.88 \pm 0.32b	1.26 \pm 0.25b	2.75 \pm 0.56a
Dimethyl trisulfide	9.26	973	972	0.09 \pm 0.01 ns	0.07 \pm 0.02 ns	ND	0.08 \pm 0.02 ns
2-Methyl-3-octanone	11.41	986	984	1.21 \pm 0.04a	1.24 \pm 0.1a	1.44 \pm 0.4a	1.66 \pm 0.1a
Ethyl hexanoate	11.32	991	974	0.11 \pm 0.01 ns	0.06 \pm 0.02 ns	ND	0.02 \pm 0.01 ns
Ethanol	11.54	1002	982	1.9 \pm 0.3d	5.6 \pm 0.1a	6.5 \pm 0.2a	3.5 \pm 0.4c
(Z)-3-Hexen-1-ol acetate	11.87	1014	1010	2.12 \pm 0.02 ns	2.08 \pm 0.04 ns	2.11 \pm 0.1 ns	ND
Ethyl acetate	12.46	1021	1008	ND	0.13 \pm 0.02b	0.29 \pm 0.04a	ND
(Z)-3-Hexen-1-ol propanoate	13.27	1108	1091	ND	ND	0.13 \pm 0.04a	ND
Nonanal	14.32	1112	1105	0.18 \pm 0.02a	0.13 \pm 0.15b	0.11 \pm 0.01b	0.15 \pm 0.03a
Methyl disulfide	15.55	1134	1106	ND	ND	ND	0.05 \pm 0.01 ns
Cis-3-hexenyl iso-butyrate	16.01	1151	1145	0.22 \pm 0.01 ns	ND	ND	ND
1,2-Dimethoxy-benzene	16.47	1156	1126	ND	ND	ND	ND
Isothiocyanates	16.72	1162	1134	0.23 \pm 0.04a	0.11 \pm 0.02b	ND	0.31 \pm 0.04a
Naphthalene	17.05	1189	1185	ND	0.02 \pm 0.01 ns	ND	0.05 \pm 0.02 ns
Butyl isothiocyanate	17.63	1221	1202	ND	0.03 \pm 0.02b	ND	ND
N-Valeric acid cis-3-hexenyl ester	18.54	1246	1243	0.08 \pm 0.03b	ND	0.22 \pm 0.01a	ND
Tridecane	19.24	1282	1300	0.10 \pm 0.03 ns	0.12 \pm 0.04 ns	ND	ND
1-Tridecene	20.82	1299	1304	0.21 \pm 0.02a	0.23 \pm 0.06a	0.27 \pm 0.03a	ND
2-Methylbutyl isothiocyanate	21.32	1372	1351	ND	0.16 \pm 0.01b	0.31 \pm 0.02a	0.26 \pm 0.02a
Tetradecane	22.55	1409	1400	ND	0.03 \pm 0.04 ns	0.08 \pm 0.05 ns	0.05 \pm 0.01 ns
Trans- β -ionone	24.26	1501	1489	ND	ND	0.12 \pm 10.01a	0.02 \pm 0.01b
Hexadecane	25.44	1612	1600	0.13 \pm 0.01a	0.14 \pm 0.02a	0.18 \pm 0.01a	0.03 \pm 0.01b
Heptadecane	26.87	1713	1702	0.1 \pm 0.05a	0.12 \pm 0.04a	0.15 \pm 0.16a	0.18 \pm 0.06a
Total content							
Hydrocarbon				0.33 ns	0.41 ns	0.43 ns	0.35 ns
Alcohols				2.89b	8.12a	7.47a	5.79a
Esters				2.76a	2.41a	3.06a	0.59b
Sulfides				6.89b	5.98b	1.62c	8.43a
Acids				0.23a	0.11b	0.28a	ND
Others				8.24a	7.55a	3.6b	7.15a

^a RT and RI are the retention time and retention index of the volatile compounds, respectively. RE is the abbreviation for reference. The analyses were identified by comparison with the NIST 11 database. The symbol a, b, c, and d show significant differences according to the independent sample *t*-test ($P < 0.05$) for each volatile compound. 'ns' indicates no significant difference. ND indicates that the volatile compound was not detected. '—' indicates that the retention time of this volatile compound is before the retention time of C6 *n*-alkanes.

all the yellowed samples. Butyl isothiocyanate and (*Z*)-3-hexen-1-ol acetate were the unique volatile esters in the control and MT samples, respectively. In the MeJA samples, (*Z*)-3-hexen-1-ol propanoate was a unique volatile ester component. Furthermore, we found that the different treatments had little effect on the contents of hydrocarbons, alcohols, and aldehydes in broccoli.

3.3 Effect of different treatments on the taste

As shown in Table 2, when the control samples began to turn yellow, the level of sweetness and bitterness increased by 446.1% and 349.4%, respectively, whereas there was no significant difference ($P < 0.05$) in the other flavors. When the samples turned yellow from green, compared to the control samples, the sourness and sweetness of the MeJA-treated



Table 2 Effects of different treatments on broccoli flavor on the day of harvesting and turning yellow^a

Flavor compounds	Treatments			
	Fresh samples	Control	MeJA	MT
Sweetness	-2.39 ± 0.23c	8.27 ± 0.04b	17.39 ± 0.11a	-2.03 ± 0.04c
Sourness	-46.38 ± 0.15b	-44.29 ± 0.08b	-23.44 ± 0.14a	-45.24 ± 0.15b
Saltiness	31.02 ± 0.27 ns	25.12 ± 0.11 ns	31.22 ± 0.18 ns	30.66 ± 0.27 ns
Bitterness	-0.81 ± 0.11c	2.02 ± 0.22b	-12.2 ± 0.02d	8.32 ± 0.07a
Astringency	1.02 ± 0.03b	1.74 ± 0.07b	0.22 ± 0.10b	7.29 ± 0.08a
Umami	14.23 ± 0.23b	14.91 ± 0.02b	12.4 ± 0.09b	31.33 ± 0.07a

^a Values represent the means ± standard deviation of three replicates of six broccoli heads ($n = 9$). Different letters in the different rows indicate significant differences under different treatments ($P < 0.05$). 'ns' indicates no significant difference.

Table 3 Effects of different treatments on the contents of different bioactive compounds in broccoli and on the antioxidant activity on the day of harvesting and turning yellow^a

Bioactive compounds	Treatments			
	Fresh samples	Control	MeJA	MT
Flavonoids (mg g ⁻¹)	13.40 ± 2.33c	18.01 ± 2.82b	23.62 ± 3.07a	15.02 ± 2.14c
V _C (mg/100g)	49.36 ± 3.42a	36.72 ± 3.15 b	10.45 ± 2.15 c	45.25 ± 3.22 a
Carotenoids (mg/100g)	0.38 ± 0.09c	0.62 ± 0.08 b	0.68 ± 0.09 b	1.71 ± 0.13 a
Sulforaphane (mg/100g)	130.5 ± 4.69a	97.5 ± 4.75 b	128.2 ± 3.67 a	96.8 ± 3.79 b
DPPH (100%)	18.3 ± 0.99c	24.1 ± 1.26 b	25.10 ± 1.47 b	39.2 ± 1.23 a

^a Values represent the means ± standard deviation of three replicates of six broccoli heads ($n = 9$). Different letters in different rows indicate significant differences under different treatments ($P < 0.05$). 'ns' indicates no significant difference.

samples increased by 47.1% and 110.3%, whereas the bitterness was significantly decreased ($P < 0.05$) by 710.2%. For the MT samples, the level of bitterness, astringency, and umami increased by about 50.5%, 319.1%, and 110.1%, respectively, compared to that in the samples without any treatment, whereas the sweetness was significantly decreased ($P < 0.05$) by 124.2%.

3.4 Effect of different treatments on the bioactive compounds

Broccoli is a good source of bioactive compounds with antioxidant, immuno-regulatory, and anti-aging effects. We detected the content of flavonoids, V_C, carotenoids, and sulforaphane in broccoli heads on the day of harvesting and turning yellow after storage at 20 °C. As shown in Table 3, when the control samples changed from green to yellow, the sulforaphane and V_C content decreased by 25.3% and 25.6%, whereas the contents of flavonoids and carotenoids increased by 34.4% and 63.2%. Compared to the control samples, MeJA treatment significantly reduced ($P < 0.05$) the V_C content by 71.5%, and increased the sulforaphane and flavonoids content by 31.5% and 30.5%. The contents of V_C and carotenoids in the MT-treated samples was significantly increased ($P < 0.05$) by 23.2% and 175.8%, respectively whereas those of other bioactive compounds tested in this study were not significantly different from the control samples. We also observed that the MT treatment prevented the decrease of V_C and carotenoids, but not of the flavonoids.

Moreover, in the MeJA treated samples, the content of sulforaphane and flavonoids was significantly accumulated ($P < 0.05$), but the normal level of V_C could not be maintained.

3.5 Effect of different treatments on the antioxidant activity

The DPPH radical scavenging activity is used to measure the antioxidant ability in plants. As shown in Table 3, compared to the fresh samples after harvest, the DPPH radical scavenging activity of all the samples was increased gradually accompanied by the yellowing of broccoli heads. These results were in agreement with those of other studies on broccoli.²⁵ Moreover, the DPPH in the control samples was significantly higher than that in the fresh samples ($P < 0.05$), and was about 62.7% lower than that in the MT samples. There was no significant difference ($P < 0.05$) in DPPH between the control and MeJA samples when the buds began to turn yellow.

4 Discussion

Sensory quality, including color, aroma, flavor, and tissue morphology, is one of the important factors for customers in choosing fruits and vegetables. The broccoli heads are composed of tightly clustered buds. The freshly picked broccoli samples were green, whereas the buds stored at 20 °C without treatment began to turn yellow on day 2 during the shelf life. During the storage, the permeability of different cell membranes would change, resulting in the disintegration of the



chloroplast membrane and deformation of the thylakoid membrane structure. The chlorophyll attached to the thylakoid membrane would thus be released and consumed by extensive action of the chlorophyll degrading enzymes. Moreover, because of the transformation of plastids, chloroplasts and leucoplasts are converted to chromoplasts, which store a large number of carotenoids. In this way, while the green pigment (chlorophyll) is consumed, a large amount of yellow pigment (carotenoids) accumulates, which in turn causes the yellowing of broccoli.²⁶ It has previously been shown that the yellowing of plants was closely related to reactive oxygen species and cell membrane components.²⁷

In this study, the MeJA-treated broccoli heads maintained a high level of sensory quality on day 4, whereas the control samples turned completely yellow. Previous studies have suggested that MeJA induced the production of ethylene to accelerate the yellowing process. However, the effect of MeJA treatment is related to factors such as treatment concentration and the maturity of fruits and vegetables, and this view has been confirmed in the relevant research results of apple quality.²⁸ Simultaneously, MeJA treatment could maintain the surface color of the fruit while reducing the rate of quality change during storage. Feng *et al.*²⁹ found that MeJA treatment could prevent the mango chilling without accelerating coloration. Dong and Cai³⁰ proved that MeJA application significantly increased chlorophyll content, chlorophyll fluorescence parameter F_v/F_m and F_v/F_0 in rice leaves under drought conditions. Moreover, the slightly delayed yellowing process in the MeJA treatment might be due to the enhanced adaptation of broccoli under the unsuitable storage temperature used in the study. MeJA plays an important role as a signal molecule in plant-induced responses and defense mechanisms. Evidence³¹ suggests that when plants are under stress, MeJA rapidly accumulated to a high content, in response to the adversity, and thereby, participating in and regulating many physiological and biochemical processes of the plant for improving the resistance to adversity. The functions of MT, including its antioxidant, anti-inflammatory properties, and the capacity to modulate mitochondrial homeostasis, are linked to the redox status of cells and tissues. MT is important for the physiological regulation of cellular homeostasis.³² In addition, Szafrńska *et al.*³³ studied the ultrastructure of *Vigna Radiata* roots and found that melatonin effectively protected the structure of plastids (chloroplast, chromoplast, leucoplast), which could prove that MT was helpful in maintaining the morphology of the chloroplast and thylakoid membranes. Thus, the yellowing process of the MT-treated broccoli samples was remarkably delayed.

Aroma is an important index that reflects the flavor, maturity, and commodity quality of fruits and vegetables. The pleasant aroma is an important factor in attracting the consumers and in enhancing the market competitiveness of the product. Aroma is determined by the content, composition, and odor threshold of the different volatile components. Broccoli is characterized by sulfurous volatile compounds, which are similar to that in other cruciferous plants. Furthermore, esters and aldehydes provide fruity and grassy aroma, respectively, whereas alcohols are often accompanied with the spoiling

odors. Dimethyl disulfide, disulfide dimethyl, dimethyl trisulfide, hexanal, *cis*-3-hexen-1-ol, and ethanol acetate were identified as the main volatile components in broccoli (Table 1). This result was similar to that obtained by Annelie *et al.*³⁴ in broccoli. As per our findings, sulfides are the main volatile components of cruciferous vegetables. The GC-MS analysis revealed that there was no significant change in the content of sulfides when the untreated sample was yellowed, which might be because of the short storage time during which the change in total sulfide content could not be registered. The treatment of broccoli samples with MeJA could significantly reduce the unique sulfide volatiles, allowing broccoli to maintain fresh aroma even when they started to turn yellow. This might be due to the differences in the levels of dimethyl disulfide in the MeJA-treated samples. The MT treatment increased the production of volatile sulfides in the yellowed broccoli and was accompanied with a full-bodied odor because of deterioration of the lipid membranes in the cells as well as because of the loss of intracellular compartmentalization due to prolonged storage, which resulted in enzymatic reactions.³⁵ In addition, MT might promote the accumulation of the free amino acid, *S*-methyl-L-cysteine sulfide, which is the main source of volatile sulfide compounds.

When the control samples turned yellow, the types of aldehydes and esters as well as their contents were not significantly changed. In the MeJA-treated samples, when the buds just lost their green color, the content of (*E*)-2-hexenal, which is used to evaluate the volatile content of plants, was increased. It might be that MeJA treatment promoted the accumulation of aldehydes,³⁶ which provided fresh grass aroma to broccoli. In addition, 2-hexenal was the main component of leaf volatiles, which had been effectively increased by MT treatment during yellowing. MT induced the synthesis of the precursor substances of 2-hexenal volatile compounds, which were mainly formed by the aldol condensation pathway.³⁷ Moreover, the MT treatment resulted in a sharp decrease in the ester content of the yellowing samples, which was different from that in the control and MeJA samples. This might be related to the reduction in the types of ester present in such samples.

A survey conducted by the Regional Fruit and Vegetables Economic Committee found evidence that flavor is one of the main reasons for some consumers rarely or never purchasing cauliflower.³⁸ Some consumers were very sensitive to the unique sulfide taste of Brassica vegetables, which gives a pungent odor and bitter taste to these vegetables.¹² Unlike other taste analysis instruments used throughout the world, the electronic tongue has a real taste analysis system, which matches the tastes in humans.³⁹ In our study, we detected the changes in sweetness, sourness, saltiness, bitterness, astringency, and umami of broccoli heads during yellowing and assessed the effects of different treatments on these changes. The results were obtained by an electronic tongue detector in the form of unitless numerical values, where a larger number indicated a more intense flavor. We found that the untreated samples had a large increase in sweetness and bitterness when they began to turn yellow. This might be related to the accumulation of carbohydrates in the broccoli samples after the harvest. At this time, the flavor of broccoli was better than that at the time of harvest.



Different treatments have influence on the specific flavor of *Brassica* vegetables, because of the enzymatic reaction products which play the significant roles in the creation of flavors.⁴⁰ Previous studies have shown that the products of hydrolysis of glucosinolates play a key role in determining the flavor of *Brassica* vegetables.⁴¹ The relationship between myrosinase and the feeling of taste was investigated by Engel *et al.*³⁸ They found that the products of the hydrolysis of glucosinolates could explain the feeling of bitterness. In our study, a high level of sulforaphane was observed in the MeJA samples when they began to turn yellow. As a hydrolysate of glucosinolates, sulforaphane affect the binding of bitter taste receptors to precursor substances, thereby, reducing the bitterness. On the other hand, MeJA effectively induced the accumulation of sugars and increased the sweetness.⁴² The decrease in bitterness and the added sweetness resulted in broccoli having a better flavor when they began to turn yellow. These findings were especially interesting considering the interactions of taste molecules and potential masking of bitterness by the sweet taste. The complex role of isothiocyanates in the perception of flavor of *Brassica* vegetables was attributed to the interaction with bitterness taste receptors.⁴³ The content of isothiocyanates in the MT-treated samples was higher than that in the MeJA samples, which provided the material basis for bitterness receptors, and increased the bitter flavor of the yellowed samples. As a flavor compound, isothiocyanates also contributed to the aroma of broccoli. Moreover, the improvement in the bitterness, astringency, and umami of broccoli samples treated with MT might be related to the high content of volatile sulfides when broccoli turned yellow from green. The above contents confirmed the close relationship between the volatile components and flavor.

Broccoli is rich in various nutrients, which is one of the important factors that attract consumers. Flavonoids and V_C have indisputable antioxidant, and cancer and chronic disease fighting properties, and carotenoids are the main source of vitamin A in the body. We found that even if the control samples started to turn yellow, the contents of flavonoids, and carotenoids continued to increase because of the shorter storage times. There are large amounts of oxidases in broccoli and we inferred that it might be due to an increase in the oxidase activity during the yellowing process, resulting in different degrees of loss of V_C content in the MeJA samples. However, when the MT-treated broccoli samples began to turn yellow, we observed increased levels of carotenoids and V_C due to the effective inhibition of oxidation and scavenging of free radicals by MT. Besides, the effect of MT was indirectly demonstrated by the increase in DPPH in the broccoli samples, accompanying the yellowing process. In our study, flavonoids were not affected by MeJA and MT. It might be possible that the signal transduction and free radicals do not directly participate in the biosynthetic pathway of flavonoids.

Sulforaphane is an important secondary metabolite in broccoli, which is widely known for its superior antioxidant properties. When broccoli lost the green color, the sulforaphane level in the control samples was drastically reduced. The reduction in sulforaphane content might be a result of

environmental discomfort and the depletion of glucosinolates. Pérez-Balibrea *et al.*⁴⁴ found that the synthesis and content of sulforaphane were directly influenced by the exogenous treatments. MeJA stimulated the production of secondary metabolites in many plant species. In our study, the sulforaphane level in the MeJA-treated samples was significantly higher ($P < 0.05$) than that in the control samples. MeJA induced the synthesis of glucosinolates and affected the content of sulforaphane in broccoli. It significantly improved the sulforaphane content, as was also reported by Dombrecht.⁴⁵ In addition, Halkier and Gershenzon⁴⁶ indicated that MeJA stimulated the biosynthesis of indoles, which was due to the anabolic effect of jasmonate compounds and aliphatic amino acids, and effected the accumulation of sulforaphane. However, MT did not have a similar effect, showing no difference in the sulforaphane content when compared to the control samples of yellowed broccoli.

MeJA was first discovered and isolated from the essential oil of *Jasminum grandiflorum* L. by Demole *et al.*,⁴⁷ which was a flavor compound. Thus, the source of MeJA is safer than other plant hormones. The significance of MeJA for higher plant metabolism is beyond doubt, and it also has a positive contribution to human health. A large number of reports indicate that MeJA has an anti-cancer effect in the clinic, and the effect was significant.^{48,49} Kniazhanski *et al.*⁵⁰ found that MeJA was effective against cervical carcinoma cell lines. Furthermore, MeJA is the involvement of ROS in MeJA-induced apoptosis, which is the anti-cancer mechanism.⁵¹ In our study, the MeJA content in treatment ($0.5 \mu\text{mol L}^{-1}$) was extremely trace, which not only ensured food safety, but also increased the commercial value of anti-cancer in broccoli to some extent. MT is an indole hormone secreted by the pineal gland, which physiological functions include inducing sleep, regulating the endocrine system, maintaining stable internal environment, enhancing immunity, anti-aging and so on.⁵² MT mediated by specific receptors, which regulates the biological rhythm of sleep awakening and plays a role in sedation and induces sleep, so that it is called physiological hypnotic agent.⁵³ In the regulation of melatonin, it has obvious therapeutic effects on sleep disorders which caused by sleep delay syndrome and abnormal time difference. Arendt *et al.*⁵⁴ analyzed a number of clinical findings and found that the recommended melatonin intake was 0.5–5 mg for each time. The concentration of MT used in our study was $100 \mu\text{mol L}^{-1}$, and the MT content was much lower than the daily recommended limit. Therefore, normal intake of broccoli would not affect the body.

5 Conclusions

In our study, the buds of untreated broccoli began to turn yellow on day 2 when stored at 20°C . The MeJA and MT treatments effectively delayed the yellowing time of broccoli by 2 and 4 days, respectively. When the control samples began to lose their green color, the substantial increase in sweetness and bitterness was accompanied by an increase in various bioactive compounds, except for sulforaphane; however, there was no significant change in the content of volatile components. Compared to the control samples, the MeJA treatment mainly



contributed to the accumulation of sulforaphane and flavonoids, to sweetness and sourness, and to the relief from the special odor of sulfides. However, the MT treatment tended to improve the antioxidant activity and the increase of bitterness, astringency, umami, and volatile sulfides. These results showed that the flavor, aroma, and nutrients of broccoli that began to turn yellow was changed to some extent, and the MeJA and MT treatment could significantly regulate the content of bioactive compounds while enhancing the sensory quality. In addition, we believe that maintenance of the original flavor of fruits and vegetables should no longer be our main focus. We should continue to explore strategies to improve the distasteful flavor of fruits and vegetables. The MeJA and MT treatments can contribute to the improvement of flavor in broccoli, whatever its form might be.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the National Key R&D Program of China (no. 2016YFD0400103).

References

- R. B. Jones, C. L. Frisina, S. Winkler, M. Imsic and R. B. Tomkins, *Food Chem.*, 2010, **123**, 237–242.
- G. Yuan, B. Sun, J. Yuan and Q. Wang, *Food Chem.*, 2010, **118**, 774–781.
- R. R. Dedolph, S. H. Wittwer, V. Tuli and D. Gilbert, *Plant Physiol.*, 1962, **37**, 509–512.
- K. M. Ku, E. H. Jeffery and J. A. Juvik, *J. Sci. Food Agric.*, 2014, **94**, 2090–2096.
- C. Wasternack and B. Hause, *Ann. Bot.*, 2013, **111**, 1021–1058.
- M. K. Kang, E. H. Jeffery and J. A. Juvik, *J. Agric. Food Chem.*, 2013, **61**, 9623–9631.
- C. Rodriguez, J. C. Mayo, R. M. Sainz, I. Antolín, F. Herrera and V. Martín, *J. Pineal Res.*, 2004, **36**, 1–9.
- S. Park, D. Lee, H. Jang, Y. Byeon, Y. Kim and K. Back, *J. Pineal Res.*, 2013, **54**, 258–263.
- M. M. Posmyk, M. BańAbusta, M. Wiczorek, E. Sliwinska and K. M. Janas, *J. Pineal Res.*, 2009, **46**, 214–223.
- M. M. Posmyk, H. Kuran, K. Marciniak and K. M. Janas, *J. Pineal Res.*, 2010, **45**, 24–31.
- M. N. Wiczorek, M. Walczak, M. Skrzypczak-Zielińska and H. H. Jeleń, *Crit. Rev. Food Sci. Nutr.*, 2017, **56**, 100–145.
- J. Lv, J. Wu, J. Zuo, L. Fan, J. Shi and L. Gao, *Food Chem.*, 2017, **216**, 225–233.
- V. Valpuesta and M. A. Botella, *Trends Plant Sci.*, 2004, **9**, 573–577.
- M. V. Eberhardt, Y. L. Chang and H. L. Rui, *Nature*, 2000, **405**, 903–948.
- M. V. Eberhardt, K. Kobira, A. S. Keck, J. A. Juvik and E. H. Jeffery, *J. Agric. Food Chem.*, 2005, **53**, 7421.
- Y. Topcu, A. Dogan, Z. Kasimoglu, H. Sahin-Nadeem, E. Polat and M. Erkan, *Plant Physiol. Biochem.*, 2015, **93**, 56–65.
- R. Molaykumar, J. Lekhraj, I. Seiichiro and T. Tojiro, *Food Chem.*, 2009, **114**, 263–269.
- F. Xu, X. Chen, J. Peng, X. Wang, J. Wang and Y. Zheng, *Eur. Food Res. Technol.*, 2012, **235**, 793–800.
- J. J. Gills, E. H. Jeffery, N. V. Matusheski, R. C. Moon, D. D. Lantvit and J. M. Pezzuto, *Cancer Lett.*, 2006, **236**, 72–79.
- A. GliszczyńskaSwięto, E. Ciska, K. PawlakLemańska, J. Chmielewski, T. Borkowski and B. Tyrakowska, *Food Addit. Contam.*, 2006, **23**, 1088–1098.
- A. Rybarczyk-Plonska, M. K. Hansen, A. B. Wold, S. F. Hagen, G. I. A. Borge and G. B. Bengtsson, *Postharvest Biol. Biotechnol.*, 2014, **98**, 82–89.
- P. Stepnowski, K. H. Blotevogel and B. Jastorff, *Int. Biodeterior. Biodegrad.*, 2004, **53**, 127–132.
- H. Liang, Q. P. Yuan, H. R. Dong and Y. M. Liu, *J. Food Compos. Anal.*, 2006, **19**, 473–476.
- T. Hatano, H. Kagawa, T. Yasuhara and T. Okuda, *Chem. Pharm. Bull.*, 1988, **36**, 2090–20134.
- P. Jin, D. Yao, F. Xu, H. Wang and Y. Zheng, *Food Chem.*, 2015, **172**, 705–709.
- M. S. Tian, *J. Am. Soc. Hortic. Sci.*, 1994, **119**, 276–281.
- J. D. Miller, R. N. Arteca and E. J. Pell, *Plant Physiol.*, 1999, **120**, 1015–1024.
- X. Fan, J. P. Mattheis and J. K. Fellman, *J. Am. Soc. Hortic. Sci.*, 1998, **123**, 421–425.
- F. Lei, Y. Zheng, Y. Zhang, F. Wang, L. Zhang and L. Zhaoxin, *J. Agric. Food Chem.*, 2000, **48**, 515–519.
- T. Dong and K. Cai, *Environ. Sci. Technol.*, 2009, **14**, 156–187.
- J. J. Cheong and Y. D. Choi, *Trends Genet.*, 2003, **19**, 409–413.
- Y. Aguilera, T. Herrera, V. Benítez, S. M. Arribas, L. D. P. Al and R. M. Esteban, *Food Chem.*, 2015, **170**, 203–211.
- K. Szafrńska, S. Glińska and K. M. Janas, *Biol. Plant.*, 2012, **57**, 91–96.
- A. Jacobsson, A. Tim Nielsen and I. Sjöholm, *J. Agric. Food Chem.*, 2004, **52**, 1607–1614.
- D. Zhao, J. Tang and X. Ding, *LWT-Food Sci. Technol.*, 2007, **40**, 439–447.
- M. Yu, L. Shen, B. Fan, D. Zhao, Y. Zheng and J. Sheng, *Postharvest Biol. Biotechnol.*, 2009, **54**, 153–158.
- E. Ortner and M. Granvogl, *J. Agric. Food Chem.*, 2017, **41**, 221–261.
- E. Engel, C. Baty, D. Le Corre, I. Souchon and N. Martin, *J. Agric. Food Chem.*, 2002, **50**, 6459–6467.
- A. Rudnitskaya, L. M. Schmidtke, A. Reis, M. R. Domingues, I. Delgado and B. Debus, *Food Chem.*, 2017, **229**, 20–27.
- F. S. Hanschen, R. Klopsch, T. Oliviero, M. Schreiner, R. Verkerk and M. Dekker, *Sci. Rep.*, 2017, **56**, 712–761.
- R. Kubec, A. Veronika Drhová and J. Velišek, *J. Agric. Food Chem.*, 1998, **46**, 4334–4340.
- L. Yu, *Postharvest Biol. Biotechnol.*, 2016, **113**, 8–16.
- D. Zabararas, M. Roohani, R. Krishnamurthy, M. Cochet and C. Delahunty, *Food Funct.*, 2013, **4**, 592–601.



- 44 S. Pérez-Balibrea, D. A. Moreno and C. García-Viguera, *Food Chem.*, 2011, **125**, 348–354.
- 45 B. Dombrecht, G. P. Xue, S. J. Sprague, J. A. Kirkegaard, J. J. Ross and J. B. Reid, *Plant Cell*, 2007, **19**, 2225–2245.
- 46 B. A. Halkier and J. Gershenzon, *Annu. Rev. Plant Biol.*, 2006, **57**, 303–333.
- 47 E. Demole, E. Lederer and D. Mercier, *Helv. Chim. Acta*, 1962, **45**, 512–567.
- 48 S. Cohen and E. Flescher, *Phytochemistry*, 2009, **70**, 1600–1609.
- 49 E. Flescher, *Cancer Lett.*, 2007, **245**, 1–10.
- 50 T. Kniazhanski, A. Jackman, A. Heyfets, P. Gonen, E. Flescher and L. Sherman, *Cancer Lett.*, 2008, **271**, 34–46.
- 51 N. Goldin, A. Heyfets, D. Reischer and E. Flescher, *J. Bioenerg. Biomembr.*, 2007, **39**, 51–57.
- 52 Y. H. Wu and D. F. Swaab, *Sleep Med.*, 2007, **8**, 623–636.
- 53 G. C. Brainard, J. P. Hanifin, J. M. Greeson, B. Byrne, G. Glickman and E. Gerner, *J. Neurosci.*, 2001, **21**, 6405–6412.
- 54 J. Arendt, E. J. Van Someren, R. Appleton, D. J. Skene and T. Akerstedt, *Clin. Med.*, 2008, **8**, 381–383.

