
Nanoscale Advances

Volume 5 Number 10 21 May 2023 Pages 2667-2834



ISSN 2516-0230



REVIEW ARTICLE



Nanoscale Advances



REVIEW

View Article Online



Cite this: Nanoscale Adv., 2023, 5,

Received 10th August 2022 Accepted 27th January 2023

DOI: 10.1039/d2na00534d

rsc.li/nanoscale-advances

The toxicity of nanoparticles and their interaction with cells: an in vitro metabolomic perspective

Mohammad Awashra ** and Piotr Mtynarz ** b

Nowadays, nanomaterials (NMs) are widely present in daily life due to their significant benefits, as demonstrated by their application in many fields such as biomedicine, engineering, food, cosmetics, sensing, and energy. However, the increasing production of NMs multiplies the chances of their release into the surrounding environment, making human exposure to NMs inevitable. Currently, nanotoxicology is a crucial field, which focuses on studying the toxicity of NMs. The toxicity or effects of nanoparticles (NPs) on the environment and humans can be preliminary assessed in vitro using cell models. However, the conventional cytotoxicity assays, such as the MTT assay, have some drawbacks including the possibility of interference with the studied NPs. Therefore, it is necessary to employ more advanced techniques that provide high throughput analysis and avoid interferences. In this case, metabolomics is one of the most powerful bioanalytical strategies to assess the toxicity of different materials. By measuring the metabolic change upon the introduction of a stimulus, this technique can reveal the molecular information of the toxicity induced by NPs. This provides the opportunity to design novel and efficient nanodrugs and minimizes the risks of NPs used in industry and other fields. Initially, this review summarizes the ways that NPs and cells interact and the NP parameters that play a role in this interaction, and then the assessment of these interactions using conventional assays and the challenges encountered are discussed. Subsequently, in the main part, we introduce the recent studies employing metabolomics for the assessment of these interactions in vitro.

Introduction

Nanomaterials are defined as materials with at least one dimension smaller than 100 nm, while nanotechnology is defined as the understanding and manipulation of matter at dimensions in the range of 1 to 100 nm, where unique phenomena enable novel applications.2 Nanotechnology introduces many potential health, environmental, and industrial

^aDepartment of Chemistry and Materials Science, School of Chemical Engineering, Aalto University, 02150 Espoo, Finland. E-mail: mohammad.awashra@aalto.fi ^bDepartment of Biochemistry, Molecular Biology and Biotechnology, Faculty of Chemistry, Wroclaw University of Science and Technology, Wroclaw, Poland. E-mail: piotr.mlynarz@pwr.edu.pl



Mohammad Awashra started pursuing his doctoral degree in the Microfabrication Group (MFG) at Aalto University starting in 2022. His research is focused on studying the cellrepellent behaviour of superhydrophobic surfaces with nano and micro features. He received his M.Sc. in Chemical Nano-Engineering in 2021 from Aix-Marseille University during the Erasmus Mundus Joint Master

Program (CNE). He received his Bachelor of Chemistry from Birzeit University in 2019.



Prof. Piotr Młynarz received his PhD in Chemistry in 2001 from the Faculty of Chemistry, University of Wrocław and joined the Faculty of Chemistry Wrocław University of Science and Technology in 2002. He is a member of the Department of Biochemistry, Molecular Biology and Biotechnology, and Head of the Bioanalytical group (2019). His research is mainly focused on the use of metabolomics

methods in the field of civilization disease diagnostics and cell metabolism.

Review Nanoscale Advances

benefits^{3,4} and its applications are widespread in daily life, transforming society.⁵ For example, its applications in the food industry range from agriculture⁶ to food processing and packaging.⁷ Furthermore, nanotechnology is applied in drug delivery,⁸ imaging, diagnostics, cosmetics,⁹ clothing,¹⁰ transportation,¹¹ biofuels,¹² and biosensors.¹³ Therefore, human exposure to nanomaterials (NMs) nowadays is highly probable. Depending on the type of product in which nanoparticles (NPs) are used, exposure may occur through inhalation, dermal, and oral pathways. Among them, inhalation is considered as the most significant exposure route for airborne NPs.¹⁴

Due to their high surface to volume ratio, high reactivity, and tunable characteristic properties, NMs exhibit great benefits such as enhanced targeting and imaging techniques.¹⁵ However, NMs may also cause some potential risks to human health and the environment.¹⁶ Given that human and environmental exposure to NMs are inevitable, nanotoxicity research is attracting increasing attention.17 In the last decade, the number of research studies on the toxicity of different types of NMs has increased dramatically. NMs may affect human health in several ways such as inflammation18 and heart problems.19,20 Thus, to understand the mechanisms of these effects, more investigations in the nanotoxicology field are necessary at the cellular and sub-cellular levels. The scope of nanotoxicity depends on many parameters that are related to the NM itself such as its size, shape, chemical composition, and coating²¹ or the exposed cell type or tissue.22

When exposed to NPs, the cell can be affected via several routes, including a decrease in cell viability and proliferation,23 inflammatory response, production of cytokines, 24,25 oxidative stress,26,27 generation of reactive oxygen species (ROS),28-30 cell membrane damage,29 mitochondrial damage,28,31 cell cycle dysregulation,²⁷ DNA damage,³² genotoxicity,²⁴ lipid peroxidation, changes in cell morphology,33 apoptosis29 or necrosis,25 and metabolic changes.30 To study the cytotoxicity of NPs, many conventional assays and biomarkers are used. For example, the cell viability and proliferation can be investigated using tetrazolium-based assays such as MTT,34 MTS,35 and WST-1.36 Alternatively, the cell inflammatory response can be investigated by measuring inflammatory biomarkers, such as IL-8, IL-6, and tumor necrosis factor, using ELISA.37 Moreover, for cell membrane integrity, lactate dehydrogenase (LDH) and trypan blue exclusion assays can be used,37 and for cell metabolism, the Alamar Blue assay is frequently used.38 However, although these assays afford general information about the cytotoxicity of NPs, they do not give molecular information about the mechanism of their cytotoxicity.38-41 Moreover, NPs can interfere with the conventional assays, and thus the use of more than one assay is important. In general, most of the studies on the cytotoxicity of NPs mainly use the conventional (phenotypic) tests and assays. Alternatively, some studies used other techniques based on the change in epigenome, transcriptome, proteome, or metabolome (omics techniques) induced by NPs.39 These techniques are beneficial to study the effect of NPs on cells at the molecular level and explain the results of conventional essays. Among them, metabolomics is one of the most powerful bioanalytical strategies, enabling a picture of the metabolites of an organism in the

course of a biological process to be obtained, which is the omics technique of interest in this review.^{40,41} The introduction of NPs in a cell line may cause a change in the levels of certain metabolites, which may give a clue on their effect on cells. During the past decade, many *in vitro* studies have used metabolomics to investigate the cytotoxicity of NPs on different cell lines.

In this review, the ways NPs and cells interact and the effects of the NP parameters on their interaction are discussed, followed by an overview of the cytotoxicity of different NPs in *in vitro* models, focusing on the use of metabolomics as a tool to identify the mechanisms and molecular information of their cytotoxicity.

2. Cellular uptake of NPs

The cytotoxic effects of NPs usually originate from their presence inside cells. ⁴² However, many applications of NMs in biomedicine require their entry in the cell to achieve their goal. Therefore, to further understand the cytotoxicity mechanisms of NPs on the cell and its metabolism, it is important to first understand the cellular uptake mechanisms of NPs. This will also aid the design of environmentally safer NMs with enhanced cellular targeting and uptake properties for therapeutic purposes. ⁴³

When immersed in a biological fluid, NPs are exposed to a different medium than that employed for their synthesis. This will force the NPs to interact with the surrounding medium, which may alter their physical and chemical properties.⁴⁴ To stabilize themselves, NPs tend "to catch" the surrounding biomolecules (proteins, lipids, *etc.*) and form a biomolecular corona or protein corona (in case they are surrounded by proteins only), which may alter their identity.⁴⁵

NPs may be taken up by the cell in an energy-independent process, such as simple diffusion or translocation. However, most NP uptake pathways are energy dependent *via* endocytosis. Endocytosis is the formation of vesicles from the cell plasma membrane to take up substances such as particles, nutrients, and dead cells from the extracellular to the intracellular environment.⁴⁶ Endocytosis is described in two categories, *i.e.*, phagocytosis and pinocytosis.

2.1 Phagocytosis

Phagocytosis is the cellular uptake of particulates $(0.5-10~\mu m)$ in the plasma-membrane envelope. It is known as a host defence mechanism, engulfing and internalizing cargos such as particles, dead cells, and cell debris. This mechanism is a ligand-induced process, where NPs are engulfed by adsorbing opsonins, followed by their interaction with complement receptors on the cell surface (see Fig. 1).

2.2 Pinocytosis

Pinocytosis is the cellular uptake of extracellular fluids and dissolved solutes.⁴⁹ It can be divided into macropinocytosis, clathrin- and caveolae-independent endocytosis, and receptor-mediated endocytosis. The latter is classified as clathrin-dependent endocytosis and caveolae-dependent endocytosis based on the proteins involved in the pathway.⁵⁰

Nanoscale Advances Review




Fig. 1 Nanoparticle internalization via phagocytoses.

2.2.1 Macropinocytosis. This mechanism involves cytoskeleton rearrangements that induce the formation of membrane ruffles, which fold back, resulting in the formation of large

intracellular vacuoles $(0.1–5~\mu m)^{51}$ referred to as macropinosomes (see Fig. 2). Macropinocytosis is actin-dependent endocytosis, while it is independent of clathrin and membrane receptors. ⁵²




Fig. 2 Pinocytosis internalization mechanisms. (1) Macropinocytosis. (2) Clathrin-mediated endocytosis. (3) Caveolae-mediated endocytosis.

Review Nanoscale Advances

2.2.2 Clathrin-mediated endocytosis. Clathrin-mediated endocytosis is the main process for the internalization of many NPs, which is used by all eukaryotic cells to internalize small particles and nutrients such as cholesterol. When the plasma membrane is rich in clathrin and ligand-receptor complexes start to form on the cell membrane surface, a cage of clathrin starts to form around the vesicle, resulting in vesicles with a diameter of 100-150 nm (see Fig. 2).43,49,53

2.2.3 Caveolae-mediated endocytosis. Caveolae are bulbshaped invaginations in the plasma membrane, which are 50-80 nm in size. These vesicles are coated by caveolin and cavin and detached from the membrane by dynamin, which is a 100 kDa GTPase (see Fig. 2).43,54,55


Role of physicochemical properties of NPs in cellular uptake and cytotoxicity

It is important to consider the physicochemical properties (size, shape, surface functionalization, surface chemistry, chemical composition, concentration, etc.) of NPs in their design for biomedical or other applications. The interactions of NPs with the cell membrane and organelles can significantly be altered at the bio-nano interface by these physicochemical properties, consequently changing the cellular uptake and nanotoxicity of the NPs. Therefore, before starting to assess the biological

responses of NPs, thorough and proper characterization of the physicochemical properties of their core and surface should be performed.⁵⁶ In this part, we mainly focus on the effect of the size, shape, and surface chemistry of NPs on their cytotoxicity and cellular uptake (see Fig. 3). The effect of the NP core composition is not discussed here given that the surface characteristics are more important than the bulk characteristics in this context.

3.1 Size

The size of NPs plays an important role in both their cellular uptake and cytotoxicity. Thus, it is considered a key factor when designing NPs for biomedical application. Due to the fact that NPs possess a size between atoms and bulk materials, they lie on the critical transition zone between two different worlds. 57,58 It is worthy to mention that the original (primary) size of NPs differs from their hydrodynamic size in biological media. 59 This is mainly because of the formation of a biomolecular corona and the aggregation of the NPs. In this case, the aggregation of NPs can be prevented by manipulating the balance of attractive and repulsive forces. 60 For instance, Fe3O4 NPs can be stabilized with citrate, preventing their aggregation due to electrostatic repulsion.61 However, due to the formation of a biomolecular corona and the different ionic strengths of biological solutions compared to water, NPs may have new surface identity. Wei et al.38 performed a cytotoxicity study on the different sizes of TiO₂ (5 and 200 nm) and Al₂O₃ (10 and 50 nm) NPs and observed






Fig. 3 Physicochemical properties of NPs.

Nanoscale Advances Review

the formation of aggregates in solution form when the NPs were suspended in cell medium without serum, where the sizes of all the NPs became 8-388-fold larger than their original sizes due to the higher ionic strength of the medium compared to water. Upon the addition of serum, the hydrodynamic sizes of the NPs decreased to only 1.6-10 folds larger than their original sizes. This is because the formation of the protein corona around the NPs prevented them from aggregating due to steric repulsion. The authors found that the smaller NPs (in terms of primary size, rather than hydrodynamic size) for both TiO2 and Al2O3 had higher cytotoxicity and much greater decrease in cell metabolic activity.

When studying the NP-cell membrane interaction mechanism dependence on the size of NPs, it was found that it has a strong influence. Specifically, large NPs (>60 nm) may cause steric hindrance, which prevents their interaction with the cell membrane.62 Conversely, NPs smaller than the cut-off size of receptor diffusion (<30 nm) may not recruit enough cell membrane receptors in the interaction region to overcome the elastic recoil force, preventing membrane wrapping from occurring.63 Moreover, the membrane receptors are known to form clusters that are 10-50 nm in size. Thus, a 50 nm NP, for example, needs to interact with only one receptor cluster, while a 500 nm NP must interact with several clusters simultaneously. This makes the internalization of the 50 nm NP energetically more favourable than the 500 nm NP.44

In general, smaller-sized NPs have been reported to have higher cellular uptake and higher cytotoxicity. For instance, Dong et al.64 reviewed 76 carefully chosen literature reports that included in vitro studies of the size-dependent cytotoxicity of amorphous silica NPs (aSiO₂ NPs) and found that 76% of these papers showed that smaller-sized aSiO2 NPs exhibited greater cytotoxicity. However, it is important to consider that the cell type plays a role in this process given that it depends on the predominant pathway of cellular uptake in each different cell. 65,66

For some NPs, the higher the cellular uptake of NPs, the greater their cytotoxicity.67 Nonetheless, there are some exceptions, where the cytotoxicity of NPs is independent of their cellular uptake. In these cases, the cytotoxicity is induced by sources other than amount of toxicant, including the NP high surface area, instability, and ion release. Gliga et al.68 found that 10 nm silver NPs (AgNPs) are more toxic to the human lung BEAS-2B cell line than other NPs with higher uptake ratios due to the release of more Ag⁺.

3.2 Shape

The shape of NPs can be controlled by manipulating the experimental conditions during their synthesis, such as supersaturation, reducing agents, temperature, surfactants, and secondary nucleation.69 There are many different shapes and geometries of NPs, such as spherical, rod, flower, star, disc, cubic, prismatic, and needle-like structures. The aspect ratio (AR), which is the proportion between width and height of NPs, is used to compare different shapes of NPs. For example, spherical AuNPs have an AR of 1, while Au nanorods (AuNRs) have a higher AR.

It was proven that the cellular uptake and cytotoxicity of NPs are affected by the AR of NPs. Given that AuNPs are common in many biomedical applications, many studies investigated their shape-dependent cellular uptake and cytotoxicity. For instance, Woźniak et al. 70 compared the in vitro cytotoxicity profiles of different shapes and sizes of bare (non-coated) AuNPs in cancer (HeLa) and normal (HEK293T) cell lines. They found that Au nanospheres (AuNS) and AuNRs had higher cytotoxicity than star-, flower- and prism-shaped AuNPs. However, the sizes of these different AuNPs shapes also differed. Specifically, the AuNSs and AuNRs had smaller sizes (10 nm and 38 \times 16 nm, respectively), while the flower-, prism-, and star-shaped AuNPs had larger sizes (~370 nm, ~160 nm, and ~240 nm, respectively). Thus, their sizes may also play a crucial role in this cytotoxicity tendency, given that smaller NPs are known to have higher cellular uptake and aggregation rate inside the cell, which explains the observed cytotoxicity.

3.3 Surface charge

NPs can have negative, positive, or neutral surface charge depending on their surface functional groups.71 The surface charge can affect the NP-cell membrane interactions, protein corona, and consequently the cellular uptake of NPs.72 Therefore, it is one of the most important physicochemical properties to control when designing NPs for biomedical applications. Generally, reports have shown that charged NPs have higher cellular uptake than neutral NPs.63 The cell membrane is negatively charged due to the anionic head group of phospholipids and the existence of some carbohydrates, such as sialic acid.73 Considering this, cationic NPs, in most nonphagocytic cells, are taken up by the cells to a greater extent than anionic NPs. However, in some cases, anionic NPs have greater cellular uptake in phagocytic cells.74,75 The surface charge of NPs can also tune their cellular uptake pathway. For instance, Untener et al.76 reported that positively charged AuNRs had a higher extent of internalization compared to their negatively charged counterparts. It was found that cationic AuNRs were taken up through macropinocytosis and clathrin-mediated endocytosis, while anionic AuNRs were internalized through macropinocytosis and caveolae-related mechanisms.

The cytotoxicity of NPs is also, as expected, affected by their surface charge. Similar to the dependence of the cellular uptake of NPs on their surface charge, in nonphagocytic cells, charged NPs were found to be more cytotoxic than their neutral counterparts, with the positively charged NPs, in most cases, being more cytotoxic than negatively charged NPs.74 Moreover, the surface charge of NPs does not only affect their cytotoxicity level but also their mechanisms. A study by Schaeublin et al.77 showed that although both charged and neutral AuNPs were taken up in similar amounts and caused cell morphology disruption and decreased cell viability through ROS generation in a human keratinocyte cell line (HaCaT) model, only charged NPs caused significant mitochondrial stress. This suggested that the surface charge of AuNPs can affect the mechanism of cell death. Further investigations on mitochondrial-mediated toxicity revealed that neutral AuNPs did not affect the




Fig. 4 Schematic representation of the mitochondrial membrane: different NP surface charges induce different mechanisms of cell death.⁷⁷ The neutrally charged NP (*) does not disrupt the mitochondrial membrane potential, and therefore apoptosis is not activated. The positively charged NP (+) disrupts the slight negative charge on the cytosolic side of the outer membrane, leading to a disruption in the mitochondrial membrane potential. The disruption damages the membrane and proteins, such as caspase activators, leak into the cytosol. The negatively charged NP (-) increases the negative charge on the outer membrane, which leads to a disruption in the mitochondrial membrane potential. The mitochondria compensate by releasing calcium ions that were stored in the matrix of the mitochondria. The spike in calcium induced apoptosis.⁷⁷

mitochondrial outer membrane potential, which has a slight negative charge, and thus apoptosis was not initiated, and the authors suggested that necrosis may be the cell death mechanism in this case. However, charged AuNPs affected this membrane in different ways. On the one hand, cationic AuNPs accumulated on the mitochondrial outer membrane due to its slight negative charge, which eventually damaged the membrane and caused the release of apoptotic proteins such as caspase-3 inducing mitochondrial-mediated apoptosis. On the other hand, anionic AuNPs increased this slight negative charge on the outer membrane, which forced the mitochondria, trying to adjust this potential disruption, to release the positively charged calcium ions into the cytosol, inducing calcium-evoked apoptosis (see Fig. 4).

3.4 Hydrophobicity

It has been shown that the hydrophobicity of NPs can affect the protein binding, cellular uptake, and cytotoxicity of NPs.78-82

The hydrophobicity and hydrophilicity of NPs can originate from the core or the functionalities of the NPs. In a recent systematic simulation study, Li et al.78 showed that changing the spikes of virus-like NPs (VLP) significantly altered the cellular uptake efficiency, while the effect of the core hydrophobicity of VLP was secondary. This study reported that VLP with hydrophobic or amphiphilic spikes were internalized more efficiently than that with hydrophilic spikes.

Generally, when keeping the other properties of NPs such as surface charge constant, their hydrophobicity has a positive trend with their cytotoxicity.74 Muthukumarasamyvel et al.81 controlled the hydrophobicity of dicationic amphiphilestabilized AuNPs by conjugating the dicationic functionality with different numbers and locations of H and OH groups. The authors observed increasing anticancer or cytotoxicity properties with an increase in the surface hydrophobicity of the NPs against A549 lung cancer cells.

(co) BY This article

Table 1 Recent studies highlighting the influence of the physicochemical properties of NPs on their cellular uptake and cytotoxicity

Property	Parameter	NPs	Cell lines	Uptake mechanism	Cytotoxicity	Highlights	Ref.
Size	(5 and 200 nm)	TiO ₂	A549	I	ROS generation ↑ Nutrient denletion	Smaller primary-sized NPs are more	38
Size	10, 40, 75 nm	Ag	BEAS-2B	Clathrin, caveolin/lipid raft, macropinocytosis and phagocytosis	Ag release (Trojan horse) DNA damage	10 nm NPs are the most cytotoxic	89
Size Shape Corona	Spheres (15 nm-NP1, 45 nm-NP2, and 80 nm-NP3), rods (33 × 10 nm-NR), stars (15 nm-NS)	Au	SMCC-7721 GES-1 4T1	Endocytosis (depends on corona)	Cell viability ↓	NS and NR are much more cytotoxic than the three spherical Au NPs. Cellular uptake in the order NP3 > NP2 \geq NP1 \sim NS	88
Size Shape Corona	Spheres (different sizes and coating), cubes, rods, prisms	Au	PC3	Endocytosis	Membrane damage Cell death	Increased uptake of smaller particles. AuNS are the most cytotoxic, followed by AuNPr, while both AuNR and AuNC are not toxic	68
Size Shape	Spheres (10 nm), flowers (370 nm), rods (41 nm), prisms (160 nm), stars (~240 nm)	Au	НеLа НЕК293Т	Endocytosis	Cell viability ↓	Au nanospheres and nanorods are more cytotoxic than star, flower and prism AuNPs	70
Shape	Rods $(L = 39 \text{ nm}, W = 18 \text{ nm})$, stars (215 nm) , spheres (6.3 nm)	Αu	hFOB 1.19 143B MG63 hTERT-HPNE	Endocytosis & phagocytosis	Mitochondrial dysfunction Membrane damage, apoptosis	Au nanostars are the most cytotoxic to the three cell lines while AuNPs spheres are the least cytotoxic	06
Shape	Spherical and needle-like	PLGA- PEG	HepG2 HeLa	Endocytosis	DNA damage, membrane damage, apoptosis	Spherical NPs have higher cellular uptake while needle-like NPs have greater cytotoxicity	91
Shape	Rods Flakes	$\mathrm{Al}_2\mathrm{O}_3$	Rat ASTs	Phagocytosis	ROS generation, inflammatory response, metabolism changes, apoptosis	Nanorods have significantly greater cytotoxicity than nanoflakes against rat astrocytes	92
Surface charge	Positive, negative, neutral	Au	НаСаТ	Endocytosis	ROS generation, oxidative stress, mitochondrial stress, apoptosis, or necrosis	All three NPs generated significant ROS levels, but only charged NPs caused mitochondrial stress. Charged NPs caused cell death through apoptosis, while neutral NPs caused it through necrosis	77
Surface charge	Positive & negative with different zeta potentials	Polymeric	L929	I	Cell viability ↓	Cationic NPs are more cytotoxic that anionic NPs. As absolute zeta potential increases, eytotoxicity increases	84
Surface charge Hydrophobicity	Positive/negative charge density and different hydrophobicity	Au	A549 HEK293	Endocytosis	ROS generation Apoptosis	Positive trend in the cytotoxicity of NPs over their surface hydrophobicity	82
Hydrophobicity	Three dicationic amphiphile-stabilized AuNPs	Au	A549	I	ROS generation Apoptosis	Positive trend in the cytotoxicity of NPs over their surface hydrophobicity	81

Review Nanoscale Advances

3.5 Surface functionalization

Changing the ligands on the surface of NPs will mostly tune the previous parameters (surface charge and hydrophobicity), which affects the protein corona, cellular uptake, and cytotoxicity of the NPs.77,83,84 However, the specific functionalities on the surface of NPs can be useful for targeting purposes. Here, overexpressed or unique receptors on the cell membrane are targeted by functionalizing the NPs with a complementary aptamer, protein, or antibody, which can specifically bind to the cell receptors. Tao et al.85 targeted cervical cancer cells through folic acid (FA)-poly(ethylene glycol)-b-poly(lactide-co-glycolide) blended NPs, which enhanced the efficacy of cancer chemotherapy through the targeted-delivery of anticancer drugs.

Lund et al.86 showed that AuNPs functionalized with 50% PEG-NH₂/50% glucose had an eighteen-fold higher internalization rate than NPs functionalized with either PEG-NH2 or glucose alone due to their different organization patterns. Alternatively, Yeh et al.87 studied the role of ligand coordination of two quantum dots (QDs) on their cytotoxicity. The authors found that monothiol-functionalized QDs had greater levels of cytotoxicity compared to dithiol-functionalized QDs in HeLa cell lines. However, the monothiol-functionalized QDs had a higher charge density, and thus it is difficult to tell if this tendency is solely related to the ligand coordination or charge density.

Studying the dependency of cellular uptake and cytotoxicity on a certain physicochemical property of NPs can be very

complex. For instance, changing their surface charge may lead to a change in hydrophobicity, hydrodynamic size, and protein corona. Furthermore, this may be done by changing the functionalities and coating of the NPs.73-77

Table 1 summarizes some recent studies exemplifying the effect of the physicochemical properties of NPs on their cellular uptake and cytotoxicity.

Cytotoxicity assessment

In vitro cytotoxicity of NPs is assessed using cell models. Although this assessment does not replace the in vivo evaluation of their cytotoxicity, it represents a screening bridge between the investigation of the quality and in vivo application of materials.56,93 Herein, we focus on the in vitro assessment of nanotoxicity. In the case of in vivo assessment, readers are encouraged to read the wholistic review by Kumar et al.94 Many in vitro assays are used to investigate or measure the cytotoxicity of NPs. These assays can be categorized to five main categories including cell viability and proliferation, ROS generation, cell stress, cell morphology phenotyping, and cell-NP uptake assays.56 Fig. 5 demonstrates some pathways of the effect of NPs on cells.

4.1 Cell viability and proliferation

Cell viability assays focus on investigating the cell metabolic and mitochondrial enzymes, such as lactate




Fig. 5 Various modes of action of NPs on cells.

Nanoscale Advances Review

dehydrogenase (LDH), an enzyme that regulates pyruvate and lactate levels through nicotinamide adenine dinucleotide (NAD) oxidation.93,95 Tetrazolium salts can react with the mitochondrial dehydrogenase enzymes. This reaction leads to the cleavage of the tetrazolium ring and conversion of these salts into a colored formazan form, which can be detected using colorimetry-spectroscopy. The detected activity of these enzymes is an indication of the cell viability. One of the most commonly used tetrazolium salts for assessing the cytotoxicity 3-(4,5-dimethylthiazol-2-yl)-2,5diphenyltetrazolium bromide (MTT) assay. 96 The other tetrazolium salts used include 3-(4,5-dimethylthiazol-2-yl)-5-(3phenyl)-2H-tetrazolium carboxymethoxyphenyl)-2-(4-sulfo (MTS), iodonitrotetrazolium (INT), and 4-[3-(4-iodophenyl)-2-(4nitrophenyl)-2H-5-tetrazolio]-1,3-benzene disulfonate (WST-1), which different to the previous salts, produce the watersoluble formazan. Other colorimetric/fluorimetric cytotoxicity assays are also used, for example, neutral red, trypan blue, lactate dehydrogenase (LDH), mitochondrial membrane potential (MMP), and Alamar Blue (resazurin) assays.

Many types of interference between NPs and cell viability assays have been reported. One way is the adsorption of the mitochondrial activity-related proteins on the NP surfaces. This may lead to the enzyme denaturation, giving false results of the cell viability profiles.⁹⁷ For instance, Stueker *et al.*⁹⁸ used

molecular dynamics simulation to investigate the effect of LDH enzyme binding on functionalized AuNPs. The authors observed that the dynamics of the side chains of the enzyme were largely constrained in all four active sites. Another way of interference is that the light absorbance spectra of the NPs can interfere with the absorption window of the assay, leading to false colorimetric measurements.99 For example, Díaz et al.100 reported that five NPs (magnetic iron/graphite, magnetite/silica, bare and poly(ethylene glycol)(PEG)-ylated silica, and magnetite/FAU zeolite) in culture medium after 72 h (in the absence of cells) showed absorbance at the same wavelength (525 nm) used in the MTT assay. This absorbance increases with the NP concentration, depending on their type. The third way of interference is that NPs may interact with the assay reagents. For instance, Hoshino et al. 101 reported that cysteamine-coated quantum dots catalytically reduced MTT to formazan without cellular metabolism taking place (see Fig. 6).

4.2 ROS generation and oxidative stress

Reactive oxygen species (ROS) are a type of unstable molecule (free radicals) that contain oxygen and can easily react with the other molecules in cells. The ROS include the superoxide anion $(O_2$, hydrogen peroxide (H_2O_2) , and hydroxyl radical (HO). ROS are normally produced by cells at certain levels to maintain




Fig. 6 Three ways of NPs interference with MTT cell viability test. (a) NPs in the absence of cells showed light absorbance at the same wavelength of MTT assay (525 nm in this study). The absorbance of the 5 NPs was measured at a concentration of 32 μ g mL⁻¹. Data was obtained from ref. 100 (b) NPs can catalyze the reduction of the MTT (or another test agent) to its colored (or fluorescent) form without the existence of cell enzymes. (c) NPs may adsorb and denature the cell enzymes that reduce the MTT dye to its colored form, giving false results.

Review

regular metabolism and homeostasis, which are considered as critical signalling molecules in cell proliferation and survival.99,102 However, they may be produced through interactions with exogenous sources such as NPs. If this event produces excessive ROS that the cellular antioxidant defense system (enzymatic antioxidants such as glutathione (GSH) peroxidases) cannot handle, oxidative stress is triggered. 102,103 This may lead to the destruction of organelles and biomolecules, including triggering membrane damage, lipid peroxidation, DNA damage, protein damage, apoptosis, necrosis, and inflammatory response, leading to many diseases such as cancer, diabetes, neurodegenerative, and cornea diseases. 103,104

NPs can generate ROS by acting as a catalyst in ROS generation reactions. For instance, Higashi et al.105 reported the catalytic generation of ROS by AuNPs and showed that this reaction can be controlled by changing conditions such as the type, concentration, and pH of the NP solution.

ROS detection can be performed by the direct measurement of ROS levels or the measurement of their oxidative damage or other outcomes. 106 Some direct methods for the detection of ROS are fluorescein-compound-based tests and electron paramagnetic resonance (EPR). The reactive fluorescein probes 2',7'difluorescein-diacetate (DCFH-DA) and dichlorodihydrofluorescein diacetate (H2 DCFDA) are non-fluorescent; however, when they are exposed to the cell cytosol enzymes, they get hydrolysed. Then, the cellular ROS oxidize them into a highly fluorescent compound, dichlorofluorescein (DCF), yielding an optical ROS concentration-dependent response, which can be measured using fluorescence microscopy or flow cytometry.107 Alternatively, indirect approaches for the detection of ROS include many assays that depend on the stimulated oxidative effect of the ROS. One approach is by measuring the enzymatic or non-enzymatic antioxidants levels. 106 Oxidative stress can also be assessed by measuring the oxidative damage of the cell biomolecules. These damaged biomolecules include proteins, lipids, and DNA and can be detected by measuring the protein carbonyl content, 108,109 malondialdehyde levels, 110,111 and 8-oxo-2'-deoxyguanosine (8-OdG) lesion, 112-114 respectively. Other genotoxicity assays include the comet, Ames, micronucleus, and chromosome aberration assays. 115

During the course of measuring NP-induced ROS generation and oxidative stress, NP-assay interferences may occur. 116,117 In colorimetric- and fluorimetric-dependent assays, NPs may interact with the final form of the dyes in a way that alters, by enhancing or reducing, the absorbance or fluorescence of the dye. For example, Aranda et al. 116 observed the quenching effect of several NPs on the dye fluorescence emission in the DCFH-DA assay, which was correlated with the cellular uptake of the NPs. The authors suggested a threshold concentration of NPs at which their oxidative effect can be detected, and they proposed that changing the experimental conditions can reduce this interference. Conversely, Pfaller et al. 117 reported the dye fluorescence enhancement of the DCFH-DA assay in the presence of Au or Fe₂O₃ NPs. This confirms that both scenarios (quenching and enhancement) may occur due to NP-probe interactions during colorimetric- and fluorimetric assays.

4.3 Inflammatory response

The inflammatory response induced by NPs in a cell line can be measured by detecting the produced inflammatory biomarkers. Macrophages and other cells release many cytokines, which play a crucial role in cell communication in the immune system by, for instance, promoting inflammation. Interleukins (ILs), such as IL-1β, IL-6, IL-8, and IL-10, in addition to other cytokines, such as tumor necrosis factor TNF-α and granulocytemacrophage colony-stimulating factor (GM-CSF), play a central role in inflammation regulation. The expression of these biomarkers can be assayed to determine the inflammatory response caused by NPs. ELISA (enzyme-linked immunosorbent assay) or western blotting, and electrophoretic mobility shift assays (EMSAs) or real-time polymerase chain reaction (RT-PCR) systems are used for the measurement of cytokines and the related genetic expressions, respectively. 118-123

NPs were reported to induce an inflammatory response in different cell lines. Many studies used conventional assays to measure this response.119-121 However, these assays can also interfere with NPs during the measurement of inflammatory response in cell lines. Some inflammatory cytokines were reported to be adsorbed on the NP surface, causing interference (Fig. 7a). Guadagnini et al.122 investigated the interferences of different NPs with some in vitro cytotoxicity assays. The authors reported that Fe₂O₃, TiO₂, and SiO₂ NPs significantly adsorbed




Fig. 7 Interference of NPs with ELISA assay. (a) NPs can adsorb different cytokines on their surface. (b) IL-8 were quantified by ELISA after 24 h of incubation with NPs at 75 μg cm⁻² after elimination of particles by centrifugation. Results (n = 6) are expressed as % of control (cytokine incubated in the absence of NPs). *Significantly different from the control (p < 0.05 ANOVA followed by Dunnett's test). (b) Is reproduced from ref. 122 with permission from Taylor & Francis. Copyright 2015.

Nanoscale Advances Review

IL-6, IL-8, and GM-CSF cytokines on their surfaces at different levels at a NP concentration of 75 μ g cm⁻². Fig. 7b shows that all the studied NPs adsorbed the IL-8 cytokine except PLGA-PEO NPs, which surprisingly increased the apparent level of cytokines, probably due to the stabilization of the peptides and their protection from proteolysis. In the case of other NPs, the level of adsorption depends on the NPs and the cytokine studied. OC-Fe₃O₄ NPs are the most cytokine-adsorbing NPs tested given that cytokines could not be detected in the supernatants. Furthermore, Piret et al. 123 observed a high inter-laboratory variability for the ELISA assay for IL1-β and TNF-α measurements and they suggested that testing of NP-cytotoxicity assay interferences should be always performed. Readers should kindly refer to ref. 122 for more information about the interference between different assay and NPs and some solutions to this problem.

4.4 Apoptosis and necrosis

Apoptosis is a programmed cell death pattern, 124 while necrosis is an unprogrammed cell death. 125 Both patterns of cell death can be an outcome of NP treatment. 126-131 Nickel ferrite (NiFe₂O₄),¹²⁶ TiO₂,¹²⁷ Fe₂O₃,¹²⁸ hydroxyapatite,¹²⁹ and Ag¹³⁰ NPs induced apoptosis in A549, BEAS-2B, ECV304, C6, and HepG2 cell lines, respectively. Alternatively, Reus et al. 131 reported dosedependent cell necrosis induced by SiO₂ NPs in BALB/c 3T3 cell line. Apoptotic cell death is mostly non-inflammatory, while necrotic cell death can be inflammatory. 132 Both pathways are extremes, and many cases are a complex combination of both. For instance, Kumar et al. 133 observed that AgNPs caused cell death in L-929 fibroblast cell lines in association with both necrosis and apoptosis. The cell death pathway is controlled by many parameters such as the surface charge, concentration, and exposure time of NPs. Schaeublin et al.77 reported that charged AuNPs caused cell death through apoptosis, while neutral AuNPs caused it through necrosis (see Fig. 4).

Many assays are used to detect apoptosis and necrosis. Phosphatidylserine (PS) migration to the extracellular side of the cell membrane and caspase activation into initiator and effector enzymes are two events that accompany apoptosis and can be used as markers to detect it. Externalized PS on the surface of the cell can be detected using fluorescein isothiocyanate (FITC)-labelled Annexin-V. Annexin-V specifically binds to the exposed PS on the cell surface in the early apoptotic cells, and then can be measured via flow cytometry or fluorescent microscopy. Alternatively, the membrane-impermeable propidium iodide (PI) dye exclusion assay is used for the identification of cellular necrosis. PI binds to DNA in the nucleus and stains it only when the cell membrane integrity is lost (which is an event that accompanies necrosis). Thus, a combination of the above-mentioned assays can determine the pattern of cell death.¹³⁴ For instance, Vafaei et al.¹³⁵ used the Annexin V-FITC/PI staining kit to study the apoptotic efficacy of zinc-phosphate NPs (ZnPNPs) against the MCF-7 breast cancer cell line. The untreated cells with NPs showed a live cell (Annexin V-FITC-/PI-) percentage of 98.6%. Conversely, after exposure to ZnPNPs, the apoptotic cell (Annexin V-FITC+/PI-)




Fig. 8 Evaluation of apoptosis and necrosis activities in MCF-7 cells using Annexin-V/PI staining. (Left) Untreated cells (control) and (Right) cells treated with ZnPNPs. Reproduced from ref. 135, with permission. Copyright @ 2020, Springer Science Business Media, LLC, part of Springer Nature.

ratio increased from 0.190% to 44.8% and the necrotic cell (Annexin V-FITC+/PI+) percentage increased to 1.34% (see Fig. 8).

Flow cytometry-based assays have negligible NP interferences. 133 Bancos et al. 136 reported that SiO2 NPs have low or no interference with flow cytometry assays. However, other colorimetric and fluorimetric-based assays face the same problems mentioned in the previous sections.

Metabolomics for the cytotoxicity assessment of NPs

In general, most studies on the cytotoxicity of NPs use the conventional (phenotypic) assays. However, many of these assays, as mentioned before, have been reported to interfere with the NPs because of their color, fluorescence, chemical activity, light scattering, etc. Thus, to precisely reveal the cytotoxicity of NPs, it is necessary to use a combination of more than two assays. This involves testing for NP interferences and eliminating them by changing experimental conditions or comparing the results of two similar tests, which is a complex and time-consuming process. However, many reports only used one or two cytotoxicity assays and ignored any potential interference with NPs.137 In addition, even though the conventional cytotoxicity assays can reveal that a certain cytotoxicity outcome happened, these assays are limited in terms of detecting the molecular information that caused this event.

The current toxicological assays need to be updated and new tools should be incorporated progressively in this field. 138 A more advanced and emerging approach to study the toxicity of particles is the "omics" technique, which is based on the change in epigenome, transcriptome, proteome, genome, lipidome, and metabolome profiles introduced by internal or external stimuli. In increasing number of studies are using this approach to investigate the in vitro and in vivo toxicity induced by NPs. The determination of new targets and biomarkers for NP toxicity is one of the strengths of the omics technique. Moreover, the omics technique has high sensitivity, which is useful because of the low levels of environmental exposure to NPs that sometimes cannot be detected using the conventional

assays.39 Another strength is that unlike the conventional assays, the omics technique has low or no interferences with NPs.39,122

In the field of toxicology, the most related omics discipline is metabolomics. 139 Metabolomics, one of the newest in the omics era, is an emerging field, which is broadly defined as the comprehensive measurement of all metabolites and lowmolecular-weight molecules in a biological specimen (tissues, cells, fluids, or organisms),40 and is one of the most powerful bioanalytical strategies that allow a picture of the changes of metabolites levels of an organism to be obtained during the course of a biological process either as a footprint (analysis of extracellular metabolites) or fingerprint (analysis of intracellular metabolites).41 The detailed analysis of low molecular weight compounds provided by nuclear magnetic resonance (NMR) spectroscopy or mass spectrometry (MS), besides the analysis performed by the powerful chemometric software (MetaboAnalyst),140 provides an accurate and quick detection and comparison of many types of chemical entities including carbohydrates, amino acids, nucleotides, lipids, steroids, fatty acids, and their derivatives, which are produced by cell metabolism.141

Currently, metabolomics is applied in many fields such as disease fingerprinting, biotechnology, environmental and plant research, toxicology and safety research, clinical medicine, and pharmacology. 139,142,143 Our group has been investigating the metabolic changes in serum, urine, and feces induced by different diseases such as lung cancer and diabetes, or other stimuli such as kidney transplant.144-146 Due to the non-invasive sampling in the metabolomic approach, the relatively low number of metabolites (compared to transcripts and proteins), and good level of knowledge about the role of most metabolites, metabolomics provides a well-grounded and precise methodology to investigate the biochemical effects and toxicity of NPs, 139,147 and it can present insight into the genotype and phenotype changes with a biological response. 48 Moreover, single-cell metabolomics is achievable today, making it possible to determine phenotypic heterogeneity among individual cells.149 Many cellular activities such as intercellular signal transduction, energy transfer, cell proliferation, and differentiation occur at the metabolite molecular level and are regulated by the presence and level of specific metabolites. Furthermore, metabolites are the end result of the expression of functional genome, transcriptome, and proteome (see Fig. 9). 150,151 This indicates that metabolomics can detect many NP cytotoxicity outcomes and reveal the molecular information behind these events even at low levels of NP exposure and with no interferences. Therefore, it is a great tool in nanotoxicology, which is being applied to reveal the effect and toxicity of NPs in many fields including environmental and agricultural fields152-154 and cancer research.155 Metabolomics can help in better understanding of the transition from in vitro to in vivo systems of NP




Fig. 9 Overview of the connection of the main omics-sciences: genomics, transcriptomics, proteomics, and metabolomics. Metabolomics represents the final output of cellular processes.

Nanoscale Advances Review

toxicity and its effect given that it is applied in both types of experiment. Furthermore, metabolomics can be combined with other omics techniques to provide a more comprehensive understanding of the effects of NPs on cells. 158-160

When comparing NMR- and MS-based metabolomics, generally, NMR has lower sensitivity than MS, and thus it is considered more suitable to analyze extracellular metabolites (exometabolome), which is done by the analysis of the cell culture media. Alternatively, the more sensitive MS techniques are more suitable for the analysis of relatively low levels of intracellular metabolites (endometabolome), especially when isolated from a limited number of cells. However, both analysis techniques are complementary and should be used simultaneously to maximize the metabolic window.

This emerging technique has not yet been widely applied for the investigation of NP cytotoxicity in *in vitro* systems and more research needs to be done on different NPs and cell lines. In this section, we focus on the metabolic changes induced by different NPs in different cells *in vitro*. The workflow of a metabolomics experiment is demonstrated in Fig. 10. This review does not go into detail on the workflow of metabolomics. In this case, for a detailed demonstration of how metabolomic workflows generate data, the reader is directed to read the following reviews and book chapters.^{39,160–166}

5.1 AuNPs

Gold nanoparticles (AuNPs) are very common in the biomedical field. AuNPs have many unique properties such as ease of synthesis, tunable size, ease of surface modification, surface plasmon resonance (SPR), and X-ray attenuation. This makes them the center of attention in many applications, including the growing field of nanomedicine, biosensors, targeted drug delivery, radiation therapy, photothermal therapy, biomedical imaging, and cancer diagnostics and therapeutics. The size of the size of

Metabolomics is used in several studies to assess the cytotoxicity of AuNPs and reveal their molecular information. Au nanorods (AuNRs) are one example of AuNPs that have strong absorption in the near-infrared spectral region and can be used in tumor thermal therapy (hyperthermia), and also in targeted tumor therapy. Wang *et al.*¹⁶⁸ observed, using conventional assays, that AuNRs have a unique influence on cell viability by causing the death of cancer cells (A549 cell line), while having negligible effect on normal cells (16HBE and MSC cell lines). The authors showed that AuNRs were released from the lysosome of cancer cells, and then translocated into the mitochondria, causing oxidative stress by the production of ROS. Alternatively, the normal cells had more intact lysosomes, and thus the AuNRs were not released in the cell cytoplasm.




Fig. 10 Metabolomics workflow for NMR- or MS-based metabolomics. DA: discriminant analysis; PCA: principal component analysis; PLS: partial least squares; and OPLS: orthogonal partial least squares.

However, the molecular information during this cellular translocation was unclear. Later, the same group, 169 used a metabonomic approach, a subset of metabolomics, 170 by applying ¹H NMR and multivariate data analysis, to study the metabolic change with time during the exposure of A595 and 16HBE cell lines to AuNRs. The authors found that both cell lines had intracellular disruption by the reduction of lactate levels and by causing oxidative stress. However, the normal cells resisted this oxidative stress by de novo GSH synthesis, unlike the cancer cells, which did not trigger this pathway, causing severe damage of their mitochondria (see Fig. 11). The metabonomic study further indicated the downregulation of nucleosides and nucleotides in the cancer cells, indicating cell death. Alternatively, the amino acid levels were upregulated in the normal cells, indicating cell stress. This study shows the usefulness of metabolomics in revealing the molecular information of the effect of NPs on cells, after conventional assays played the role of a general scanner for these effects.

Metabolomics can help in identifying biomarkers for NP cytotoxicity. For example, Xu et al. 171 investigated the potential harmful effects of AuNRs on male reproduction by studying the metabolic change in spermatocyte-derived cells (GC-2) and Sertoli (TM-4) cell line after exposure to 10 nM of AuNRs. Employing metabolomics, the authors observed a strong downregulation in glycine levels in TM-4 cells, while there was

no significant change in GC-2 cells. To identify what may accompany this reduction of glycine (potential biomarker), high content screening (HCS) and JC staining were used, and it was found that AuNRs decreased the membrane permeability and mitochondrial membrane potential of TM-4 cells. Moreover, the authors observed a disruption in the mRNA and protein levels of blood-testis barrier (BTB) factors using RT-PCR and western blot. Then, to confirm that glycine is a biomarker for these events, the authors repeated the experiments after adding glycine to the medium and noticed that the cells recovered from the previous harmful effects. This experiment reveals that glycine can be recognized as a biomarker to the changes in membrane permeability, mitochondrial membrane potential, and blood-testis barrier (BTB) factors in further similar experiments.

Huang et al. 172 observed that spherical AuNPs (20 nm) were not cytotoxic against the human dermal fibroblast (HDF) cell line. The authors combined bioinformatics with metabolomics to determine the molecular information of this toxicity resistance. Firstly, they detected that 29, 30 and 27 metabolites were differentially expressed in HDFs after 4, 8, and 24 h treatment with AuNPs, respectively. Among them, only six metabolites were determined to be key metabolites using bioinformatics techniques including expression pattern analysis and metabolic pathway analysis using MetaboAnalyst online tool. The key




Fig. 11 Summary of various metabolic responses of A549 (Right) and 16HBE (Left) cells to AuNR exposure. Metabolites in red or blue represent a significant increase or decrease in their levels, respectively, in the AuNR-treated groups compared with the non-treated groups. (This figure has been reprinted from ref. 169 with permission. Copyright © 2013, Elsevier Ltd).

Summary of AuNP-induced perturbation of metabolic pathways and their biological impact on different cell lines Table 2

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
AuNRs	15×58	CTAB	A549 16HBE	50 μΜ 12, 24, 48 h	¹ H NMR	Amino acid ↑ in 16HBE nucleosides and nucleotides ↓ in A549	Oxidative stress and cell death in A549	169
AuNRs	11×42	N/A	GC-2 TM-4	10 nM 24 h	GC-TOF-MS	Glycine ↓ in TM-4 Amino acid Metabolic disruption ↑	Cell and mitochondrial membrane disruption blood-testis barrier (BTB)	171
AuNRs	13.2×55.7	PSS PDDAC PEI	A549 16HBE	50 μΜ 12, 24, 48 h	¹ H NMR GC-FID/MS	Glucose ↑ Pyruvate ↑ Lactate ↑	Oxidative stress and cell death in A549	177
AuNRs	$14-16 \times 61-78$	Phospholipid PEG	MCF-7	0.05, 0.1 nM 4 h	LC-MS	Purine Pyrimidine GSH Amino acid J	Dysfunction in TCA cycle Reduction in glycolytic activity Imbalance of the redox state	178
Au	18	Citrate PSS PVP	HepG2	PSS, PVP - 0.25 nM Cit - 0.5 nM 3 h	LC/GC-MS (endo) ¹ H NMR (exo)	Uniform decrease in intracellular metabolite levels	Loss of cell membrane integrity	173
Au	30	N/A	Caco-2	59 μg mL ⁻¹ 72 h	LC-HRMS/MS	 5, 30 - amino acid ↑ 5 - TCA pathway ↓ 30 - glvcolvsis ↓ 	5 – small molecule biochemistry, cellular assembly and organization, and cellular growth and proliferation 30 – cellular degeneration and cell morphology	175
Au	5.67 5.90 5.90	CeO ₂ Chitosan Chitosan	HeLa HeLa	20 µg mL ⁻¹ 24, 48, 72 h 20 µg mL ⁻¹	¹ H NMR ALSOFAST – ¹ H, ¹³ C- HSQC of ¹³ C-labeled metabolites	CGO ₂ – pyruvate †, lactate † Chitosan – lactate † Chitosan – GSH †, UDP-NAG ↓, and alteration of glucose metabolism	Coop, and an analogic respiration Chitosan – aerobic respiration Chitosan – antioxidant effect	179
	5.65	CeO ₂		48 h		CeO_2 – no detected metabolites	CeO ₂ – antioxidant effect of this particle is lower and less related with the labelled glucose metabolism	
Au	5.65	CeO ₂	RBCs PMNs PBMCs	20 μg mL ⁻¹ 24 h	¹H NMR	Chitosan – reduced GSH ↑ CeO ₂ – amino acid ↑	Chitosan – antioxidant effect CeO ₂ – antioxidant effect of this particle is lower Lower AuChi toxicity compared with AuCeO ₂ PMN has higher pronounced antioxidant impact than PRMC	181
Au	ro	2-Mercapto-1- methylimidazole	SH-SY5Y	100 ng mL ⁻¹ 1, 2, 4, 6 h	¹H NMR	Glutamine, glutamate, leucine, tyrosine, PC/GPC and alanine	Oxidative stress Immune response Antioxidant mechanism (restore the initial state)	182
Au	20	Citrate	HDFs	200 μM 4, 8, 24 h	LC/MS	dSH ↑	Anti-oxidative stress mechanism	172

Review

metabolic pathway was identified to be the GSH pathway with GSH as the key metabolite. Subsequently, these results were verified and it was found that the increase in GSH levels after AuNP treatment may be the reason behind the toxicity resistance behaviour of the cells, given that GSH can trigger an oxidative stress protection mechanism that helps in avoiding cytotoxicity. This reveals that GSH can be considered as a biomarker for oxidative stress resistance.

Lindeque *et al.*¹⁷³ used MS metabolomics to study the effect of citrate-, poly-(sodium styrene sulfonate)-, and polyvinylpyrrolidone (PVP)-capped AuNPs on the intracellular metabolites of HepG2 cells. Surprisingly, after 3 h of treatment, a holistic depletion of intracellular metabolites was observed for all the capped AuNPs. Usually, metabolic changes result in the upregulation of the metabolite levels because of secondary pathways, clearance issues, and reduced enzyme functionality.¹⁷⁴ Firstly, the authors suggested that a loss of cell membrane integrity happened, but the exometabolomic data, measured using the NMR technique, was not consistent with this reasoning. Subsequently, they hypothesized that the AuNPs bind to the intracellular metabolites with or without replacing the surface coatings.

Gioria et al.175 combined proteomics and metabolomics to gain a further understanding of the effects of two sizes, i.e., 5 and 30 nm, of AuNPs on the human colon adenocarcinoma Caco-2 cell line. The proteome and metabolome are directly interconnected and influence each other given that the protein levels can change the metabolic profile of a cell system and vice versa. Genomics and transcriptomics were excluded from this study due to their restricted value given that they provide limited information about phenotyping. The authors used liquid chromatography high-resolution tandem mass spectrometry (LC-HRMS/MS) and two-dimensional gel electrophoresis (2DE) to obtain qualitative and quantitative data of deregulated metabolites and proteins, respectively. Subsequently, the data was combined and interpreted using systems biology analysis. After 72 h of exposure to AuNPs, 61 proteins and 35 metabolites in the cell extract were identified to be up-/ down-regulated. The internalization mechanism was found to be endocytosis due to the downregulation of the SH3GL1 and EAA1 proteins, which are involved in the endocytic pathway. The smaller-sized AuNPs caused a greater number of deregulated proteins and metabolites due to their higher internalization in the cells. Concerning metabolomics, the metabolite propionylcarnitine (C-3 carnitine) and glycine levels increased upon exposure to AuNPs, which indicates apoptosis. This study further reported the accumulation of GSH in both 5 and 30 nm AuNP-treated cells, which indicates that an antioxidative mechanism occurred as a self-defense system against oxidative stress. These results were confirmed using fluorescence microscopy analysis, where the over-expression of Annexin-V and nuclear fragmentation induced by AuNPs were evident, emphasizing that apoptosis occurred.

Omics technology together with complementary methods not only offer a promising tool in nanotoxicology to understand the molecular mechanisms of NP toxicity, but they also enhance the development and design of nano-drugs. For instance, Ali et al. 176 combined MS-based metabolomics and proteomics results through network analysis to better understand the molecular mechanism of AuNR photo-thermal therapy in the human oral squamous cell carcinoma (HSC-3) cell line. The results showed an upregulation in phenylalanine, which is considered an outcome of apoptosis pathways, indicating the good photo-thermal therapy efficiency of the AuNRs. Table 2 summarizes the studies that used the metabolomics technique to assess the effect of AuNPs in vitro on different cell lines.

5.2 AgNPs

Silver nanoparticles (AgNPs) have various interesting biological properties and are known for their well-reported antibacterial activity. They have a wide range of applications including cosmetics, textiles, and biomedical products. Also, their therapeutic application as antiviral and anticancer drugs is expected to be further expanded. Regarding the use of AgNPs as potential drug carriers for cancer therapy from proteogenomic and metabolomic perspectives, the reader is directed to the review by Raja *et al.* AgNPs have been shown to influence different cells causing apoptosis, lipid peroxidation, and DNA damage. 187–190

One of the advantages of metabolomics is that it is capable of detecting early biochemical events and metabolic changes even during the absence of a significant cytotoxic response by conventional assays. Carrola et al. 191 studied the effect of citratestabilized 30 nm AgNPs on the human epidermis keratinocyte (HaCaT) cell line after 48 h of exposure at two concentrations, *i.e.*, 40 $\mu g \text{ mL}^{-1}$ (close to IC₅₀ = 38.7 \pm 2.5 $\mu g \text{ mL}^{-1}$) and 10 μg mL⁻¹ (no significant cell viability loss). Using NMR-based metabolomics, the authors observed that most metabolic changes happened at the lower concentration, which allowed the detection of early biochemical events, including upregulated GSH-based antioxidant protection, downregulated tricarboxylic acid (TCA) cycle activity, energy depletion, and cell membrane modification. In a similar study, 192 NMR metabolomics was used to assess the metabolic effects of two types of coated AgNPs towards the human hepatoma (HepG2) cell line and significant metabolome changes were observed at a subtoxic concentration of AgNPs. These changes include energy production, antioxidant defence system, protein degradation, and lipid metabolism pathways, suggesting that the cells have metabolism-mediated protective mechanisms against AgNPs. In the third study by this group, 193 they investigated the effect of size and surface chemistry of AgNPs on the metabolic change caused in the HaCaT cell line. The authors used citrate-coated AgNPs with a diameter of 10, 30, and 60 nm, and 30 nm AgNPs coated with citrate, polyethylene glycol (PEG), or bovine serum albumin (BSA). It was found that the largest NPs and the PEG-coated NPs had the least impact on cell metabolism and viability, which is the expected tendency, as mentioned before in Section 3. Furthermore, Carrola et al.194 used NMR metabolomics to characterize the responses of RAW 264.7 macrophages to subtoxic concentrations of AgNPs (30 nm) and ionic silver (Ag+). They observed that the exposure to AgNPs caused a downregulation in intracellular glucose utilization, possibly

Nanoscale Advances Review

due to the reprogramming of the TCA cycle towards anaplerotic fuelling and production of anti-inflammatory metabolites. Also, an upregulation in the synthesis of GSH was observed, enabling the cells to control the ROS levels. In contrast, macrophages exposed to Ag⁺ at equivalent subtoxic concentrations showed reduced metabolic activity, lower ability to counterbalance ROS generation, and alterations in membrane lipids. This indicates that the ionic form of silver has a greater effect on the cells and is one of the sources of AgNP cytotoxicity.

Huang et al. 172 compared the effect of AgNPs and AuNPs, and showed that while AuNPs had no cytotoxicity, AgNPs induced grade 1 cytotoxicity after HDF cells were exposed to them for 72 h. Using metabolomics, the citrate cycle pathway was determined to be the key metabolic pathways in the AgNPtreated cells with malic acid as the key metabolite. Thus, the mechanism of AgNP cytotoxicity is by the upregulation of citric acid content, which indicated the inhibition of malic acid synthesis, influencing the production of ATP (mitochondrial dysfunction) and inhibiting cell proliferation, leading to cytotoxicity (see Fig. 12). Conversely, AuNPs were not cytotoxic due to the triggering of the antioxidant defence system by the upregulation of GSH. Kim et al.195 used high-resolution magic angle spinning (HR-MAS) NMR-based metabolomics to study the cytotoxicity of AgNPs against human Chang liver cells. The authors observed the depletion of GSH, lactate, taurine, and

glycine levels, while most amino acids, choline analogues, and pyruvate were upregulated by the AgNPs. It is probable that the downregulation of GSH induced the conversion of lactate and taurine to pyruvate.

The effect of AgNPs was also studied on non-mammalian cells such as yeast and unicellular alga. Babele et al. 196 studied the effect of 1.0 mg L⁻¹ of 50-100 nm-sized AgNPs, prepared using aqueous gooseberry extract, on yeast Saccharomyces cerevisiae cells. Untargeted ¹H NMR-based metabolomics revealed a several-fold increase or decrease in the levels of 55 different metabolites, including the ones involved in amino acid metabolism, glycolysis, and tricarboxylic acid (TCA) cycle, organic acids, nucleotide metabolism, urea cycle, and lipids metabolism. The authors noticed a reduced level of GSH, which indicates that oxidative stress occurred, leading to the strong cytotoxicity of AgNPs to the yeast cells. Qu et al. 197 investigated the effect of AgNPs on the performance of Chlorella vulgaris F1068 unicellular green alga in phosphorus assimilation (phosphorus removal by algae-based biotechnology). Using MS-based metabolomics, the authors observed the inhibition of algal assimilation. AgNPs disturbed the metabolic responses related to the phosphorus assimilation by reducing the levels of guanine, glutamine, alanine, and aspartic acid and increasing the levels of succinic acid. The NPs also inhibited phospholipid metabolism by the




Fig. 12 Comparison of the metabolic changes induced due to the interactions between AuNPs or AgNPs with HDFs cells. While AgNPs (Right) induced cytotoxicity in the HDF cells, the effect of AuNPs (Left) was suppressed by an antioxidant mechanism.¹⁷²

Table 3 Summary of AgNP-induced perturbation of metabolic pathways and their biological impact on different cell lines

	,							
NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
Ag	20	N/A	HDFs	200 µM 4, 8, 24 h	LC/MS	Citric acid ↑	Oxidative stress and cell death	172
Ag	30	Citrate	НаСаТ	10, 40 $\mu g \; m L^{-1}$ 48 h	¹ H NMR	GSH↑ TCA↓	Antioxidant protection Cell membrane modification Energy depletion	191
Ag	Cit – 29	Citrate	HepG2	Cit – 6.4, 11.0 μg mL ⁻¹	$^{1}\mathrm{H}\ \mathrm{NMR}$	TCA ↑	Metabolism-mediated protective mechanisms	192
	GS - 33	Biogenic (GS)		GS – 5.4, 14.0 μ g mL ⁻¹		Pyruvate use ↑	Protein degradation	
Ag	10 30 60	Citrate PEG BSA	НаСаТ	40 mg mL ⁻¹ 48 h	¹ H NMR	Glycolysis ↓ Energy production ↓	Oxidative stress Largest NP and PEG-NPs have the lowest impact of cell metabolism	193
Ag	30	Citrate	RAW 264.7	23.2, 35.3 µg mL ⁻¹ 24 h	¹H NMR	Intracellular glucose ↓ TCA GSH ↑ Anti-inflammatory metabolites ↑	ROS/RNS levels control	194
Ag	15	N/A	A549	38.6 $\mu g m L^{-1}$ 1, 6, 24 h	DIMS	Amino acid ↑ Clycolysis ↓ GSH ↓	Oxidative stress Apoptosis	199
Ag	5-10	N/A	Human Chang liver cell	N/A	HRMAS- ¹ H NMR	GSH, lactate, taurine, and glycine ↓ Amino acid, choline analogues, pyruvate ↑	Mitochondria-involved apoptosis DNA breaks Lipid membrane peroxidation Protein carbonylation	195
Ag	8.69	N/A	HT29	25 $\mu g m L^{-1}$ 12 h	UPLC Q-TOF MS	Nicotinic acid↑ ATP↓	Mitochondrial dysfunction Membrane damage Inhibit cancer proliferation	200
Ag	50-100	N/A	Yeast S. cerevisiae	$1.0~\mu \mathrm{g~mL}^{-1}$ $3~\mathrm{h}$	¹H NMR	Reduced GSH ↑ TCA ↓ Glycolysis ↓ Amino acid ↓ Urea cycle	Oxidative stress	196
Ag	15	N/A	Alga <i>C. vulgaris</i> F1068	0.09, 0.2 μg mL ⁻¹ 148 h	GC-TOF-MS	Glycerol-3-phosphate ↓ Myo-inositol ↓ Serine ↑	Oxidative stress Membrane damage Inhibition of the algal assimilation (66,2% reduction)	197
Ag	20	Citrate	Alga S. obliquus	1, 10, 100 $\mu g \; L^{-1}$ 148 h	GC-QTOF-MS	Carbohydrates; p-galactose, sucrose, and p-fructose ↑ Amino acids as glycine GSH ↑ TCA internution	Growth inhibition Cell wall damage Oxidative stress	201
Ag	20	Citrate	Alga P. malhamensis	40.7, 1000 $\mu g L^{-1}$ 2, 24 h	TC-MS	Amino acid	Photosynthesis and photorespiration disruption Oxidative stress	202

Table 3 (Contd.)

NP	Size [nm]	NP Size [nm] Coating Cell	Cell	Dose/exposure time	Analytical platform	Dose/exposure time Analytical platform Perturbed metabolic pathway	Biological effect	Ref.
Ag	23.4	PVP	Cyanobacteria M gernoinosa	$0.075,0.15~{ m mg}{ m L}^{-1}$	TC-MS	Nucleotides Fatty acids Amino acids; arginine and proline Cellular stress	Cellular stress	203
				96 h		ndole alkaloid biosynthesis ↑ Phospholipid metabolism ↑	ROS generation Damage to photosynthesis and	
Ag.	I	N/A	Chlorella mrenoidosa	0.5, 5, and $10 \text{ mo } \text{L}^{-1}$	TC-MS	Amino acid	cenular memoranes Oxidative stress	198
			Fyriance	0-72 h 1 or 3 repeated exposure		Carbohydrate	Membrane damage	

downregulation of glycerol-3-phosphate and myo-inositol and upregulation of serine. Furthermore, GSH metabolism was affected by the NPs, which induced oxidative stress in the alga cells (upregulation of glycine). Cao *et al.*¹⁹⁸ showed that the effect of AgNPs on *Chlorella pyrenoidosa* can be altered by the number of repeated exposures. In this study, NP single exposure had a greater impact on the *C. pyrenoidosa* metabolome than repeated exposure. Table 3 summarizes the studies that used the metabolomics technique to assess the effect of AgNPs *in vitro* on different cell lines.

5.3 TiO₂ NPs

Micro-titania (titanium oxide, TiO2) particles are known as biologically inert in humans, enabling their use in many products such as cosmetics and pharmaceuticals. 204,205 Nano-titania (TiO₂ NPs) are also used as additives in many products such as sunscreen products, paints, printing ink, rubber, paper, sugar, cement, toothpaste, film, biomedical ceramics, implanted biomaterials, antimicrobial plastic packaging, and self-cleaning sanitary ceramicss. 206 However, TiO2 NPs can enter the body via inhalation, ingestion, and dermal contact and they have been shown to exert significant toxic effects, such as cell metabolic change,206 chronic pulmonary inflammation,207 and proinflammatory effects in cells.208 Raja et al.209 reviewed the microenvironmental influence of TiO2 NP-induced mechanical stimuli on tumor cells and showed using the omics analysis that the exposure of cancer cells to TiO₂ NPs caused gene mutations, protein alterations, and metabolite changes.

Chen et al.210 observed mitochondrial dysfunction caused by TiO₂ NPs in a macrophage (RAW) cell line and primary mouse bone marrow-derived macrophages (BMDM) using a combination of metabolomics, lipidomics, and proteomics. The targeted UPLC-MS-based metabolomic analysis revealed a significant upregulation in the production of COX-2 metabolites including PGD2, PGE2, and 15dPGJ2, indicating an inflammatory response in macrophages. The authors also used GC-MS-based metabolic flux analysis, which is a technique that uses MS to track the fate of stable isotope tracers (e.g., ¹³C-glucose and ¹⁵Nglutamine), allowing the investigation of the contribution of specific metabolic pathways to the prevailing levels of specific metabolites, 211 to measure the metabolic flux in the tricarboxylic acid (TCA) cycle using 13C-labelled glutamine. They observed a downregulation in TCA cycle metabolism and ATP production caused by significant mitochondrial dysfunction after the exposure of macrophages to TiO₂ NPs. In a similar study, Tucci et al.206 studied the response of the human keratinocyte HaCaT cell line after exposure to 10-100 nm TiO2 NPs and found that the NPs were only present in the phagosomes of the cells without their internalization in any other cytoplasmic organelle. Specifically, "268" metabolites were detected using GC/LC-MSbased metabolomics, of which 85 metabolites were found to be significantly altered at 100 µg mL⁻¹ dose of NPs. As stated in other studies, TiO2 NPs have shown significant and rapid effects on mitochondrial function by altering energy metabolism and anabolic pathways. However, they did not affect the cell cycle phase distribution or cell death.

Review Nanoscale Advances

Jin et al. 212 used GC/TOFMS-based metabolomics to study the metabolic changes in L929 cells and their corresponding culture media induced by 5 nm-TiO2 NPs. At concentrations higher than 100 μg mL⁻¹, the NPs caused a depletion in the cellular carbohydrate metabolism (the major biochemical metabolism pathway) after causing energy metabolism disruption, pentose phosphate pathway inhibition, nicotinamide metabolism block, mitochondria damage, and oxidative stress activation. Bo, Jin, Liu et al.213 again used GC/TOFMS-based metabolomics to study the change in amino acid levels in L929 cells after they were exposed to TiO2 NPs. The study revealed that seven metabolic pathways among the regulated pathways were significantly altered including 12 amino acids, i.e., L-α-alanine, β-alanine, glycine, L-aspartate, L-methionine, Lcysteine, glutamate, L-pyroglutamate, L-asparagine, L-glutamine, S-adenosyl methionine, and L-lysine.

In dental science, the use of TiO₂ NPs as an additive to glass ionomer cements is known to improve their mechanical and antibacterial properties. However, the study by Garcia-Contreras et al.214 showed that these NPs may induce proinflammation in human gingival fibroblast (HGF) cells. Nevertheless, the molecular mechanism of the pro-inflammatory action of TiO2 NPs on these cells was still unclear. MS metabolomics was used to reveal the mechanism of this proinflammatory action by the treatment of HGF cells with IL-1b alone or in combination with TiO2 NPs.215 A total of 109 metabolites was successfully identified and quantified by CE/ TOFMS. Most amino acids levels were downregulated at high concentrations of TiO2 NPs, while ophthalmate, α-aminoadipate, kynurenine, and β-alanine were upregulated. The activation of the urea cycle, polyamine, S-adenosylmethionine, and GSH synthetic pathways was stronger than that of the other pathways. The intracellular levels of urea cycle metabolites were downregulated significantly in the presence of both IL-1b/TiO₂

NPs. In conclusion, ornithine was downregulated, which led to an immediate decline in putrescine. That latter is used to synthesize spermidine, which has anti-inflammatory properties. Thus, the reduction of this polyamine level accelerated the inflammation in HGF cells upon exposure to a combination of IL-1b/TiO2 NPs.

Kitchin et al.216 studied the effect of four different TiO2 NPs (in addition to two CeO2 NPs) on human liver HepG2 cells. Using LC/GC MS-based metabolomics, five out of the six NPs were found to cause a significant downregulation in GSH concentration. The authors observed a decrease in the GSH system in GSH precursors (glutamate and cysteine), GSH itself, and GSH metabolites (the gamma-glutamyl condensation products, glutamine, alanine, valine, 5-oxoproline, and cysteine-GSH). Among the 265 metabolites detected, the reduction in GSH was the largest deregulation. This indicates that the NPs are acting via an oxidative stress mode, which is a consistent biochemical effect of NPs.

Metabolomics can help to better understand the transition from in vitro to in vivo systems of NPs toxicity given that it can be applied in both types of experiment. For example, Cui et al.217 employed LC-MS-based metabolomics to investigate the effect of four metal oxide NPs, including TiO2 NPs, in vitro on human bronchial epithelial (BEAS-2B) cell line, and in vivo on mouse model after lung exposure. Their study showed that in vitro metabolomic findings can effectively reveal the biochemical effects in vivo, and that LC-MS-based metabolomics is sensitive enough to detect the tiny metabolomic changes when conventional cytotoxicity assays cannot detect any significant effect. Fig. 13a shows the workflow of this study. BEAS-2B cells were exposed to the four studied NPs, and then the metabolomics experiment was performed in vitro. This was followed by validation in vitro by enzymatic assays, in vivo using a mouse model after lung exposure to respective NPs, and finally by cellular




Fig. 13 Untargeted metabolomic analysis was used to reveal the effect of exposure of two different doses (12.5 and 25 μ g mL⁻¹) of ZnO, SiO₂, TiO₂, and CeO₂ NPs on the metabolism of human bronchial epithelial cells (BEAS-2B). (a) Schematic diagram of the study workflow. (b) Hierarchical cellular stress responses based on metabolomics and functional assays. The cells were maintained in the healthy state at the tier 1 stage. At an intermediate level of cellular stress (tier 2), the exposure to SiO₂, TiO₂, and CeO₂ NPs altered several metabolic pathways and induced inflammation. At a high level of cellular stress (tier 3), ZnO NPs significantly affected toxicity and DNA damage related metabolic pathways. Only a short list of significantly altered pathways is presented due to the limited space. Adapted with permission from ref. 217 Copyright 2019, the American Chemical Society.

Open Access Article. Published on 30 2023. Downloaded on 01-11-2025 11:40:01.

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Table 4 Summary of TiO₂ NP-induced perturbation of metabolic pathways and their biological impact on different cells

)	,	2)						
NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
TiO_2	10	N/A	RAW264.7	10, 100 $\mu \mathrm{g} \; \mathrm{mL}^{-1}$	UPLC-MS	$COX-2$; PGD_2 , PGE_2 , 15 d- PGJ_2	Mitochondrial dysfunction	210
			ВМDМ	24 h	GC-MS for ¹³ C-labelled glutamine (flux analysis)	TCA↓ ATP↓	Inflammatory response	
${ m TiO}_2$	10–100	N/A	НаСаТ	$5, 50, 100 \ \mathrm{\mu g \ mL}^{-1}$ 24 h	GC-MS LC/MS/MS	Acetyl-CoA GSH ↓	Oxidative stress Mitochondrial function	206
						Acetyl-carnitine Glycolysis ↓ Pentose phosphate pathway ↓ Nucleotide ⊥	disruption	
${ m TiO}_2$	N/A	N/A	L929	$30~\mu \mathrm{g~mL}^{-1}$	GC-TOF-MS	Carbohydrate metabolism and TCA ↓	Metabolism changes	150
				48 h		Glycolysis↓ Fatty acid↓ Purine metabolism⊥		
${ m TiO}_2$	ις	N/A	L929	100, 200 $\mu\mathrm{g~mL}^{-1}$	GC-TOF-MS	Carbohydrate metabolism and TCA ↓	Mitochondria damage	212
				48 h		Pentose phosphate pathway↓ Nicotinamide metabolism block	Oxidative stress	
TiO_2	23	N/A	L929	$100~\mathrm{\mu g~mL^{-1}}$	GC-TOF-MS	Amino acid ↑	Oxidative stress	213
				48 h		Carbohydrate↑ Nucleotide↑	Energy damage Inhibition of DNA and RNA synthesis	
TiO_2	18	N/A	HGF	0.2, 0.8, 3.2 mM + II8 3 ng mI1	CE-TOF-MS	Amino acid ↓	Inflammation	215
				24 h		Urea cycle ↓ GSH synthesis ↓ Polyamine ↓		
TiO_2	8–142	N/A	HepG2	3, 30 $\mu \mathrm{g} \ \mathrm{mL}^{-1}$	GC-MS	GSH↓ Fafty acid	Oxidative stress	216
TiO_2	21	N/A	BEAS-2B	$12.5, 25 \mathrm{\mu g mL^{-1}}$	LC-MS	S1P pathway ↑	Inflammation	217
				6 h		Folate cycle ↑ Fatty acid oxidation GSH ↓ Lipid ↑	Oxidative stress DNA damage	
${ m TiO}_2$	21	N/A	RLE-6TN	$0.1, 1, 10 \text{ µg cm}^{-2}$ 24 h	HPLC-MS/MS FIA-MS/MS	Lipids↑ Amino acid Biogenic amines	Oxidative stress	231
TiO_2	21	N/A	RLE-6TN NR8383	1, 2.5, 5, 10, μg cm ⁻² 24, 48 h	HPLC-MS	Amino acid Biogenic amine Lipid	Oxidative stress	230
TiO_2	21	N/A	RLE-6TN	0.1 –50 ${ m \mu g~cm^{-2}}$	HPLC-MS	esh	Oxidative stress	232

Table 4 (Contd.)

NP	Size [nm] Coating	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
			NR8383	24, 48 h		Amino acid Biogenic amine Linid		
TiO_2	42	N/A	E. coli	$2.5-10 \; \mathrm{\mu g \; mL^{-1}}$ 3 h	GC-TOF-MS	Polyamine; putrescine ↓ Amino acid: glvcine ↑	Oxidative stress	233
TiO_2	8–37	N/A	E. coli	10, 100 ppm 3 h	¹H NMR	TCA † Amino acid † ATP † Fatty acid † Polvamine: putrescine †	Cell membrane damage Oxidative stress	234
TiO_2	30	N/A	P. polycephalum macroplasmodium	9, 15 18 mg m L^{-1} 72 h	GC-MS	Amino acid GSH Nucleotide Polyamine	Oxidative stress ROS imbalance	235
TiO ₂ /CdS/ 10/30/40 ZnS	10/30/40	N/A	Bacillus subtilis	$0.1953,3.125~{ m mg~mL}^{-1}$	LC/MS	Carbonyarac Lipid ↓ Biomolecules synthesis ↓ ATP ↓	Membrane damage ROS generation	236

function assays. The TiO2 NPs significantly altered the metabolic pathways of sphingosine-1-phosphate, fatty acid oxidation, folate cycle, inflammation/redox, and lipid metabolism, inducing inflammation. In addition, this effect was dosedependent for some metabolites. Fig. 13b shows the altered metabolites and effect of the four studied metal oxide (MO_r) NPs and their numbers, respectively.

Metabolomics is also applied in many in vivo nanotoxicity studies.218,219 For instance, Chen et al. performed three recent studies of TiO2 NP toxicity in vivo using MS-based metabolomics, once in rats by feces metabolite analysis,220 and then screened for urine221 and serum222 biomarkers in human workers exposed to these NPs in factories. This group also performed another metabolomics study using rat serum after subchronic oral exposure of TiO2 NPs.223 Han et al.224 used MSbased metabolomics to study the influence of TiO2 NPs on the fecal metabolome in rats after oral administration for 90 days. Åslund et al.225 used NMR-based metabolomics to assess the effects of 5 nm-TiO2 NPs on Eisenia fetida earthworms and observed metabolic changes related to oxidative stress. Eight years later, Zhu et al. 226 used transcriptomics besides metabolomics to investigate the same effect of TiO2 NPs on the same earthworm and noticed that the antioxidant system and metabolic profiles of the earthworms were significantly affected. Ratnasekhar et al. 227 used MS-based metabolomics to investigate the effects of TiO2 NPs on the soil nematode Caenorhabditis elegans. The results indicated the disruption of the tricarboxylic acid (TCA) cycle, arachidonic acid metabolism, and glyoxylate dicarboxylate metabolism pathways. For more about the in vivo metabolic effects of NPs including Ag, TiO2, and carbon-based NPs on organisms (plants, aquatic, and terrestrial invertebrates), the reader is kindly referred to the chapter by Farré and Jha.165

Metabolomics reveals the global responses that cannot be observed by conventional toxicity endpoints, leading to an effective assessment of the effects of NPs in the environment, in vivo, and in vitro. Metabolomics has also been used to reveal the metabolite corona that is surrounding TiO2 NPs. 228,229 Table 4 summarizes the studies that used the metabolomics technique to assess the effect of TiO2 NPs in vitro on different cells.

5.4 SiO₂ NPs

The annual global production of SiO₂ NPs is reported to exceed 1.5 million tons, making SiO2 NPs one of the most widely used NPs in the industrial manufacturing, drug delivery, cancer therapy, and biotechnological fields.39 This widespread is due to their biocompatibility, stability, and other unique properties compared with their bulk.237

Although SiO2 NPs have been shown to have different cytotoxic effects on cells, the molecular mechanism of this cytotoxicity still needs to be explored using novel analytical techniques, such as metabolomics. Huang et al.238 used MSbased metabolomics to reveal the molecular information of the effect of SiO₂ NPs on the human fetal lung fibroblast MRC-5 cell line. The authors observed NP dose-dependent changes in the metabolic profiles of the cells. As the dose increased, there

Nanoscale Advances Review

was a downregulation in the amino acid and GSH levels together with an upregulation in urea and phospholipid concentrations, causing oxidative stress and energy metabolism disturbance. Feng et al.237 used NMR-based metabolomics to study the effects of 0.01 or 1.0 mg mL⁻¹ of hydrophilic SiO₂ NPs on the human cervical adenocarcinoma (HeLa) cell line. They studied both the intracellular and extracellular metabolome changes. In the early stage of NP exposure, no clear dose-effect of the HeLa cell metabolome was observed, which implied that the cellular stress-response regulated the metabolic variations in the HeLa cells. Afterwards, the NPs induced cell membrane modification, catabolism of carbohydrate and protein, and a stress response. The toxicological effects induced by highdosage SiO2 NPs could be derived from the elevated levels of ATP and ADP, the utilization of glucose and amino acids and the production of metabolic end-products such as glutamate, glycine, lysine, methionine, phenylalanine, and valine. Irfan et al.239 used conventional assays and NMR-based extracellular metabolomics to study the effect of fumed SiO2 NPs on human lung A549 cells. The authors observed an upregulation in the extracellular glucose, lactate, phenylalanine, histidine, and tyrosine levels in a time- and NP dose-dependent manner. There was also an increase in intracellular ROS and cell membrane damage at 4 h and a loss of cell viability after 48 h observed by conventional assays.

A few metabolomics studies compared the in vitro and in vivo outcomes of SiO2 NP treatments. For instance, Chatterjee et al.240 used NMR-based untargeted-metabolomics to study the effect of amorphous SiO2 NPs on the human hepatoma HepG2 cell line and mice liver (Fig. 14). Firstly, this study determined the altered metabolites in the cells and mice liver using OPLS-DA analysis (Fig. 14a and d, respectively). Subsequently, the selected significantly altered metabolites were determined (Fig. 14b and e), followed by pathway analysis using the MetaboAnalyst 3.0 software (Fig. 14c and f). In both in vitro and in vivo systems, the perturbation of GSH metabolism and the depletion of the GSH pool were detected after aSiO2 NP treatment. Moreover, the in vitro results were further supported by the in vivo data, specifically for metabolite profiling and pathway analysis, were there were 8 common altered metabolic pathways in the two systems. This study revealed that the major causes of aSiO2 NP-mediated hepatotoxicity were the suppression of GSH metabolism and oxidative stress. In a similar study, Bannuscher et al. 230 studied the responses of rat lung epithelial cells (RLE-6TN) and alveolar macrophages (NR8383) (in vitro) to four well-selected SiO2 NPs, differing in structure, size, and surface charge, and compared the results to in vivo responses in rat lung tissues. The authors observed a cell-specific time- and concentration-dependent changes in vitro and identified several biomarker candidates such as Asp, Asn, Ser, Pro, spermidine, putrescine, and LysoPCaC16:1 in vitro, and then verified them in vivo.

It was proven that SiO₂ NP exposure inevitably induces damage to the respiratory system, however, knowledge of its mode of action and metabolic interactions with the cells is limited. Zhao et al.241 performed a study to reveal the molecular information of the metabolic responses of the lung bronchial epithelial BEAS2B cell line after SiO2 NP exposure, using MSbased metabolomics. They revealed that even with low cytotoxicity, SiO2 NPs still caused global metabolism disruption. Specifically, five metabolic pathways were significantly perturbed; in particular, oxidative stress, as confirmed by GSH depletion, mitochondrial dysfunction-related GSH metabolism, and pantothenate and coenzyme A (CoA) biosynthesis. The identified key metabolites were GSH, glycine, beta-alanine, cysteine, cysteinyl-glycine, and pantothenic acid. Oxidative DNA damage and cell membrane disintegration were detected by observing elevated 8-oxo-2'-deoxyguanosine (8-OdG) and decreased phospholipids levels.

Several studies compared the effect of SiO₂ NPs on cells to other NPs using metabolomics and other omics techniques. For example, Karkossa et al.231 used targeted metabolomics and global proteomics to compare the effect of SiO2 NPs with different particle sizes, surface charges, and hydrophobicity to the effect of TiO2, graphene oxide (GO), phthalocyanine blue, phthalocyanine green, and Mn₂O₃ NPs on RLE-6TN alveolar epithelial cells. Alternatively, Cui et al.217 used MS-based metabolomics to reveal the significantly altered metabolites and metabolic pathways in human bronchial epithelial cells and a mouse model exposed to four different types of metal oxide NPs (SiO₂, ZnO, TiO₂, and CeO₂) at both high (25 μ g mL⁻¹) and low (12.5 μ g mL⁻¹) doses (see Fig. 13). Table 5 summarizes the studies that used metabolomics technique to assess the effect of SiO2 NPs in vitro on different cells.

5.5 ZnO NPs

Zinc oxide NPs are gaining increasing attention due to their unique properties, especially their optical and electronic properties. Also, they can be prepared using a variety of methods and in a range of different morphologies.246 This makes them the third highest global production volume among metalcontaining NMs247 and excellent for a broad range of applications, including optoelectronic devices (light-emitting diodes (LEDs), laser diodes, solar cells, and photodetectors), electronic devices (transistors),246 and active compounds in sunscreens, drug delivery, biomedical engineering, food additives, and cosmetics.248 It was shown that human exposure to these engineered NPs can cause health problems for both consumers and industry workers, making it important to further investigate in their toxicity and improve their safety when they are used and produced.217,249

The respiratory tract is the primary route of exposure to airborne NPs such as ZnO. Thus, it is common to use the bronchial epithelial BEAS-2B cell line as an in vitro model to study the toxicity of these NPs. For instance, Lim et al.247 used this cell line to perform an MS-based metabolomics study to reveal the effect of ZnO NPs on the respiratory system. The authors revealed ROS-mediated cell death associated with mitochondrial dysfunction and interference in regulating energy metabolism. This was concluded after observing a significant decrease in the levels of amino acids (valine, tryptophan, lysine, proline, threonine, glycine, serine, glutamic acid, and aspartic acid) and TCA intermediate metabolites




Fig. 14 Global metabolomics and pathway analysis in aSiO₂ NP-exposed HepG2 cells (a-c) and Institute for Cancer Research (ICR) mice (d-f). OPLS-DA score plot from the NMR spectra of metabolomes from HepG2 cells treated with 100 mg L^{-1} aSiO₂ NPs for 24 h (a) and from ICR mice liver treated with 50 mg kg^{-1} aSiO₂ NPs for 24 h. Selected significantly altered global metabolites level in HepG2 cells (b) and in ICR mice liver (e) after treatment with the NPs. Pathway-based enrichment analysis performed by MetaboAnalyst 3.0 with significant altered metabolites (>1.5 fold) in HepG2 cells (c) and ICR mice liver (f) after treatment with the NPs. This figure has been reproduced from ref. 240 with permission from Elsevier B.V., Copyright 2018

(citrate) (Fig. 15a). These results indicate that ZnO NPs can be seriously harmful to human health if they were inhaled.

Although Zn is a key micronutrient for plants, a high dose of this metal is toxic to plants either in the nano or other forms.

Salehi et al.250 used UHPLC-QTOF metabolomics to study the effect of ZnO NPs and bulk ZnSO4 on bean plants (Phaseolus vulgaris L). The results indicated the unique NP-related toxic effects of ZnO in beans compared to the ionic forms of Zn. Two

Open Access Article. Published on 30 2023. Downloaded on 01-11-2025 11:40:01.

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Table 5 Summary of SiO₂ NP-induced perturbation of metabolic pathways and their biological impact on different cells

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
SiO_2	45	FBS Tris(2,2'-bipyridyl)- dichlororuthenium(11)	MRC-5	$2.5, 10, 40, 80 \ \mathrm{\mu g \ mL}^{-1}$ 24 h	GC-MS LC-MS	Amino acid ↓ GSH↓	Oxidative stress Energy metabolism disturbance	238
		hexahydrate				Urea cycle ↑ Phospholipid ↑	Membrane damage Cell death	
SiO_2	20	N/A	НеГа	$0.01, 1~\mathrm{mg~mL}^{-1}$	HRMAS ¹ H NMR	Lipid ↑	Catabolism of carbohydrate and	237
				6, 48 h		Carbohydrate and protein metabolism Amino acid	protein Cell membrane modification Oxidative stress	
SiO_2	7–14	N/A	A549	10, 25, 50, 100 μg mL ⁻¹ 4, 12, 24, 48 h	¹ H NMR	Glucose † Lactate † Dhenylolnine †	Oxidative stress	239
SiO_2	57.7	N/A	BEAS-2B	2 , 10, 50 $\mu\mathrm{g}~\mathrm{mL}^{-1}$ 24 h	UPLC-MS	GSH ↓ 8-OdG ↑	Oxidative stress DNA damage	241
						ATP \\ NRF2 \\ Pantothenate and CoA	dysfunction Cell membrane dis- integrity	
SiO_2	16	N/A	BEAS-2B	12.5, 25 µg mL ⁻¹ 6 h	LC-MS	Stopy mit. Stopy To S	Inflammation Oxidative stress DNA damage	217
SiO_2	10-30	N/A	HepG2	30 $\mu g m L^{-1}$ 72 h	GC-MS LC-MS	Lipid↑ GSH ⊥	Oxidative stress	242
SiO_2	20–50	N/A	НерG2	$100 \mathrm{\mu g \ mL}^{-1}$ 24 h	¹ H NMR	GSH↓ Acetate↓ Choline↑ Biotin merabolism	Oxidative stress	240
SiO_2	15, 15, 8, 40	None, amino, none, none	RLE-6TN NR8383	1, 2.5, 5, 10, 50 μg cm ⁻² 24, 48 h	HPLC-MS	Amino acid Biogenic amine Linid	Oxidative stress	230
SiO_2	15, 15, 15, 40, 8, 8, 8	None, phosphate, amino, none, none, 2% TMS, 3% TMS	RLE-6TN	$10 \ \mathrm{\mu g \ cm^{-2}}$ 24 h	HPLC-MS/MS FIA-MS/MS	Amino acid Biogenic amine Lipid	Oxidative stress Apoptosis	231
SiO ₂	8, 15, 40	N/A	NR8383	$2.5, 5, 10 \ \mu \mathrm{g} \ \mathrm{cm}^{-2}$ $24 \ \mathrm{h}$	HPLC-MS	Amino acid Biogenic amine Phosphatidylcholine	Oxidative stress Mitochondrial dysfunction DNA damage Cell death	243

Review

		Adenosine triphosphate 🔱						
	Cell viability	↑ HSĐ						
	Cell death	p-Glucose ↑		24 h				
245	Oxidative stress	Adenine ↑	UHPLC-MS	$25, 50 \ \mathrm{\mu g \ mL^{-1}}$	L-02	N/A	58	\mathcal{O}_2
		TCA						
		ATP ↓						
		Lactate ↑		24, 48, 72 h				
244	Inflammation	Glycolysis ↑	1 H NMR	10, 500 $\mu g m L^{-1}$	RAW 264.7	N/A	100 - 125	\mathcal{O}_2
		Lipid						
		Biogenic amine						
		Amino acid		24, 48 h	NR8383	none		
230	Oxidative stress	GSH	HPLC-MS	$1-50 \ \mathrm{\mu g \ cm^{-2}}$	RLE-6TN	None, amino, none,	8, 15, 15, 40	\mathcal{O}_2
Ref.	Biological effect	pathway	platform	Dose/exposure time	Cell	Coating	Size [nm]	<u></u>
		Perturbed metabolic	Analytical					

similar studies of the effect of ZnO NPs have been done on tomato and cucumber.^{251,252} Wan et al.²⁵³ performed a metabolomics analysis to reveal the effect of ZnO NPs on salt tolerance in the Sophora alopecuroides plant. Moreover, He et al. 254 elucidated toxicodynamic differences at the molecular scale between ZnO NPs and ZnCl₂ in Enchytraeus crypticus, a model species in soil ecotoxicology, using non-targeted metabolomics. They found that the number of altered metabolites after Zn²⁺ exposure was larger than the number of altered metabolites after ZnO NP exposure, indicating the higher toxicity of the Zn ionic form (Fig. 15b and c). For more information about nanotechnology in agriculture and the effect of metallic-, metal oxide-, and carbonbased-NPs on plants, the reader is advised to read the review by Paramo et al.255 and review by Majumdar et al.256

The toxic effects of a NP may be reduced by applying coexposure with another NP. For instance, Wu et al.257 studied the combined effects of graphene oxide (GO) and ZnO NPs on human A549 cells using NMR-based metabolomics. PLS-DA analysis showed that the control and GO-alone exposure groups overlapped, indicating a low effect of 10 mg L^{-1} GO on the metabolome profiles. In contrast, ZnO NP-alone exposure significantly altered the metabolome profiles in A549 cells. A total of 14 altered metabolites was shared in the ZnO NP-alone and the co-exposure with GO groups. However, the levels of fold changes of the 14 shared metabolites were lower in the coexposure group than that in the ZnO NP-alone group. This tendency indicates that GO alleviated the toxicity induced by ZnO NPs in the cellular metabolism by reducing or blocking their internalization in the cells (Fig. 15d-f). Table 6 summarizes the studies that used metabolomics technique to assess the effect of ZnO NPs in vitro on different cells.

5.6 Other metal- and metal oxide-NPs

5.6.1 Cobalt ferrite (CoFe₂O₄) NPs. Cobalt ferrite (CoFe₂O₄) NPs have interesting properties, such as mechanical hardness, excellent chemical stability, high anisotropy, paramagnetism, and coercivity.259 Oliveira et al.260 studied the cytotoxic effect, cellular uptake, and metabolomic effect of CoFe₂O₄ NPs on the HeLa and HaCaT cell lines. This study revealed, using NMR-based metabolomics, that although the uptake of NPs at 2 mg mL⁻¹ caused low cytotoxicity, it significantly impacted the cell metabolism. Both cell lines shared stress-related metabolic changes such as upregulation in alanine and creatine. A downregulation in fumarate level was present in HeLa cells treated with the NPs. Given that this metabolite is associated with cell proliferation and tumor growth, it was concluded that CoFe₂O₄ NPs can inhibit tumorigenesis.

5.6.2 Copper oxide (CuO) NPs. Copper oxide (CuO) NPs have been used in heat transfer fluids, semiconductors, and intrauterine contraceptive devices.261 Human exposure to CuO NPs is rapidly increasing, and thus reliable toxicity test systems are urgently needed. It was shown that CuO NPs are more toxic than their microparticles (MPs). To reveal the mechanism of this toxicity, Murgia et al. 262 used MS-based metabolomics to the study the effect of CuO micro- and nano-particles against




Fig. 15 [(a) Comparison of ZnO NPs with PS NPs]. 247 (a) Summary comparing the relevant metabolic responses of BEAS-2B cells to ZnO and PS_LD NP exposure. Arrow in orange or blue represents a significant increase or decrease in the ZnO or PS_LD NP exposure, respectively, compared with the non-treated groups. This figure is adapted with permission from ref. 247 Copyright 2019, Taylor & Francis. [(b and c) Comparison of ZnO NPs with Zn²⁺]. 254 (b) Proportion of significantly changed metabolites in different categories after 2 (2 d) and 7 (7 d) days exposure to ZnO NPs and ZnCl₂. (c) Edwards-Venn diagram of the total number of significantly changed metabolites. The total numbers of significantly changed metabolites in ZnO_2 d (d = days), ZnO_7 d, ZnCl₂_2 d, and ZnCl₂_7 d groups were 99, 121, 128, and 183, respectively. The altered metabolites were obtained by conducting PLS-DA analyses for each Zn-exposed group vs. the matched control group (VIP > 1 and p < 0.05). Note: the metabolites identified from the positive ion mode and negative ion mode were merged together. (b and c) Are reprinted with permission from ref. 254 Copyright 2020, the American Chemical Society. [(d-f) Comparison of ZnO NPs solo exposure to their co-exposure with GO]. Schematic illustration of A549 cells ZnO NPs solo exposure vs. co-exposure with GO. The GO sheets reduce the cytotoxicity of ZnO NPs by blocking their internalization into the A549 cells. (e) Influence of GO on the bioavailability of ZnO NPs. The Zn concentrations were normalized by the protein concentrations. All data expressed as the mean \pm SD. All differences were identified by one-way ANOVA followed by Tukey post hoc test. * Indicates p-value. (f) Interaction network of metabolites in ZnO NPs solo and co-exposure groups. (d-f) Are reproduced from ref. 257 with permission from The Royal Society of Chemistry, Copyright 2019.

Table 6 Summary of ZnO NP-induced perturbation of metabolic pathways and their biological impact on different cells

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
ZnO	60	N/A	BEAS-2B	10 μg mL ⁻¹	HPLC-MS/MS	Amino acid ↓	Mitochondrial dysfunction	247
				24 h	GC-MS/MS	TCA ↓	Cell death ROS	
ZnO	22.6	N/A	BEAS-2B	12.5, 25 μg mL ⁻¹ 6 h	LC-MS	Amino acid ↓ Nucleotides ↓ Lipid ↓ Toxic metabolites ↑	Inflammation Oxidative stress DNA damage High cytotoxicity	217
ZnO- GO	ZnO – 50	N/A	A549	ZnO – 20 μg mL^{-1}	¹ H NMR	TCA ↓	Membrane damage	257
				GO – 10 μg mL ⁻¹ 24 h		GSH ↓ Choline Amino acid Carbohydrate	Oxidative stress Energy metabolism disruption GO reduced the impact of nano-ZnO	
ZnO	42 34	None Triethoxycaprylsilane	A549	10, 15 μg mL ⁻¹ 1, 6, 24 h	DIMS	GSH ↓ Amino acid	Oxidative stress Apoptosis	199
ZnO	71	N/A	E. coli	0.025–0.2 μg mL ⁻¹ 3 h	GC-TOF-MS	Amino acid; glycine	Oxidative stress	233
ZnO	<70	N/A	Yeast S. cerevisiae BY4741	10 μg mL ⁻¹ 3 h	¹ H NMR	Amino acid TCA ↓ GSH ↓ Glycolysis ↓ Fatty acid ↓ Purine and pyrimidine ↓	DNA and protein damage Oxidative stress Antioxidation Energy metabolism disruption	258

human bone marrow mesenchymal stem cells (hBMMSCs). It was found that the MPs increased the levels of serine, glyceric acid, and succinic acid, while glutamine was the only discriminant metabolite for the class of samples treated with NPs. This proves that ROS formation is the active mode of action in NP treatment, providing the first step toward the understanding of the mechanism of toxicity of CuO NP-treated cells. Wang et al. 263 compared the effect of CuO NPs, MPs, and Cu ions on microalga Chlorella vulgaris after 5 days exposure using global metabolomics. A total of 75 differentiated metabolites was identified. Most metabolic pathways perturbed after CuO NP exposure were shared by that after CuO MP and Cu ion exposure. Only one difference between metabolic responses to particles and that to ions was observed, which is the accumulation of fatty acid oxidation products, i.e., particles caused higher fold changes at 1 mg L⁻¹ and lower fold changes at 10 mg L⁻¹ compared with ions. This indicates the significant role of dissolved Cu ions on the toxicity of CuO NPs and MPs. Kruszka et al.264 compared the effect of Cu and CuO NPs on the secondary metabolism of Hypericum perforatum L. cell suspension cultures and found that metal NPs induce higher metabolic changes than their counterpart metal oxide NPs. Table 7 summarizes the studies that used the metabolomics technique to assess the effect of other metal/metal oxide NPs in vitro on different cells.

5.7 Carbon-based NPs

5.7.1 Graphene. Graphene has attracted significant attention due to its unique and novel properties, which has promising applications in different fields, including biomedical engineering, tissue engineering, and biosensors. However, graphene-based drug delivery systems and other biomedical applications are associated with challenges related to the safety of carbon NMs for clinical use. Many groups have investigated the cytotoxicity of graphene. In this case, although the conventional *in vitro* toxicity assays of graphene yielded contradictory results, Jiao *et al.*²⁸⁶ used the metabolomics approach to investigate the metabolic responses on graphene-treated HepG2 and detected twelve metabolites as potential biomarkers. The authors also determined three KEGG pathways including arginine and proline metabolism, purine metabolism, and glycophospholipid metabolism.

Adamson *et al.*²⁸⁷ studied the metabolic change caused in macrophages by graphene nanoplatelets. The number of compounds changed following exposure to graphene was determined to be both concentration and time dependent. The identified metabolites are related to several metabolism pathways, such as GSH metabolism, pantothenate and CoA biosynthesis, sphingolipid metabolism, purine metabolism, arachidonic acid metabolism and others. Graphene oxide (GO) also has some biomedical applications but a greater

Summary of metal/oxide NM NP-induced perturbation of metabolic pathways and their biological impact on different cells

Table 7

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
CuO	30	N/A	hBMMSCs	$2.51~\mu \mathrm{g}~\mathrm{mL}^{-1}$	GC-MS	Serine Glyceric acid Succinic acid Glutamine	ROS formation	262
CuO	20-80	N/A	НерG2	3 μg mL ⁻¹ 72 h	GC-MLC-MS	S-Adenosylhomocysteine ↑ Lysolipids ↑ Sphingolipids ↑ S-Adenosylmethionine ↓	Oxidative stress	216
CuO	28	N/A	A549	10 μg mL ⁻¹ 0, 1, 3, 6, 12, 24 h	HPLC-MS	GST ↓ MNA ↑ GSH ↓ GPC ↑ Aprile a seid ↑	Oxidative stress Hypertonic stress Apoptosis	265
CuO	<50	N/A	HCF-116	2.5, 5 μg mL ⁻¹ 24 h	LC-QToF-MS	Anning acts Triacylglycerols ↑ Phosphatidylcholines ↑ Ceramides ↑ GSSG ↑	Autophagy Oxidative stress	266
CuO	15-50	N/A	Alga C. vulgaris	1, 10 $\mu { m g \ mL}^{-1}$ 120 h	LC-QToF-MS	Fatty acid ↑ GSH ↓ Phosphatidylcholines ↓ Phosphatidylchowerols ↑	Oxidative stress Osmotic stress ROS formation Membrane damage	263
CeO ₂	5-50	N/A	HepG2	$30,100~\mu \mathrm{g}~\mathrm{mL}^{-1}$ 72 h	GC-MS LC-MS	S-Adenosylhomocysteine ↑ Lipids ↑ UDP-glucuronate ↓ S-Adenosylmethionine ↓	Oxidative stress	242
CeO_2	58	N/A	HepG2	3, 30 $\mu g m L^{-1}$ 72 h	GC-MS UPLC/MS/MS	GSII↓ GSBI↓ Lipids↓ Fattv acid	Oxidative stress	216
CeO_2	4.7	None, or Zr	A549	128 $\mu g m L^{-1}$ 1 6 24 h	DIMS	racy aca Cysteine↑ ~-Glutamyleysteine↑	Oxidative stress	207
CeO_2	50×6.7	N/A	BEAS-2B	1, 5, 27 m 12.5, 25 µg mL ⁻¹ 6 h	IC-MS	7 Statemy 23 Sente S1P pathway ↑ Folate cycle ↑ GSH ↓	Apoposas Inflammation Oxidative stress DNA damage	217
CeO_2	210	N/A	HuH-7	100 $\mu g \; m L^{-1}$ 24 h	HPLC-MS	Lipid Histamine ↑ Hydroxyproline ↑ Proline ↓	Genotoxicity	267

Table 7 (Contd.)

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
CeO ₂	4–5	PVP	Alga C. reinhardtii	0.029, 0.144, 0.7280, 400, 2000, 10 000 μg 1-1	LTQ-FT-MS	Only at 10 000 $\mu g \; L^{-1}$	Photosynthesis decrease	268
				72 h		Pyruvate ↑ GSH ↑ Purine ↑	Energy metabolism disruption	
$\mathrm{CoFe_2O_4}$	27	N/A	НеLа НаСаТ	$\begin{array}{c} 2 \text{ mg mL}^{-1} \\ 24 \text{ h} \end{array}$	¹ H NMR	Pyrimidine ↓ Alanine ↑ Creatine ↑ GSH ↑	Metabolic change HeLa – inhibition of tumorigenesis	260
$\mathrm{CoFe_2O_4}$	Core-9 Full-50	(RITC)-SiO ₂	HEK293	$0.1, 1.0 \ \mathrm{Hg \ mL^{-1}}$ 12 h	GC-MS/MS	HeLa – fumarate, lactate ↓ Glucose ↓ Amino acid Fatty acid Polyamine	ROS generation Glucose metabolic dysfunction	30
$\mathrm{CoFe}_2\mathrm{O}_4$	Core-9 Full-50	$(RITC)$ -SiO $_2$	HEK293	$1.0~ m \mu g~mL^{-1}$	GC-MS/MS	Organic acid Glutamic acid ↑ Krebs cycle	Mitochondrial damage ROS generation	156
$\mathrm{CoFe_2O_4}$	Core-9	(RITC)-SiO ₂	HEK293	$1.0~\mu \mathrm{g~mL}^{-1}$	GC-MS/MS	AIP ↓ Lipid ↓	Membrane fluidity decreased	269
$\mathrm{CoFe_2O_4}$	Full-50 Core-9	(RITC)-SiO ₂	BV2 murine	$0.1,1.0~\mu{ m g~mL}^{-1}$	GC-MS/MS	ATP ↓ GSH	Cell movement decreased ROS generation	270
$\mathrm{Mn_2O_3}$	5 5	N/A	nncrognar cens RLE-6TN	12 n 10 μg cm ⁻² 24 h	HPLC-MS/MS FIA-MS/MS	ICA No significant metabolic changes	Innanimatory response Oxidative stress Apoptosis	231
$\mathrm{Mn_2O_3}$	50	N/A	NR8383	$2.5, 5, 10 \mu \mathrm{g cm^{-2}}$ $24 \mathrm{h}$	HPLC-MS	Amino acid Biogenic amine	High cytotoxicity Oxidative stress Mitochondrial dysfunction	243
$\mathrm{Al}_2\mathrm{O}_3$	64	N/A	нве	100, 500 $\mu \mathrm{g \; mL}^{-1}$ 24 h	GC/TOF-MS	Phosphatidylcholine p-Glutamic acid ↑ Succinic acid ↑	DNA damage & cell death Mitochondria-dependent apoptosis Oxidative stress	271
γ -Al $_2$ O $_3$	[20–30 ×	N/A	ASTs	$125~ m \mu g~mL^{-1}$	UHPLC-MS/	3-Methylhistidine↓ TCA↓ Amino acid	ROS generation ROS generation	92
	$\begin{bmatrix} 20-30 \\ [20-30 \times \\ 100-200 \end{bmatrix}$			72 h	CIVI	Lipid Purines, and pyrimidines	Oxidative stress Inflammation	
$\mathrm{Fe}_2\mathrm{O}_3$	$\begin{bmatrix} 16-44 \times \\ 45-172 \end{bmatrix}$	N/A	THP-1	$20100~\mu\mathrm{g~mL}^{-1}$	LC-ToF-MS	Sphingolipids	Apoptosis Inflammation	272
	$\begin{bmatrix} 20 - 1/3 \end{bmatrix}$ $\begin{bmatrix} 20 - 53 \times 88 - 322 \end{bmatrix}$			24 h		Ceramides Sphingosine-1-phosphates Glucosylceramide	Energy metabolism disruption Cell proliferation Autophagy	

Table 7 (Contd.)

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
Fe ₃ O ₄ FeO. mix	19.4	N/A	BEAS-2B	0.003, 0.03, 0.3 μg mI _. ⁻¹ 24 h	DI-MS	No significant change	Low cytotoxicity	273
ZrO_2	31.9	N/A	MC3T3-E1	100 µg mL ⁻¹ 24 h 48 h	LC-ToF-MS	Phospholipids 👃	Oxidative stress	274
$\mathrm{MoS}_2\mathrm{-Ag}$	Ag - 18	Chitosan	Yeast S. cerevisiae	20 $\mu g L^{-1} N-Ag$, 1 $mg L^{-1} CS-MoS_2$ 24 h	GC-TOF-MS	Amino acid Organic acid	Oxidative stress lower than Ag alone Membrane stress more than	275
MoS_2	5–10	N/A	Mesophyll protoplasts	50 µg mL ⁻¹ 0.5-3 h	GC-MS	Fatty acid ATP ↓ 3-(4-Hydroxyphenyl) propionic acid ↑ Carechin	Ag alone ROS generation Mitochondrial dysfunction	276
MoS_2	N/A	N/A	E. coli	1, 10, 100, 1000 µg mL ⁻¹ 6 h	GC-TOF-MS	Glycine, serine, and threonine Urea cycle	Antimicrobial activity LDH release	277
MoS_2	Thickness - 4	Chitosan	HDFs	25, 100 μg mL ⁻¹ 72 h	GC-TOF-MS	Tyruvac Amino acid GSH ↓ TCA ↓	Mode Sentration Membrane damage ROS generation DNA damage Inflammation Anontosis	278
WS_2	100-4000	N/A	A549 - THP-1	50, 100 $\mu \mathrm{g} \ \mathrm{mL}^{-1}$ 24 h	GC-MS	Amino acid Glycolysis ↑ Pyruvate	Trepression of the Physical Structure of Physical Cell migration Cell migration	279
Fe	50	N/A	Mesophyll protoplasts	$50 \ \mathrm{\mu g \ mL}^{-1}$ $0.5-3 \ \mathrm{h}$	GC-MS	ATP↑ Phenvlalanine	Photosynthesis enhancement	276
Fe	9.9	N/A	Front process S. aureus E. coli	$^{100}\mathrm{kg}\mathrm{mL}^{-1}$ 0.5 h	¹H NMR UHPLC-HRMS	A-Linolenic acid Lipoic acid metabolism Phenylalanine Tyrosine, and tryptophan biosynthesis	No significant effect on bacterial metabolism	280
Bismuth	<100	Carboxyl	H. pylori	100 $\mu g \; m L^{-1}$ 24 h	$^{1}\mathrm{H}\ \mathrm{NMR}$	TCA Nucleotide Amino acid	Antimicrobial activity Release of some metabolites to	281
Cu	5.3	N/A	S. aureus E. coli	$100~{ m \mu g~mL^{-1}}$ 0.5 h	¹H NMR UHPLC-HRMS	Amino sugar Nucleotide sugar	Blocking cell metabolism	280
$Cu(OH)_2$	140	N/A	HepG2	$25 \mu \mathrm{g \ mL}^{-1}$ 48 h	UHPLC-MS/ MS	Glycolysis↑ Lipid↑ TCA↓	Energetic stress Oxidative stress	282
$S(\alpha - \text{ or } \beta - \text{SNPs})$	10 50	PEG	A. niger (MTCC- 10180)	4 mg mL $^{-1}$ 16 mg mL $^{-1}$ 48 h	GC-MS	TCA ↓	Antimicrobial activity Oxidative phosphorylation	283

Table 7(Contd.)

Ref.	280	284	ion 285	
Biological effect	Blocking cell metabolism Cell death	Mitochondrial dysfunction	Energy metabolism disruption	
Perturbed metabolic pathway	Pentose phosphate TCA Amino sugar	Nucleotide sugar Glycolysis ↑ Pentose phosphate ↑ Lipid ↑	TCA↓ TCA↓ Glycolysis↑ Fatty acid↑	Lipiu
Analytical platform	¹H NMR UHPLC-HRMS	GC-MS	GC-MS LC-MS/MS	
Dose/exposure time	$100~\mathrm{\mu g~mL}^{-1}$ 0.5 h	8, 80 µg mL ⁻¹ 48 h	100, 200, 500 µg mL ⁻¹ 48 h	
Cell	S. aureus E. coli	HUVECs	HUVEC	
Coating	N/A	N/A	N/A	
Size [nm] Coating	15	43.8	489 × 11	
Ð.	u-Fe	pd-n	$^{13}\mathrm{C}_{2}$	

understanding of its cytotoxicity and efficiency as a drug carrier is needed. Raja *et al.*²⁸⁸ used NMR-based metabolomics to assess the metabolic effect of GO nanosheets on MCF-7 breast cancer cells. The treatment affected arginine metabolism, proline metabolism, and aminoacyl-tRNA biosynthesis, including anabolism and catabolism. Moreover, GO increased the number of metabolic disturbances in cancer steroids in a dose-dependent manner.

5.7.2 Carbon black NPs (CBNPs). Carbon black NPs (CBNPs) are the core component of fine particulate matter in the atmosphere, which make its exposure to the respiratory system easy. It was reported that CBNPs can induce inflammation, oxidative stress, and changes in cell signalling and gene expression in mammalian cells and organs. Hou et al.289 used MS-based metabolomics to reveal this mechanism in A549 cells. Their study identified a total of 32 differential metabolites between the CBNP exposure and control groups. The pathway analysis showed that the metabolic changes were involved in tricarboxylic acid (TCA) cycle, alanine, aspartate, glutamate, and histidine metabolism. This suggests that CBNPs act by affecting the normal process of energy metabolism and disturbing several vital signalling pathways in the cells, finally leading to cell apoptosis. Other studies performed in vivo experiments and assessed the effect of carbon-based NMs on the ecosystem by studying some models such as earthworms. For instance, Xu et al.290 studied the impacts of three carbon NMs, i.e., carbon black (CB), reduced graphene oxide (RGO), and single-wall carbon nanotubes (SWCNTs), on Eisenia fetida, an early warning soil invertebrate for pollution events. They concluded that the soil environmental risk of C-NMs was related to their particle morphology, contributing to a comprehensive understanding of nano-agriculture. Table 8 summarizes the studies that used the metabolomics technique to assess the effect of other C-based NPs in vitro on different cells.

5.8 Polymeric NPs

Polymeric NPs such as polystyrene (PS) are gaining considerable attention because of their growing accumulation in the environment and the high probability of human and animal exposure. Therefore, more research must be done to increase our understanding of their potential effects. Kim et al. 303 studied the metabolic effects of PS NPs on human epithelial colorectal cells (Caco-2). The authors designed two methods to investigate the exposure of Caco-2 cells to NPs, where the first is by exposing cells to a high concentration of 50 nm PS NPs for 24 h (acute), and the second is by exposing them to a relatively lower concentration for over 1.5 months (chronic). The biological assays were performed using specific NP concentrations, which were 10 and 80 $\mu g \text{ mL}^{-1}$ for the acute model and 0.1 $\mu g \text{ mL}^{-1}$ for the chronic exposure model. After acute exposure, untargeted metabolic profiling was performed and the change in lipid metabolic pathways determined, including steroid and arachidonic acid metabolism. Alternatively, chronic exposure induced relatively minor changes. However, there was still a potential effect on fatty acid biosynthesis, indicating that acute and chronic exposure to PS NPs may disturb lipid

Table 8 Summary of C-based NP-induced perturbation of metabolic pathways and their biological impact on different cells

9		5			, , , , , , , , , , , , , , , , , , , ,			
NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
9	N/A	N/A	Нер <u>б</u> 2	$25~{ m \mu g~mL}^{-1}$	UPLC-MS	Arginine and proline Purine Glycophospholipid Urea cycle	Membrane damage Protein damage	286
g	$2000\times2000\times12$	N/A	RAW 264.7	50, 100 μg mL ⁻¹ 1, 3 h	HPLC-MS	GSH GSH Sphingolipid CoA	Mitochondrial membrane potential CD36-dependent response	287
_o	N/A	N/A	НаСаТ	5 μg mL ⁻¹ 168 h	¹H NMR	Furine Fumarate † Glycerophosphocholine † Pyruvate ↓ Phosphocreatine ↓ Phosphocholine	ROS generation Mitochondrial damage Cell death Cell migration	291
g	N/A	Carboxyl	A549	0.01, 0.1 μg mL ⁻¹ 48 h	LC-MS	Anisophoranic Anino acid Organic acids Glycerophospholipid Glycerolinids	ROS generation Apoptosis	292
05	N/A	N/A	НаСаТ	5 μg mL ⁻¹ 168 h	¹H NMR	Fumarate ↓ Alanine ↑ Pyruvate ↑ Phosphocreatine ↑ Glycerophosphocholine ↑	ROS generation Mitochondrial damage Cell death Cell migration	291
09	N/A	N/A	MCF-7	20, 40, 60 μg mL ⁻¹ 24 h	¹ H NMR	Arginine Proline GSH TCA	Catabolism Therapeutic activity	288
05	300	PEG	Saos-2	75 μg mL ⁻¹ 24 h	¹H NMR	Amino acid ↓ Taurine ↓ Creatine ↓ Phospbocholine ↑	Proliferation delay Oxidative stress	293
OĐ	N/A	N/A	RLE-6TN	$10 \ \mu \mathrm{g} \ \mathrm{cm}^{-2}$ 24 h	HPLC-MS/MS FIA-MS/MS	Amino acid Biogenic amine Linid	Oxidative stress Apoptosis	231
09	N/A	N/A	NR8383	2.5, 5, 10 µg cm ⁻² 24 h	HPLC-MS	No significant alteration	Oxidative stress Mitochondrial dysfunction DNA damage Cell death	243
OĐ	$500-5000 \times 0.8-1.2$	N/A	Alga <i>C. vulgaris</i>	0.01 – $10~\mu \mathrm{g~mL}^{-1}$ 96 h	GC-MS	Alkanes, lysine, octadecadienoic acid and valine	ROS generation Oxidative stress	294
MWCNT	$500-2000 \times 8-15 \times 3-5$	None Hydroxyl Carboxyl	BEAS-2B HepG2	2–39 μg mL ⁻¹ 24 h	¹H NMR	Choline Betaine Succinate	Oxidative stress Inflammation Profibrosis DNA damage	295

Table 8 (Contd.)

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
MWCNT	[(1-3) or (0.05-0.2) $10^4 \times 20-30$ or $8-15$]	Hydroxyl	BEAS-2B HepG2	$6-45 \ \mu g \ mL^{-1}$ 24 h	¹H NMR	TCA GSH Amino acid	Oxidative stress Inflammation Profibrosis	296
MWCNT	846 imes 11	N/A	HBEC-3KT	$0.96, 1.92 \text{µg cm}^{-2}$ 2, 13 weeks	FT-MS/MS	TG, SM, Cer, PE, Chol↑	ROS generation Apoptosis	297
SWCNT	$500-3000 \times 1-2$	Carboxyl	Alga C. vulgaris	0.01–10 µg mL ⁻¹ 96 h	GC-MS	Alkanes, lysine, octadecadienoic acid and valine	ROS generation Oxidative stress	294
CNT	N/A	Carboxyl	A549	0.01, 0.1 μg mL ⁻¹ 48 h	I.C-MS	Amino acid Organic acids Glycerophospholipid	Energy metabolism dysfunction ROS generation Apoptosis	292
C_{60}	100	Carboxyl	A549	0.01, 0.1 μg mL ⁻¹ 48 h	LC-MS	Amino acid Organic acids Glycerophospholipid Glycerolinide	Energy metabolism dysfunction	292
C_{60}	120–150	N/A	Alga S. obliquus	$0.1, 1 \ \mu g \ mL^{-1}$ 168 h	GC-Q-TOF-MS	TCA Sucrose, p-glucose, malic	Growth inhibition Photosynthesis decrease	298
Chlack	30–50	N/A	A549	70 μg mL ⁻¹ 48 h	UHPLC-Q-TOF-MS	TCA Alanine Aspartate Glutamate Histidine ↑	Inflammation Oxidative stress Energy and signalling pathway disruption Apoptosis	289
C_{black}	14	None Benzo $[a]$	EA.hy926	$0.001-1 \ \mathrm{\mu g \ mL^{-1}}$ 72 h	GC-MS	Amino acid Organic acid Fatty acid	Enhanced cell proliferation Cell motility Cell migration increase	299
Ccellulose	2-40	N/A	V69V	10, 100 $\mu\mathrm{g~mL}^{-1}$ 6 h	LC-MS	Fatty acid	Cell viability Metabolic activity disruption	300
Cyclodextrin	0.5-1	Folate	MDA-MB-231	1:16 (v/v) 24 h	LC-Q-TOF-MS	Hexose metabolism Glycolysis TCA	Apoptosis	301
Glycyrrhizin (GL-BSA- Lut-NPs)	225.3	Luteolin	Bel-7402	0.375, 0.75, 1.25, 2.5, 5 mg mL ⁻¹	¹ H NMR	Lactate, 3-hydroxybutyrate, lipid, tyrosine, and β -glucose \downarrow	Antitumor effect	302
		BSA		24 h		Glutamate, asparagine, choline, and creatine ↑	Apoptosis	

Table 9 Summary of polymeric NP-induced perturbation of metabolic pathways and their biological impact on different cells

	Samuely of postmicion			the partition of the process of the constant o				
NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
NG NG	7.0	4 /N	RFAS-7R	10 50 119 mJ ⁻¹	HPLC-MS/MS	Amino seid ↑	Anti-oxidative	747
2	1	41/14		24 55 (51			protection	ì
				24 h	GC-MS/MS	TCA ↑	Autophagy	
						Lactate ↑	Perturbed energetic metabolism	
PS	50	N/A	Caco-2	Acute [10, 80 μg mL ⁻¹ – 24 h]	UPLC-MS	Acute – lipid; steroid and arachidonic acid	Metabolic disturbance	303
				Chronic $[0.1 \ \mu g \ mL^{-1} - 45 \ days]$		Chronic – fatty acid biosynthesis	ROS generation	
PS	50	Amino	Cyanobacterium	$2.5, 4 \mathrm{\mu g mL^{-1}}$	GC-ToF-MS	Amino acid ↓	Oxidative stress	305
			S. elongatus	48 h		GSH↓ o-Phosphoethanolamine↑	Membrane damage ROS generation	
PLGA	100-120	None	A549	$92~{ m \mug~mL}^{-1}$	LC-MS	Nucleotide	Energy metabolism	306
		MTX	THP-1	24 h		TCA ↓	More effective on	
						Glycolysis ↓ GSH	cancer cell than free	
PLGA	100–125	A/N	RAW 264.7	$10.500 \text{ ug mL}^{-1}$	¹ H NMR	Glycolysis ↑	Inflammation	244
		<u> </u>		24, 48, 72 h		Lactate ↑	Energy metabolism	
				,			disruption	
PLGA	250.90	L-Carnitine	CFs	$2~{ m mg~mL}^{-1}$	GC-MS	Amino acid ↑	Very effective drug	307
				144 h		2-Ketoisocaproic ↑ Glucose ↑	carrier for amino acid	
DIIA	246 71	N/A	DC12	V/N	MO-TOE-MS	Amino seid	Cell differentiation	304
	218.57	U /V		12, 24, 36 h	UHPLC-MS	Carbohydrate	enhancement	*
,				7		Lipid		
PLLA (LJ@AA)	150	PEG–amino acid	MCF7	$1 \mathrm{\mu g mL^{-1}}$ 48 h	UPLC-MS/MS	Amino acid ↓	Cancer inhibition	157
PET	10-80	N/A	Caco-2	$30 \mu \mathrm{g} \mathrm{mL}^{-1}$ 48 h	¹H NMR	Glucose ↓ Lactate ↓ Alanine ↑	Oxidative stress	308
Silk	$\frac{10}{0-125}$	N/A	RAW 264.7	10, 500 μg mL ⁻¹ 24, 48, 72 h	¹H NMR	Glycolysis↑ Lactate↑ Pyrnyafe	Inflammation	244
Strigol1/albumin/ chitosan	5-10	N/A	HepG2	9.24–92.4 nM 48 h	LC-MS-MS	Spermine and spermidine ↑ Glutamine ↓	Apoptosis Anti-carcinogenic effect	309
Platicur-NCs	100	Chitosan	НеГа	Dark – 75 μM	$^1\mathrm{H}$ NMR	rumararc ↓ Light – glutamine, acetate, olnose ↑	ROS generation	310
		Chitosan-pectin		Light – 0.05 µM		Dark – lactate, creatine,	Oxidative stress	
				24 h		SST ↑ HSD	Apoptosis	
6OCaproβ Cyclodextrin	15 3	N/A	MCF-7	N/A 48 h	Q-TOF-LC MS	Serine biosynthesis Estrogen biosynthesis Phospholipid biosynthesis	Apoptosis	311

(Contd. Table 9

NP	Size [nm]	Coating	Cell	Dose/exposure time	Analytical platform	Perturbed metabolic pathway	Biological effect	Ref.
Lipo-lysosomes	97.88	L-Carnitine	CFs	2 mg mL ⁻¹ 144 h	GC-MS	Amino acid † 2-Ketoisocaproic ↓ 3-Phenyllactic acid ↓ 2-Hydroxyhuturate ↑	Very effective drug carrier for the synthesis of unsaturated fatty	307
Cationic liposomes	15 2.2	N/A	L02	116 μg mL ⁻¹ 24 h	UHPLC-Q-TOF- MS	Sphingolipid Fatty acid TCA GSH Methionine	Ros generation Oxidative stress Energy metabolism disruption	312
Chloroquine Dendrimer	23	Curcumin	Parasite P. falciparum	0.1 –2.5 $\mu \mathrm{g~mL}^{-1}$ 48 h	¹ H NMR	rymmame Pyrimidine TCA↑ GSH↑	Effective drug carrier	313
Lambda phage like	16	Fluorescein-5- maleimide Trastuzumab	SKBR3	150 nM 2 µМ 3 h	UHPLC-MS	TCA ↓ Glycolysis Fatty acid synthesis Protein synthesis	DNA damage Oxidative stress Mitochondrial dysfunction	314

homeostasis. They also confirmed the changes in the expression levels of lipid transcriptional regulatory factor coding genes, namely, sterol regulatory element-binding transcription factors 1 and 2. The total fatty acid composition was further studied to verify metabolic disturbance by chronic exposure. Su et al. 304 investigated the effect of poly(L-lactic acid) (PLLA) nanofibers on PC12 cell differentiation at the metabolic level. Many differential metabolites were identified and two pathways and three metabolites critical to PC12 cell differentiation were influenced by the nanofibers. Table 9 summarizes the studies that used the metabolomics technique to assess the effect of other polymeric NPs in vitro on different cells.

Conclusions

Currently, it is obvious for many researchers that nanotechnology provides countless benefits, and consequently its demand is increasing daily. It is very important to assess the safety of every NP that is being produced and to test its beneficial and disadvantageous effects. Understanding the interactions between NPs with cells and how NPs are internalized in cells are the first step to assess their toxicity. Here, not only the type of the NP matters, but also its physicochemical properties such as size, shape, and surface properties. It was proven that NPs with different properties have different effect on cells. Some sizes of NPs are not toxic, but others are severely harmful to cells. In general, conventional assays are the most used strategy to assess the effect of NPs on cells. However, these assays were found to interfere with NPs, giving false results in some cases, and they are unable to reveal the molecular information of the toxicity or effect of NPs. Thus, an increasing number of researchers are heading towards the use of other analytical techniques. Metabolomics is a powerful technique that provides a full picture of the toxicity of NPs by analyzing the functionality of an existing living system by measuring the metabolic change induced by NPs. Unlike conventional assays, this tool does not interfere with NPs and provides information at the molecular level about the toxicity of NPs. Furthermore, it forms an additional bridge that connects the in vitro with the in vivo models, as proven by several references. It was shown in this review that NPs can harm the cell through different ways, including cell viability and proliferation perturbation, inflammatory response, oxidative stress, ROS generation, and cell death via apoptosis or necrosis. Moreover, using metabolomics, NPs were shown to perturb the metabolic pathways of cells, including the TCA cycle, DNA and protein synthesis, glycolysis, glutathione, and amino acid pathways. Thus far, metabolomics has been used in many studies to assess the effects of different NPs on living organisms. However, more research needs to be done to identify and validate specific biomarkers of the effects of NPs on cells. Reaching this point will introduce a huge step in determining the toxicity of NPs and how to avoid or multiply this toxicity. This will help in designing better NPs for biomedical applications and producing safer NPs for industrial applications. Nevertheless, long-term targeted studies should also be performed to fill many gaps in this field. Also, the combination of metabolomics with other techniques is

required in some cases to provide a bigger picture on the events occurring in the cell.

Author contributions

Mohammad Awashra: literature search, writing, figures designing and creating. Piotr Mlynarz: review idea, reviewing, and editing.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors thank Wroclaw University of Science and Technology subsidy for Faculty of Chemistry for financial support and CNE Erasmus Mundus Program for managing the main author master thesis. Fig. 1-7a, 9, 10, 12 and 15a, d, and table of entry figure are created with https://BioRender.com.

References

- 1 X. Cheng, in Nanolithography, Elsevier, 2014, pp. 348-375.
- 2 J. Hulla, S. Sahu and A. Hayes, Nanotechnology, Hum. Exp. Toxicol., 2015, 34, 1318-1321.
- 3 P. F. Wright, Potential risks and benefits of nanotechnology: perceptions of risk in sunscreens, Med. J. Aust., 2016, 204, 369-370.
- 4 C. S. C. Santos, B. Gabriel, M. Blanchy, O. Menes, D. García, M. Blanco, N. Arconada and V. Neto, Industrial applications of nanoparticles - a prospective overview, Mater. Today: Proc., 2015, 2, 456-465.
- 5 X. He, H. Deng and H. Hwang, The current application of nanotechnology in food and agriculture, J. Food Drug Anal., 2019, 27, 1-21.
- 6 L. F. Fraceto, R. Grillo, G. A. de Medeiros, V. Scognamiglio, G. Rea and C. Bartolucci, Nanotechnology in agriculture: which innovation potential does it have?, Front. Environ. Sci., 2016, 4(20), DOI: 10.3389/fenvs.2016.00020.
- 7 C. Chellaram, G. Murugaboopathi, A. A. John, R. Sivakumar, S. Ganesan, S. Krithika and G. Priya, Significance of manotechnology in food industry, APCBEE Proc., 2014, 8, 109-113.
- 8 M. J. Mitchell, M. M. Billingsley, R. M. Haley, M. E. Wechsler, N. A. Peppas and R. Langer, Engineering precision nanoparticles for drug delivery, Nat. Rev. Drug Discovery, 2021, 20, 101-124.
- 9 S. Raj, U. Sumod, S. Jose and M. Sabitha, Nanotechnology in cosmetics: Opportunities and challenges, J. Pharm. BioAllied Sci., 2012, 4, 186.
- 10 A. Javed, M. Azeem, J. Wiener, M. Thukkaram, J. Saskova and T. Mansoor, Ultrasonically assisted in situ deposition ZnO nano particles on cotton fabrics multifunctional textiles, Fibers Polym., 2021, 22, 77-86.

11 J. Mathew, J. Joy and S. C. George, Potential applications of nanotechnology in transportation: a review, I. King Saud Univ., Sci., 2019, 31, 586-594.

- 12 P. T. Sekoai, C. N. M. Ouma, S. P. du Preez, P. Modisha, N. Engelbrecht, D. G. Bessarabov and A. Ghimire, Application of nanoparticles in biofuels: an overview, Fuel, 2019, 237, 380-397.
- 13 X. Zhang, Q. Guo and D. Cui, Recent advances in nanotechnology applied to biosensors, Sensors, 2009, 9,
- 14 P. J. Borm, D. Robbins, S. Haubold, T. Kuhlbusch, H. Fissan, K. Donaldson, R. Schins, V. Stone, W. Kreyling, Lademann, J. Krutmann, D. Warheit E. Oberdorster, The potential risks of nanomaterials: a review carried out for ECETOC, Part. Fibre Toxicol., 2006, 3, 11.
- 15 A. B. Asha and R. Narain, in Polymer Science and Nanotechnology, Elsevier, 2020, pp. 343-359.
- 16 P. C. Ray, H. Yu and P. P. Fu, Toxicity and Environmental Risks of Nanomaterials: Challenges and Future Needs, J. Environ. Sci. Health, Part C: Environ. Carcinog. Ecotoxicol. Rev., 2009, 27, 1-35.
- 17 T. A. Saleh, Nanomaterials: Classification, properties, and environmental toxicities, Environ. Technol. Innovation, 2020, 20, 101067.
- 18 A. Seaton, L. Tran, R. Aitken and K. Donaldson, Nanoparticles, human health hazard and regulation, J. R. Interface, 2010, 7(suppl_1), DOI: Soc., rsif.2009.0252.focus.
- 19 H. Kan, D. Pan and V. Castranova, Engineered nanoparticle exposure and cardiovascular effects: the role of a neuronalregulated pathway, Inhalation Toxicol., 2018, 30, 335-342.
- 20 V. v. Chrishtop, A. Y. Prilepskii, V. G. Nikonorova and V. A. Mironov, Nanosafety vs. nanotoxicology: adequate animal models for testing in vivo toxicity of nanoparticles, Toxicology, 2021, 462, DOI: 10.1016/j.tox.2021.152952.
- 21 M. F. Hallock, P. Greenley, L. DiBerardinis and D. Kallin, Potential risks of nanomaterials and how to safely handle materials of uncertain toxicity, J. Chem. Health Saf., 2009, **16**, 16–23.
- 22 S. K. Sohaebuddin, P. T. Thevenot, D. Baker, J. W. Eaton and L. Tang, Nanomaterial cytotoxicity is composition, size, and cell type dependent, Part. Fibre Toxicol., 2010, 7, 22.
- 23 A. A. Alshatwi, P. V. Subbarayan, E. Ramesh, A. A. Al-Hazzani, M. A. Alsaif and A. A. Alwarthan, Aluminium nanoparticles induce mitochondrial-mediated oxidative stress and alter the expression of antioxidant enzymes in human mesenchymal stem cells, Food Addit. Contam., Part A, 2013, 30, 1-10.
- 24 V. A. Senapati, A. Kumar, G. S. Gupta, A. K. Pandey and A. Dhawan, ZnO nanoparticles induced inflammatory response and genotoxicity in human blood cells: a mechanistic approach, Food Chem. Toxicol., 2015, 85,
- 25 X. Yuan, W. Nie, Z. He, J. Yang, B. Shao, X. Ma, X. Zhang, Z. Bi, L. Sun, X. Liang, Y. Tie, Y. Liu, F. Mo, D. Xie, Y. Wei

- and X. Wei, Carbon black nanoparticles induce cell necrosis through lysosomal membrane permeabilization and cause subsequent inflammatory Theranostics, 2020, 10, 4589-4605.
- 26 D. R. Nogueira, C. M. B. Rolim and A. A. Farooqi, Nanoparticle induced oxidative stress in cancer cells: adding new pieces to an incomplete jigsaw puzzle, Asian Pac. J. Cancer Prev., 2014, 15, 4739-4743.
- 27 V. Ramalingam, S. Revathidevi, T. Shanmuganayagam, L. Muthulakshmi and R. Rajaram, Biogenic gold nanoparticles induce cell cycle arrest through oxidative stress and sensitize mitochondrial membranes in A549 lung cancer cells, RSC Adv., 2016, 6, 20598-20608.
- 28 P. P. Fu, Q. Xia, H.-M. Hwang, P. C. Ray and H. Yu, Mechanisms of nanotoxicity: Generation of reactive oxygen species, J. Food Drug Anal., 2014, 22, 64-75.
- 29 L. Zhang, L. Wu, Y. Mi and Y. Si, Silver nanoparticles induced cell apoptosis, membrane damage of Azotobacter vinelandii and Nitrosomonas europaea via generation of reactive oxygen species, Bull. Environ. Contam. Toxicol., 2019, 103, 181-186.
- 30 T. H. Shin, C. Seo, D. Y. Lee, M. Ji, B. Manavalan, S. Basith, S. K. Chakkarapani, S. H. Kang, G. Lee, M. J. Paik and C. B. Park, Silica-coated magnetic nanoparticles induce glucose metabolic dysfunction in vitro via the generation of reactive oxygen species, Arch. Toxicol., 2019, 93, 1201-
- 31 H. Zhao, L. Chen, G. Zhong, Y. Huang, X. Zhang, C. Chu, L. Chen and M. Wang, Titanium dioxide nanoparticles induce mitochondrial dynamic imbalance and damage in HT22 Cells, J. Nanomater., 2019, 2019, 1-16.
- 32 J. Duan, Y. Yu, Y. Li, Y. Yu, Y. Li, X. Zhou, P. Huang and Z. Sun, Toxic effect of silica nanoparticles on endothelial cells through DNA damage response via Chk1-dependent G2/M checkpoint, PLoS One, 2013, 8, e62087.
- 33 A. Shvedova, V. Castranova, E. Kisin, D. Schwegler-Berry, A. Murray, V. Gandelsman, A. Maynard and P. Baron, Exposure to carbon nanotube material: assessment of nanotube cytotoxicity using human keratinocyte cells, J. Toxicol. Environ. Health, Part A, 2003, 66, 1909-1926.
- 34 A. Magrez, S. Kasas, V. Salicio, N. Pasquier, J. W. Seo, M. Celio, S. Catsicas, B. Schwaller and L. Forró, Cellular toxicity of carbon-based nanomaterials, Nano Lett., 2006, 6, 1121-1125.
- 35 H. A. Jeng and J. Swanson, Toxicity of metal oxide nanoparticles in mammalian cells, J. Environ. Sci. Health, Part A: Toxic/Hazard. Subst. Environ. Eng., 2006, 41, 2699-2711.
- 36 J. Blacklock, A. Vetter, A. Lankenau, D. Oupický and H. Möhwald, Tuning the mechanical properties of bioreducible multilayer films for improved cell adhesion and transfection activity, Biomaterials, 2010, 31, 7167-7174.
- 37 H. Bahadar, F. Maqbool, K. Niaz and M. Abdollahi, Toxicity of nanoparticles and an overview of current experimental models, Iran. Biomed. J., 2016, 20, 1-11.

- 38 Z. Wei, L. Chen, D. M. Thompson and L. D. Montoya, Effect of particle size on *in vitro* cytotoxicity of titania and alumina nanoparticles, J. Exp. Nanosci., 2014, 9, 625-638.
- 39 E. Fröhlich, Role of omics techniques in the toxicity testing of nanoparticles, J. Nanobiotechnol., 2017, 15, 84.
- 40 C. B. Clish, Metabolomics: an emerging but powerful tool for precision medicine, Molecular Case Studies, 2015, 1, a000588.
- 41 A. Klassen, A. T. Faccio, G. A. B. Canuto, P. L. R. da Cruz, H. C. Ribeiro, M. F. M. Tavares and A. Sussulini, in Advances in Experimental Medicine and Biology, 2017, pp.
- 42 T. Peng, C. Wei, F. Yu, J. Xu, Q. Zhou, T. Shi and X. Hu, Predicting nanotoxicity by an integrated machine learning and metabolomics approach, Environ. Pollut., 2020, 267, 115434.
- 43 S. Behzadi, V. Serpooshan, W. Tao, M. A. Hamaly, Y. Alkawareek, E. C. Dreaden, D. Brown, A. M. Alkilany, O. C. Farokhzad and M. Mahmoudi, Cellular uptake of nanoparticles: journey inside the cell, Chem. Soc. Rev., 2017, 46, 4218-4244.
- 44 Ž. Krpetić, S. Anguissola, D. Garry, P. M. Kelly and K. A. Dawson, in Advances in Experimental Medicine and Biology, 2014, pp. 135-156.
- 45 D. Nierenberg, A. R. Khaled and O. Flores, Formation of a protein corona influences the biological identity of nanomaterials, Rep. Practical Oncol. Radiother., 2018, 23, 300-308.
- 46 D. Manzanares and V. Ceña, Endocytosis: the nanoparticle and submicron nanocompounds gateway into the cell, Pharmaceutics, 2020, 12, 371.
- 47 S. Gordon, Phagocytosis: an immunobiologic process, Immunity, 2016, 44, 463-475.
- 48 E. Colucci-Guyon, F. Niedergang, B. J. Wallar, J. Peng, A. S. Alberts and P. Chavrier, A role for mammalian diaphanous-related formins in complement receptor (CR3)-mediated phagocytosis in macrophages, Curr. Biol., 2005, 15, 2007-2012.
- 49 M. Mahmoudi, K. Azadmanesh, M. A. Shokrgozar, W. S. Journeay and S. Laurent, Effect of nanoparticles on the cell life cycle, Chem. Rev., 2011, 111, 3407-3432.
- 50 L. Kou, J. Sun, Y. Zhai and Z. He, The endocytosis and intracellular fate of nanomedicines: implication for rational design, Asian J. Pharm. Sci., 2013, 8, 1-10.
- 51 K. Hirota and H. Ter, in Molecular Regulation of Endocytosis, 2012, DOI: 10.5772/45820.
- 52 J. P. Lim and P. A. Gleeson, Macropinocytosis: an endocytic pathway for internalising large gulps, Immunol. Cell Biol., 2011, 89, 836-843, DOI: 10.1038/icb.2011.20.
- 53 I. H. Pastan and M. C. Willingham, Receptor-mediated endocytosis of hormones in cultured cells, Annu. Rev. Physiol., 1981, 43, 239-250.
- 54 O. Kovtun, V. A. Tillu, N. Ariotti, R. G. Parton and B. M. Collins, Cavin family proteins and the assembly of caveolae, J. Cell Sci., 2015, 128(7), 1269-1278, DOI: 10.1242/jcs.167866.

- 55 P. Lajoie and I. R. Nabi, Lipid rafts, caveolae, and their endocytosis, Int. Rev. Cell Mol. Biol., 2010, 282(C), 135-163, DOI: 10.1016/S1937-6448(10)82003-9.
- 56 C. F. Jones and D. W. Grainger, In vitro assessments of nanomaterial toxicity, Adv. Drug Delivery Rev., 2009, 61(6), 438-456, DOI: 10.1016/j.addr.2009.03.005.
- 57 M. Zhu, G. Nie, H. Meng, T. Xia, A. Nel and Y. Zhao, Physicochemical properties determine nanomaterial cellular uptake, transport, and fate, Acc. Chem. Res., 2013, 46(3), 622-631, DOI: 10.1021/ar300031y.
- 58 C. Huang, X. Chen, Z. Xue and T. Wang, Effect of structure: a new insight into nanoparticle assemblies from inanimate to animate, Sci. Adv., 2020, 6(20), eaba1321, DOI: 10.1126/ sciadv.aba1321.
- 59 L. Li, W. S. Xi, O. Su, Y. Li, G. H. Yan, Y. Liu, H. Wang and A. Cao, Unexpected size effect: the interplay between different sized nanoparticles in their cellular uptake, Small, 2019, 15(38), 1901687, DOI: 10.1002/smll.201901687.
- 60 S. Shrestha, B. Wang and P. Dutta, Nanoparticle processing: understanding and controlling aggregation, Adv. Colloid Interface Sci., 2020, 279, 102162, DOI: 10.1016/ j.cis.2020.102162.
- 61 C. Hui, C. Shen, T. Yang, L. Bao, J. Tian, H. Ding, C. Li and H.-J. Gao, Large-scale Fe 3 O 4 nanoparticles soluble in water synthesized by a facile method, J. Phys. Chem. C, 2008, 112, 11336-11339.
- 62 P. Sabourian, G. Yazdani, S. S. Ashraf, M. Frounchi, S. Mashayekhan, S. Kiani and A. Kakkar, Effect of physico-chemical properties of nanoparticles on their intracellular uptake, Int. J. Mol. Sci., 2020, 21, 8019.
- 63 K. Kettler, K. Veltman, D. van de Meent, A. van Wezel and A. J. Hendriks, Cellular uptake of nanoparticles as determined by particle properties, experimental conditions, and cell type, Environ. Toxicol. Chem., 2014, 33(3), 481-492, DOI: 10.1002/etc.2470.
- 64 X. Dong, Z. Wu, X. Li, L. Xiao, M. Yang, Y. Li, J. Duan and Z. Sun, The size-dependent cytotoxicity of amorphous silica nanoparticles: a systematic review of in vitro studies, Int. J. Nanomed., 2020, 15, 9089-9113, DOI: 10.2147/IJN.S276105.
- 65 B. D. Chithrani and W. C. W. Chan, Elucidating the mechanism of cellular uptake and removal of proteincoated gold nanoparticles of different sizes and shapes, Nano Lett., 2007, 7(6), 1542-1550, DOI: 10.1021/nl070363y.
- 66 B. D. Chithrani, A. A. Ghazani and W. C. W. Chan, Determining the size and shape dependence of gold nanoparticle uptake into mammalian cells, Nano Lett., 2006, **6**(4), 662–668, DOI: **10.1021/nl0523960**.
- 67 H. Sun, C. Jiang, L. Wu, X. Bai and S. Zhai, Cytotoxicityrelated bioeffects induced by nanoparticles: the role of surface chemistry, Front. Bioeng. Biotechnol., 2019, 7, 414, DOI: 10.3389/fbioe.2019.00414.
- 68 A. R. Gliga, S. Skoglund, I. Odnevall Wallinder, B. Fadeel and H. L. Karlsson, Size-dependent cytotoxicity of silver nanoparticles in human lung cells: The role of cellular uptake, agglomeration and Ag release, Part. Fibre Toxicol., 2014, 11(1), 11, DOI: 10.1186/1743-8977-11-11.

- 69 Z. Wu, S. Yang and W. Wu, Shape control of inorganic nanoparticles from solution, Nanoscale, 2016, 8(3), 1237-1259, DOI: 10.1039/c5nr07681a.
- 70 A. Woźniak. A. Malankowska, G. Nowaczyk. B. F. Grześkowiak, K. Tuśnio, R. Słomski, A. Zaleska-Medynska and S. Jurga, Size and shape-dependent cytotoxicity profile of gold nanoparticles for biomedical applications, J. Mater. Sci.: Mater. Med., 2017, 28(6), 92, DOI: 10.1007/s10856-017-5902-y.
- 71 A. Asati, S. Santra, C. Kaittanis and J. M. Perez, Surfacecharge-dependent cell localization and cytotoxicity of cerium oxide nanoparticles, ACS Nano, 2010, 4(9), 5321-5331, DOI: 10.1021/nn100816s.
- 72 S. Jeon, J. Clavadetscher, D. K. Lee, S. v. Chankeshwara, M. Bradley and W. S. Cho, Surface charge-dependent cellular uptake of polystyrene nanoparticles, Nanomaterials, 2018, 8(12), 1028, DOI: 10.3390/ NANO8121028.
- 73 R. Schauer, Sialic acids as regulators of molecular and cellular interactions, Curr. Opin. Struct. Biol., 2009, 19, 507-514, DOI: 10.1016/j.sbi.2009.06.003.
- 74 E. Fröhlich, The role of surface charge in cellular uptake and cytotoxicity of medical nanoparticles, Int. J. Nanomed., 2012, 7, 5577-5591, DOI: 10.2147/IJN.S36111.
- 75 C. He, Y. Hu, L. Yin, C. Tang and C. Yin, Effects of particle size and surface charge on cellular uptake and biodistribution of polymeric nanoparticles, Biomaterials, 2010, 31(13), 3657-3666. DOI: 10.1016/ j.biomaterials.2010.01.065.
- 76 E. A. Untener, K. K. Comfort, E. I. Maurer, C. M. Grabinski, D. A. Comfort and S. M. Hussain, Tannic acid coated gold nanorods demonstrate a distinctive form of endosomal uptake and unique distribution within cells, ACS Appl. Mater. Interfaces, 2013, 5(17), 8366-8373, DOI: 10.1021/ am402848q.
- 77 N. M. Schaeublin, L. K. Braydich-Stolle, A. M. Schrand, M. Miller, J. Hutchison, J. J. Schlager and S. M. Hussain, Surface charge of gold nanoparticles mediates mechanism of toxicity, Nanoscale, 2011, 3(2), 410-420, DOI: 10.1039/c0nr00478b.
- 78 J. Li, J. Wang, Y. Yan, J. Zhang and Z. Li, Understanding the role of hydrophobicity arrangement in cellular uptake of synthetic virus-like nanoparticles, Chem. Phys. Lett., 2021, 766, 138336, DOI: 10.1016/j.cplett.2021.138336.
- 79 X. Bai, S. Wang, X. Yan, H. Zhou, J. Zhan, S. Liu, V. K. Sharma, G. Jiang, H. Zhu and B. Yan, Regulation of cell uptake and cytotoxicity by nanoparticle core under the controlled shape, size, and surface chemistries, ACS Nano, 2020, 14(1), 289-302, DOI: 10.1021/acsnano.9b04407.
- 80 Y. Li, X. Chen and N. Gu, Computational investigation of interaction between nanoparticles and membranes: hydrophobic/hydrophilic effect, J. Phys. Chem. B, 2008, 112(51), 16647–16653, DOI: 10.1021/jp8051906.
- 81 T. Muthukumarasamyvel, G. Rajendran, D. Santhana Panneer, J. Kasthuri, K. Kathiravan and N. Rajendiran, Role of surface hydrophobicity of dicationic amphiphilestabilized gold nanoparticles on A549 lung cancer cells,

ACS Omega, 2017, 2(7), 3527–3538, DOI: 10.1021/acsomega.7b00353.

- 82 H. Sun, Y. Liu, X. Bai, X. Zhou, H. Zhou, S. Liu and B. Yan, Induction of oxidative stress and sensitization of cancer cells to paclitaxel by gold nanoparticles with different charge densities and hydrophobicities, *J. Mater. Chem. B*, 2018, **6**(11), 1633–1639, DOI: **10.1039/c7tb03153j**.
- 83 S. Jambhrunkar, Z. Qu, A. Popat, J. Yang, O. Noonan, L. Acauan, Y. Ahmad Nor, C. Yu and S. Karmakar, Effect of surface functionality of silica nanoparticles on cellular uptake and cytotoxicity, *Mol. Pharm.*, 2014, 11(10), 3642– 3655, DOI: 10.1021/mp500385n.
- 84 X. R. Shao, X. Q. Wei, X. Song, L. Y. Hao, X. X. Cai, Z. R. Zhang, Q. Peng and Y. F. Lin, Independent effect of polymeric nanoparticle zeta potential/surface charge, on their cytotoxicity and affinity to cells, *Cell Proliferation*, 2015, 48(4), 465–474, DOI: 10.1111/cpr.12192.
- 85 W. Tao, J. Zhang, X. Zeng, D. Liu, G. Liu, X. Zhu, Y. Liu, Q. Yu, L. Huang and L. Mei, Blended Nanoparticle System Based on Miscible Structurally Similar Polymers: A Safe, Simple, Targeted, and Surprisingly High Efficiency Vehicle for Cancer Therapy, *Adv. Healthcare Mater.*, 2015, 4(8), 1203–1214, DOI: 10.1002/adhm.201400751.
- 86 T. Lund, M. F. Callaghan, P. Williams, M. Turmaine, C. Bachmann, T. Rademacher, I. M. Roitt and R. Bayford, The influence of ligand organization on the rate of uptake of gold nanoparticles by colorectal cancer cells, *Biomaterials*, 2011, 32(36), 9776–9784, DOI: 10.1016/ j.biomaterials.2011.09.018.
- 87 Y. C. Yeh, K. Saha, B. Yan, O. R. Miranda, X. Yu and V. M. Rotello, The role of ligand coordination on the cytotoxicity of cationic quantum dots in HeLa cells, *Nanoscale*, 2013, 5(24), 12140–12143, DOI: 10.1039/c3nr04037b.
- 88 L. Ding, C. Yao, X. Yin, C. Li, Y. Huang, M. Wu, B. Wang, X. Guo, Y. Wang and M. Wu, Size, shape, and protein corona determine cellular uptake and removal mechanisms of gold nanoparticles, *Small*, 2018, 14(42), 1801451, DOI: 10.1002/smll.201801451.
- 89 C. Carnovale, G. Bryant, R. Shukla and V. Bansal, Identifying trends in gold nanoparticle toxicity and uptake: size, shape, capping ligand, and biological corona, *ACS Omega*, 2019, 4(1), 242–256, DOI: 10.1021/acsomega.8b03227.
- 90 K. P. Steckiewicz, E. Barcinska, A. Malankowska, A. Zauszkiewicz-Pawlak, G. Nowaczyk, A. Zaleska-Medynska and I. Inkielewicz-Stepniak, Impact of gold nanoparticles shape on their cytotoxicity against human osteoblast and osteosarcoma in in vitro model. Evaluation of the safety of use and anti-cancer potential, *J. Mater.* Sci.: Mater. Med., 2019, 30(2), 22, DOI: 10.1007/s10856-019-6221-2.
- 91 B. Zhang, P. Sai Lung, S. Zhao, Z. Chu, W. Chrzanowski and Q. Li, Shape dependent cytotoxicity of PLGA-PEG nanoparticles on human cells, *Sci. Rep.*, 2017, 7(1), 7315, DOI: 10.1038/s41598-017-07588-9.

- 92 L. Dong, S. Tang, F. Deng, Y. Gong, K. Zhao, J. Zhou, D. Liang, J. Fang, M. Hecker, J. P. Giesy, X. Bai and H. Zhang, Shape-dependent toxicity of alumina nanoparticles in rat astrocytes, *Sci. Total Environ.*, 2019, 690, 158–166, DOI: 10.1016/j.scitotenv.2019.06.532.
- 93 D. T. Savage, J. Z. Hilt and T. D. Dziubla, In vitro methods for assessing nanoparticle toxicity, *Methods Mol. Biol.*, 2019, **1894**, 1–29, DOI: **10.1007/978-1-4939-8916-4** 1.
- 94 V. Kumar, N. Sharma and S. S. Maitra, In vitro and in vivo toxicity assessment of nanoparticles, *Int. Nano Lett.*, 2017, 7(4), 243–256, DOI: 10.1007/s40089-017-0221-3.
- 95 J. M. Veranth, in *Nanoscience and Nanotechnology*, John Wiley & Sons, Inc., Hoboken, NJ, USA, 2008, pp. 261–286.
- 96 C. M. Sayes, K. L. Reed and D. B. Warheit, Assessing toxicology of fine and nanoparticles: comparing in vitro measurements to in vivo pulmonary toxicity profiles, *Toxicol. Sci.*, 2007, 97(1), 163–180, DOI: 10.1093/toxsci/kfm018.
- 97 X. Han, R. Gelein, N. Corson, P. Wade-Mercer, J. Jiang, P. Biswas, J. N. Finkelstein, A. Elder and G. Oberdörster, Validation of an LDH assay for assessing nanoparticle toxicity, *Toxicology*, 2011, 287(1–3), 99–104, DOI: 10.1016/j.tox.2011.06.011.
- 98 O. Stueker, V. A. Ortega, G. G. Goss and M. Stepanova, Understanding interactions of functionalized nanoparticles with proteins: a case study on lactate dehydrogenase, *Small*, 2014, **10**(10), 2006–2021, DOI: **10.1002/smll.201303639**.
- 99 B. Drasler, P. Sayre, K. G. Steinhäuser, A. Petri-Fink and B. Rothen-Rutishauser, In vitro approaches to assess the hazard of nanomaterials, *NanoImpact*, 2017, 8, 99–116, DOI: 10.1016/j.impact.2017.08.002.
- 100 B. Díaz, C. Sánchez-Espinel, M. Arruebo, J. Faro, E. de Miguel, S. Magadán, C. Yagüe, R. Fernández-Pacheco, M. R. Ibarra, J. Santamaría and Á. González-Fernández, Assessing methods for blood cell cytotoxic responses to inorganic nanoparticles and nanoparticle aggregates, *Small*, 2008, 4(11), 2025–2034, DOI: 10.1002/ smll.200800199.
- 101 A. Hoshino, K. Fujioka, T. Oku, M. Suga, Y. F. Sasaki, T. Ohta, M. Yasuhara, K. Suzuki and K. Yamamoto, Physicochemical properties and cellular toxicity of nanocrystal quantum dots depend on their surface modification, *Nano Lett.*, 2004, 4(11), 2163–2169, DOI: 10.1021/nl048715d.
- 102 P. D. Ray, B. W. Huang and Y. Tsuji, Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling, *Cell. Signalling*, 2012, 24, 981–990, DOI: 10.1016/j.cellsig.2012.01.008.
- 103 L. He, T. He, S. Farrar, L. Ji, T. Liu and X. Ma, Antioxidants maintain cellular redox homeostasis by elimination of reactive oxygen species, *Cell. Physiol. Biochem.*, 2017, 44(2), 532–553, DOI: 10.1159/000485089.
- 104 M. Nita and A. Grzybowski, The role of the reactive oxygen species and oxidative stress in the pathomechanism of the age-related ocular diseases and other pathologies of the anterior and posterior eye segments in adults, *Oxid. Med.*

- Cell. Longevity, 2016, 2016, 3164734, DOI: 10.1155/2016/ 3164734.
- 105 Y. Higashi, J. Mazumder, H. Yoshikawa, M. Saito and E. Tamiya, Chemically regulated ROS generation from nanoparticles for enzyme-free electrochemiluminescent immunosensing, Anal. Chem., 2018. **90**(9), 5773-5780. DOI: 10.1021/ acs.analchem.8b00118.
- 106 M. Katerji, M. Filippova and P. Duerksen-Hughes, Approaches and methods to measure oxidative stress in clinical samples: research applications in the cancer field, Oxid. Med. Cell. Longevity, 2019, 2019, 1279250, DOI: 10.1155/2019/1279250.
- 107 B. Kalyanaraman, V. Darley-Usmar, K. J. A. Davies, P. A. Dennery, H. J. Forman, M. B. Grisham, G. E. Mann, K. Moore, L. J. Roberts and H. Ischiropoulos, Measuring reactive oxygen and nitrogen species with fluorescent probes: challenges and limitations, Free Radical Biol. Med., 2012, 52, 1-6. DOI 10.1016/ j.freeradbiomed.2011.09.030.
- 108 P. Whongsiri, S. Phoyen and C. Boonla, Oxidative stress in urothelial carcinogenesis: Measurements of protein carbonylation and intracellular production of reactive oxygen species, Methods Mol. Biol., 2018, 1655, 109-117, DOI: 10.1007/978-1-4939-7234-0_9.
- 109 M. Jeszka-Skowron, T. Podgórski and B. Czarczyńska-Goślińska, Determination of Antioxidant Biomarkers in Biological Fluids, Springer, Cham, 2021, DOI: 10.1007/978-3-030-61879-7_11.
- 110 A. G. Fantel, Reactive oxygen species in developmental toxicity: review and hypothesis, Teratology, 1996, 53, 196-217, DOI: 10.1002/(SICI)1096-9926(199603)53:3<196::AID-TERA7>3.0.CO;2-2.
- 111 H. Ohkawa, N. Ohishi and K. Yagi, Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction, Anal. Biochem., 1979, 95(2), 351-358, DOI: 10.1016/0003-2697(79)90738-3.
- 112 M. Pylväs, U. Puistola, L. Laatio, S. Kauppila and P. Karihtala, Elevated serum 8-OHdG is associated with poor prognosis in epithelial ovarian cancer, Anticancer Res., 2011, 31(4), 1411-1415.
- 113 S. Toyokuni, T. Tanaka, Y. Hattori, Y. Nishiyama, A. Yoshida, K. Uchida, H. Hiai, H. Ochi and T. Osawa, Quantitative immunohistochemical determination of 8hydroxy-2'- deoxyguanosine by a monoclonal antibody N45.1: its application to ferric nitrilotriacetate-induced renal carcinogenesis model, Lab. Invest., 1997, 76(3), 365-
- 114 C.-C. Chiou, P.-Y. Chang, E.-C. Chan, T.-L. Wu, K.-C. Tsao and J. T. Wu, Urinary 8-hydroxydeoxyguanosine and its analogs as DNA marker of oxidative stress: development of an ELISA and measurement in both bladder and prostate cancers, Clin. Chim. Acta, 2003, 334, 87-94, DOI: 10.1016/S0009-8981(03)00191-8.
- 115 N. Golbamaki, B. Rasulev, A. Cassano, R. L. Marchese Robinson, E. Benfenati, J. Leszczynski and T. D. Cronin, Genotoxicity of metal oxide

- nanomaterials: review of recent data and discussion of possible mechanisms, Nanoscale, 2015, 7, 2154-2198, DOI: 10.1039/c4nr06670g.
- 116 A. Aranda, L. Sequedo, L. Tolosa, G. Quintas, E. Burello, J. v. Castell and L. Gombau, Dichloro-dihydro-fluorescein diacetate (DCFH-DA) assay: a quantitative method for oxidative stress assessment of nanoparticle-treated cells, Toxicol. in Vitro, 2013, 27(2), 954-963, DOI: 10.1016/ j.tiv.2013.01.016.
- 117 T. Pfaller, R. Colognato, I. Nelissen, F. Favilli, E. Casals, D. Ooms, H. Leppens, J. Ponti, R. Stritzinger, V. Puntes, D. Boraschi, A. Duschl and G. J. Oostingh, The suitability of different cellular in vitro immunotoxicity and genotoxicity methods for the analysis of nanoparticleinduced events, Nanotoxicology, 2010, 4(1), 52-72, DOI: 10.3109/17435390903374001.
- 118 S. Kinoshita, S. Akira and T. Kishimoto, A member of the C/ EBP family, NF-IL6 beta, forms a heterodimer and transcriptionally synergizes with NF-IL6, Proc. Natl. Acad. Sci. U. S. A., 1992, 89, 1473-1476, DOI: 10.1073/ pnas.89.4.1473.
- 119 F. Yu, X. Zhang, L. Gao, H. Xue, L. Liu, S. Wang, S. Chen and L. Huang, LncRNA loc105377478 promotes NPs-Nd2O3induced inflammation in human bronchial epithelial cells through the ADIPOR1/NF-kB axis, Ecotoxicol. Environ. Saf., 2021, 208, 111609, DOI: 10.1016/ j.ecoenv.2020.111609.
- 120 S. Tada-Oikawa, M. Eguchi, M. Yasuda, K. Izuoka, A. Ikegami, S. Vranic, S. Boland, L. Tran, G. Ichihara and Ichihara, Functionalized surface-charged nanoparticles induce pro-inflammatory responses, but are not lethal to caco-2 cells, Chem. Res. Toxicol., 2020, 33(5), 1226-1236, DOI: 10.1021/acs.chemrestox.9b00478.
- 121 H. Lin, G. Fu, Q. Yu, Z. Wang, Y. Zuo, Y. Shi, L. Zhang, Y. Gu, L. Qin and T. Zhou, Carbon black nanoparticles induce HDAC6-mediated inflammatory responses in 16HBE cells, Toxicol. Ind. Health, 2020, 36(10), 759-768, DOI: 10.1177/0748233720947214.
- 122 R. Guadagnini, B. Halamoda Kenzaoui, L. Walker, G. Pojana, Z. Magdolenova, D. Bilanicova, M. Saunders, L. Juillerat-Jeanneret, A. Marcomini, A. Huk, M. Dusinska, L. M. Fjellsbo, F. Marano and S. Boland, Toxicity screenings of nanomaterials: Challenges due interference with assay processes and components of classic in vitro tests, Nanotoxicology, 2015, 9(S1), 13-24, DOI: 10.3109/17435390.2013.829590.
- 123 J. P. Piret, O. M. Bondarenko, M. S. P. Boyles, M. Himly, A. R. Ribeiro, F. Benetti, C. Smal, B. Lima, A. Potthoff, M. Simion, E. Dumortier, P. E. C. Leite, L. B. Balottin, J. M. Granjeiro, A. Ivask, A. Kahru, I. Radauer-Preiml, U. Tischler, A. Duschl, C. Saout, S. Anguissola, A. Haase, A. Jacobs, I. Nelissen, S. K. Misra and O. Toussaint, Pan-European inter-laboratory studies on a panel of in vitro cytotoxicity pro-inflammation and assays for nanoparticles, Arch. Toxicol., 2017, 91(6), 2315-2330, DOI: 10.1007/s00204-016-1897-2.

124 S. Elmore, Apoptosis: A Review of Programmed Cell Death, Toxicol. Pathol., 2007, 35, 495-516, DOI: 10.1080/ 01926230701320337.

- 125 P. R. Ruffolo, The pathogenesis of necrosis. i. correlated light and electron microscopic observations of the myocardial necrosis induced by the intravenous injection of papain, Am. J. Pathol., 1964, 45, 741-756.
- 126 M. Ahamed, M. J. Akhtar, M. A. Siddiqui, J. Ahmad, J. Musarrat, A. A. Al-Khedhairy, M. S. AlSalhi and S. A. Alrokayan, Oxidative stress mediated apoptosis induced by nickel ferrite nanoparticles in cultured A549 cells, Toxicology, 2011, 283(2-3), 101-108, DOI: 10.1016/ j.tox.2011.02.010.
- 127 E. J. Park, J. Yi, K. H. Chung, D. Y. Ryu, J. Choi and K. Park, Oxidative stress and apoptosis induced by titanium dioxide nanoparticles in cultured BEAS-2B cells, Toxicol. Lett., 2008, 180(3), 222-229, DOI: 10.1016/j.toxlet.2008.06.869.
- 128 M. T. Zhu, Y. Wang, W. Y. Feng, B. Wang, M. Wang, H. Ouyang and Z. F. Chai, Oxidative stress and apoptosis induced by iron oxide nanoparticles in cultured human umbilical endothelial cells, J. Nanosci. Nanotechnol., 2010, 10, 8584-8590, DOI: 10.1166/jnn.2010.2488.
- 129 J. Xu, P. Xu, Z. Li, J. Huang and Z. Yang, Oxidative stress and apoptosis induced by hydroxyapatite nanoparticles in C6 cells, J. Biomed. Mater. Res., 2012, 100 A(3), 738-745, DOI: 10.1002/jbm.a.33270.
- 130 Y. Xue, T. Zhang, B. Zhang, F. Gong, Y. Huang and M. Tang, Cytotoxicity and apoptosis induced by silver nanoparticles in human liver HepG2 cells in different dispersion media, J. Appl. Toxicol., 2016, 36(3), 352-360, DOI: 10.1002/ jat.3199.
- 131 T. L. Reus, B. H. Marcon, A. C. C. Paschoal, I. R. S. Ribeiro, M. B. Cardoso, B. Dallagiovanna and A. M. de Aguiar, Dosedependent cell necrosis induced by silica nanoparticles, Toxicol. in Vitro, 2020, 63, 104723, DOI: 10.1016/ j.tiv.2019.104723.
- 132 W. Fiers, R. Beyaert, W. Declercq and P. Vandenabeele, More than one way to die: apoptosis, necrosis and reactive oxygen damage, Oncogene, 1999, 18, 7719-7730, DOI: 10.1038/sj.onc.1203249.
- 133 G. Kumar, H. Degheidy, B. J. Casey and P. L. Goering, Flow cytometry evaluation of in vitro cellular necrosis and apoptosis induced by silver nanoparticles, Food Chem. Toxicol., 2015, 85, 45-51, DOI: 10.1016/j.fct.2015.06.012.
- 134 I. Vermes, C. Haanen, H. Steffens-Nakken and C. Reutellingsperger, A novel assay for apoptosis Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V, J. Immunol. Methods, 1995, 184(1), 39-51, DOI: 10.1016/ 0022-1759(95)00072-I.
- 135 S. Vafaei, S. A. Sadat Shandiz and Z. Piravar, Zincphosphate nanoparticles as a novel anticancer agent: an in vitro evaluation of their ability to induce apoptosis, Biol. Trace Elem. Res., 2020, 198(1), 109-117, DOI: 10.1007/s12011-020-02054-6.
- 136 S. Bancos, D.-H. Tsai, V. Hackley, J. L. Weaver and K. M. Tyner, Evaluation of viability and proliferation

- profiles on macrophages treated with silica nanoparticles in vitro via plate-based, flow cytometry, and coulter counter assays, ISRN Nanotechnol., 2012, 2012, 454072, DOI: 10.5402/2012/454072.
- 137 K. J. Ong, T. J. MacCormack, R. J. Clark, J. D. Ede, V. A. Ortega, L. C. Felix, M. K. M. Dang, G. Ma, H. Fenniri, J. G. C. Veinot and G. G. Goss, Widespread nanoparticle-assay interference: implications nanotoxicity testing, PLoS One, 2014, 9(3), e90650, DOI: 10.1371/journal.pone.0090650.
- 138 B. Naseer, G. Srivastava, O. S. Qadri, S. A. Faridi, R. U. Islam and K. Younis, Importance and health hazards of nanoparticles used in the food industry, Nanotechnol. Rev., 2018, 7(6), 623-641, DOI: 10.1515/ntrev-2018-0076.
- 139 T. Ramirez, M. Daneshian, H. Kamp, F. Y. Bois, M. R. Clench, M. Coen, B. Donley, S. M. Fischer, R. Ekman, E. Fabian, C. Guillou, J. Heuer, H. T. Hogberg, H. Jungnickel, H. C. Keun, G. Krennrich, Krupp, A. Luch, F. Noor, E. Peter, B. Riefke, M. Seymour, N. Skinner, L. Smirnova, E. Verheij, S. Wagner, T. Hartung, B. van Ravenzwaay and M. Leist, T4 Report* Metabolomics in toxicology and preclinical research, ALTEX, 2013, 30(2), 209-225, DOI: 10.14573/ altex.2013.2.209.
- 140 Z. Pang, J. Chong, G. Zhou, D. A. de Lima Morais, L. Chang, M. Barrette, C. Gauthier, P.-É. Jacques, S. Li and J. Xia, MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional insights, Nucleic Acids Res., 2021, 49, W388-W396, DOI: 10.1093/nar/gkab382.
- 141 J. R. Idle and F. J. Gonzalez, Metabolomics, Cell Metab., 2007, 6, 348-351, DOI: 10.1016/j.cmet.2007.10.005.
- 142 D. F. Gomez-Casati, M. I. Zanor and M. v. Busi, Metabolomics in plants and humans: applications in the prevention and diagnosis of diseases, BioMed Res. Int., 2013, 2013, 792527, DOI: 10.1155/2013/792527.
- 143 V. Masutin, C. Kersch and S. Schmitz-Spanke, A systematic review: metabolomics-based identification of altered metabolites and pathways in the skin caused by internal and external factors, Exp. Dermatol., 2022, 700-714, DOI: 10.1111/exd.14529.
- 144 S. Deja, I. Porebska, A. Kowal, A. Zabek, W. Barg, K. Pawelczyk, I. Stanimirova, M. Daszykowski, Korzeniewska, R. Jankowska and P. Mlynarz, Metabolomics provide new insights on lung cancer staging and discrimination from chronic obstructive pulmonary disease, J. Pharm. Biomed. Anal., 2014, 100, 369-380, DOI: 10.1016/j.jpba.2014.08.020.
- 145 S. Deja, E. Barg, P. Młynarz, A. Basiak and E. Willak-Janc, 1H NMR-based metabolomics studies of urine reveal differences between type 1 diabetic patients with high and low HbAc1 values, J. Pharm. Biomed. Anal., 2013, 83, 43-48, DOI: 10.1016/j.jpba.2013.04.017.
- Mika, W. Wojtowicz, A. Ząbek, P. Młynarz, M. Chmielewski, T. Sledzinski and P. Stepnowski, Application of nuclear magnetic resonance spectroscopy for the detection of metabolic disorders in patients with

- moderate kidney insufficiency, J. Pharm. Biomed. Anal., 2018, 149, 1-8, DOI: 10.1016/j.jpba.2017.10.037.
- 147 Q. Bu, G. Yan, P. Deng, F. Peng, H. Lin, Y. Xu, Z. Cao, T. Zhou, A. Xue, Y. Wang, X. Cen and Y. L. Zhao, NMRbased metabonomic study of the sub-acute toxicity of titanium dioxide nanoparticles in rats after oral administration, Nanotechnology, 2010, 21(12), 125105, DOI: 10.1088/0957-4484/21/12/125105.
- 148 K. Grintzalis, T. N. Lawson, F. Nasser, I. Lynch and M. R. Viant, Metabolomic method to detect a metabolite amino-functionalized polystyrene corona nanoparticles, Nanotoxicology, 2019, 13(6), 783-794, DOI: 10.1080/17435390.2019.1577510.
- 149 I. Lanekoff, V. v. Sharma and C. Marques, Single-cell metabolomics: where are we and where are we going?, Curr. Opin. Biotechnol., 2022, 75, 102693, DOI: 10.1016/ j.copbio.2022.102693.
- 150 Y. Liu, Y. Cheng, T. Chen, Y. Zhang, X. Wang, A. Zhao, W. Jia, Y. Bo and C. Jin, GC/TOFMS Analysis of Endogenous Metabolites in Mouse Fibroblast Cells and Its Application in TiO2 Nanoparticle-Induced Cytotoxicity Study, Chromatographia, 2012, 75, 1301-1310, DOI: 10.1007/s10337-012-2315-4.
- 151 A. E. Steuer, L. Brockbals and T. Kraemer, Metabolomic strategies in biomarker research-new approach for indirect identification of drug consumption and sample manipulation in clinical and forensic toxicology?, Front. Chem., 2019, 7, DOI: 10.3389/fchem.2019.00319.
- 152 C. Deng, Q. Tang, Z. Yang, Z. Dai, C. Cheng, Y. Xu, X. Chen, X. Zhang and J. Su, Effects of iron oxide nanoparticles on phenotype and metabolite changes in hemp clones (Cannabis sativa L.), Front. Environ. Sci. Eng., 2022, 16, 134, DOI: 10.1007/s11783-022-1569-9.
- 153 A. Bouyahya, N. el Omari, M. Hakkour, N. el Menyiy, T. Benali, D. Kulikov, M. Karpukhin, M. A. Shariati, B. Venkidasamy, M. Thiruvengadam and I. Chamkhi, A review on transcriptomic and metabolomic responses of plants to nanopollution, Environ. Sci. Pollut. Res., 2022, 29, 22913-22929, DOI: 10.1007/s11356-022-18659-4.
- 154 Z. Chen, Z. Guo, J. Niu, N. Xu, X. Sui, H. A. Kareem, M. U. Hassan, M. Yan, Q. Zhang, Z. Wang, F. Mi, J. Kang, J. Cui and Q. Wang, Phytotoxic effect and molecular mechanism induced by graphene towards alfalfa (Medicago sativa L.) by integrating transcriptomic and metabolomics analysis, Chemosphere, 2022, 290, 133368, DOI: 10.1016/j.chemosphere.2021.133368.
- 155 D. Kumar, I. Mutreja, K. Chitcholtan and P. Sykes, Cytotoxicity and cellular uptake of different sized gold nanoparticles in ovarian cancer cells, Nanotechnology, 2017, 28(47), 475101, DOI: 10.1088/1361-6528/aa935e.
- 156 W. Shim, M. J. Paik, D. T. Nguyen, J. K. Lee, Y. Lee, J. H. Kim, E. H. Shin, J. S. Kang, H. S. Jung, S. Choi, S. Park, J. S. Shim and G. Lee, Analysis of changes in gene expression and metabolic profiles induced by silicacoated magnetic nanoparticles, ACS Nano, 2012, 6(9), 7665-7680, DOI: 10.1021/nn301113f.

- 157 L. Kou, X. Jiang, Y. Tang, X. Xia, Y. Li, A. Cai, H. Zheng, H. Zhang, V. Ganapathy, O. Yao and R. Chen, Resetting amino acid metabolism of cancer cells by ATB0,+-targeted nanoparticles for enhanced anticancer therapy, Bioact. Mater., 2022, 15-28,DOI: j.bioactmat.2021.07.009.
- 158 T. H. Shin, D. Y. Lee, H.-S. Lee, H. J. Park, M. S. Jin, M.-J. Paik, B. Manavalan, J.-S. Mo and G. Lee, Integration of metabolomics and transcriptomics in nanotoxicity studies, BMB Rep., 2018, 51, 14-20.
- 159 A. Machuca, E. Garcia-Calvo, D. S. Anunciação and J. L. Luque-Garcia, Integration of transcriptomics and metabolomics to reveal the molecular mechanisms underlying rhodium nanoparticles-based photodynamic cancer therapy, Pharmaceutics, 2021, 13(10), 1629, DOI: 10.3390/pharmaceutics13101629.
- 160 T. H. Shin, S. Nithiyanandam, D. Y. Lee, D. H. Kwon, J. S. Hwang, S. G. Kim, Y. E. Jang, S. Basith, S. Park, J. S. Mo and G. Lee, Analysis of nanotoxicity with integrated omics and mechanobiology, Nanomaterials, 2021, 11(9), 2385, DOI: 10.3390/nano11092385.
- 161 M. Lv, W. Huang, Z. Chen, H. Jiang, J. Chen, Y. Tian, Z. Zhang and F. Xu, Metabolomics techniques for nanotoxicity investigations, Bioanalysis, 2015, 7, 1527-1544, DOI: 10.4155/bio.15.83.
- 162 F. Ahmad, X. Wang and W. Li, Toxico-metabolomics of engineered nanomaterials: progress and challenges, Adv. Funct. Mater., 2019, **29**, 1904268, DOI: **10.1002**/ adfm.201904268.
- 163 W. Feng, Omic" Techniques for Nanosafety, Toxicol. Nanomater., 2016, DOI: 10.1002/9783527689125.ch12.
- 164 L. K. Schnackenberg, J. Sun and R. D. Beger, Metabolomics techniques in nanotoxicology studies, Methods Mol. Biol., 2012, 926, 141-156, DOI: 10.1007/978-1-62703-2-1_10.
- 165 M. Farré and A. N. Jha, in Environmental Metabolomics, Elsevier, 2020, pp. 259-281.
- 166 M. Mortimer, Y. Wang and P. A. Holden, Molecular Mechanisms of Nanomaterial-Bacterial Interactions Revealed by Omics-The Role of Nanomaterial Effect Level, Front. Bioeng. Biotechnol., 2021, 9, DOI: 10.3389/ fbioe.2021.683520.
- 167 E. Boisselier and D. Astruc, Gold nanoparticles in nanomedicine: preparations, imaging, therapies and toxicity, Chem. Soc. Rev., 2009, 38(6), 1759-1782, DOI: 10.1039/b806051g.
- 168 L. Wang, Y. Liu, W. Li, X. Jiang, Y. Ji, X. Wu, L. Xu, Y. Qiu, K. Zhao, T. Wei, Y. Li, Y. Zhao and C. Chen, Selective targeting of gold nanorods at the mitochondria of cancer cells: Implications for cancer therapy, Nano Lett., 2011, 11(2), 772-780, DOI: 10.1021/nl103992v.
- 169 L. Zhang, L. Wang, Y. Hu, Z. Liu, Y. Tian, X. Wu, Y. Zhao, H. Tang, C. Chen and Y. Wang, Selective metabolic effects of gold nanorods on normal and cancer cells and their application in anticancer drug screening, Biomaterials, 34(29), 7117-7126, DOI: 10.1016/ j.biomaterials.2013.05.043.

170 J. J. Ramsden, Metabolomics and Metabonomics, in *Computational Biology*, 2009, pp. 1–6.

- 171 B. Xu, M. Chen, X. Ji, Z. Mao, X. Zhang, X. Wang and Y. Xia, Metabolomic profiles delineate the potential role of glycine in gold nanorod-induced disruption of mitochondria and blood-testis barrier factors in TM-4 cells, *Nanoscale*, 2014, 6(14), 8265–8273, DOI: 10.1039/c4nr01035c.
- 172 Y. Huang, X. Lü, R. Chen and Y. Chen, Comparative study of the effects of gold and silver nanoparticles on the metabolism of human dermal fibroblasts, *Regener. Biomater.*, 2020, 7(2), 221–232, DOI: 10.1093/RB/RBZ051.
- 173 J. Z. Lindeque, A. Matthyser, S. Mason, R. Louw and C. J. F. Taute, Metabolomics reveals the depletion of intracellular metabolites in HepG2 cells after treatment with gold nanoparticles, *Nanotoxicology*, 2018, 12(3), 251– 262, DOI: 10.1080/17435390.2018.1432779.
- 174 C. J. Reinecke, G. Koekemoer, F. H. van der Westhuizen, R. Louw, J. Z. Lindeque, L. J. Mienie and I. Smuts, Metabolomics of urinary organic acids in respiratory chain deficiencies in children, *Metabolomics*, 2012, 8(2), 264–283, DOI: 10.1007/s11306-011-0309-0.
- 175 S. Gioria, J. L. Vicente, P. Barboro, R. la Spina, G. Tomasi, P. Urbán, A. Kinsner-Ovaskainen, R. François and H. Chassaigne, A combined proteomics and metabolomics approach to assess the effects of gold nanoparticles in vitro, *Nanotoxicology*, 2016, 10(6), 736–748, DOI: 10.3109/17435390.2015.1121412.
- 176 M. R. K. Ali, Y. Wu, T. Han, X. Zang, H. Xiao, Y. Tang, R. Wu, F. M. Fernández and M. A. El-Sayed, Simultaneous Time-Dependent Surface-Enhanced Raman Spectroscopy, Metabolomics, and Proteomics Reveal Cancer Cell Death Mechanisms Associated with Gold Nanorod Photothermal Therapy, J. Am. Chem. Soc., 2016, 138(47), 15434–15442, DOI: 10.1021/jacs.6b08787.
- 177 Z. Liu, L. Wang, L. Zhang, X. Wu, G. Nie, C. Chen, H. Tang and Y. Wang, Metabolic characteristics of 16HBE and A549 cells exposed to different surface modified gold nanorods, *Adv. Healthcare Mater.*, 2016, 5(18), 2363–2375, DOI: 10.1002/adhm.201600164.
- 178 L. A. Dahabiyeh, N. N. Mahmoud, M. A. Al-Natour, L. Safo, D. H. Kim, E. A. Khalil and R. Abu-Dahab, Phospholipid-gold nanorods induce energy crisis in mcf-7 cells: Cytotoxicity evaluation using lc-ms-based metabolomics approach, *Biomolecules*, 2021, 11(3), 364, DOI: 10.3390/biom11030364.
- 179 J. R. Herance, H. García, P. Gutiérrez-Carcedo, S. Navalón, A. Pineda-Lucena and M. Palomino-Schätzlein, A translational approach to assess the metabolomic impact of stabilized gold nanoparticles by NMR spectroscopy, *Analyst*, 2019, 144(4), 1265–1274, DOI: 10.1039/c8an01827h.
- 180 M. P. Schätzlein, J. Becker, D. Schulze-Sünninghausen, A. Pineda-Lucena, J. R. Herance and B. Luy, Rapid twodimensional ALSOFAST-HSQC experiment for metabolomics and fluxomics studies: application to a 13C-enriched cancer cell model treated with gold

- nanoparticles, *Anal. Bioanal. Chem.*, 2018, **410**(11), 2793–2804, DOI: **10.1007/s00216-018-0961-6**.
- 181 M. Palomino-Schätzlein, H. García, P. Gutiérrez-Carcedo, A. Pineda-Lucena and J. R. Herance, Assessment of gold nanoparticles on human peripheral blood cells by metabolic profiling with 1H-NMR spectroscopy, a novel translational approach on a patient-specific basis, *PLoS One*, 2017, 12(12), e0189748, DOI: 10.1371/journal.pone.0189748.
- 182 D. Paris, D. Melck, A. Longo, S. Napoletano, G. Carotenuto, L. Nicolais, A. Motta and E. Vitale, Metabolic response of SH-SY5Y cells to gold nanoparticles by NMR-based metabolomics analyses, *Biomed. Phys. Eng. Express*, 2016, 2(4), 045003, DOI: 10.1088/2057-1976/2/4/045003.
- 183 M. K. Rai, S. D. Deshmukh, A. P. Ingle and A. K. Gade, Silver nanoparticles: The powerful nanoweapon against multidrug-resistant bacteria, *J. Appl. Microbiol.*, 2012, 112, 841–852, DOI: 10.1111/j.1365-2672.2012.05253.x.
- 184 S. Galdiero, A. Falanga, M. Vitiello, M. Cantisani, V. Marra and M. Galdiero, Silver nanoparticles as potential antiviral agents, *Molecules*, 2011, 16, 8894–8918, DOI: 10.3390/ molecules16108894.
- 185 L. Wei, J. Lu, H. Xu, A. Patel, Z. S. Chen and G. Chen, Silver nanoparticles: Synthesis, properties, and therapeutic applications, *Drug Discovery Today*, 2015, **20**(5), 595–601, DOI: **10.1016/j.drudis.2014.11.014**.
- 186 G. Raja, Y. K. Jang, J. S. Suh, H. S. Kim, S. H. Ahn and T. J. Kim, Microcellular environmental regulation of silver nanoparticles in cancer therapy: a critical review, *Cancers*, 2020, 12, 664, DOI: 10.3390/cancers12030664.
- 187 D. Krzyzanowski, M. Kruszewski and A. Grzelak, Differential action of silver nanoparticles on abcb1 (Mdr1) and abcc1 (mrp1) activity in mammalian cell lines, *Materials*, 2021, 14(12), 3383, DOI: 10.3390/ ma14123383.
- 188 P. Paciorek, M. Zuberek and A. Grzelak, Products of lipid peroxidation as a factor in the toxic effect of silver nanoparticles, *Materials*, 2020, 13(11), 2460, DOI: 10.3390/MA13112460.
- 189 K. Sikorska, I. Gradzka, I. Wasyk, K. Brzóska, T. M. Stepkowski, M. Czerwińska and M. K. Kruszewski, The impact of Ag nanoparticles and CdTe quantum dots on expression and function of receptors involved in amyloid-β uptake by BV-2 microglial cells, *Materials*, 2020, 13(14), 3227, DOI: 10.3390/ma13143227.
- 190 R. Rajan, P. P. Huo, K. Chandran, B. Manickam Dakshinamoorthi, S. il Yun and B. Liu, A review on the toxicity of silver nanoparticles against different biosystems, *Chemosphere*, 2022, 292, 133397, DOI: 10.1016/j.chemosphere.2021.133397.
- 191 J. Carrola, V. Bastos, J. M. P. Ferreira De Oliveira, H. Oliveira, C. Santos, A. M. Gil and I. F. Duarte, Insights into the impact of silver nanoparticles on human keratinocytes metabolism through NMR metabolomics, *Arch. Biochem. Biophys.*, 2016, 589, 53–61, DOI: 10.1016/ j.abb.2015.08.022.

- 192 J. Carrola, R. J. B. Pinto, M. Nasirpour, C. S. R. Freire, A. M. Gil, C. Santos, H. Oliveira and I. F. Duarte, NMR metabolomics reveals metabolism-mediated protective effects in liver (HepG2) cells exposed to subtoxic levels of silver nanoparticles, J. Proteome Res., 2018, 17(4), 1636-1646, DOI: 10.1021/acs.jproteome.7b00905.
- 193 J. Carrola, V. Bastos, I. Jarak, R. Oliveira-Silva, E. Malheiro, A. L. Daniel-da-Silva, H. Oliveira, C. Santos, A. M. Gil and I. F. Duarte, Metabolomics of silver nanoparticles toxicity in HaCaT cells: structure-activity relationships and role of ionic silver and oxidative stress, Nanotoxicology, 2016, 10(8), 1105-1117, DOI: 10.1080/17435390.2016.1177744.
- 194 J. Carrola, V. Bastos, A. L. Daniel-da-Silva, A. M. Gil, C. Santos, H. Oliveira and I. F. Duarte, Macrophage Metabolomics Reveals Differential Metabolic Responses to Subtoxic Levels of Silver Nanoparticles and Ionic Silver, Eur. J. Inorg. Chem., 2020, 2020(19), 1867-1876, DOI: 10.1002/ejic.202000095.
- 195 S. Kim, S. Kim, S. Lee, B. Kwon, J. Choi, J. W. Hyun and S. Kim, Characterization of the effects of silver nanoparticles on liver cell using HR-MAS NMR spectroscopy, Bull. Korean Chem. Soc., 2011, 32(6), 2021-2026, DOI: 10.5012/bkcs.2011.32.6.2021.
- 196 P. K. Babele, A. K. Singh and A. Srivastava, Bio-inspired silver nanoparticles impose metabolic and epigenetic toxicity to saccharomyces cerevisiae, Front. Pharmacol., 2019, 10, DOI: 10.3389/fphar.2019.01016.
- 197 R. Qu, Q. Xie, J. Tian, M. Zhou and F. Ge, Metabolomics reveals the inhibition on phosphorus assimilation in Chlorella vulgaris F1068 exposed to AgNPs, Sci. Total Environ., 2021, 770(20), 145362, DOI: 10.1016/ j.scitotenv.2021.145362.
- 198 M. Cao, X. Huang, F. Wang, Y. Zhang, B. Zhou, H. Chen, R. Yuan, S. Ma, H. Geng, D. Xu, C. Yan and B. Xing, Transcriptomics and metabolomics revealed biological response of Chlorella pyrenoidesato single and repeated exposures of AgNPs at different concentrations, Environ. Sci. Technol., 2021, 55(23), 15776-15787, DOI: 10.1021/acs.est.1c04059.
- 199 S. Dekkers, T. D. Williams, J. Zhang, J. Zhou, R. J. Vandebriel, L. J. J. de La Fonteyne, E. R. Gremmer, S. He, E. J. Guggenheim, I. Lynch, F. R. Cassee, W. H. de Jong and M. R. Viant, Multi-omics approaches confirm metal ions mediate the main toxicological pathways of metal-bearing nanoparticles in lung epithelial A549 cells, Environ. Sci.: Nano, 2018, 5(6), 1506-1517, DOI: 10.1039/ c8en00071a.
- 200 J. Cao, X. Qin and Z. Li, Synthesis of Silver Nanoparticles from the Polysaccharide of Farfarae Flos and Uncovering Its Anticancer Mechanism Based on the Cell Metabolomic Approach, J. Proteome Res., 2022, 21, 172-181, DOI: 10.1021/acs.jproteome.1c00668.
- 201 P. Wang, B. Zhang, H. Zhang, Y. He, C. N. Ong and J. Yang, Metabolites change of Scenedesmus obliquus exerted by AgNPs, J. Environ. Sci., 2019, 76, 310-318, DOI: 10.1016/ j.jes.2018.05.017.

- 202 W. Liu, S. Majumdar, W. Li, A. A. Keller and V. I. Slaveykova, Metabolomics for early detection of stress in freshwater alga Poterioochromonas malhamensis exposed to silver nanoparticles, Sci. Rep., 2020, 10(1), 20563, DOI: 10.1038/ s41598-020-77521-0.
- 203 J. L. Zhang, Z. P. Zhou, Y. Pei, Q. Q. Xiang, X. X. Chang, J. Ling, D. Shea and L. Q. Chen, Metabolic profiling of silver nanoparticle toxicity in Microcystis aeruginosa, Environ. Sci.: Nano, 2018, 5(11), 2519-2530, DOI: 10.1039/
- 204 G. A. Hart and T. W. Hesterberg, In vitro toxicity of respirable-size particles of diatomaceous earth and crystalline silica compared with asbestos and titanium dioxide, J. Occup. Environ. Med., 1998, 40(1), 29-42, DOI: 10.1097/00043764-199801000-00008.
- 205 B. K. Bernard, M. R. Osheroff, A. Hofmann and J. H. Mennear, Toxicology and carcinogenesis studies of dietary titanium dioxide-coated mica in male and female fischer 344 rats, J. Toxicol. Environ. Health, 1990, 29(4), 417-429, DOI: 10.1080/15287399009531402.
- 206 P. Tucci, G. Porta, M. Agostini, D. Dinsdale, I. Iavicoli, K. Cain, A. Finazzi-Agró, G. Melino and A. Willis, Metabolic effects of TIO2 nanoparticles, a common component of sunscreens and cosmetics, on human keratinocytes, Cell Death Dis., 2013, 4(3), e549, DOI: 10.1038/cddis.2013.76.
- 207 G. Oberdörster, J. Ferin, R. Gelein, S. C. Soderholm and J. Finkelstein, Role of the alveolar macrophage in lung injury: studies with ultrafine particles, Environ. Health Perspect., 1992, 97, 193-199, DOI: 10.1289/ehp.9297193.
- 208 K. Peters, R. E. Unger, C. J. Kirkpatrick, A. M. Gatti and E. Monari, Effects of nano-scaled particles on endothelial cell function in vitro: studies on viability, proliferation and inflammation, J. Mater. Sci.: Mater. Med., 2004, 15, 321-325, DOI: 10.1023/B:JMSM.0000021095.36878.1b.
- 209 G. Raja, S. Cao, D. H. Kim and T. J. Kim, Mechanoregulation of titanium dioxide nanoparticles in cancer therapy, Mater. Sci. Eng., C, 2020, 107, 110303, DOI: 10.1016/j.msec.2019.110303.
- 210 Q. Chen, N. Wang, M. Zhu, J. Lu, H. Zhong, X. Xue, S. Guo, M. Li, X. Wei, Y. Tao and H. Yin, TiO2 nanoparticles cause mitochondrial dysfunction, activate inflammatory responses, and attenuate phagocytosis in macrophages: a proteomic and metabolomic insight, Redox Biol., 2018, 15, 266-276, DOI: 10.1016/j.redox.2017.12.011.
- 211 S. Radenkovic, I. Vuckovic and I. R. Lanza, Metabolic flux analysis: moving beyond static metabolomics, Trends Biochem. Sci., 2020, 45, 545-546, DOI: 10.1016/ j.tibs.2020.02.011.
- 212 C. Jin, Y. Liu, L. Sun, T. Chen, Y. Zhang, A. Zhao, X. Wang, M. Cristau, K. Wang and W. Jia, Metabolic profiling reveals disorder of carbohydrate metabolism in mouse fibroblast cells induced by titanium dioxide nanoparticles, J. Appl. Toxicol., 2013, 33(12), 1442-1450, DOI: 10.1002/jat.2808.
- 213 Y. Bo, C. Jin, Y. Liu, W. Yu and H. Kang, Metabolomic analysis on the toxicological effects of TiO2 nanoparticles in mouse fibroblast cells: from the perspective of

perturbations in amino acid metabolism, Toxicol. Mech. Methods. 2014, 24(7), 461-469. DOI: 10.3109/ 15376516.2014.939321.

- 214 R. Garcia-Contreras, R. J. Scougall-Vilchis, R. Contreras-Bulnes, Y. Kanda, H. Nakajima and H. Sakagami, Induction of prostaglandin E2 production by TiO2 nanoparticles in human gingival fibroblast, In Vivo, 2014, 28(2), 217-222.
- 215 R. Garcia-Contreras, M. Sugimoto, N. Umemura, M. Kaneko, Y. Hatakeyama, T. Soga, M. Tomita, R. J. Scougall-Vilchis, R. Contreras-Bulnes, H. Nakajima and H. Sakagami, Alteration of metabolomic profiles by titanium dioxide nanoparticles in human gingivitis model, Biomaterials, 2015, 57, 33-40, DOI: 10.1016/ j.biomaterials.2015.03.059.
- 216 K. T. Kitchin, E. Grulke, B. L. Robinette and B. T. Castellon, Metabolomic effects in HepG2 cells exposed to four TiO₂ and two CeO₂ nanomaterials, Environ. Sci.: Nano, 2014, 1(5), 466-477, DOI: 10.1039/c4en00096j.
- 217 L. Cui, X. Wang, B. Sun, T. Xia and S. Hu, Predictive metabolomic signatures for safety assessment of metal oxide nanoparticles, ACS Nano, 2019, 13(11), 13065-13082, DOI: 10.1021/acsnano.9b05793.
- 218 S. Jia, M. I. Setyawati, M. Liu, T. Xu, J. Loo, M. Yan, J. Gong, S. H. Chotirmall, P. Demokritou, K. W. Ng and M. Fang, Association of nanoparticle exposure with serum metabolic disorders of healthy adults in printing centers, J. Hazard. Mater., 2022, 432, 128710, DOI: 10.1016/ j.jhazmat.2022.128710.
- 219 Q. Zhang, X. Chang, X. Wang, H. Zhan, Q. Gao, M. Yang, H. Liu, S. Li and Y. Sun, A metabolomic-based study on disturbance of bile acids metabolism induced by intratracheal instillation of nickel oxide nanoparticles in rats, Toxicol. Res. (Camb), 2021, 10(3), 579-591, DOI: 10.1093/toxres/tfab039.
- 220 Z. Chen, S. Han, D. Zhou, S. Zhou and G. Jia, Effects of oral exposure to titanium dioxide nanoparticles on gut microbiota and gut-associated metabolism: in vivo, Nanoscale, 2019, 11(46), 22398-22412, DOI: 10.1039/ c9nr07580a.
- 221 Z. Chen, S. Han, J. Zhang, P. Zheng, X. Liu, Y. Zhang and G. Jia, Exploring urine biomarkers of early health effects occupational exposure to titanium dioxide nanoparticles using metabolomics, Nanoscale, 2021, 13(7), 4122-4132, DOI: 10.1039/d0nr08792k.
- 222 Z. Chen, S. Han, J. Zhang, P. Zheng, X. Liu, Y. Zhang and G. Jia, Metabolomics screening of serum biomarkers for occupational exposure of titanium dioxide nanoparticles, Nanotoxicology, 2021, 15(6), 832-849, DOI: 10.1080/ 17435390.2021.1921872.
- 223 Z. Chen, S. Han, D. Zhou, P. Zheng, S. Zhou and G. Jia, Serum metabolomic signatures of Sprague-Dawley rats administration of titanium nanoparticles, NanoImpact, 2020, 19, 100236, DOI: 10.1016/j.impact.2020.100236.
- 224 S. Han, Z. J. Chen, D. Zhou, P. Zheng, J. H. Zhang and G. Jia, Effects of titanium dioxide nanoparticles on fecal

- metabolome in rats after oral administration for 90 days, Beijing Daxue Xuebao, Yixueban, 2020, 52(3), 457-463, DOI: 10.19723/j.issn.1671-167X.2020.03.010.
- 225 M. L. Whitfield Åslund, H. McShane, M. J. Simpson, A. J. Simpson, J. K. Whalen, W. H. Hendershot and G. I. Sunahara, Earthworm sublethal responses to titanium dioxide nanomaterial in soil detected by ¹H NMR metabolomics, Environ. Sci. Technol., 2012, 46(2), 1111-1118, DOI: 10.1021/es202327k.
- 226 Y. Zhu, X. Wu, Y. Liu, J. Zhang and D. Lin, Integration of transcriptomics and metabolomics reveals the responses of earthworms to the long-term exposure of TiO2 nanoparticles in soil, Sci. Total Environ., 2020, 719, 137492, DOI: 10.1016/j.scitotenv.2020.137492.
- 227 C. Ratnasekhar, M. Sonane, A. Satish and M. K. R. Mudiam, Metabolomics reveals the perturbations in the metabolome of Caenorhabditis elegans exposed to titanium dioxide nanoparticles, Nanotoxicology, 2015, 9(8), 994–1004, DOI: 10.3109/17435390.2014.993345.
- 228 J. Kurepa, T. E. Shull and J. A. Smalle, Metabolomic analyses of the bio-corona formed on TiO2 nanoparticles incubated with plant leaf tissues, J. Nanobiotechnol., 2020, 18(1), 28, DOI: 10.1186/s12951-020-00592-8.
- 229 A. J. Chetwynd, W. Zhang, J. A. Thorn, I. Lynch and R. Ramautar, The nanomaterial metabolite corona determined using a quantitative metabolomics approach: a pilot study, Small, 2020, 16(21), 2000295, DOI: 10.1002/ smll.202000295.
- 230 A. Bannuscher, B. Hellack, A. Bahl, J. Laloy, H. Herman, M. S. Stan, A. Dinischiotu, A. Giusti, B.-C. Krause, J. Tentschert, M. Roşu, C. Balta, A. Hermenean, M. Wiemann, A. Luch and A. Haase, Metabolomics profiling to investigate nanomaterial toxicity in vitro and in vivo, Nanotoxicology, 2020, 14, 807-826.
- 231 I. Karkossa, A. Bannuscher, B. Hellack, A. Bahl, S. Buhs, P. Nollau, A. Luch, K. Schubert, M. von Bergen and A. Haase, An in-depth multi-omics analysis in RLE-6TN rat alveolar epithelial cells allows for nanomaterial categorization, Part. Fibre Toxicol., 2019, 16(1), 38, DOI: 10.1186/s12989-019-0321-5.
- 232 I. Karkossa, A. Bannuscher, B. Hellack, W. Wohlleben, J. Laloy, M. S. Stan, A. Dinischiotu, M. Wiemann, A. Luch, A. Haase, M. von Bergen and K. Schubert, Nanomaterials induce different levels of oxidative stress, depending on the used model system: comparison of in vitro and in vivo effects, Sci. Total Environ., 2021, 801, 149538, DOI: 10.1016/j.scitotenv.2021.149538.
- 233 K. Pathakoti, M. Manubolu and H. min Hwang, Mechanistic insights into TiO2 and ZnO nanoparticleinduced metabolic changes in Escherichia coli under solar simulated light irradiation, Water, Air, Soil Pollut., 2020, 231(1), 16, DOI: 10.1007/s11270-019-4388-2.
- 234 M. Planchon, T. Leger, O. Spalla, G. Huber and R. Ferrari, Metabolomic and proteomic investigations of impacts of titanium dioxide nanoparticles on Escherichia coli, PLoS DOI: 10.1371/ One, 2017, 12(6), e0178437, journal.pone.0178437.

- 235 Z. Zhang, Z. C. Liang, J. H. Zhang, S. L. Tian, J. le Qu, J. N. Tang and S. de Liu, Nano-sized TiO₂ (nTiO₂) induces metabolic perturbations in Physarum polycephalum macroplasmodium to counter oxidative stress under dark conditions, Ecotoxicol. Environ. Saf., 2018, 154, 108-117, DOI: 10.1016/j.ecoenv.2018.02.012.
- 236 N. Kumar, A. Mittal, M. Yadav, S. Sharma, T. Kumar, R. Chakraborty, S. Sengupta and N. S. Chauhan, Photocatalytic TiO₂/CdS/ZnS nanocomposite induces Bacillus subtilis cell death by disrupting its metabolism and membrane integrity, Indian J. Microbiol., 2021, 61(4), 487-496, DOI: 10.1007/s12088-021-00973-z.
- 237 J. Feng, J. Li, H. Wu and Z. Chen, Metabolic responses of HeLa cells to silica nanoparticles by NMR-based metabolomic analyses, Metabolomics, 2013, 9(4), 874-886, DOI: 10.1007/s11306-013-0499-8.
- 238 S. M. Huang, X. Zuo, J. J. E. Li, S. F. Y. Li, B. H. Bay and C. N. Ong, Metabolomics studies show dose-dependent toxicity induced by SiO2 nanoparticles in MRC-5 human fetal lung fibroblasts, Adv. Healthcare Mater., 2012, 1(6), 779-784, DOI: 10.1002/adhm.201200114.
- 239 A. Irfan, M. Cauchi, W. Edmands, N. J. Gooderham, J. Njuguna and H. Zhu, Assessment of Temporal Dose-Toxicity Relationship of Fumed Silica Nanoparticle in Human Lung A549 Cells by Conventional Cytotoxicity and 1H-NMR-Based Extracellular Metabonomic Toxicol. Sci., 2014, 138, 354-364.
- 240 N. Chatterjee, J. Jeong, D. Yoon, S. Kim and J. Choi, Global metabolomics approach in in vitro and in vivo models reveals hepatic glutathione depletion induced by amorphous silica nanoparticles, Chem.-Biol. Interact., 2018, 293, 100-106, DOI: 10.1016/j.cbi.2018.07.013.
- 241 X. Zhao, A. Abulikemu, S. Lv, Y. Qi, J. Duan, J. Zhang, R. Chen, C. Guo, Y. Li and Z. Sun, Oxidative stress- and mitochondrial dysfunction-mediated cytotoxicity by silica nanoparticle in lung epithelial cells from metabolomic perspective, Chemosphere, 2021, 275, 129969, DOI: 10.1016/j.chemosphere.2021.129969.
- 242 K. T. Kitchin, S. Stirdivant, B. L. Robinette, B. T. Castellon and X. Liang, Metabolomic effects of CeO₂, SiO₂ and CuO metal oxide nanomaterials on HepG2 cells, Part. Fibre Toxicol., 2017, 14(1), 50, DOI: 10.1186/s12989-017-0230-4.
- 243 A. Bannuscher, I. Karkossa, S. Buhs, P. Nollau, K. Kettler, M. Balas, A. Dinischiotu, B. Hellack, M. Wiemann, A. Luch, M. von Bergen, A. Haase and K. Schubert, A multi-omics approach reveals mechanisms nanomaterial toxicity and structure-activity relationships in alveolar macrophages, Nanotoxicology, 2020, 14(2), 181-195, DOI: 10.1080/17435390.2019.1684592.
- 244 R. Saborano, T. Wongpinyochit, J. D. Totten, B. F. Johnston, F. P. Seib and I. F. Duarte, Metabolic reprogramming of macrophages exposed to silk, poly(lactic-co-glycolic acid), and silica nanoparticles, Adv. Healthcare Mater., 2017, 6(14), 1601240, DOI: 10.1002/adhm.201601240.
- 245 Y. Zhu, Y. Zhang, Y. Li, C. Guo, Z. Fan, Y. Li, M. Yang, X. Zhou, Z. Sun and J. Wang, Integrative proteomics and metabolomics approach to elucidate metabolic

- dysfunction induced by silica nanoparticles hepatocytes, J. Hazard. Mater., 2022, 434, 128820.
- 246 A. B. Djurišić, X. Chen, Y. H. Leung and A. Man Ching Ng, ZnO nanostructures: Growth, properties and applications, J. Mater. Chem., 2012, 22(14), 6526-6535, DOI: 10.1039/ c2jm15548f.
- 247 S. L. Lim, C. T. Ng, L. Zou, Y. Lu, J. Chen, B. H. Bay, H. M. Shen and C. N. Ong, Targeted metabolomics reveals differential biological effects of nanoplastics and nanoZnO in human lung cells, Nanotoxicology, 2019, 13(8), 1117-1132, DOI: 10.1080/17435390.2019.1640913.
- 248 A. A. Keller, W. Vosti, H. Wang and A. Lazareva, Release of engineered nanomaterials from personal care products throughout their life cycle, J. Nanopart. Res., 2014, 16(7), 2489, DOI: 10.1007/s11051-014-2489-9.
- 249 J. Heim, E. Felder, M. N. Tahir, A. Kaltbeitzel, U. R. Heinrich, C. Brochhausen, V. Mailänder, W. Tremel and J. Brieger, Genotoxic effects of zinc oxide nanoparticles, Nanoscale, 2015, 7(19), 8931-8938, DOI: 10.1039/c5nr01167a.
- 250 H. Salehi, N. de Diego, A. Chehregani Rad, J. J. Benjamin, M. Trevisan and L. Lucini, Exogenous application of ZnO nanoparticles and ZnSO4 distinctly influence the metabolic response in Phaseolus vulgaris L., Sci. Total Environ., 2021, 778, 146331, DOI: j.scitotenv.2021.146331.
- 251 S. Li, J. Liu, Y. Wang, Y. Gao, Z. Zhang, J. Xu and G. Xing, Comparative physiological and metabolomic analyses revealed that foliar spraying with zinc oxide and silica nanoparticles modulates metabolite profiles in cucumber (Cucumis sativus L.), Food Energy Secur., 2021, 10(1), e269, DOI: 10.1002/fes3.269.
- 252 L. Sun, Y. Wang, R. Wang, R. Wang, P. Zhang, Q. Ju and J. Xu, Physiological, transcriptomic, and metabolomic analyses reveal zinc oxide nanoparticles modulate plant growth in tomato, Environ. Sci.: Nano, 2020, 7(11), 3587-3604, DOI: 10.1039/d0en00723d.
- 253 J. Wan, R. Wang, H. Bai, Y. Wang and J. Xu, Comparative physiological and metabolomics analysis reveals that single-walled carbon nanohorns and ZnO nanoparticles affect salt tolerance in: Sophora alopecuroides, Environ. Sci.: Nano, 2020, 7(10), 2968-2981, DOI: 10.1039/ d0en00582g.
- 254 E. He, R. Qiu, X. Cao, L. Song, W. J. G. M. Peijnenburg and H. Qiu, Elucidating toxicodynamic differences at the molecular scale between ZnO nanoparticles and ZnCl₂ in Enchytraeus crypticus via nontargeted metabolomics, Environ. Sci. Technol., 2020, 45(6), 3487-3498, DOI: 10.1021/acs.est.0c00663.
- 255 L. A. Paramo, A. A. Feregrino-Pérez, R. Guevara, S. Mendoza and K. Esquivel, Nanoparticles in agroindustry: applications, challenges, and trends, toxicity, Nanomaterials, 2020, 10, 1654, DOI: 10.3390/ nano10091654.
- 256 S. Majumdar and A. A. Keller, Omics to address the opportunities and challenges of nanotechnology in

agriculture, Crit. Rev. Environ. Sci. Technol., 2021, 51(22), 2595-2636, DOI: 10.1080/10643389.2020.1785264.

- 257 B. Wu, J. Wu, S. Liu, Z. Shen, L. Chen, X. X. Zhang and H. q. Ren, Combined effects of graphene oxide and zinc oxide nanoparticle on human A549 cells: bioavailability, toxicity and mechanisms, Environ. Sci.: Nano, 2019, 6(2), 635-645, DOI: 10.1039/C8EN00965A.
- 258 P. Kumar Babele, Zinc oxide nanoparticles impose metabolic toxicity by de-regulating proteome and metabolome in Saccharomyces cerevisiae, Toxicol. Rep., 2019, 6, 64-73, DOI: 10.1016/j.toxrep.2018.12.001.
- 259 S. Amiri and H. Shokrollahi, The role of cobalt ferrite magnetic nanoparticles in medical science, Mater. Sci. Eng., C, 2013, 33, 1-8, DOI: 10.1016/j.msec.2012.09.003.
- 260 A. B. B. Oliveira, F. R. de Moraes, N. M. Candido, I. Sampaio, A. S. Paula, A. de Vasconcellos, T. C. Silva, A. H. Miller, P. Rahal, J. G. Nery and M. F. Calmon, Metabolic effects of cobalt ferrite nanoparticles on cervical carcinoma cells and nontumorigenic keratinocytes, J. Proteome Res., 2016, 15(12), 4337-4348, DOI: 10.1021/acs.jproteome.6b00411.
- 261 V. Aruoja, H. C. Dubourguier, K. Kasemets and A. Kahru, Toxicity of nanoparticles of CuO, ZnO and TiO2 to microalgae Pseudokirchneriella subcapitata, Sci. Total Environ., 2009, 407(4), 1461-1468, j.scitotenv.2008.10.053.
- 262 A. Murgia, L. Mancuso, C. Manis, P. Caboni and G. Cao, GC-MS metabolomics analysis of mesenchymal stem cells treated with copper oxide nanoparticles, Toxicol. Mech. Methods, 2016, 26(8), 611-619, DOI: 10.1080/ 15376516.2016.1220654.
- 263 L. Wang, X. Huang, W. Sun, H. Z. Too, A. K. C. Laserna and S. F. Y. Li, A global metabolomic insight into the oxidative stress and membrane damage of copper oxide nanoparticles and microparticles on microalga Chlorella vulgaris, Environ. Pollut., 2020, 258, 113647, DOI: 10.1016/ j.envpol.2019.113647.
- 264 D. Kruszka, R. K. Selvakesavan, P. Kachlicki and G. Franklin, Untargeted metabolomics analysis reveals the elicitation of important secondary metabolites upon treatment with various metal and metal oxide nanoparticles in Hypericum perforatum L. suspension cultures, Ind. Crops Prod., 2022, 178, 114561, DOI: 10.1016/j.indcrop.2022.114561.
- 265 M. S. P. Boyles, C. Ranninger, R. Reischl, M. Rurik, R. Tessadri, O. Kohlbacher, A. Duschl and C. G. Huber, Copper oxide nanoparticle toxicity profiling using untargeted metabolomics, Part. Fibre Toxicol., 2016, 13(1), 49, DOI: 10.1186/s12989-016-0160-6.
- 266 N. G. Chavez Soria, D. S. Aga and G. E. Atilla-Gokcumen, Lipidomics reveals insights on the biological effects of copper oxide nanoparticles in a human colon carcinoma cell line, Mol. Omics, 2019, 15(1), 30-38, DOI: 10.1039/ c8mo00162f.
- 267 B. C. Krause, F. L. Kriegel, V. Tartz, H. Jungnickel, P. Reichardt, A. V. Singh, P. Laux, M. Shemis and Luch, Combinatory effects of cerium dioxide

- nanoparticles and acetaminophen on the liver-a case study of low-dose interactions in human HuH-7 cells, Int. J. Mol. Sci., 2021, 22(13), 6866, DOI: 10.3390/ijms22136866.
- 268 N. S. Taylor, R. Merrifield, T. D. Williams, J. K. Chipman, J. R. Lead and M. R. Viant, Molecular toxicity of cerium oxide nanoparticles to the freshwater Chlamydomonas reinhardtii is associated with supraenvironmental exposure concentrations, Nanotoxicology, 2016, **10**(1), 32-41, DOI: 10.3109/17435390.2014.1002868.
- 269 T. H. Shin, A. A. Ketebo, D. Y. Lee, S. Lee, S. H. Kang, S. Basith, B. Manavalan, D. H. Kwon, S. Park and G. Lee, Decrease in membrane fluidity and traction force induced silica-coated magnetic nanoparticles, Nanobiotechnol., 2021, 19(1), 21, DOI: 10.1186/s12951-020-00765-5.
- 270 T. H. Shin, B. Manavalan, D. Y. Lee, S. Basith, C. Seo, M. J. Paik, S.-W. Kim, H. Seo, J. Y. Lee, J. Y. Kim, A. Y. Kim, J. M. Chung, E. J. Baik, S. H. Kang, D.-K. Choi, Y. Kang, M. M. Mouradian and G. Lee, Silica-coated magnetic-nanoparticle-induced cytotoxicity is reduced in microglia by glutathione and citrate identified using integrated omics, Part. Fibre Toxicol., 2021, 18, 42.
- 271 X. Li, C. Zhang, X. Zhang, S. Wang, Q. Meng, S. Wu, H. Yang, Y. Xia and R. Chen, An acetyl-L-carnitine switch on mitochondrial dysfunction and rescue in the metabolomics study on aluminum oxide nanoparticles, Part. Fibre Toxicol., 2016, 13(1), 4, DOI: 10.1186/s12989-016-0115-y.
- 272 X. Cai, J. Dong, J. Liu, H. Zheng, C. Kaweeteerawat, F. Wang, Z. Ji and R. Li, Multi-hierarchical profiling the structureactivity relationships of engineered nanomaterials at nano-bio interfaces, Nat. Commun., 2018, 9(1), 4416, DOI: 10.1038/s41467-018-06869-9.
- 273 C. Guo, R. J. M. Weber, A. Buckley, J. Mazzolini, S. Robertson, J. M. Delgado-Saborit, J. Z. Rappoport, J. Warren, A. Hodgson, P. Sanderson, J. K. Chipman, M. R. Viant and R. Smith, Environmentally relevant iron oxide nanoparticles produce limited acute pulmonary effects in rats at realistic exposure levels, Int. J. Mol. Sci., 2021, 22(2), 556, DOI: 10.3390/ijms22020556.
- 274 M. Ye, L. Wang, Z. Wu and W. Liu, Metabolomic profiling of ZrO 2 nanoparticles in MC3T3-E1 cells, IET Nanobiotechnol., 2021, 15, 687-697.
- 275 Q. Yang, L. Zhang, A. Ben, N. Wu, Y. Yi, L. Jiang, H. Huang and Y. Yu, Effects of dispersible MoS2 nanosheets and Nano-silver coexistence on the metabolome of yeast, Chemosphere, 2018, **198**, 216-225, DOI: 10.1016/ j.chemosphere.2018.01.140.
- 276 A. Wang, Q. Jin, X. Xu, A. Miao, J. C. White, J. L. Gardea-Torresdey, R. Ji and L. Zhao, High-throughput screening for engineered nanoparticles that enhance photosynthesis using mesophyll protoplasts, J. Agric. Food Chem., 2020, 68(11), 3382-3389, DOI: 10.1021/acs.jafc.9b06429.
- 277 N. Wu, Y. Yu, T. Li, X. Ji, L. Jiang, J. Zong and H. Huang, Investigating the influence of MoS2 nanosheets on E. coli from metabolomics level, PLoS One, 2016, 11(12), e0167245, DOI: 10.1371/journal.pone.0167245.

Nanoscale Advances

- 278 Y. Yu, N. Wu, Y. Yi, Y. Li, L. Zhang, Q. Yang, W. Miao, X. Ding, L. Jiang and H. Huang, Dispersible MoS₂ nanosheets activated TGF-β/Smad pathway and perturbed the metabolome of human dermal fibroblasts, ACS Biomater. Sci. Eng., 2017, 3(12), 3261-3272, DOI: 10.1021/ acsbiomaterials.7b00575.
- 279 P. Yuan, X. Hu and Q. Zhou, The nanomaterial-induced bystander effects reprogrammed macrophage immune function and metabolic profile, Nanotoxicology, 2020, 14(8), 1137-1155, DOI: 10.1080/17435390.2020.1817598.
- 280 T. G. Chatzimitakos and C. D. Stalikas, Qualitative Alterations of Bacterial Metabolome after Exposure to Metal Nanoparticles with Bactericidal Properties: A Comprehensive Workflow Based on 1H NMR, UHPLC-HRMS, and Metabolic Databases, I. Proteome Res., 2016, 15(9), 3322-3330, DOI: 10.1021/acs.jproteome.6b00489.
- 281 P. Nazari, R. Dowlatabadi-Bazaz, M. R. Mofid, M. R. Pourmand, N. E. Daryani, M. A. Faramarzi, Z. Sepehrizadeh and A. R. Shahverdi, The antimicrobial effects and metabolomic footprinting of carboxyl-capped bismuth nanoparticles against helicobacter pylori, Appl. Biochem. Biotechnol., 2014, 172(2), 570-579, DOI: 10.1007/ s12010-013-0571-x.
- 282 X. Li, Y. Qin, L. Kong, X. Yan, W. Zhang, C. J. Martyniuk, X. Wang and B. Yan, Metabolomic and bioenergetic responses of human hepatocellular carcinoma cells following exposure to commercial copper hydroxide nanopesticide, Environ. Sci.: Nano, 2022, 9, 589-605.
- 283 S. R. Choudhury, A. Mandal, M. Ghosh, S. Basu, D. Chakravorty and A. Goswami, Investigation of orthorhombic antimicrobial physiology of monoclinic nanoallotropes of sulfur at the interface of transcriptome and metabolome, Biotechnol., 2013, 97(13), 5965-5978, DOI: 10.1007/s00253-013-4789-x.
- 284 D. Zhang, X. Li, X. Xie, W. Zheng, A. Li, Y. Liu, X. Liu, R. Zhang, C. Deng, J. Cheng, H. Yang and M. Gong, Exploring the Biological Effect of Biosynthesized Au-Pd Core-Shell Nanoparticles through an Untargeted Metabolomics Approach, ACS Appl. Mater. Interfaces, 2021, 13, 59633-59648.
- 285 D. Zhang, W. Zheng, X. Li, A. Li, N. Ye, L. Zhang, Y. Liu, X. Liu, R. Zhang, M. Wang, J. Cheng, H. Yang and M. Gong, Investigating the effect of Ti₃C₂ (MXene) nanosheet on human umbilical vein endothelial cells via a combined untargeted and targeted metabolomics approach, Carbon, 2021, 178, 810-821, DOI: 10.1016/ j.carbon.2021.04.023.
- 286 G. Jiao, X. Li, N. Zhang, J. Qiu, H. Xu and S. Liu, Metabolomics study on the cytotoxicity of graphene, RSC Adv., 2014, 4(84), 44712-44717, DOI: 10.1039/c4ra06312k.
- 287 S. X. F. Adamson, R. Wang, W. Wu, B. Cooper and J. Shannahan, Metabolomic insights of macrophage responses to graphene nanoplatelets: Role of scavenger receptor CD36, PLoS One, 2018, 13(11), e0207042, DOI: 10.1371/journal.pone.0207042.

- 288 G. Raja, V. Selvaraj, M. Suk, K. T. Suk and T. J. Kim, Metabolic phenotyping analysis of graphene oxide exposures in nanosheets breast cancer metabolomics profiling techniques, Process Biochem., 2021, 104, 39-45, DOI: 10.1016/j.procbio.2021.02.016.
- 289 L. Hou, S. Guan, Y. Jin, W. Sun, Q. Wang, Y. Du and R. Zhang, Cell metabolomics to study the cytotoxicity of carbon black nanoparticles on A549 cells using UHPLC-Q/ TOF-MS and multivariate data analysis, Sci. Total 698, 134122, DOI: 10.1016/ Environ., 2020, j.scitotenv.2019.134122.
- 290 K. Xu, X. Wang, C. Lu, Y. Liu, D. Zhang and J. Cheng, Toxicity of three carbon-based nanomaterials earthworms: Effect of morphology on biomarkers, cytotoxicity, and metabolomics, Sci. Total Environ., 2021, 777, 146224, DOI: 10.1016/j.scitotenv.2021.146224.
- 291 J. Frontiñán-Rubio, M. Victoria Gómez, C. Martín, M. González-Domínguez, M. Durán-Prado and E. Vázquez, Differential effects of graphene materials on the metabolism and function of human skin cells, Nanoscale, 2018, 10(24), 11604-11615, DOI: 10.1039/ c8nr00897c.
- 292 D. Zhang, L. Zhang, W. Zheng, F. Wu, J. Cheng, H. Yang and Investigating biological M. Gong, effects multidimensional carboxylated carbon-based nanomaterials on human lung A549 cells revealed via non-targeted metabolomics approach, Nanotechnology, 2021, 32(1), 015704, DOI: 10.1088/1361-6528/abb55b.
- 293 M. Cicuéndez, J. Flores, H. Oliveira, M. T. Portolés, M. Vallet-Regí, M. Vila and I. F. Duarte, Metabolomic response of osteosarcoma cells to nanographene oxidemediated hyperthermia, Mater. Sci. Eng., C, 2018, 91, 340-348, DOI: 10.1016/j.msec.2018.05.057.
- 294 X. Hu, S. Ouyang, L. Mu, J. An and Q. Zhou, Effects of graphene oxide and oxidized carbon nanotubes on the cellular division, microstructure, uptake, oxidative stress, and metabolic profiles, Environ. Sci. Technol., 2015, 49(18), 10825-10833, DOI: 10.1021/acs.est.5b02102.
- 295 N. Chatterjee, J. Yang, D. Yoon, S. Kim, S. W. Joo and J. Choi, Differential crosstalk between global DNA methylation and metabolomics associated with cell type specific stress response by pristine and functionalized MWCNT, Biomaterials, 2017, 115, 167-180, DOI: 10.1016/ j.biomaterials.2016.11.005.
- 296 N. Chatterjee, J. Yang, S. Kim, S. W. Joo and J. Choi, Diameter size and aspect ratio as critical determinants of uptake, stress response, global metabolomics and epigenetic alterations in multi-wall carbon nanotubes, DOI: Carbon, 2016, 108, 529-540, 10.1016/ j.carbon.2016.07.031.
- 297 S. Phuyal, M. Kasem, O. Knittelfelder, A. Sharma, d. M. Fonseca, V. Vebraite, S. Shaposhnikov, Slupphaug, V. Skaug and Characterization of the proteome and lipidome profiles of human lung cells after low dose and chronic exposure to multiwalled carbon nanotubes, Nanotoxicology, 2018, 12(2), 138-152, DOI: 10.1080/17435390.2018.1425500.

298 C. Du, B. Zhang, Y. He, C. Hu, Q. X. Ng, H. Zhang, C. N. Ong and ZhifenLin, Biological effect of aqueous C60 aggregates on Scenedesmus obliquus revealed by transcriptomics and non-targeted metabolomics, J. Hazard. Mater., 2017, 324, 221-229, DOI: 10.1016/j.jhazmat.2016.10.052.

- 299 M. Pink, N. Verma, A. W. Rettenmeier and S. Schmitz-Spanke, Integrated proteomic and metabolomic analysis to assess the effects of pure and benzo[a]pyrene-loaded carbon black particles on energy metabolism and motility in the human endothelial cell line EA.hy926, Arch. Toxicol., 2014, 88(4), 913-934, DOI: 10.1007/s00204-014-1200-3.
- 300 H. Zanin, L. M. Hollanda, H. J. Ceragioli, M. S. Ferreira, Machado, M. Lancellotti, R. R. Catharino, V. Baranauskas and A. O. Lobo, Carbon nanoparticles for gene transfection in eukaryotic cell lines, Mater. Sci. Eng., C, 2014, 39(1), 359-370, DOI: 10.1016/j.msec,2014.03.016.
- 301 I. Varol, O. Kaplan, N. Erdoğar, S. Öncül, T. T. Nielsen, A. Ercan, E. Bilensoy and M. Çelebier, Q-TOF LC/MSbased untargeted metabolomics approach to evaluate the effect of folate-conjugated cyclodextrins on triple-negative breast cancer cells, Curr. Pharm. Anal., 2021, 17(10), 1272-1281, DOI: 10.2174/1573412917999201020205745.
- 302 Y. Yang, H. Liang, P. Zeng, W. Fu, J. Yu, L. Chen, D. Chai, Wen, A. Chen and Y. Zhang, characterization, and anti-hepatocellular carcinoma effect of glycyrrhizin-coupled bovine serum albumin-loaded luteolin nanoparticles, *Pharmacogn. Mag.*, 2022, **18**(77), 216-225, DOI: 10.4103/pm.pm_34_21.
- 303 S. J. Kim, N. P. Long, C. W. Jung, N. H. Anh, J. E. Min, H. M. Kim and S. W. Kwon, Exposure to nano-polystyrene induces metabolic alteration in lipid homeostasis in Caco-2, Environ. Sci.: Nano, 2021, 8(5), 1408-1424, DOI: 10.1039/d1en00145k.
- 304 X. Su, Y. Huang, R. Chen, Y. Zhang, M. He and X. Lü, Metabolomics analysis of poly(l-lactic acid) nanofibers' performance on PC12 cell differentiation, Regener. Biomater., 2021, 8(4), rbab031, DOI: 10.1093/rb/rbab031.
- 305 L. J. Feng, J. W. Li, E. G. Xu, X. D. Sun, F. P. Zhu, Z. Ding, H. Tian, S. S. Dong, P. F. Xia and X. Z. Yuan, Short-term exposure to positively charged polystyrene nanoparticles causes oxidative stress and membrane destruction in cyanobacteria, Environ. Sci.: Nano, 2019, 6(10), 3072-3079, DOI: 10.1039/c9en00807a.
- 306 M. A. Al-Natour, A. Alazzo, A. M. Ghaemmaghami, D. H. Kim and C. Alexander, LC-MS metabolomics comparisons of cancer cell and macrophage responses to methotrexate and polymer-encapsulated methotrexate, Int. J. Pharm.: X, 2019, 1, 100036, DOI: 10.1016/ j.ijpx.2019.100036.

- 307 M. Yaşacan, A. Erikçi, C. C. Eylem, S. Y. Çiftçi, E. Nemutlu, K. Ulubayram and İ. Eroğlu, Polymeric nanoparticle versus liposome formulations: comparative physicochemical and metabolomic studies as l-carnitine delivery systems, AAPS PharmSciTech, 2020, 21(8), 308, DOI: 10.1208/s12249-020-01852-4.
- 308 D. Magrì, M. Veronesi, P. Sánchez-Moreno, V. Tolardo, T. Bandiera, P. P. Pompa, A. Athanassiou and D. Fragouli, PET nanoplastics interactions with water contaminants and their impact on human cells, Environ. Pollut., 2021, 271, 116262, DOI: 10.1016/j.envpol.2020.116262.
- 309 A. L. Al-Malki, A. Bakkar, E. A. Huwait, E. K. Barbour, K. O. Abulnaja, T. A. Kumosani and S. S. Moselhy, Strigol1/albumin/chitosan nanoparticles decrease cell viability, induce apoptosis and alter metabolomics profile in HepG2 cancer cell line, Biomed. Pharmacother., 2021, 142, 111960, DOI: 10.1016/j.biopha.2021.111960.
- 310 F. de Castro, V. Vergaro, M. Benedetti, F. Baldassarre, L. del Coco, M. M. Dell'anna, P. Mastrorilli, F. P. Fanizzi and Ciccarella, Visible Light-Activated Water-Soluble Platicur Nanocolloids: Photocytotoxicity and Metabolomics Studies in Cancer Cells, ACS Appl. Bio 6836-6851, Mater., 2020, 3(10), DOI: 10.1021/ acsabm.0c00766.
- 311 A. Ercan, M. Çelebier, G. Varan, S. Öncül, M. Nenni, O. Kaplan and E. Bilensoy, Global omics strategies to investigate the effect of cyclodextrin nanoparticles on MCF-7 breast cancer cells, Eur. J. Pharm. Sci., 2018, 123, 377-386, DOI: 10.1016/j.ejps.2018.07.060.
- 312 J. Yu, J. Chen, H. Zhao, J. Gao, Y. Li, Y. Li, J. Xue, A. Dahan, D. Sun, G. Zhang and H. Zhang, Integrative proteomics and metabolomics analysis reveals the toxicity of cationic liposomes to human normal hepatocyte cell line L02, 2018, **14**(5), 362-372, DOI: **10.1039**/ Mol. Omics, c8mo00132d.
- 313 T. Elmi, M. Shafiee Ardestani, F. Hajialiani, M. Motevalian, M. Mohamadi, S. Sadeghi, Z. Zamani and F. Tabatabaie, Novel chloroquine loaded curcumin based anionic linear globular dendrimer G2: A metabolomics study on Plasmodium falciparum in vitro using 1H NMR spectroscopy, Parasitology, 2020, 147(7), 747-759, DOI: 10.1017/S0031182020000372.
- 314 A. Catala, M. Dzieciatkowska, G. Wang, A. Gutierrez-Hartmann, D. Simberg, K. C. Hansen, A. D'Alessandro and C. E. Catalano, Targeted intracellular delivery of Trastuzumab using designer phage lambda nanoparticles alters cellular programs in human breast cancer cells, ACS Nano, 2021, 15(7), 11789-11805, DOI: 10.1021/ acsnano.1c02864.