

# ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## COMMUNICATION

## Controllable aggregation-induced emission based on tetraphenylethylene-functionalized pillar[5]arene via host-guest recognition

Jie Wu,<sup>a</sup> Shu Sun,<sup>a, b</sup> Xiaoqing Feng,<sup>a</sup> Jianbing Shi,<sup>b</sup> Xiao-Yu Hu<sup>\*a</sup> and Leyong Wang<sup>\*a</sup><sup>5</sup> Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX


DOI: 10.1039/b000000x

A novel TPE-functionalized pillar[5]arene (TPEP5) was successfully synthesized, and the motion of the TPE motif was restricted via pillararene-based host-guest recognition-mediated cross-linking, resulting in the efficient “turn-on” of fluorescence emission based on AIE mechanism.

Molecules with aggregation-induced emission (AIE)<sup>1</sup> or aggregation-induced enhanced emission (AIEE)<sup>2</sup> characteristics have provided a promising platform for design and creation of efficient light emitters ranging from optical materials to sensors, owing to the enhanced emission in their aggregate or solid-state forms.<sup>3</sup> Since the first AIE molecule reported by Tang *et al* in 2001,<sup>1a</sup> a wide variety of AIE molecules have been synthesized based on the mechanism of restriction of intramolecular motions (RIM).<sup>1b, 3b</sup> Among various AIE molecules, tetraphenylethylene (TPE) and its functionalized derivatives can be readily obtained via facile synthetic transformations, which are non-emissive in molecularly dissolved state, but enhanced fluorescence emission could be achieved in both the aggregated form and the solid state.<sup>4</sup> In contrast to aggregation-caused quenching (ACQ) effect of conventional organic luminophores, TPE-based AIE-active materials are demonstrated to have improved efficiency and sensitivity as chemosensors, bio-probes, and solid-state emitters and have already shown practical applications in these fields.<sup>5</sup> For example, Tang and co-workers synthesized a peptide-conjugated TPE derivative, which could be used as live-cell-permeable, fluorescent light up probe for real-time cell apoptosis imaging.<sup>6</sup> In addition, Zhang *et al.* demonstrated that the TPE derivatives containing adenine or thymine moieties could be used as “turn on” chemosensors for selective detection of Ag<sup>+</sup> and Hg<sup>2+</sup> ions.<sup>5c</sup>

Recently, non-covalent interactions such as host-guest recognition have been proved to be an efficient strategy to restrict the intramolecular motions of TPE molecule, concomitantly accompanied with the turn-on of fluorescence emission via the AIE mechanism.<sup>7</sup> For example, Liu and co-workers integrated the concept of AIE with the specific host-guest supramolecular recognition between K<sup>+</sup> ions and crown ether moieties to develop effective fluorometric K<sup>+</sup> probes.<sup>7a</sup> Considering the unique structure and interesting host-guest chemistry of pillararene, which can form supramolecular inclusion complex with various kinds of linear guests,<sup>8</sup> the grafting of pillararenes onto the periphery of TPE can provide a novel strategy for fabricating various functional AIE luminogens and achieving the fluorescent

detection of various types of guest compounds mediated by the pillararene-based host-guest interactions.<sup>9</sup> Herein, we designed and for the first time successfully synthesized TPE-functionalized pillar[5]arene (TPEP5) by attaching four DMPillar[5]arene (DMP5) groups onto the periphery of TPE (Scheme 1). It was found that TPEP5 dissolved in CHCl<sub>3</sub>/acetone solution with negligible fluorescence emission, whereas, upon addition of the guest molecule (G1), TPEP5 could be effectively induced to aggregate due to the pillararene-based host-guest recognition-mediated cross-linking via the formation of TPEP5⊃G1 (1:2 molar ratio) inclusion complex, which concomitantly resulting in the “turn-on” of fluorescence emission based on the AIE mechanism. Moreover, the fluorescence “turn-off” was found upon the gradual addition of adiponitrile (G3, a competitive guest), which was easily visualized by naked eye. Thus, this novel supramolecular system based on TPEP5⊃G1 complex creates unique possibilities to fabricate novel types of pillararene-based fluorescent probes.




**Scheme 1** Schematic illustration of the construction of the luminescent supramolecular aggregates based on aggregation-induced emission (AIE) of pillar[5]arene-functionalized tetraphenylethylene (TPEP5), induced by host-guest recognition-mediated cross-linking between G1 and pillar[5]arene moieties, and its application in the detection of adiponitrile.

The TPE-functionalized pillar[5]arene (TPEP5) was prepared by attaching four DMPillar[5]arene (DMP5) groups onto the periphery of TPE through the alkyne-azide Click reaction (Scheme S1, ESI†).<sup>10</sup> To the best of our knowledge, this is the first example of successful synthesis of TPE-functionalized pillar[5]arene (TPEP5), which could be used to fabricate functional luminescent supramolecular aggregates induced by host-guest recognition-mediated cross-linking between G1 and pillar[5]arene moieties based on the AIE mechanism.

The complexation between TPEP5 and G1 was initially

investigated by  $^1\text{H}$  NMR spectroscopy as shown in Fig. 1. The proton NMR spectra of **TPEP5**, **G1**, and a mixture of **TPEP5** and 4 equiv. of **G1** showed that this complexation system is a fast-exchanging process on the proton NMR time scale. As can be seen from Fig. 1b, after complexation the peaks of phenyl protons  $\text{H}_4$ ,  $\text{H}_5$ ,  $\text{H}_6$ , methylene and methoxyl protons  $\text{H}_{10}$ ,  $\text{H}_{11}$ ,  $\text{H}_{12}$  from pillar[5]arene, and triazole protons  $\text{H}_1$  on **TPEP5** shifted downfield slightly. The proton signals derived from  $\text{H}_d$ ,  $\text{H}_e$ ,  $\text{H}_f$ ,  $\text{H}_g$ , and  $\text{H}_a$  of **G1** shifted upfield remarkably due to the shielding effect of the electron-rich cavities of pillar[5]arene on **TPEP5**. While, no obvious change was observed for the protons  $\text{H}_b$  and  $\text{H}_c$  on **G1**. The above results revealed that the pillar[5]arene motifs on **TPEP5** were fully threaded by the guest **G1** with the protons  $\text{H}_d$ ,  $\text{H}_e$ ,  $\text{H}_f$ ,  $\text{H}_g$  and  $\text{H}_a$  in the pillar[5]arene cavities and other protons  $\text{H}_b$  and  $\text{H}_c$  out of the cavities. In addition, the  $^1\text{H}$  NMR spectrum of a mixture of model compound DMPillar[5]arene (**DMP5**) and 0.5 equiv. of **G1** was also investigated and similar complexation-induced chemical shift changes were observed (Fig. S21). Moreover, 2D NOESY experiment was also performed to study the host-guest complexation between **DMP5** and **G1** (Fig. S23). NOE correlation signals were observed between protons  $\text{H}_1$  on **DMP5** and  $\text{H}_d$ ,  $\text{H}_e$ ,  $\text{H}_f$ ,  $\text{H}_g$  on **G1**, as well as protons  $\text{H}_2/\text{H}_3$  on **DMP5** and  $\text{H}_d$ ,  $\text{H}_e$ ,  $\text{H}_f$ ,  $\text{H}_g$  on **G1**, which also confirmed the above threading binding mode.




**Fig. 1**  $^1\text{H}$  NMR spectra ( $\text{CDCl}_3/\text{acetone-}d_6$  (1:8, v/v), 300 MHz, 298 K) of (a) 5 mM **G1**; (b) 5 mM **TPEP5** and 20 mM **G1**; (c) 5 mM **TPEP5**.

Further investigation of the complex stoichiometry between **TPEP5** and **G1** was carried out by Job's plot method using model compound **DMP5**, and the result indicated a 2:1 stoichiometry complex was formed between **DMP5** and **G1** (Fig. S22), which further confirmed our envision that the guest molecule **G1** could served as a cross-linker to bind with two molecules of **DMP5** motifs (Fig. S31), leading to the aggregation of **TPEP5** and form supramolecular network (for details, see ESI $^\dagger$ , Fig S34).<sup>8d</sup> In order to investigate the binding affinity of pillar[5]arene-**G1** recognition motif, model compounds **DMP5** and **G2** (1-(4-bromobutyl)-4-((4-methoxyphenoxy)methyl)-1H-1,2,3-triazole, analogues of **G1**), were applied for the  $^1\text{H}$  NMR titration experiments, where the association constant ( $K_a$ ) of the formation of 1:1 **DMP5** $\rightarrow$ **G2** complex was calculated to be  $(7.30 \pm 0.49) \times 10^2 \text{ M}^{-1}$  ( $\text{CDCl}_3/\text{acetone-}d_6$ , Fig. S26-S28).

Considering the AIE feature of TPE core in **TPEP5** and based on the above established novel **DMP5** $\rightarrow$ **G1** (2:1 molar ratio) supramolecular inclusion complex, we envisage that the addition


of **G1** will induce the aggregation of **TPEP5**, which concomitantly resulting in the "turn-on" of fluorescence emission based on the AIE mechanism. Thus the fluorescence properties of such host-guest recognition-induced aggregation of **TPEP5** were further investigated. In preliminary experiments, we found that the choice of solvents also played important roles in observing the host-guest recognition-induced AIE. Lots of efforts were therefore made in the initial stage, searching for an appropriate solvent system, and finally,  $\text{CHCl}_3/\text{acetone}$  (1/8, v/v) was selected as the best solvent system for such supramolecular aggregation (for details, see Fig. S25). It was found that **TPEP5** dissolved in  $\text{CHCl}_3/\text{acetone}$  (1/8, v/v) showed negligible fluorescence emission due to the efficient nonradiative annihilation caused by the intramolecular rotation of the phenyl rings in TPE core of **TPEP5**<sup>1a</sup> (Fig 2, Inset A). However, when **G1** was added into the above **TPEP5** solution, the fluorescence emission increased gradually due to the formation of supramolecular network and the rotation of phenyl rings in the TPE core of **TPEP5** is restricted. As shown in Fig. 2, a dramatic emission enhancement was observed when 16.0 equiv. of **G1** was added, and this fluorescence enhancement can be easily distinguished by naked eye when illuminating the solution with UV light (365 nm) as indicated in the inset of Fig. 2, which further supporting the proposed AIE mechanism. Moreover, the quantum yield of **TPEP5** with 8.0 equiv. **G1** was determined to be 12.3%, measured by using quinine sulfate in 0.1 M  $\text{H}_2\text{SO}_4$  (quantum yield = 54.6%) as the standard (Fig. S32).



**Fig. 2** Fluorescence spectral changes of **TPEP5** (0.04 mM) upon gradual addition of **G1** (0 – 0.64 mM) in  $\text{CHCl}_3/\text{acetone}$  (1/8, v/v) ( $\lambda_{\text{ex}} = 330 \text{ nm}$ ). The inset shows the photographs of the solution of **TPEP5** in the (A) absence and (B) presence of **G1** (0.64 mM) under UV light (365 nm) illumination at 298 K.


According to the previous report, dinitrile compounds show very strong binding affinities with pillar[5]arene based on the cooperative multiple hydrogen bond and dipole-dipole interactions.<sup>8f</sup> Hence, we envisioned that the complexation between **TPEP5** and **G1** could also be destroyed after the addition of size-fit dinitrile, such as adiponitrile, resulting in the fluorescence "turn-off" of the above supramolecular **TPEP5** $\rightarrow$ **G1** system. To investigate the fluorescence sensing effect of the above **TPEP5** $\rightarrow$ **G1** system for adiponitrile, fluorescence titration experiments were performed by adding different concentrations of adiponitrile (**G3**) to the **TPEP5** $\rightarrow$ **G1** system in  $\text{CHCl}_3/\text{acetone}$  (1/8, v/v). As shown in Fig. 3, significant quenching of the fluorescence intensity was observed upon the gradual addition of adiponitrile, which could also be easily visualized by naked eye when illuminating the solution with UV light (365 nm). For the quenching of the fluorescence, a possible reason is that after addition of the complete guest **G3**, a more stable inclusion complex **TPEP5** $\rightarrow$ **G3** was formed (Fig S29-S30), which could

not lead to the cross-linking of **TPEP5** due to the fact that **G3** can bind with only one molecule of **DMP5**, generating a simple 1:1 inclusion complex. Therefore, **TPEP5** could not be induced to aggregate and result in the fluorescence “turn-off”.



**Fig. 3** (a) Fluorescence quenching of a solution of **TPEP5** (0.04 mM) and **G1** (0.64 mM) upon gradual addition of adiponitrile (**G3**, 0 – 20.48 mM)  $\text{CHCl}_3/\text{acetone}$  (1/8, v/v) ( $\lambda_{\text{ex}} = 330 \text{ nm}$ ). The Inset shows the photographs of the solution of **TPEP5** and **G1** in the (B) absence and (C) presence of adiponitrile (20.48 mM) under UV light (365 nm) illumination at 298 K. (b) The plot of fluorescence intensity against the concentration of **G3**.

Furthermore, transmission electron microscopy (TEM) was also used to provide further insight into the size and shape of the supramolecular aggregates formed from **TPEP5** and **G1**. As shown in Fig. 4, spherical aggregates with a diameter of  $\sim 2 \mu\text{m}$  were observed for the supramolecular aggregates formed in  $\text{CHCl}_3/\text{acetone}$  solution (Fig. 4a and 4b). Moreover, the dynamic light scattering (DLS) measurements showed that different size distributions were observed and the mean size of the above aggregates was about  $2 \mu\text{m}$  in diameter (Fig. S33), which was in good agreement with the above TEM results. Therefore, the above results further confirmed the formation of large sized supramolecular aggregates *via* host-guest recognition-mediated cross-linking.



**Fig. 4** TEM images: (a) TEM image of **TPEP5**⊃**G1** complex; (b) enlarged TEM image of (a). Samples were prepared by placing one drop of the  $\text{CHCl}_3/\text{acetone}$  solution of the mixtures of **TPEP5** with 4 equiv. **G1** onto a carbon-coated copper grid.

## Conclusions

In summary, a novel TPE-functionalized pillar[5]arene (**TPEP5**) was successfully synthesized by incorporating four pillar[5]arene groups onto the periphery of TPE through the alkyne-azide Click reaction. The formation of **TPEP5**⊃**G1** (1:2 molar ratio) supramolecular inclusion complex based on host-guest interactions led to the effective aggregation of **TPEP5**, resulting in the “turn-on” of fluorescence emission based on the AIE mechanism. Moreover, fluorescence “turn-off” could be observed upon further addition of adiponitrile due to the competitive host-guest complexation. In addition, DLS and TEM images confirmed the formation of large sized spherical aggregates due to the host-guest recognition-induced cross-linking. Therefore, this novel supramolecular system offers a new opportunity for the fabrication of novel types of pillararene-based

AIE luminogens. Future work will focus on the design and synthesis of highly efficient and selective pillararene-based functional AIE materials.

## Notes and references

- <sup>a</sup> Key Laboratory of Mesoscopic Chemistry of MOE, Center for Multimolecular Organic Chemistry, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210093, China. Fax: +86 025 83317761; Tel: +86 025 83592529; E-mail: huxy@nju.edu.cn; lywang@nju.edu.cn.
  - <sup>b</sup> College of Materials Science and Engineering, Beijing Institute of Technology, Beijing 100081, China.
- † Electronic Supplementary Information (ESI) available: [Synthesis and characterization data; determination of the association constants; fluorescence spectra and DLSI study]. See DOI: 10.1039/b000000x/
- ‡ We gratefully thank the financial support of the National Basic Research Program of China (2014CB846000) and the National Natural Science Foundation of China (No. 91227106, 21202083).
- III Jie Wu and Shu Sun contributed equally to this work.
- (a) J. Luo, Z. Xie, J. W. Y. Lam, L. Cheng, H. Chen, C. Qiu, H. S. Kwok, X. Zhan, Y. Liu, D. Zhu and B. Z. Tang, *Chem. Commun.*, 2001, 1740-1741; (b) Y. Hong, J. W. Y. Lam and B. Z. Tang, *Chem. Soc. Rev.*, 2011, **40**, 5361-5388.
  - B.-K. An, S.-K. Kwon, S.-D. Jung and S. Y. Park, *J. Am. Chem. Soc.*, 2002, **124**, 14410-14415.
  - (a) K. Hatano, H. Saeki, H. Yokota, H. Aizawa, T. Koyama, K. Matsuoka and D. Terunuma, *Tetrahedron Lett.*, 2009, **50**, 5816-5819; (b) Y. Hong, J. W. Y. Lam and B. Z. Tang, *Chem. Commun.*, 2009, 4332-4353; (c) M. Wang, G. Zhang, D. Zhang, D. Zhu and B. Z. Tang, *J. Mater. Chem.*, 2010, **20**, 1858-1867.
  - H. Tong, Y. Hong, Y. Dong, H. Tong, J. W. Y. Lam, Z. Li, Z. Guo, Z. Guo and B. Z. Tang, *Chem. Commun.*, 2006, 3705-3707.
  - (a) Y. Dong, J. W. Y. Lam, A. Qin, J. Liu, Z. Li, B. Z. Tang, J. Sun and H. S. Kwok, *Appl. Phys. Lett.*, 2007, **91**, -; (b) H. Tong, Y. Hong, Y. Dong, M. Häußler, Z. Li, J. W. Y. Lam, Y. Dong, H. H. Y. Sung, I. D. Williams and B. Z. Tang, *J. Phys. Chem. B*, 2007, **111**, 11817-11823; (c) L. Liu, G. Zhang, J. Xiang, D. Zhang and D. Zhu, *Org. Lett.*, 2008, **10**, 4581-4584; (d) Y. Hong, M. Häußler, J. W. Y. Lam, Z. Li, K. K. Sin, Y. Dong, H. Tong, J. Liu, A. Qin, R. Renneberg and B. Z. Tang, *Chem. – Eur. J.*, 2008, **14**, 6428-6437; (e) M. Wang, X. Gu, G. Zhang, D. Zhang and D. Zhu, *Anal. Chem.*, 2009, **81**, 4444-4449; (f) Q. Chen, N. Bian, C. Cao, X.-L. Qiu, A.-D. Qi and B.-H. Han, *Chem. Commun.*, 2010, **46**, 4067-4069; (g) T. Sanji, K. Shiraishi, M. Nakamura and M. Tanaka, *Chem. – Asian J.*, 2010, **5**, 817-824.
  - H. Shi, R. T. K. Kwok, J. Liu, B. Xing, B. Z. Tang and B. Liu, *J. Am. Chem. Soc.*, 2012, **134**, 17972-17981.
  - (a) X. Wang, J. Hu, T. Liu, G. Zhang and S. Liu, *J. Mater. Chem.*, 2012, **22**, 8622-8628; (b) G. Liang, J. W. Y. Lam, W. Qin, J. Li, N. Xie and B. Z. Tang, *Chem. Commun.*, 2014, **50**, 1725-1727.
  - (a) T. Ogoshi, S. Kanai, S. Fujinami, T.-a. Yamagishi and Y. Nakamoto, *J. Am. Chem. Soc.*, 2008, **130**, 5022-5023; (b) M. Xue, Y. Yang, X. Chi, Z. Zhang and F. Huang, *Acc. Chem. Res.*, 2012, **45**, 1294-1308; (c) T. Ogoshi, *J. Incl. Phen. Macro. Chem.*, 2012, **72**, 247-262; (d) X.-Y. Hu, X. Wu, S. Wang, D. Chen, W. Xia, C. Lin, Y. Pan and L. Wang, *Polym. Chem.*, 2013, **4**, 4292-4297; (e) C. Li, K. Han, J. Li, Y. Zhang, W. Chen, Y. Yu and X. Jia, *Chem. – Eur. J.*, 2013, **19**, 11892-11897; (f) X. Shu, S. Chen, J. Li, Z. Chen, L. Weng, X. Jia and C. Li, *Chem. Commun.*, 2012, **48**, 2967-2969; (g) X.-B. Hu, Z. Chen, G. Tang, J.-L. Hou and Z.-T. Li, *J. Am. Chem. Soc.*, 2012, **134**, 8384-8387; (h) Y. Yao, X. Chi, Y. Zhou and F. Huang, *Chem. Sci.*, 2014, DOI: 10.1039/C4SC00585F; (i) X. Shu, J. Fan, J. Li, X. Wang, W. Chen, X. Jia and C. Li, *Org. Biomol. Chem.*, 2012, **10**, 3393-3397.
  - During we preparing our manuscript, Huang and co-works reported the host-guest complexation induced emission based on water-soluble pillar[6]arene (**WP6**) and TPE ammonium salt, the intramolecular rotation of the TPE motif were hampered upon the addition of **WP6**. P. Wang, X. Yan and F. Huang, *Chem. Commun.*, 2014, **50**, 5017-5019.
  - J. Wang, J. Mei, R. Hu, J. Z. Sun, A. Qin and B. Z. Tang, *J. Am. Chem. Soc.*, 2012, **134**, 9956-9966.