



Cite this: *Sustainable Food Technol.*,  
2026, 4, 411

## Development and characterization of probiotic processed cheese

Deepak Kumar,<sup>a</sup> Sangita Ganguly,<sup>ID</sup> \*<sup>a</sup> Santosh Tambde,<sup>a</sup> Yogesh Khetra,<sup>ID</sup> <sup>a</sup>  
P. Narender Raju,<sup>ID</sup> <sup>a</sup> and Rashmi Hogarehalli Mallappa<sup>b</sup>

Processed cheese is a popular variety of cheese and is generally devoid of lactic organisms. The incorporation of probiotic organisms into processed cheese will provide functionality and health beneficial properties to the product, along with an increased market demand. Four strains of the probiotic organism *Lactiplantibacillus plantarum*, namely, Lp1, Lp6, Lp7 and Lp9, were subjected to thermal stability testing, and based on the maximum thermal stability, *L. plantarum* (Lp9) was selected for further experimentation. Based on the moisture level (46%) and probiotic count, an inoculum of 14% (w/w) was selected for probiotic processed cheese preparation. Probiotic processed cheese was prepared by incorporating an optimized level of probiotic inoculum in normal processed cheese at 60 °C. Probiotic processed cheese had a sensory and compositional quality comparable to that of the control cheese. The product had lower meltability, hardness and cohesiveness than the control cheese. Probiotic processed cheese exhibited improved bio-functional attributes in terms of antioxidant and ACE-inhibitory activity compared with the control cheese. Probiotic processed cheese had an initial probiotic count of 7.66 log CFU g<sup>-1</sup>, which remained stable in the processed cheese matrix during 35 days of refrigerated storage.

Received 14th July 2025  
Accepted 18th November 2025

DOI: 10.1039/d5fb00375j

rsc.li/susfoodtech

### Sustainability spotlight

Processed cheese is a popular variety of cheese and is generally devoid of lactic organisms. The incorporation of probiotic organisms into processed cheese will provide more functionality and health beneficial properties to the product, along with an increased market demand. This study describes a sustainable way to incorporate probiotic organisms into processed cheese for enhanced health beneficial properties for promoting good health and improved lifestyle. This approach contributes to sustainable food innovation by enhancing processed cheese with health promoting properties without compromising shelf stability or processing efficiency.

## 1 Introduction

Functional foods are those that provide health benefits beyond basic nutrition, such as preventing diseases and supporting health management,<sup>1</sup> and fermented dairy products are a popular type of functional food since they are believed to offer more health benefits than regular milk. The use of fermented dairy products containing probiotic bacteria as functional foods has increased due to new scientific research showing their ability to prevent diseases.

Probiotics are live microorganisms that, upon administration in adequate amounts, confer health benefits to the host.<sup>2</sup> Probiotic microorganisms offer several health benefits, including immune modulation, cholesterol reduction, and cancer prevention.<sup>3,4</sup> Some probiotics commonly used in various fermented products, including cheese, are *Lactobacillus*

and *Bifidobacterium* spp.<sup>5</sup> To provide health benefits to the host, the concentration of probiotic bacteria should be 7–8 log CFU g<sup>-1</sup> or mL of the product.<sup>6</sup>

Cheeses are dairy products having a strong potential to serve as delivery matrices for probiotic microorganisms due to their inherent physical and chemical nature. Cheese has comparatively low acidity, high pH, high buffering capacity and solid consistency. Cheeses are nutrient dense foods with a low oxygen content compared with *dahi* or yogurt. These inherent properties of cheese provide protection to the probiotic organisms during storage and gastrointestinal transit.<sup>7</sup> Hard variety of cheese like Cheddar, may provide certain benefits over yoghurt-type products regarding a delivery medium of viable probiotic, such as the low acidity and the high-fat content as compared to yoghurt environment. Further, the texture of cheese may offer an encapsulation-type protection to the microorganisms during passage through the GI tract.<sup>8</sup> Several soft cheeses, like whey Cheese,<sup>9</sup> white cheese,<sup>10</sup> and cottage cheese, have been used as carriers of live probiotic organisms.<sup>11</sup> Processed cheese is a product obtained by blending natural cheese with emulsifying salts and dairy and non-dairy ingredients. This mixture is

<sup>a</sup>Dairy Technology Division, ICAR-National Dairy Research Institute, Karnal, Haryana, India. E-mail: sangitandri@gmail.com; Fax: +91-184-2250042; Tel: +91-184-2259248

<sup>b</sup>Dairy Bacteriology Section, ICAR-National Dairy Research Institute, SRS, Bengaluru, India



subjected to a heat treatment and continuous mixing to achieve a uniform consistency suitable for extended storage. Processed cheese differs from conventional cheese since it is not obtained from traditional processes, involving traditional lactic fermentation of milk or direct acidification of milk using food grade organic acids. Processed cheese is widely used in various food applications, including sandwiches, burgers, pizzas, and ready-to-eat meals. Its unique properties, such as high meltability, smoothness, and convenience, make it a popular ingredient in the food industry.<sup>12</sup> Some researchers have tried to incorporate beneficial probiotic organisms in 'requeijao cremoso' type processed cheese and reported that the fusion stage is the most appropriate for the incorporation of organisms.<sup>13</sup> In another study, probiotic processed cheese was prepared by the addition of spores of *B. coagulans* ATCC7050. The sensory quality of probiotic processed cheese was reduced with an increment in spore count, and the viability of the probiotic organism was maintained at an optimal level during 60 days of storage.

Processed cheese is extremely popular in the Indian market and is mostly used for direct consumption. It is generally devoid of live microorganisms due to its production techniques. However, if probiotic organisms can be incorporated into processed cheese, this could lead to increased consumer acceptability and market demand, leading to categorization of the product as a functional food with added health benefits. To the best of our knowledge, no studies have been conducted in the area of the incorporation of live probiotic organisms into processed cheese. Considering all the aspects, a study is conducted for the development of probiotic processed cheese with the following objectives: (a) to develop probiotic processed cheese, and (b) to characterize the developed probiotic processed cheese, including compositional, functional, rheological, textural, sensory, microbial and bio-functional quality.

## 2 Materials and methods

### 2.1 Materials

Raw chilled cow milk (fat –4–4.2% and solid-not-fat –8.2–8.3%) and fresh as well as three-month-old matured cheddar cheese were collected from the Experimental Dairy Plant, National Dairy Research Institute, Karnal, Haryana, India. Skim milk powder (SMP) was obtained from Modern Dairies Ltd, Karnal, Haryana, India. All chemicals used during the investigation were obtained from M/s Sigma Aldrich Ltd, USA. All microbiological media were purchased from M/s Himedia, Mumbai, Maharashtra, India. Probiotic organisms, specifically *Lactiplantibacillus plantarum* four strains Lp1, Lp6, Lp7 and Lp9, were collected from the Molecular Biology Unit at the National Dairy Research Institute in Karnal, Haryana, India. Alpha Packaging's polypropylene (PP) tubs, manufactured in Surat, Gujarat, India, were procured from a local supplier in Karnal, Haryana, India.

### 2.2 Maintenance, preservation, and propagation of cultures

Freeze-dried cultures were activated and sub-cultured using the methodology described by Sameer *et al.*<sup>14</sup> *Lactobacilli* stock

cultures were stored at –20 °C in De Man, Rogosa and Sharpe (MRS) broth with 20% glycerol. For cheese production, the chosen organism was cultured in milk.

**2.2.1 Screening probiotic organisms.** The selection process for probiotic microorganisms is based on their ability to survive at different temperatures and their thermal stability.<sup>15</sup> About 50 g of probiotic culture was placed inside a polypropylene sample container. The container was then immersed in a water bath maintained at temperatures of 60, 65, and 70 °C for 5 minutes. After that, the samples were heat-treated and cooled to room temperature, followed by plating of probiotics on MRS agar, and incubated for 24–48 hours at a temperature of 37 °C. After calculating the count of probiotics, the survival rate was determined using the formula given by Guo *et al.*:<sup>16</sup>

$$\text{Survival rate(\%)} = \frac{\log \text{CFU } N_1}{\log \text{CFU } N_0} \times 100,$$

where  $N_1$  denotes the probiotic count after exposure to thermal treatment and  $N_0$  denotes the initial probiotic count.

**2.2.2 Selection of probiotic inoculum levels.** The appropriate level of probiotic inoculum was determined based on the percentage of moisture and probiotic count of the final product. To comply with the Food Safety and Standards Authority of India<sup>17</sup> legal regulations, processed cheese should have a maximum moisture content of 47%. Further, a probiotic count of 7–8 log CFU g<sup>-1</sup> is required to qualify as a probiotic product.

**2.2.3 Preparation of probiotic inoculum (probiotic curd).** The probiotic inoculum was prepared using the protocol described below (Fig. 1). The probiotic curd was prepared by inoculating about 1% culture that had been activated in skim milk to yield probiotic inoculum. At 37 °C, a 1% culture was inoculated into pasteurized standardised (16% TS) cow milk. The cow milk was standardized to a suitable solid (16% TS) level by the addition of a calculated amount of skimmed milk powder. The milk was then incubated for 14 hours at 37 °C. The



Fig. 1 Flow diagram for the preparation of probiotic inoculum.



acidity, probiotic count, and total solids of the probiotic inoculum were measured after 14 hours of incubation.

### 2.3 Preparation of probiotic processed cheese (PPC)

The control and probiotic processed cheeses were prepared in a process cheese kettle with a maximum capacity of 50 kg situated at the experimental dairy plant of ICAR-NDRI, Karnal. In one batch, about 20–25 kg of processed cheese was prepared. The manufacturing process of control processed cheese involves the use of old (*i.e.*, three months old) and young cheddar cheeses. About 25% old cheddar cheese (38.7% moisture) and 75% young cheddar cheese (39.35% moisture) were used for the preparation of the control processed cheese. Shredded cheddar cheese was blended and subjected to thermal processing at 90 °C for 5 minutes and mixed with trisodium citrate at a rate of 2% dissolved in a calculated quantity of water (76.61 ml per kg of processed cheese). Probiotic processed cheese was prepared using old and young cheddar cheese and probiotic curd (inoculum) with 16% total solids. A lower amount of water was required for probiotic processed cheese preparation as the water content of the curd was considered. The probiotic curd was added at a selected rate at 60 °C, followed by thorough mixing using a high speed automated stirrer.

### 2.4 Analysis of probiotic processed cheese

**2.4.1 Compositional analysis.** The moisture, lipid, and ash content of control and probiotic processed cheese were analysed using standard procedures.<sup>18</sup> The KELPLUS CLASSIC-DX VATS (B) was used to estimate the total protein content of cheese by applying the Micro-Kjeldahl method.<sup>19</sup> To calculate the lactose content of cheese, the protein, ash, and fat contents were subtracted from the total solid content. The titratable acidity was determined following the standard method.<sup>19</sup> The water activity of the cheese was measured using an Aqua Lab water activity meter (Decagon Devices Inc., Washington, USA).

**2.4.2 Analysis of functional characteristics.** Meltability of cheeses was estimated using a modified Schreiber test developed by Gunasekaran and AK.<sup>20</sup> The oiling-off properties of the cheese were measured using the modified method described by Wang and Sun.<sup>21</sup>

**2.4.3 Textural attributes.** The textural properties of processed cheese samples were evaluated using Texture Analyzer TAXT2i (M/s Stable Micro Systems, Godalming, Surrey, UK) equipped with a 25 Kg load cell. Cylindrical processed cheese samples (16 mm × 17.5 mm) were stored at 5 °C, and a two-bite compression test was performed for the cheese using the P-75 probe attachment to measure textural attributes. The test conditions were as follows: mode: measure force in compression; option: return to start; Pre-test speed: 2.0 mm s<sup>-1</sup>; test speed: 1.0 mm s<sup>-1</sup>; post-test speed: 2.0 mm s<sup>-1</sup>; Force: 0.10 N; trigger type: auto; data acquisition rate: 200.00 PPS. All analyses were completed in triplicate for both control and processed cheeses.<sup>22</sup>

**2.4.4 Rheological analysis.** Processed cheese samples were evaluated for rheological attributes using a controlled stress

rheometer (MCR52, Anton Paar, GmbH, Germany) with Rheoplus/32, ServiceV.3.61 software, parallel plate geometry, PP-50 (1.002° inclination), with a gap of 1.5 mm. The temperature of 20 °C was controlled using a Peltier control system throughout the study. Rheological analysis was performed soon after the product was manufactured. The measurement was conducted at 20 °C, and samples were kept for an equilibration period of 60 s before actual testing. The volumes of the prepared control and probiotic processed cheese samples were individually placed on the plate and completely covered, and the excess sample was trimmed off. A strain sweep test was conducted in advance to ensure that all rheological measurements were carried out within the linear viscoelastic region. The experiment was carried out using a frequency sweep test. A frequency sweep was conducted by applying frequencies in descending order from 100 Hz to 0.01 Hz at 20 °C using 0.1% strain amplitude. The strain eventually chosen was 0.5, which is well within the linear region. Both  $G'$  and  $G''$  were obtained from the results of this test. A temperature sweep was performed at a constant frequency of 10 Hz and a constant amplitude strain of 0.1% (obtained from LVE), with the temperature varying from 20 to 90 °C at 3 °C min<sup>-1</sup> using a Peltier heating element. The storage modulus ( $G'$ ), the loss modulus ( $G''$ ), and loss tangent ( $\tan \delta$ ) were determined. All measurements were performed in triplicate.

**2.4.5 Sensory analysis.** Ten trained panellists comprising staff and scientists of the Dairy Technology Division, National Dairy Research Institute, Karnal, Haryana, India, evaluated the sensory parameters of control and probiotic processed cheeses.<sup>23</sup> The sensory evaluation analysis was conducted in a sensory evaluation laboratory with separate booths. The training was conducted using market-processed cheese samples, including a control processed cheese sample, to familiarize the panellists with the evaluation criteria. The samples were served in the sensory laboratory at a controlled temperature (20 ± 2 °C) under normal light. The panellist selection criteria were based on a non-smoker, aged between 25 and 60 years, non allergic to dairy products, and availability to participate in the sensory analysis during testing time (11.00–11.30 am; 3.00–3.30 pm). The panellists assessed various attributes of cheese, such as flavor, body and texture, colour and appearance, and sliceability (maximum score: total sensory score-100; colour and appearance-10; spreadability-15; texture-30 and flavour-45). The panellists were allowed to use water and bland crackers for palate cleansing between the samples. An exemption from ethical committee approval was obtained to conduct a sensory study for dairy and food products developed at the ICAR-National Dairy Research Institute, Karnal. However, the study was approved by the Institute of ICAR-NDRI, Karnal, and information regarding the product was provided to each panellist prior to conducting the study. Consent was obtained from each panellist prior to their participation in the study.

**2.4.6 Colour profile analysis.** The colour profiles of control and probiotic processed cheeses were measured using the reflectance spectroscopy method with the reflectance meter, colour flex (Hunter Lab, Reston, Virginia, USA). Universal Software (Version 4.10) was used for colour profile analysis. Before



conducting the test, the instrument requires calibration with standard black glass and white tile, as specified by the manufacturer. The light source was a dual beam xenon flash lamp. Data were obtained in terms of  $L^*$  [lightness, ranges from 0 (black) to 100 (white)],  $a^*$  [redness, ranges from +60 (red) to -60 (green)], and  $b^*$  [yellowness, ranges from +60 (yellow) to -60 (blue)] from the software in values of the international colour system.<sup>24</sup>

**2.4.7 Bio-functional activities of control and probiotic processed cheeses.** Water-soluble extracts of cheese samples were prepared by employing a method developed by Kuchroo and Fox.<sup>25</sup> ABTS and DPPH methods were used to determine the antioxidant activity of water-soluble extracts.<sup>26</sup> Angiotensin-converting enzyme (ACE) inhibitory activity was measured using the modified method by Hernandez *et al.*<sup>27</sup>

**2.4.8 Viability of probiotic organisms.** The probiotic count of probiotic-processed cheese was determined using the procedure described by Meira *et al.*<sup>28</sup> with MRS medium, and the plates were incubated at 37 °C for 48 h and expressed as log CFU g<sup>-1</sup>. PPC was stored in polypropylene tubs at a refrigeration temperature of 7 °C. The viability of probiotic organisms was estimated during refrigerated storage at every 5-day interval until the end of the sensory shelf life.

## 2.5 Statistical analysis

The statistical analyses were performed using IBM SPSS 20 (version 20; IBM Corporation, USA). A two-tailed paired *t*-test was used to determine the significance of the difference between the two treatments, while an analysis of variance (ANOVA) was employed to examine the significance of the difference between more than two treatments at the 5% level of significance. Three replicates were used for all experimental analyses for probiotic and control processed cheeses.

# 3 Results and discussion

## 3.1 Screening of probiotic strain

Four probiotic organisms of *Lactobacillus* spp., Lp1, Lp6, Lp7, and Lp9, were checked for thermal stability at different time-temperature conditions to select the most suitable culture for the development of probiotic processed cheese. Among the four strains, the Lp9 strain showed a higher survival rate of 89.19%

and 83.83% when exposed to thermal temperatures of 60 and 65 °C for 5 minutes, respectively (Table 1). When exposed to a temperature of 60 °C for 5 minutes, Lp9 had the highest number of viable probiotic counts (*i.e.*, 9.49 log CFU mL<sup>-1</sup>), which was significantly greater ( $p < 0.05$ ) than the other strains. However, when all strains were subjected to a temperature exposure of 70 °C for 5 min, there was a complete absence of live probiotic organisms. Strain Lp9 showed maximum thermal resistance in the experiment; therefore, it was selected for further experiments. The *L. plantarum* Lp9 is an indigenous strain that displayed a high survival at a low pH and bile and demonstrated health-promoting properties in terms of anti-oxidative, antibacterial, and cholesterol lowering properties with a potential for exploitation in functional food development.<sup>29</sup> De Angelis *et al.*<sup>30</sup> conducted a study on the heat stress tolerance and responsiveness of different strains of *L. plantarum*. The researchers found that the stationary-phase cells of *L. plantarum* DPC2739 had decimal reduction times (*D* value) of 32.9 and 14.7 s at 60 °C and 72 °C, respectively, in sterile milk. Parente *et al.*<sup>31</sup> studied the heat stress behavior of *L. plantarum* and demonstrated that stationary-phase cells exhibited greater resistance to heat stress compared to cells in the exponential growth phase. De Angelis and Gobbetti<sup>32</sup> stated that during the growth phase, cells undergo an adaptive process where certain genes are activated to handle various stress conditions, such as nutrient depletion, acidity, and heat. This adaptive mechanism enables cells to express a broad stress response, resulting in the development of more resilient cells capable of surviving unfavorable growth environments.

**3.1.1 Standardization of inoculum level.** Probiotic inoculum was prepared by inoculating about 1% culture that had been activated in skim milk. At 37 °C, a 1% culture was inoculated into pasteurized standardised (16%) cow milk. The milk was then incubated for 14 hours at 37 °C (Fig. 1). The probiotic inoculum prepared using standardized cow milk had a probiotic count of 10.38 log CFU g<sup>-1</sup> with an acidity of 0.82% LA (lactic acid). Processed cheese was prepared using cheddar cheese and probiotic inoculum, which was added to the processed cheese at 60 °C at levels of 10, 12, 14, and 15% (w/w). The results indicated that at all levels of 10–15% inoculum, the probiotic count was sufficiently higher (7.54–7.91 log CFU g<sup>-1</sup>) to consider the product as probiotic; however, the moisture content of cheese varies significantly (Table 2). The moisture

Table 1 Screening of probiotic strains<sup>a</sup>

Probiotic organisms	Thermal treatments				
	Without heat treatment (log CFU mL <sup>-1</sup> )	60 °C/5 min		65 °C/5 min	
		Log CFU mL <sup>-1</sup>	% Survival	Log CFU mL <sup>-1</sup>	% Survival
Lp1	10.54 <sup>a</sup> ± 0.01	9.30 <sup>b</sup> ± 0.02	88.24	6.56 <sup>c</sup> ± 0.19	62.24
Lp6	10.42 <sup>a</sup> ± 0.00	9.21 <sup>b</sup> ± 0.00	88.39	8.62 <sup>c</sup> ± 0.01	82.73
Lp7	10.58 <sup>a</sup> ± 0.00	9.27 <sup>b</sup> ± 0.08	87.62	6.93 <sup>c</sup> ± 0.02	65.50
Lp9	10.64 <sup>a</sup> ± 0.05	9.49 <sup>b</sup> ± 0.13	89.19	8.92 <sup>c</sup> ± 0.01	83.83

<sup>a</sup> Mean ± S.D ( $n = 3$ ); means with different superscripts within a row differ significantly ( $p < 0.05$ ).



**Table 2** Standardization of the inoculum level of the probiotic processed cheese (PPC)<sup>a</sup>

Inoculum level (%)	Moisture of PPC (%)	Probiotic count of PPC (log cfu mL <sup>-1</sup> )
10	44.65 <sup>d</sup> ± 0.06	7.54 <sup>d</sup> ± 0.02
12	45.12 <sup>c</sup> ± 0.07	7.58 <sup>c</sup> ± 0.02
14	46.09 <sup>b</sup> ± 0.04	7.81 <sup>b</sup> ± 0.00
15	47.35 <sup>a</sup> ± 0.25	7.91 <sup>a</sup> ± 0.01

<sup>a</sup> Mean ± S.D ( $n = 3$ ); means with different superscripts within a row differ significantly ( $p < 0.05$ ).

**Fig. 2** Flow diagram for the preparation of probiotic processed cheese.

content of the processed cheese produced with a 14% (w/w) inoculum level during the experiment was observed to be 46.09%. Similarly, at the 15% inoculum level, the moisture content was higher (47.35%) and not within the legal limit. An inoculum level of 15% was not selected, as the moisture content of the processed cheese exceeded the legal limit. Considering

the appropriate moisture content (46.09%) and sufficiently high probiotic count (7.81 log CFU g<sup>-1</sup>) of the end product, an inoculum level of 14% (w/w) was selected for further study.

### 3.2 Development of probiotic processed cheese

Probiotic processed cheese was prepared as per the given protocol (Fig. 2). The selected *L.plantarum* (Lp9) was added at a 14% (w/w) level at the final stage of processed cheese preparation at a temperature of 60 °C with thorough mixing. The developed probiotic processed cheese is shown in Fig. 3A and B.

### 3.3 Characterization of probiotic processed cheese

**3.3.1 Composition and physico-chemical characterization of control and probiotic processed cheeses.** Table 3 presents the compositions of the control and probiotic processed cheeses. The moisture contents of control and probiotic processed cheese were 46.00% and 46.09%, respectively, which were non-significant ( $p > 0.05$ ). Additionally, a non-significant difference was observed in fat, protein, lactose, ash content, and pH values of the control and probiotic processed cheeses. The probiotic count of probiotic process cheese was 7.81 log CFU g<sup>-1</sup>. Silva *et al.*<sup>13</sup> developed a low sodium probiotic queijo cremoso

**Table 3** Composition, physico-chemical characteristics and the probiotic count of the control processed cheese (CPC) and probiotic processed cheese (PPC)<sup>a</sup>

Parameter	CPC	PPC
<b>Composition</b>		
Moisture (%)	46.00 ± 0.69 <sup>a</sup>	46.09 ± 0.04 <sup>a</sup>
Fat (%)	26.65 ± 0.86 <sup>a</sup>	26.33 ± 0.76 <sup>a</sup>
Protein (%)	19.06 ± 0.15 <sup>a</sup>	19.25 ± 0.03 <sup>a</sup>
Lactose (%)	4.01 ± 0.23 <sup>a</sup>	4.20 ± 0.14 <sup>a</sup>
Ash (%)	4.38 ± 0.12 <sup>a</sup>	4.14 ± 0.14 <sup>a</sup>
<b>Physico-chemical properties</b>		
pH	5.76 ± 0.05 <sup>a</sup>	5.67 ± 0.04 <sup>a</sup>
Titrate acidity (%) LA	0.97 ± 0.00 <sup>a</sup>	1.04 ± 0.01 <sup>a</sup>
Water activity	0.94 ± 0.00 <sup>a</sup>	0.96 ± 0.00 <sup>a</sup>
Probiotic count (log CFU g <sup>-1</sup> )	—	7.81 ± 0.00

<sup>a</sup> Mean ± S.D ( $n = 3$ ); means with different superscripts within a row differ significantly ( $p < 0.05$ ).

**Fig. 3** (A) Probiotic processed cheese and (B): sliced probiotic processed cheese.

processed cheese using *Lactobacillus acidophilus* with a count higher than 6 log CFU g<sup>-1</sup>. In another study, probiotic processed cheese analogues with reduced emulsifying salts were prepared using *Bacillus coagulans* spores with a count of 7–8 log CFU g<sup>-1</sup>, with a shelf life of 60 days at refrigerated storage.<sup>33</sup>

**3.3.2 Functional properties of control and probiotic processed cheeses.** It is necessary to examine the functional properties of processed cheese samples, *viz.*, meltability and oiling-off, because these properties determine the quality characteristics of processed cheese.

**3.3.2.1 Meltability.** Meltability refers to how easily cheese can flow or spread when heated.<sup>34</sup> To evaluate processed cheese quality, meltability, measured as the ratio of melted to unmelted cheese area, is a crucial parameter. In our study, a significant ( $p < 0.05$ ) difference in meltability was found between control and probiotic processed cheeses, with values of 10.24 and 9.012, respectively (Fig. 4A). The possible reason for the lower meltability of probiotic processed cheese might be due to the addition of curd. Chaudhary *et al.*<sup>35</sup> found that the meltability of processed cheese varied from 10.56 to 15.55 depending on the intact casein content, and our experiment yielded similar meltability values within this range. The incorporation of non-fat-dry milk and whey protein concentrate (WPC) into processed cheese formulations may contribute to decreased meltability in processed cheese.<sup>36,37</sup> The reason has been proposed as the heat-induced disulfide interactions, involving the formation of free sulfhydryl groups from  $\beta$ -lactoglobulins, having a marked influence on the melting attributes of the processed cheeses.<sup>36</sup> In another study, researchers heated whey protein dispersions to induce disulfide cross-linking between the whey proteins and produced polymerized cross-linked whey proteins, followed by addition into a model processed cheese system. They reported that as the level of polymerized whey proteins increased, there was a decline in the meltability of the process cheese analogs produced.<sup>38</sup> Further, an increment in mixing speed during processed cheese manufacturing is associated with a decreased meltability of the final product.<sup>39</sup> In our case, probiotic processed cheese manufacturing was associated with extensive stirring and enhanced mixing speed so that probiotic curd could mix uniformly into the processed cheese matrix, leading to a reduced meltability of the probiotic sample compared to the control one.

**3.3.2.2 Oiling-off.** There was a significant ( $p < 0.05$ ) difference in oiling-off between control and probiotic processed cheeses (Fig. 4B). The control and probiotic processed cheeses had oiling-off values of 2.30 and 2.23, respectively. The higher oiling-off value of the control compared to the probiotic cheese might be attributed to the higher amount of aged cheese in the control compared to the probiotic cheese. Chaudhary *et al.*<sup>35</sup> reported that aged cheeses provided a greater oiling-off compared to young cheddar cheeses due to increased protein breakdown and lipolysis.

Oiling off the properties of processed cheese is contributed by free oil formation and oil separation due to the leakage of fat. This occurred as a result of liquefied fat separation particularly at the surface of the cheese when the cheese is subjected to melting during the heating process.<sup>40</sup> Some studies disclosed that the occurrence of free oil formation happens due to the presence of additional proteins in the processed cheese matrix, which decreases the ability of processed cheese to maintain a uniform emulsion of fat inside the protein matrix.<sup>41</sup> The stability of processed cheese emulsions can be affected by pH. Process cheese with a higher pH had an open structure and therefore caused a weaker emulsion, resulting in oiling off.<sup>42</sup> In our case, the control had a higher pH than the probiotic-processed cheese.

#### 3.4 Textural attributes of control and probiotic processed cheeses

Textural attributes studied for control and probiotic processed cheeses include hardness, adhesiveness, springiness, cohesiveness, gumminess, chewiness, and resilience. The textural attributes of the control and probiotic cheeses are shown in Table 4.

The hardness value of probiotic processed cheese was 9.327 N, which was significantly ( $p < 0.05$ ) lower than that of the control (10.188 N). The higher hardness of the control cheese compared to its probiotic counterpart might be attributed to the addition of curd to the probiotic cheese, as curd typically possesses a weaker structure in comparison to rennet curd. Some researchers have reported that the incorporation of whey proteins in a process cheese formula may lead to an increased firmness.<sup>36,37</sup> Similarly, the addition of phytosterols at 3% and 4% levels increased the firmness of processed cheese spreads at a significant ( $p < 0.05$ ) level compared to the control.<sup>42</sup> The increased firmness of the cheese spread might be due to the




Fig. 4 (A) Meltability of the control processed cheese (CPC) and probiotic processed cheese (PPC) and (B) oiling-off of CPC and PPC.

Table 4 Textural attributes of the control (CPC) and probiotic processed cheese (PPC)<sup>a</sup>

Textural attributes	CPC	PPC
Hardness	10.19 ± 0.000 <sup>a</sup>	9.337 ± 0.003 <sup>b</sup>
Springiness	0.37 ± 0.002 <sup>a</sup>	0.20 ± 0.000 <sup>b</sup>
Cohesiveness	0.24 ± 0.002 <sup>a</sup>	0.34 ± 0.001 <sup>b</sup>
Gumminess	2.48 ± 0.021 <sup>a</sup>	3.14 ± 0.009 <sup>b</sup>
Chewiness	0.92 ± 0.001 <sup>a</sup>	0.63 ± 0.001 <sup>b</sup>

<sup>a</sup> Mean ± S.D. ( $n = 3$ ); means with different superscripts within a row differ significantly ( $P < 0.05$ ).



addition of free phytosterols, which were in powder form and occupied the free space present in the cheese spread. Further, the firmness of all the inulin-incorporated processed cheese spreads was significantly higher than that of the control.<sup>43</sup> This might be due to the high water binding capacity and higher molecular weight of inulin.<sup>44</sup> Further, the presence of MPC at a higher level in the matrix causes a significant increase in the hardness of processed cheese spread.<sup>45</sup> In our case, the addition of curd caused an increment in moisture content, leading to reduced hardness. In a study incorporation of denatured whey proteins in the cheese had resulted higher moisture and lower hardness in processed cheese as compared to UF added processed cheese.<sup>46</sup> Similarly, studies have associated the effect of the final pH of process cheese with its firmness.<sup>12</sup> It was observed that with the increment of the final pH of the processed cheese from 5.0 to 6.2, its firmness initially improved. In our case, higher pH and hardness were observed in the control cheese. The springiness values for control and probiotic cheese were 0.372 and 0.201, respectively, showing a statistically significant difference ( $p < 0.05$ ). The control had a chewiness value of 0.923, while the probiotic cheese had a chewiness value of 0.633. The chewiness of the probiotic processed cheese was significantly lower ( $p < 0.05$ ) compared to the control; this might be due to the lower hardness of the probiotic cheese. Both springiness and chewiness are firmness-dependent parameters, resulting in higher values for control cheese compared to probiotic cheese, which is consistent with findings reported by El-Aidie *et al.*<sup>47</sup> The gumminess values of the control and probiotic processed cheeses were 2.479 and 3.143, respectively. The higher gumminess values of probiotic cheese were owing to the higher cohesiveness values of probiotic cheese compared to

control cheese. The cohesiveness values of the control and probiotic processed cheeses were 0.243 and 0.337, respectively. The cohesiveness value of the probiotic cheese was significantly ( $p < 0.05$ ) higher than that of the control cheese. The disruption of the continuity of the protein matrix due to the addition of curds may lead to a softer texture, which might be the reason for the increased gumminess and cohesiveness of probiotic processed cheese. A similar result was found, in which an increment in gumminess and cohesiveness was observed in the case of the processed whey cheese.<sup>48</sup> However, with the addition of phytosterols from 0 to 4%, a sharp, steady and significant ( $p < 0.05$ ) decrease in the work of adhesion was reported in processed cheese spread.<sup>42</sup> The decrease in the work of adhesion was due to the reduction in work needed to overcome the attractive force between the surface of the product and the surface of the probe as a result of the reduction in stickiness of the product and might also be due to weak gel formation as a consequence of the addition of insoluble phytosterols. However, increasing inulin levels caused a marginal decline in the work of adhesion.<sup>43</sup>

### 3.5 Rheological attributes of the control and probiotic processed cheeses

**3.5.1 Small amplitude oscillatory rheology.** The viscoelasticity of foods can be analysed using a frequency sweep test. It measures the storage and loss modulus of food over a range of frequencies. The dynamic rheology data obtained in the experiment confirmed the viscoelastic nature of the control and probiotic processed cheeses (Fig. 5A and B). The storage modulus was higher than the loss modulus for both samples,



Fig. 5 Rheological attributes of the control and probiotic processed cheese: (A) storage modulus and (B) loss modulus.



indicating a dominant contribution of the elastic component to the product's viscoelasticity.

This behaviour is typically associated with solid viscoelastic materials.<sup>49</sup> The values observed for  $G'$  and  $G''$  of the probiotic cheese were lower than those observed for the hard cheese.<sup>50</sup> A similar tendency of viscoelasticity was observed for processed cheese.<sup>46</sup> Similarly, Lee *et al.*<sup>51</sup> observed that  $G'$  values of unheated processed cheese (at 20 °C) were higher than those of  $G''$  with increasing frequency ranging from 0.01 to 10 Hz. Hence, the investigation of the above experiments matches our results and gives a clear idea of product behaviour. The control and probiotic processed cheeses exhibited a visco-elastic behaviour during the unmelted state at 20 °C. Further, the probiotic processed cheese had a lower elastic modulus ( $G'$ ) than the control processed cheese, which might be attributed to lower hardness than the control processed cheese, as depicted in Fig. 5.

**3.5.2 Temperature sweep of the control and probiotic processed cheeses.** Temperature sweeps were performed from 20 to 90 °C at 4 °C min<sup>-1</sup>. The test combines thermal treatments and low shear. Measurements were conducted at a constant frequency of 1 Hz and a constant strain of 1%. In duplicate, these conditions ensured that the samples were tested within the linear viscoelastic range.

The probiotic processed cheese had a profile with an increase in  $G''$  and  $G'$  after the unique crossing-point, corresponding to the maximum value of  $\tan \delta$  observed at 65 °C, as shown in Fig. 6a. The viscous component  $G''$  was greater than

the elastic component  $G'$ . The pattern observed for the visco-elastic changes with temperature for processed cheese is very different from that for the other cheeses.<sup>52</sup> This profile shows good meltability of the probiotic processed cheese, which is associated with enough fat emulsification. The control processed cheese had a profile with an increase in  $G''$  and  $G'$  after the unique crossing-point, corresponding to the maximum value of  $\tan \delta$  observed at 61 °C, as illustrated in Fig. 6b. The viscous component  $G''$  was higher than the elastic component  $G'$ . The pattern of viscoelastic changes with temperature is the same for both cheeses, *i.e.*, the probiotic processed and control processed cheeses.

In our study, a decline in the  $G'$  was observed with an increment of temperature from 20 to 90 °C for the control and probiotic processed cheeses, indicating the loosening of elasticity of the cheese matrix as the temperature increased. This indicates the breaking of protein–protein interactions inside a casein network as well as the liquefaction of fat together with the distortion of fat globules, which had plasticized in the protein matrix and allowed it to flow.<sup>53</sup> The fat completely melted at 40 °C;<sup>54</sup> thus, an increment was observed in  $\tan \delta$  in the control and probiotic samples Fig. 6. An increase in  $\tan \delta$  was observed with an increment in temperature after 40 °C, exhibiting a melt-like behaviour. At this temperature, protein softening started in the cheese matrix due to the decline of specific casein–casein interactions in the structure.<sup>54,55</sup>

### 3.6 Sensory quality of products

Sensory evaluation is an important parameter for the acceptability of processed cheese. The body and texture, sliceability and overall acceptability scores were lower for probiotic processed cheese compared to the control. However, the color and flavour scores obtained for probiotic processed cheese were higher than the control. However, there were non-significant differences observed in all sensory parameters between the control processed cheese and the probiotic processed cheese (Fig. 7). Several authors have studied the effects of the incorporation of probiotic organisms into varieties of cheeses. The addition of probiotic *Bacillus coagulans* spores in the processed cheese non-significantly affected the sensory attributes of the final product.<sup>56</sup> Probiotic Minas fresh cheese containing probiotic bacteria *L. paracasei* subsp. *paracasei* had a better sensorial quality compared to the control cheese.<sup>57</sup> The effect of the addition of whey protein in the process cheese matrix has been extensively studied on its sensory qualities, which exhibited similar or enhanced sensory quality in the final product.<sup>36,38,55</sup> In our case, fermented milk had been incorporated into the probiotic processed cheese, which had contributed towards an improved flavour of the probiotic processed cheese. Fermentation of milk is known for the production of several significant aromatic and flavour compounds as metabolites.<sup>58</sup>

### 3.8 Colour profile for control and probiotic processed cheese

Colour is considered an important parameter for the sensory acceptability of a product.<sup>59</sup> In colour measurement, " $L^*$ " value




Fig. 6 (a): Temperature sweep of probiotic processed cheese. (b): Temperature sweep of the control processed cheese.





Fig. 7 Sensory analysis of the control processed cheese (CPC) and probiotic processed cheese (PPC).

represents lightness (100) and blackness (0); “ $a^*$ ” represents red (+ve) to green (–ve) hues, while “ $b^*$ ” represents yellow (+ve) to blue (–ve) hues.<sup>60</sup> The  $L^*$  values for control and probiotic processed cheeses were 70.87 and 71.07, respectively, as shown in Table 5. Both cheeses had statistically similar  $L^*$  values ( $p > 0.05$ ). The probiotic processed cheese was whiter than the control one, which could be attributed to the incorporation of probiotic curd. The ability of protein particles to scatter light in the visible spectrum can lead to a whitening effect, thereby attributing the product’s whiteness to the protein matrix.<sup>14</sup> The respective  $a^*$  values for the control and probiotic cheeses were 1.19 and 1.20, respectively. The  $a^*$  values of the probiotic and control processed cheeses were not statistically significant from each other ( $p > 0.05$ ). The  $b^*$  values for control and probiotic processed cheeses were statistically non-significant ( $p > 0.05$ ) and 24.91 and 25.83, respectively. The higher yellowness of the probiotic cheese (Table 5) might be attributed to the incorporation of probiotic curd prepared from cow milk.<sup>61</sup>

### 3.9 Bio-functional activities of the control and probiotic processed cheeses

Control and probiotic processed cheeses were evaluated for bio-functional activities, viz., *in vitro* ACE-inhibitory and antioxidant activity.

**3.9.1 ACE-inhibitory activity.** The ACE-inhibitory activities of the water-soluble extracts of the probiotic and control cheeses were 83.61% and 64.84%, respectively, as shown in Fig. 8C. Probiotic processed cheese showed significantly ( $p < 0.05$ ) higher ACE-inhibitory activity compared to the control cheese. The higher ACE-inhibitory activity of the probiotic cheese was attributed to the generation of substantial amounts of ACE-inhibitory peptides. Ong *et al.*<sup>62</sup> reported that cheddar cheese from 1 to 36 weeks of ripening showed an ACE-inhibitory activity in the range of 10–87.0% due to the production of ACE-inhibitory peptides. Several researchers had reported higher *in vitro* ACE-inhibitory properties in probiotic yoghurt.<sup>63,64</sup> The increased protein breakdowns and generation of bio-active



Table 5 Colour profile of the control (CPC) and probiotic processed cheese (PPC)<sup>a</sup>

Colour parameters	<i>L</i> *		<i>a</i> *		<i>b</i> *	
	CPC	PPC	CPC	PPC	CPC	PPC
Colour values	70.87 ± 0.33 <sup>a</sup>	71.07 ± 0.40 <sup>a</sup>	1.19 ± 0.03 <sup>a</sup>	1.20 ± 0.06 <sup>b</sup>	24.91 ± 0.04 <sup>a</sup>	25.83 ± 0.03 <sup>a</sup>

<sup>a</sup> Mean ± S.D. (*n* = 3); means with different superscripts within a row differ significantly (*p* < 0.05).

peptides during fermentation might be the major reason for the higher ACE-inhibitory attributes of probiotic fermented products.<sup>63,64</sup> Most of the reported ACE-inhibitory peptides are usually short peptides with a proline residue at the carboxyl terminal end. Proline is known to be tolerant to degradation by digestive enzymes, which may pass the peptides with the same amino acid residues from the small intestine into the blood stream.<sup>65</sup> In our case, the incorporation of probiotic curd resulted in a higher ACE-inhibitory property of the probiotic-processed cheese.

**3.9.2 Antioxidant activity.** The total antioxidant capacity (TAC) of water-soluble extracts of the probiotic and control cheeses was estimated by ABTS and DPPH free radical scavenging assay. By employing the ABTS assay, probiotic and control cheese samples displayed 85.46% and 66.94% inhibition, respectively (Fig. 8A). The total antioxidant activities of the probiotic and control processed cheeses were found to be 1.95 μM mg<sup>-1</sup> and 1.54 μM mg<sup>-1</sup>, respectively. It was observed that the probiotic processed cheese had significantly (*p* < 0.05) higher antioxidant activity in contrast to the control processed cheese. This could be attributed to the release of anti-oxidative peptides by the starter proteases. The TAC of probiotic and control cheese was found to be 20.73 μM mg<sup>-1</sup> and 5.03 μM mg<sup>-1</sup>, respectively, as determined using DPPH assay (Fig. 8B). The total antioxidant activity observed by the DPPH assay was significantly (*p* < 0.05) higher for probiotic cheese than the control cheese. Gupta *et al.*<sup>66</sup> reported that cheddar cheese was made using *L. casei* ssp. *casei* and *L. paracasei* ssp. *paracasei* as adjunct cultures displaying an antioxidant activity of up to 90% as percentage inhibition. Consumption of probiotics alone or in food has been reported to have a strain-specific antioxidant activity and a reduction in cellular damage caused by oxidation.<sup>67</sup> In milk, an increment in antioxidant activity was observed after fermentation using several probiotic strains.<sup>68</sup> The antioxidant activity of curd and yogurt is mainly derived from the hydrolysis of milk components by starter organisms.<sup>65</sup> Particularly, the hydrolysis products weighing 4–20 KDa showed a remarkably higher antioxidant activity in the antioxidant assay.<sup>69</sup> Additionally, it was observed that an increased degree of hydrolysis of milk proteins was associated with an increment in antioxidant activity, as observed by DPPH, ABTS, and FRAP radical scavenging activity.<sup>70</sup> Further, it has been reported that hydrolysis and the release of bacterial cell wall components during fermentation may lead to the release of phenolic compounds in the food matrix, resulting in an enhanced antioxidant capacity.<sup>69</sup> In our case, probiotic fermentation of milk during curd production generated bioactive compounds with

a higher antioxidant activity in probiotic curd. The incorporation of probiotic curd in processed cheese contributed towards enhanced antioxidant activity of the probiotic processed cheese compared to the control cheese.

### 3.10 Viability of probiotic organisms in process cheese

The probiotic count was found to be 7.66 log CFU g<sup>-1</sup> in fresh probiotic processed cheese. There was a significant (*p* < 0.05) increase in probiotic count during refrigerated storage (Fig. 9). The probiotic count was 9 log CFU g<sup>-1</sup> on the 20<sup>th</sup> day of refrigerated storage; thereafter, the count decreased to 8.19 log CFU g<sup>-1</sup> at the 35<sup>th</sup> day of refrigerated storage at 4 °C in PP packaging material. A population of 10<sup>6</sup>–10<sup>7</sup> CFU g<sup>-1</sup> in the final product was reported to be sufficient and effective to consider a product as a probiotic one.<sup>71</sup> Several researchers have reported various stabilities of various probiotic organisms in different cheese matrices. An increment in *L. plantarum*96 was reported



Fig. 8 (A) % ABTS inhibition of CPC and PPC. (B) % DPPH scavenging activity of CPC and PPC. (C) ACE inhibitory activity of CPC and PPC.






Fig. 9 Probiotic count of the probiotic processed cheese during refrigerated storage.

in the experimentally manufactured semi-hard cheese with low-cooking curd during a 180-day maturation period, with a final count of  $7.39 \log \text{CFU g}^{-1}$ .<sup>72</sup> The initial count of *L. acidophilus* La 05 reduced significantly ( $p < 0.05$ ) from  $7.87 \log \text{CFU g}^{-1}$ , to  $7.6 \log \text{CFU g}^{-1}$  at the end of the 12 days of refrigerated storage in buffalo milk Ricotta cheese.<sup>14</sup> Madureira *et al.*<sup>73</sup> and Meira *et al.*<sup>24</sup> observed an increment in probiotic count in Ricotta cheese during refrigerated storage. It was observed that the *L. acidophilus* Ki strain increased from  $7.15 \log \text{CFU g}^{-1}$  to  $9.39 \log \text{CFU g}^{-1}$  in the plain whey cheese matrix at the end of 28 days of storage at  $7^\circ \text{C}$ .<sup>73</sup> However, the respective organism increased from  $7.06 \log \text{CFU g}^{-1}$  to  $7.97 \log \text{CFU g}^{-1}$  in the salt-added whey cheese matrix at 28 days of storage at  $7^\circ \text{C}$ .<sup>73</sup> The count of *L. acidophilus* was increased in goat Ricotta cheese matrix from  $6.01 \pm 0.6 \log \text{CFU g}^{-1}$  to  $6.29 \pm 0.9 \log \text{CFU g}^{-1}$  at 7 days of refrigerated storage.<sup>24</sup>

## 4 Conclusion

Processed cheese is a popular variety of cheese that is mainly devoid of any live microorganisms due to its production process. The current study attempts to develop probiotic processed cheese through the incorporation of fermented milk containing live lactobacillus spp. in processed cheese emulsion at  $60^\circ \text{C}$  temperature. Probiotic processed cheese exhibited 46% moisture, 28% fat, and 19% protein, with a probiotic count of  $7.66 \log \text{CFU g}^{-1}$ . The product had acceptable sensory properties that were comparable to those of the control product. The product had lower meltability, hardness and cohesiveness compared to control cheese. Control and probiotic processed cheeses exhibited a visco-elastic behaviour during the unmelted state at  $20^\circ \text{C}$ . However, the probiotic processed cheese had a lower elastic modulus ( $G'$ ) than the control processed cheese, which might be attributed to the lower hardness of the probiotic processed cheese. The probiotic processed cheese exhibited significantly higher ACE-inhibitory and antioxidant properties as observed through ABTS and DPPH studies. The reason might be that the probiotic fermentation of milk during curd production generated bioactive compounds, which resulted in

enhanced bio-activity of probiotic processed cheese. The product had a probiotic count of  $7.66 \log \text{CFU g}^{-1}$  in fresh, which increased significantly ( $p < 0.05$ ) up to the 20<sup>th</sup> day of refrigerated storage, followed by a decline. Probiotic viability was maintained at more than  $8 \log \text{CFU g}^{-1}$  in the processed cheese matrix during 35 days of refrigerated storage.

This approach could enhance the functional attributes of processed cheese with enhanced consumer acceptability and market demand. Further, this approach contributes to sustainable food innovation by enhancing processed cheese with health promoting properties without compromising shelf stability or processing efficiency.

## Conflicts of interest

The authors declare that they have no conflicts of interest.

## Data availability

All data supporting the findings of this study are available from the corresponding author upon reasonable request.

## Acknowledgements

The authors are thankful to the Director, ICAR-NDRI, Karnal, for the funding support for this project.

## References

- 1 M. Alongi and M. Anese, *J. Funct. Foods*, 2021, **81**, 104466.
- 2 G. Reid, A. A. Gadir and R. Dhir, *Front. Microbiol.*, 2019, **10**, 444972.
- 3 M. Kechagia, D. Basoulis, S. Konstantopoulou, D. Dimitriadi, K. Gyftopoulou, N. Skarmoutsou and E. M. Fakiri, *ISRN Nutr.*, 2013, **2013**, 1–7.
- 4 R. Nagpal, H. Yadav, A. K. Puniya, K. Singh, S. Jain and F. Marotta, *Int. J. Probiotics Prebiotics*, 2007, **2**, 75.
- 5 J. Alard, V. Peucelle, D. Boutillier, J. Breton, S. Kuyille, B. Pot, S. Holowacz and C. Grangette, *Benefic. Microbes*, 2018, **9**, 317–331.
- 6 N. K. Ganguly, S. K. Bhattacharya, B. Sesikeran, G. B. Nair, B. S. Ramakrishna, H. P. S. Sachdev and R. Hemalatha, *Indian J. Med. Res.*, 2011, **134**, 22–25.
- 7 R. Karimi, A. M. Mortazavian and A. G. Da Cruz, *Dairy Sci. Technol.*, 2011, **91**, 283–308.
- 8 J. M. Castro, M. E. Tornadijo, J. M. Fresno and H. Sandoval, *BioMed Res. Int.*, 2015, **2015**, 1–11.
- 9 A. R. Madureira, M. Amorim, A. M. Gomes, M. E. Pintado and F. X. Malcata, *Food Res. Int.*, 2011, **44**, 465–470.
- 10 A. Kasimoğlu, M. Göncüoğlu and S. Akgün, *Int. Dairy J.*, 2004, **14**, 1067–1073.
- 11 L. Abadía-García, A. Cardador, S. T. Martín del Campo, S. M. Arvizu, E. Castaño-Tostado, C. Regalado-González, B. García-Almendarez and S. L. Amaya-Llano, *Int. Dairy J.*, 2013, **33**, 191–197.
- 12 R. Kapoor and L. E. Metzger, *Compr. Rev. Food Sci. Food Saf.*, 2008, **7**, 194–214.



- 13 R. Silva, T. C. Pimentel, F. Eustáquio de Matos Junior, E. A. Esmerino, M. Q. Freitas, C. S. Fávaro-Trindade, M. C. Silva and A. G. Cruz, *Food Biosci.*, 2022, **46**, 101517.
- 14 B. Sameer, S. Ganguly, Y. Khetra and L. Sabikhi, *LWT*, 2020, **121**, 108944.
- 15 X. Liu, C. P. Champagne, B. H. Lee, J. I. Boye and M. Casgrain, *Biotechnol. Res. Int.*, 2014, **2014**, 1–21.
- 16 Z. Guo, J. Wang, L. Yan, W. Chen, X. ming Liu and H. ping Zhang, *LWT–Food Sci. Technol.*, 2009, **42**, 1640–1646.
- 17 FSSAI, *FSSAI Annual Report 2020-2021*, New Delhi, 2021.
- 18 IS: SP 18, Part XI, *ISI Handbook of Food Analysis. Part XI. Dairy Products*, Indian Standards Institution, New Delhi, 1981, p. 43.
- 19 Association of Official Analytical Chemists (AOAC), *Official Methods of Analysis of the Association of Analytical Chemists*, 18th edn, AOAC, Gaithersburg, MD, USA, 2005.
- 20 S. Gunasekaran and M. M. Ak, *Cheese Rheology and Texture*, CRC Press, 2002.
- 21 H.-H. Wang and D.-W. Sun, *J. Food Eng.*, 2004, **61**, 47–55.
- 22 N. Shirashoji, J. J. Jaeggi and J. A. Lucey, *J. Dairy Sci.*, 2006, **89**, 15–28.
- 23 A. Giri, S. K. Kanawjia and M. P. Singh, *J. Food Sci. Technol.*, 2017, **54**, 2443–2451.
- 24 Q. G. S. Meira, M. Magnani, F. C. de Medeiros Júnior, R. de, C. R. do E. Queiroga, M. S. Madruga, B. Gullón, A. M. P. Gomes, M. M. E. Pintado and E. L. de Souza, *Food Res. Int.*, 2015, **76**, 828–838.
- 25 C. N. Kuchroo and P. F. Fox, *Milchwissenschaft*, 1982, **37**, 331–335.
- 26 B. Hernández-Ledesma, B. Miralles, L. Amigo, M. Ramos and I. Recio, *J. Sci. Food Agric.*, 2005, **85**, 1041–1048.
- 27 B. Hernández-Ledesma, P. J. Martín-Álvarez and E. Pueyo, *J. Agric. Food Chem.*, 2003, **51**, 4175–4179.
- 28 Q. G. S. Meira, M. Magnani, F. C. de Medeiros Júnior, R. de, C. R. do E. Queiroga, M. S. Madruga, B. Gullón, A. M. P. Gomes, M. M. E. Pintado and E. L. de Souza, *Food Res. Int.*, 2015, **76**, 828–838.
- 29 J. K. Kaushik, A. Kumar, R. K. Duary, A. K. Mohanty, S. Grover and V. K. Batish, *PLoS One*, 2009, **4**, e8099.
- 30 M. De Angelis, R. Di Cagno, C. Huet, C. Crecchio, P. F. Fox and M. Gobbetti, *Appl. Environ. Microbiol.*, 2004, **70**, 1336–1346.
- 31 E. Parente, F. Ciocia, A. Ricciardi, T. Zotta, G. E. Felis and S. Torriani, *Int. J. Food Microbiol.*, 2010, **144**, 270–279.
- 32 M. De Angelis and M. Gobbetti, in *Stress Responses of Lactic Acid Bacteria*, ed. E. Tsakalidou and K. Papadimitriou, Springer US, New York, 2011, pp. 219–249.
- 33 S. Ehsannia and M. R. Sanjabi, *J. Food Sci. Technol.*, 2016, **53**, 996–1003.
- 34 K. Muthukumarappan, Y.-C. Wang and S. Gunasekaran, *J. Dairy Sci.*, 1999, **82**, 1068–1071.
- 35 D. Chaudhary, C. T. Suresh, Y. Khetra, G. S. Meena and S. Hossain, *J. Food Sci. Technol.*, 2023, **60**, 600–608.
- 36 S. Mleko and E. A. Foegeding, *Milchwissenschaft*, 2000, **55**, 513–516.
- 37 I. Laye, T. R. Lindstrom, L. Lowry, F. I. Mei, M. Zwolfer, O. Diaz-Castillo, J. Dolande, S. Havlik and V. Rueda, *U.S. Patent*, 2004126473, 2004.
- 38 S. Mleko and E. A. Foegeding, *Milchwissenschaft*, 2001, **56**, 612–615.
- 39 S. K. G. Purna, A. Pollard and L. E. Metzger, *J. Dairy Sci.*, 2006, **89**, 2386–2396.
- 40 E. S. A. Farahat, A. G. Mohamed, M. M. El-Loly and W. A. M. S. Gafour, *Food Biosci.*, 2021, **42**, 101128.
- 41 H.-P. Bachmann, *Int. Dairy J.*, 2001, **11**, 505–515.
- 42 A. Giri, S. K. Kanawjia and A. Rajoria, *Food Chem.*, 2014, **157**, 240–245.
- 43 A. Giri, S. K. Kanawjia and Y. Khetra, *Food Bioprocess Technol.*, 2014, **7**, 1533–1540.
- 44 F. C. A. Buriti, I. A. Castro and S. M. I. Saad, *Food Chem.*, 2010, **123**, 1190–1197.
- 45 P. Salunke and L. E. Metzger, *Int. Dairy J.*, 2022, **128**, 105324.
- 46 N. S. Joshi, R. P. Jhala, K. Muthukumarappan, M. R. Acharya and V. V. Mistry, *Int. J. Food Prop.*, 2004, **7**, 519–530.
- 47 S. A. M. El-Aidie, A. M. Mabrouk, A. R. Abd-Elgawad and H.-E. M. El-Garhi, *Biocatal. Agric. Biotechnol.*, 2023, **51**, 102798.
- 48 S. E. Chatziantoniou, A. S. Thomareis and M. G. Kontominas, *Eur. Food Res. Technol.*, 2015, **241**, 737–748.
- 49 T. Kahyaoglu and S. Kaya, *Int. Dairy J.*, 2003, **13**, 867–875.
- 50 E. B. Muliawan and S. G. Hatzikiriakos, *Int. Dairy J.*, 2007, **17**, 1063–1072.
- 51 S. K. Lee, M. Huss, H. Klostermeyer and S. G. Anema, *Int. Dairy J.*, 2013, **32**, 79–88.
- 52 J. M. Reparet and Y. Noël, *Lait*, 2003, **83**, 321–333.
- 53 C. Dularia, G. S. Meena, S. Hossain, Y. Khetra and S. Arora, *Food Hydrocoll.*, 2023, **142**, 108842.
- 54 C. A. Brickley, M. A. E. Auty, P. Piraino and P. L. H. McSweeney, *J. Food Sci.*, 2007, **72**, C483–C490.
- 55 J. A. Lucey, M. E. Johnson and D. S. Horne, *J. Dairy Sci.*, 2003, **86**, 2725–2743.
- 56 S. Ehsannia and M. R. Sanjabi, *J. Food Process. Preserv.*, 2016, **40**, 667–674.
- 57 F. C. A. Buriti, J. S. da Rocha, E. G. Assis and S. M. I. Saad, *LWT–Food Sci. Technol.*, 2005, **38**, 173–180.
- 58 W. Engels, J. Siu, S. van Schalkwijk, W. Wesselink, S. Jacobs and H. Bachmann, *Foods*, 2022, **11**, 1005.
- 59 M. P. Dutra, P. C. Palhares, J. R. O. Silva, I. P. Ezequiel, A. L. S. Ramos, J. R. O. Perez and E. M. Ramos, *Small Rumin. Res.*, 2013, **115**, 56–61.
- 60 T. Sanz, A. Salvador, A. Jiménez and S. M. Fiszman, *Eur. Food Res. Technol.*, 2008, **227**, 1515–1521.
- 61 E. Šertović, Z. Sarić, M. Barać, I. Barukčić, A. Kostić and R. Božanić, *Food Technol. Biotechnol.*, 2019, **57**, 461–471.
- 62 L. Ong, A. Henriksson and N. P. Shah, *Int. Dairy J.*, 2007, **17**, 67–78.
- 63 M. Shakerian, S. H. Razavi, S. A. Ziai, F. Khodaiyan, M. S. Yarmand and A. Moayedi, *J. Food Sci. Technol.*, 2015, **52**, 2428–2433.
- 64 O. N. Donkor, A. Henriksson, T. K. Singh, T. Vasiljevic and N. P. Shah, *Int. Dairy J.*, 2007, **17**, 1321–1331.
- 65 T. Virtanen, A. Pihlanto, S. Akkanen and H. Korhonen, *J. Appl. Microbiol.*, 2007, **102**, 106–115.



- 66 A. Gupta, B. Mann, R. Kumar and R. B. Sangwan, *Int. J. Dairy Technol.*, 2009, **62**, 339–347.
- 67 Y. Wang, Y. Wu, Y. Wang, H. Xu, X. Mei, D. Yu, Y. Wang and W. Li, *Nutrients*, 2017, **9**, 521.
- 68 N. Gjorgievski, J. Tomovska, G. Dimitrovska, B. Makarijoski and M. A. Shariati, *J. Hyg. Eng. Des.*, 2014, **8**, 88–92.
- 69 J.-W. Yoon, S.-I. Ahn, J.-W. Jhoo and G.-Y. Kim, *Food Sci. Anim. Resour.*, 2019, **39**, 162–176.
- 70 B. N. P. Sah, T. Vasiljevic, S. McKechnie and O. N. Donkor, *Crit. Rev. Food Sci. Nutr.*, 2018, **58**, 726–740.
- 71 A. Talwalkar, C. W. Miller, K. Kailasapathy and M. H. Nguyen, *Int. J. Food Sci. Technol.*, 2004, **39**, 605–611.
- 72 V. Lovayová, E. Dudriková, K. Rimárová and L. Siegfried, *J. Food Sci. Technol.*, 2015, **52**, 4697–4702.
- 73 A. R. Madureira, M. S. Gião, M. E. Pintado, A. M. P. Gomes, A. C. Freitas and F. X. Malcata, *J. Food Sci.*, 2006, **70**, M160–M165.

