

Cite this: *Dalton Trans.*, 2016, **45**,
1823Received 4th August 2015,
Accepted 29th October 2015

DOI: 10.1039/c5dt02999f

www.rsc.org/dalton

Resolution of P-stereogenic P-heterocycles *via* the formation of diastereomeric molecular and coordination complexes (a review)

Péter Bagi,^{*a} Viktória Ujj,^b Mátyás Czugler,^a Elemér Fogassy^a and György Keglevich^{*a}

TADDOL derivatives and the Ca²⁺-salts of tartaric acid derivatives were found to be versatile and generally applicable resolving agents for the preparation of the enantiomers of P-stereogenic heterocyclic phosphine oxides and phosphinates *via* the formation of the corresponding diastereomeric molecular and coordination complexes. A few of the diastereomeric intermediates were characterized by single crystal X-ray crystallography to gain insights into the binding mode of the corresponding heterocyclic phosphine oxide ("guest") and the resolving agent ("host") and to study the underlying phenomenon of enantiomeric recognition.

1. Introduction

Optically active organophosphorus compounds are of great importance in organic syntheses.¹ An important application is the use of chiral P(III) derivatives (phosphorous, phosphonous and phosphinous esters/amides and especially phosphines) as P-ligands in transition metal complexes which are potential catalysts in enantioselective homogeneous catalytic transform-

ations, such as hydrogenation or hydroformylation.^{2–5} Among chiral organophosphorus compounds, the species bearing P-stereogenic center(s) are of special interest.^{6–10}

P-stereogenic organophosphorus compounds cannot be found in the natural pool of chirality in an optically active form, thus resolution and asymmetric synthesis remain the primary sources for the preparation of such compounds.^{9,11} Despite the wide variety of asymmetric syntheses,^{12–21} the resolution of racemic P-compounds is a well-established method for the preparation of the corresponding optically active organophosphorus compounds.^{9,11,21,22} In the following few paragraphs, selected examples are shown for the preparation of optically active P-stereogenic heterocycles *via* the for-

^aDepartment of Organic Chemistry and Technology, Budapest University of Technology and Economics, 1521 Budapest, Hungary.

E-mail: gkeglevich@mail.bme.hu; Fax: +36 1 4633648; Tel: +36 1 4631111/5883

^bComInnex Inc., Záhony u. 1031 Budapest, Hungary

Péter Bagi

research field includes optical resolution and the development of new catalysts.

Péter Bagi was born in Nyíregyháza (Hungary) in 1986. He graduated at the Budapest University of Technology and Economics in 2011 as a chemical engineer and earned his PhD in 2014. Currently, he is a post-doctoral fellow at the Budapest University of Technology and Economics and, besides educational activity, he has been working on the preparation and application of optically active P-stereogenic compounds. His



György Keglevich

and microwave-assisted reactions, and industrial organophosphorus syntheses. He has published 460 articles, and he serves as an editor for *Curr. Org. Chem.*, *Curr. Org. Synth.*, *Curr. Green Chem.*, *Lett. Org. Chem.* and *Lett. Drug Design Disc.*

György Keglevich was born in 1957 in Budapest, and became a chemical engineer in 1981. He obtained PhD and DSc in 1990 and 1994, respectively, and was appointed to Professor in 1996. He has been the Head of Organic Chemistry and Technology Department, Budapest University of Technology and Economics since 1999. His research interests include organophosphorus and environmentally-friendly ("green") chemistry, catalytic



mation of diastereomers. The systematization of the optical resolution of the acyclic derivatives was not our aim, as this was covered by the review of Pietrusiewicz and Zablocka, as well as by the book of Grabulosa.^{9,11}

One of the first examples for the preparation of optically active P-heterocycles *via* the formation of covalent diastereomers was described by Campbell and Way. They prepared diastereomeric amides by the reaction of an optically active dibenzophosphole oxide bearing a carboxyl functional group with 1-phenylethylamine.²³ Mathey *et al.* reported the synthesis of 1-phosphanorbornadiene-2-carboxaldehydes which were resolved *via* the corresponding diastereomeric acetals formed with (*S,S*)-1,2-diphenylethane-1,2-diol.²⁴

Tang and his co-workers elaborated the resolution of *t*-butyl-hydroxy-benzodihydrooxaphosphole oxide by reacting the racemate with menthyl chloroformate. The separation of the covalent diastereomers followed by hydrolysis afforded the enantiomerically pure *t*-butyl-hydroxy-benzodihydrooxaphosphole oxide.^{25,26} Pietrusiewicz *et al.* reported the oxidative resolution of phenylhexahydrocyclopenta[*b*]phosphole with menthyl bromoacetate. The corresponding enantiopure phenylhexahydrocyclopenta[*b*]phosphole oxide was obtained after purification and hydrolysis of the corresponding diastereomeric phosphonium salt.^{11,27}

The resolution of P-stereogenic compounds may be elaborated *via* the formation of diastereomeric salts if the corresponding organophosphorus compound is a phosphonium salt or it contains an acidic or basic functional group. Ostrogovich and Kerek elaborated the resolution of 1-(4-diethylaminophenyl)-2-phospholene oxide by forming diastereomeric salts with camphorsulfonic acid.²⁸ Kobayashi *et al.* described the optically active borane complex of 1-phenylphospholane-2-carboxylic acid prepared *via* optical resolution by applying quinine.²⁹ For the preparation of optically active phosphonium salts, the silver salt of dibenzoyl-tartaric acid, camphorsulfonic acid or menthoxy-acetic acid finds wide-spread application.

Using these resolving agents, the resolution of a few P-stereogenic phosphonium salts bearing a heterocyclic moiety is elaborated (Fig. 1).^{30–35}

Organophosphorus compounds having neither acidic nor basic functional groups may be resolved *via* the formation of diastereomeric molecular complexes. *O,O'*-Dibenzoyl-(2*R*,3*R*)-tartaric acid was suitable for the preparation of optically active phosphine oxides,^{36–43} among which a few P-heterocyclic and P-stereogenic examples can be found, such as 1-phenyl-2-phospholene oxide,¹¹ DuanPhos oxide⁴⁴ and a bis-benzophosphetane oxide (Fig. 2).⁴⁵

Exploiting the coordination ability of phosphines to transition metals, the palladium complex of phenylethylamine or naphthylethylamine is a generally applicable resolving agent for P-stereogenic phosphines *via* the formation of the corresponding diastereomeric coordination complex. In this manner several mono- or bis-P-heterocyclic phosphines could be separated into their enantiomers (Fig. 3).^{46–54}

Herein, our aim was to summarize the results obtained in the field of the resolution methods developed by our research group for the preparation of optically active five- and six-membered P-heterocyclic phosphine oxides, phosphinates and a few other derivatives (**1–8**) by applying TADDOL derivatives [(–)-**9** and (–)-**10**] or the Ca²⁺-salts of tartaric acid derivatives [(–)-**11** and (–)-**12**] as the resolving agents (Fig. 4).

2. The optical resolution of P-heterocycles

2.1. Optical resolution of five- and six-membered P-heterocycles (**1–8**) with TADDOL derivatives [(–)-**9** and (–)-**10**]

2.1.1. Resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (1a–j**) with TADDOL [(–)-**9**].** First, our aim was to elaborate the enantiomeric separation of 1-aryl- and 1-alkyl-3-



Viktória Ujj

Viktória Ujj was born in Vác (Hungary) in 1983. She graduated as a chemical engineer at the Budapest University of Technology and Economics in 2006, and obtained her PhD with her thesis on the resolution of P-chiral P-heterocycles in 2010. Then, she started working in the field of pharmaceutical research. Since 2012, she has been working as a senior research scientist at ComInnex Inc., focusing on the syntheses of heterocyclic organic compounds.



Mátyás Czugler

Mátyás Czugler was born in Tárnok (Hungary) in 1948, and graduated at the Technical University of Budapest as a chemical engineer in 1972. His research in crystallography involves mainly solid state description of organic supramolecular ensembles, often with biological importance. His DSc was defended in 2006. He led the Crystallography Department (Hungarian Academy) as academic research professor in the period of 2005–2013. In 2007, he was appointed as Adjunct Professor at TUB. His track record shows over 150 publications and experience in crystallizing and solving non-routine crystal structures.






Fig. 1 Resolution of P-heterocyclic phosphonium salts using the silver salt of camphorsulfonic acid.




Fig. 2 Resolution of P-heterocyclic phosphine oxides using O,O'-dibenzoyl-(2R,3R)-tartaric acid.

methyl-3-phospholene 1-oxides (**1a-j**). As the cyclic phosphine oxides (**1a-j**) have neither acidic nor basic functional groups, we anticipated that the resolution methods based on the for-

mation of diastereomeric molecular complexes were worth trying. In the initial screening study of the resolving agents, TADDOL [(−)-**9**] and spiro-TADDOL [(−)-**10**] seemed to be promising for the enantiomeric separation of racemic 3-phospholene oxides **1a-j**. Previously, TADDOL derivatives [(−)-**9** and (−)-**10**] were found to be excellent chiral host molecules in the resolution of racemates containing H-acceptor functional groups. To the best of our knowledge, this was the first case, when the TADDOL derivatives [(−)-**9** and (−)-**10**] were used as the resolving agents to prepare the enantiomers of P-stereogenic phosphine oxides.⁵⁵

It was the first breakthrough, when TADDOL (−)-**9** was applied in the resolution of 1-aryl- and alkyl-3-methyl-3-phospholene 1-oxide (**1a-j**).⁵⁶ According to our resolution method (Scheme 1), the racemic phospholene oxide (**1a-j**) and half equivalent of TADDOL [(−)-**9**] were dissolved in ethyl acetate. Hexane was then added to the solution to promote the precipitation of the corresponding diastereomeric molecular associates. It was found that the maximum resolving capability could be obtained by using the “half equivalent method”.^{57–60} The diastereomeric complexes formed were further purified by two recrystallizations, and the 3-phospholene oxide (**1a-j**) enantiomers were then recovered from the corresponding diastereomers by flash column chromatography (Scheme 1). In all



Elemér Fogassy

Elemér Fogassy was born in 1934 in Budapest, and graduated at the Technical University of Budapest (Hungary) as a Chemical Engineer in 1957. He earned his PhD in 1974, and defended his DSc degree in 1986. He was working for pharmaceutical factories in the period of 1957–64, then he joined the Department of Organic Chemical Technology, TUB, and was appointed as Professor in 1987.

From 2010, he is an Emeritus

Professor. He is the author or co-author of 315 papers/patents in the subject of optical resolution. He received the Academic Patent Award and the Price for Pharmaceutical Research.






Fig. 3 Resolution of P-heterocyclic phosphines using the palladium complex of phenylethylamine or naphthylethylamine.

instances, the enantiomeric purity of the given enantiomeric mixtures was determined by HPLC or GC on chiral stationary phases. The selected results are summarized in Table 1.

Applying TADDOL [(-)-9] as the resolving agent for the enantiomeric separation of 3-phospholene oxides (1), the corresponding enantiomeric excess values fell in the range of 6–97%, whereas the resolving capability values (*S*) were 0.05–0.43 (Table 1).

It was observed that the ethyl derivative (1e) could be resolved with a lower efficiency (*S*) than the propyl derivative (1f) (compare Table 1, entries 5 and 6). It was also found that the enantiomers of 3-phospholene oxides bearing a branched substituent (1g, 1i and 1j) could be prepared with lower resolving capability (*S*) than the derivatives having the normal alkyl chain (1e, 1f and 1h) (compare Table 1, entries 7, 9 and 10 with 5, 6 and 8). On the one hand, these results reveal that an adequate length of the alkyl chain is necessary for the interactions being responsible for the enantiomeric recognition between the host [(-)-9] and the guest molecules (1e–j). On the other hand, the results also show that the increasing steric bulk of the alkyl chain may hinder or weaken the non-bonding interactions between the corresponding 3-phospholene oxide (1e–j) and the TADDOL [(-)-9] molecule that leads to less efficient enantiomeric separation.

In the case of phenyl-3-phospholene oxide (1a), it was observed that kinetic or thermodynamic effects did not affect

significantly the resolving capability values obtained.⁶¹ It was found that (*S*)-aryl-3-phospholene oxides (1a–d) could be prepared with TADDOL [(-)-9] (Table 1, entries 1–4). However, among the alkyl derivatives (1e–j), the TADDOL [(-)-9] resolving agent preferred diastereomeric complex formation with the (*R*)-enantiomers in all but one instance, and the only exception was the butyl-3-phospholene oxide (1h) (Table 1, entries 5–10). The ¹H NMR spectra of the corresponding diastereomers revealed that the ratio of the corresponding 3-phospholene oxide (1a–g and 1i) and TADDOL [(-)-9] was 1 : 1 in most instances (Table 1, entries 1–7 and 9).

2.1.2. Resolution of 1-alkoxy-3-methyl-3-phospholene 1-oxides (4a–d) with TADDOL [(-)-9]. Beside the resolution of five-membered cyclic phosphine oxides (1a–j), TADDOL [(-)-9] was also suitable for the resolution of the corresponding phosphinate analogues, namely alkoxy-3-phospholene oxides (4a–d). The resolution experiments were carried out as described in section 2.1.1 (Scheme 2). The selected results are summarized in Table 2.

The corresponding optically active alkoxy-3-phospholene oxides (4a–d) were prepared with an ee above 99% in three out of four instances (Table 2, entries 2–4). Similarly to the results obtained in the case of the alkyl derivatives (1e–j), it was observed that the length and the bulkiness of the alkoxy chain influenced the overall efficiency of the resolution (*S*). The enantiomeric excess and the resolving capability (*S*) values











	Phosphine oxides or sulfide	Phosphinates or phosphoramidate
5-Membered P-heterocycles	 1	 4
	 2	
5-Membered P-heterocycles	 3	 4
	 5	
6-Membered P-heterocycles	 6	 8
	 7	
Resolving Agents	 (-)-9 or (-)-10	 (-)-11 or (-)-12
	 R¹ Me R² Me	

Fig. 4 The racemic compounds (1–8) and the resolving agents [(–)-9 and (–)-12] included in this review.




Scheme 1 General protocol for the resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (1a–j) with TADDOL [(–)-9].

were lower in the case of the ethoxy derivative (**4a**) than in that of the propoxy derivative (**4b**) (Table 2, entries 1 and 2). The (*R*)-*i*-propoxy-3-methyl-3-phospholene oxide [(*R*)-**4c**] was pre-

pared with lower resolving capability than the propoxy derivative (**4b**) despite the fact that the corresponding enantiomeric excess values were identical (Table 2, entries 2 and 3).



Entry	Y	Diastereomeric complex	ee ^a (%)	Yield ^{a,b} (%)	S ^{a,c} (–)	Abs. config. ^d	Ref.
1	Ph (1a)	(1a)·[(–)- 9]	97	44	0.43	(S)	56, 61
2	2-MeC ₆ H ₄ (1b)	(1b)·[(–)- 9]	57	49	0.28	(S)	61
3	4-MeC ₆ H ₄ (1c)	(1c)·[(–)- 9]	69	42	0.29	(S)	61
4	1-Naphthyl (1d)	(1d)·[(–)- 9]	70	42	0.29	(S)	61
5	Et (1e)	(1e)·[(–)- 9]	24	36	0.09	(R)	61
6	Pr (1f)	(1f)·[(–)- 9]	95	35	0.33	(R)	61
7	i-Pr (1g)	(1g)·[(–)- 9]	6	81	0.05	(R)	62
8 ^e	Bu (1h)	(1h)·[(–)- 9] ₂ ^g	74	23	0.17	(S)	63
9	i-Bu (1i)	(1i)·[(–)- 9]	35	14	0.05	(R)	64
10 ^f	i-Pent (1j)	(1j) ₃ ·[(–)- 9] ₂ ^g	18	38	0.07	(R)	65



rac. **4** + 0.5 eq. (-)-**9** $\xrightarrow[\text{hexane}]{\text{ethyl acetate}}$ $[\mathbf{4}_{\text{prec}}] \cdot [(-)\text{-}\mathbf{9}] \downarrow$ + **4**_{sol}


R = Et (**4a**), Pr (**4b**), *i*-Pr (**4c**),
 (1'*R*,2'*S*,5'*R*)-menthyl (**4d**)

$\downarrow$ column chromat.

4_{prec}

Entry	R	Diastereomeric complex	ee ^a (%)	Yield ^{a,b} (%)	S ^{a,c} (–)	Abs. config. ^d	Ref.
1	Et (4a)	(4a)-[(–)- 9]	44	25	0.11	(S)	61
2	Pr (4b)	(4b)-[(–)- 9]	>99	38	0.38	(R)	67
3	i-Pr (4c)	(4c)-[(–)- 9]	>99	5	0.05	(R)	61
4	Men (4d)	(4d)-[(–)- 9]	>99	25	0.25	(–) ^e	68

In all but one instance, spiro-TADDOL [(–)-**10**] was also found to be an applicable resolving agent for the preparation of optically active aryl- and alkyl-3-phospholene oxides (**1a–i**) with an ee of 25–99%, and resolving capability values of 0.10–0.55 (Table 3, entries 1–9). The i-pentyl-3-phospholene



Scheme 3 General route for the resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a–j**) with spiro-TADDOL [(–)-**10**].

Table 3 Resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a–j**) with 0.5 equiv. of spiro-TADDOL [(–)-**10**] in a mixture of ethyl acetate and hexane

Entry	Y	Diastereomeric complex	ee ^a (%)	Yield ^{a,b} (%)	S ^{a,c} (–)	Abs. config. ^d	Ref.
1	Ph (1a)	(1a)·[(–)- 10]	>99	29	0.29	(S)	56, 61
2	2-MeC ₆ H ₄ (1b)	(1b)·[(–)- 10]	>99	41	0.41	(S)	61
3	4-MeC ₆ H ₄ (1c)	(1c)·[(–)- 10]	>99	30	0.30	(S)	61
4	1-Naphthyl (1d)	(1d)·[(–)- 10]	>99	55	0.55	(S)	61
5	Et (1e)	(1e)·[(–)- 10]	58	45	0.26	(R)	61
6 ^e	Pr (1f)	(1f)·[(–)- 10] ₂ ^f	89	30	0.27	(S)	61
7	i-Pr (1g)	(1g)·[(–)- 10]	64	60	0.39	(R)	62
8 ^e	Bu (1h)	(1h)·[(–)- 10] ₂ ^f	95	52	0.49	(S)	63
9	i-Bu (1i)	(1i)·[(–)- 10]	25	39	0.10	(S)	64
10	i-Pent (1j)	(1j) ₂ ·[(–)- 10]	Rac.	—	—	—	65

^a Enantiomeric excess, yield and resolving capability obtained after recrystallizations. ^b Based on the half of the racemate that is regarded to be 100% for each antipode. ^c Resolving capability, also known as the Fogassy parameter [$S = (\text{Yield}/100) \times (\text{ee}/100)$]. ^d Determined by X-ray crystallography or CD spectroscopy. ^e 1 equiv. of spiro-TADDOL [(–)-**10**] was used. ^f The composition of the diastereomer was different from that of the predominant associate shown in Scheme 3.

oxide (**1j**) was the only five-membered P-heterocycle, where no enantiomeric discrimination could be observed (Table 3, entry 10).

The direct comparison of the results obtained with TADDOL [(–)-**9**] or spiro-TADDOL [(–)-**10**] in a mixture of ethyl acetate and hexane shows that in almost all instances the application of spiro-TADDOL [(–)-**10**] resulted in better enantiomeric separation of the corresponding 3-phospholene oxide (**1a–i**) leading to higher resolving capability values (compare Tables 1 and 3). Considering the resolving capability values, the phenyl- and propyl-3-phospholene oxides (**1a** and **1f**) were the exceptions, where the application of TADDOL [(–)-**9**] was more beneficial than that of spiro-TADDOL [(–)-**10**] (compare Table 1, entry 1 with Table 3, entry 1 and Table 1, entry 6 with Table 3, entry 6).

In all but three instances, the composition of the diastereomers was 1 : 1 [(**1**)·(–)-**10**] similarly to the results obtained with TADDOL [(–)-**9**] (compare Tables 1 and 3). In the case of the propyl-, butyl- and the i-pentyl-3-phospholene oxides (**1f**, **1h**


and **1j**) the composition of the diastereomer was different, as the given complex contained the corresponding 3-phospholene oxide (**1f**, **1h** or **1j**) and spiro-TADDOL [(–)-**10**] in a ratio of 1 : 2 or 2 : 1 (Table 3, entries 6, 8 and 10).

According to our expectation, in most instances both TADDOL derivatives [(–)-**9** or (–)-**10**] preferred diastereomeric complex formation with the same enantiomer of the corresponding aryl- or alkyl-3-phospholene oxide (**1a–e**, **1g**, **1h** and **1j**). However, in the case of the propyl- and i-butyl derivatives (**1f** and **1i**), despite the same absolute configuration and chiral environment of the resolving agents [(–)-**9** and (–)-**10**], different 3-phospholene oxide enantiomers could be prepared with TADDOL [(–)-**9**] or spiro-TADDOL [(–)-**10**] (compare Tables 1 and 3).

In this series of experiments, it was also observed that the sterically demanding alkyl chains on the 3-phospholene moiety (**1i** or **1j**) decreased the overall efficiency of the resolution with TADDOL derivatives [(–)-**9** or (–)-**10**] (Table 3, entries 9 and 10).



2.1.5. Resolution of 1-alkoxy-3-methyl-3-phospholene 1-oxides (4a-d) with spiro-TADDOL [(-)-10]. Besides the aryl- and alkyl-3-phospholene oxides (**1a-j**), their phosphinate analogues (**4a-d**) were also the subject of our systematic resolution studies with spiro-TADDOL [(-)-10]. Similarly to the cases described in sections 2.1.3 and 2.1.4, the resolution experi-



Scheme 4 General method for the resolution of 1-alkoxy-3-methyl-3-phospholene 1-oxides (**4a–d**) with spiro-TADDOL [(-)-**10**].

Table 5 Resolution of 1-alkoxy-3-methyl-3-phospholene 1-oxides (**4a–d**) with 0.5 equiv. of spiro-TADDOL [(-)-**10**] in various solvents

Entry	R	Solvent	Diastereomeric complex	ee ^a (%)	Yield ^{a,b} (%)	S ^{a,c} (–)	Abs. config. ^d	Ref.
1	Et (4a)	EtOAc–hexane	(4a)·[(-)- 10]	95	50	0.48	(<i>R</i>)	61
2	Et (4a)	acetone	(4a)·[(-)- 10] ₂ ^g	58	38	0.22	(<i>R</i>)	70
3	Pr (4b)	EtOAc–hexane	(4b)·[(-)- 10]	93	43	0.40	(<i>R</i>)	67
4 ^e	Pr (4b)	EtOH	(4b)·[(-)- 10] ₂ ^g	93	15	0.14	(<i>S</i>)	67
5	<i>i</i> -Pr (4c)	EtOAc–hexane	(4c)·[(-)- 10]	>99	37	0.37	(<i>R</i>)	61
6	<i>i</i> -Pr (4c)	acetone–hexane	(4c)·[(-)- 10]	>99	56	0.56	(<i>R</i>)	69
7	Men (4d)	EtOAc–hexane	(4d)·[(-)- 10]	90	45	0.41	(–) ^f	68

^a Enantiomeric excess, yield and resolving capability obtained after recrystallizations. ^b Based on the half of the racemate that is regarded to be 100% for each antipode. ^c Resolving capability, also known as the Fogassy parameter [$S = (\text{Yield}/100) \times (\text{ee}/100)$].⁶⁶ ^d Determined by X-ray crystallography or CD spectroscopy. ^e 1 equiv. of spiro-TADDOL [(-)-**10**] was used. ^f The sign of optical rotation. The absolute P-configuration was not determined. ^g The composition of the diastereomer was different from that of the predominant associate shown in Scheme 4.

ments were carried out in a few solvents or solvent mixtures according to the procedure described in section 2.1.3 (Scheme 4). The selected results are summarized in Table 5.

In the case of the alkoxy-3-phospholene oxides (**4a–d**), the enantiomeric excess values were in the range of 58–99%, whereas the corresponding resolving capability values were 0.14–0.56 (Table 5). The direct comparison of the result obtained with either spiro-TADDOL [(-)-**10**] or TADDOL [(-)-**9**] in a mixture of ethyl acetate and hexane revealed that the application of spiro-TADDOL [(-)-**10**] was more advantageous considering the resolving capability values (compare Table 2, entries 1–4 and Table 5, entries 1, 3, 5 and 7). Moreover, the TADDOL derivatives [(-)-**9** or (-)-**10**] formed diastereomeric molecular complexes with the same antipodes of the corresponding alkoxy-3-phospholene oxides (**4b–d**) in all but one instance. The exception was the ethoxy derivative (**4a**), in which case different antipodes of **4a** could be prepared with TADDOL [(-)-**9** or spiro-TADDOL [(-)-**10**] (compare Table 2, entry 1 and Table 5, entry 1) similarly to the case described for the two alkyl-3-phospholene oxides (**1f** and **1i**).


In two out of three cases, the solvents other than the mixture of ethyl acetate and hexane lowered the overall efficiency (*S*) of the resolution of the alkoxy-3-phospholene oxides (**4a** and **4b**) (compare Table 5, entry 1 with 2 and entry 3 with 4). Moreover, in the case of the propoxy derivative (**4b**), the solvent influenced the enantiopreference of spiro-TADDOL [(-)-**10**], and both antipodes of **4b** could be prepared with the

same resolving agent in different solvents (Table 5, entries 3 and 4).


2.1.6. Resolution of a 3-methyl-2-phospholene oxide (2), a 3-methyl-3-phospholene sulfide (3), six-membered P-heterocyclic phosphine oxides (5–7) and a six-membered phosphoramidate (8) with TADDOL derivatives [(-)-9** and (-)-**10**].** After thorough investigation of the resolution of 1-aryl-, 1-alkyl- and 1-alkoxy-3-phospholene oxides (**1** and **4**) with TADDOL and spiro-TADDOL [(-)-**9** and (-)-**10**], we were interested in investigating if these resolution methods were of more general value, so the enantiomeric separation of other five- and six-membered P-heterocyclic phosphine oxides, a phosphine sulfide and a phosphonic ester-amide (**2**, **3** and **5–8**) was also studied. The resolution of the corresponding racemic compound (**2**, **3** and **5–8**) was carried out according to the method described in section 2.1.1. The results are summarized in Schemes 5–10.

The resolution procedure applying spiro-TADDOL [(-)-**10**] as the resolving agent was also suitable for the enantiomeric separation of 1-(4-methylphenyl)-3-methyl-2-phospholene 1-oxide (**2**). The (–)-**2** enantiomer was obtained with an ee of 96% and a resolving capability of 0.35 which is in good agreement with the results obtained in the case of the corresponding 3-phospholene oxide analogue (**1c**) (compare Scheme 5 and Table 3, entry 3). This result suggests that the position of the double-bond in the phospholene moiety does not affect significantly the overall efficiency of resolution with spiro-TADDOL [(-)-**10**].⁷⁰







Scheme 5 The resolution of 1-(4-methylphenyl)-3-methyl-2-phospholene 1-oxide (**2**) with spiro-TADDOL **[(-)-10]**.



Scheme 6 The resolution of 1-phenyl-3-methyl-3-phospholene sulfide (**3**) with spiro-TADDOL **[(-)-10]**.




Scheme 7 The resolution of 6,6-dichloro-1-methyl-3-phenyl-3-phosphabicyclo[3.1.0]hexane 3-oxide (**5**) with spiro-TADDOL **[(-)-10]**.




Scheme 8 The resolution of 4-chloro-1-phenyl-1,2-dihydrophosphinine 1-oxide (**6**) with spiro-TADDOL **[(-)-10]**.





Scheme 9 The resolution of 4-chloro-5-methyl-1-phenyl-1,2,3,6-tetrahydrophosphinine 1-oxide (7) with TADDOL [(-)-9].



Scheme 10 The resolution of 6-diethylamino-dibenzo[c,e][5,6]oxaphosphorine 6-oxide (8) with spiro-TADDOL [(-)-10].

The (+)-1-phenyl-3-methyl-3-phospholene sulfide [(+)-3] was also prepared by resolution with spiro-TADDOL [(-)-10], the ee was above 99% in this case (Scheme 6). The crude product of the phospholene sulfide (3) was used as the starting material in this resolution experiment. Interestingly, less efficient enantioseparation of phosphine sulfide 3 was achieved, when purified phenyl-3-phospholene sulfide (3) was used as the racemic starting material.⁶¹

The wide applicability of TADDOL derivatives [(-)-9 and (-)-10] for the preparation of optically active five-membered phosphine oxides (1 and 2), a sulfide (3) and phosphinates (4) motivated us to extend this method to six-membered derivatives as well, such as phosphine oxides (5–7) and a phosphonic ester-amide (8). Among the six-membered P-heterocyclic phosphine oxides, the (+)-6,6-dichloro-1-methyl-3-phenyl-3-phosphabicyclo[3.1.0]hexane 3-oxide [(+)-5] bearing the dichlorocyclopropane ring and the P=O function in the position *trans* disposition could be prepared with an ee of 55% using spiro-TADDOL [(-)-10] as the resolving agent (Scheme 7).⁷¹


Racemic 4-chloro-1-phenyl-1,2-dihydrophosphinine 1-oxide (6) comprised a 3 : 1 mixture of two double-bond isomers (6A and 6B) that could not be separated. Therefore, the mixture of dihydrophosphinine oxides 6A and 6B was resolved using spiro-TADDOL [(-)-10] in a mixture of acetone–hexane. The ratio of the double-bond isomers (6A and 6B) did not change significantly during the resolution procedure. However, the

enantiomeric excess of 6A and 6B was slightly different (89% and 95%, respectively) (Scheme 8).⁷²

TADDOL [(-)-9] was also suitable for the partial resolution of 4-chloro-5-methyl-1-phenyl-1,2,3,6-tetrahydrophosphinine 1-oxide (7), whereupon (S)-7 was obtained with an ee of 65% (Scheme 9).⁷¹ Recently, X-ray crystallographic and CD spectroscopic studies have allowed us to assign (S) absolute configuration to the (-)-4-chloro-5-methyl-1-phenyl-1,2,3,6-tetrahydrophosphinine 1-oxide enantiomer [(-)-7].⁷³

A six-membered phosphonic ester-amide, the 6-diethylamino-dibenzo[c,e][5,6]oxaphosphorine 6-oxide (8) was partially resolved with spiro-TADDOL [(-)-10], and as a result, (-)-8 was obtained with an ee of 86% (Scheme 10).⁷⁰


Besides the systematic resolution experiments of the six-membered heterocyclic phosphine oxides (5–7) with TADDOL derivatives (-)-9 or (-)-10, we investigated the stereochemical outcome of the 3-methyl-1-phenyl-3-phospholene 1-oxide (1a) → 6,6-dichloro-1-methyl-3-phenyl-3-phosphabicyclo[3.1.0]hexane 3-oxide (5) → 4-chloro-1-phenyl-1,2-dihydrophosphinine 1-oxide (6) → 4-chloro-5-methyl-1-phenyl-1,2,3,6-tetrahydrophosphinine 1-oxide (7) reaction sequence described earlier by our research group.^{74,75} It was proven that the dichlorocarbene addition on the double-bond of phenyl-3-phospholene oxide (1a) involved racemization on the P-stereogenic center. However, the thermolytic ring opening of 3-phosphabicyclo[3.1.0]hexane oxide (5), and the selective reduction of the α,β-double bond of the 4-chloro-3-methyl-1-





Upon applying $\text{Ca}(\text{H-DBTA})_2$ [(–)-**11**] as the resolving agent, the corresponding enantiomeric excess values of the aryl- and alkyl-3-phospholene oxides (**1a–j**) fell in the range of 13–99%, whereas the resolving capability values (*S*) were 0.02–0.50 (Table 6). In 5 out of 10 cases, the 3-phospholene oxides (**1a**, **1b**, **1d**, **1f** and **1j**) could be prepared with an ee above 90% (Table 6, entries 1, 2, 4, 6 and 11). It is noteworthy that the resolving capability values (*S*) obtained in the instance of the aryl-3-phospholene oxides (**1a–d**) (*S* = 0.12–0.50) were significantly higher than in the case of the corresponding alkyl derivatives (**1e–j**) (*S* = 0.02–0.17) (compare Table 6, entries 1–4

To overcome these disadvantages deriving from the application of TADDOL derivatives (–)-**9** or (–)-**10**, we attempted the application of cheap and readily available tartaric acid and its derivatives. Unfortunately, none of the tartaric acid derivatives tried out could form a diastereomeric molecular complex with 3-phospholene oxides (**1** or **4**). Then, we focused on the acidic calcium salts of *O,O'*-dibenzoyl- and *O,O'*-di-*p*-toluoyl-(2*R*,3*R*)-tartaric acids [Ca(H-DBTA)₂ (–)-**11** and Ca(H-DPTTA)₂ (–)-**12**,



Scheme 12 General route for the resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a–j**) with $\text{Ca}(\text{H-DBTA})_2$ [(–)-**11**].

Table 6 Resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a–j**) with 0.25 equiv. of Ca(H-DBTA)₂ [(–)-**11**]

Entry	Y	Solvent	Diastereomeric complex	ee ^a (%)	Yield ^{a,b} (%)	S ^{a,c} (–)	Abs. config.	Ref.
1	Ph (1a)	EtOAc–EtOH ^d EtOH–water ^e	(1a) ₂ ·(–)- 11 ^g	96	52	0.50	(<i>R</i>)	80
2	2-MeC ₆ H ₄ (1b)	MeCN–EtOH ^{d,e}	(1b) ₂ ·(–)- 11	93	33	0.31	(<i>R</i>)	80
3	4-MeC ₆ H ₄ (1c)	MeCN–EtOH ^{d,e}	(1c) ₂ ·(–)- 11	44	28	0.12	(<i>S</i>)	80
4	1-Naphthyl (1d)	EtOAc–EtOH ^d EtOH–water ^e	(1d) ₂ ·(–)- 11	99	42	0.42	(<i>R</i>)	80
5	Et (1e)	EtOAc–EtOH ^{d,e}	(1e) ₂ ·(–)- 11	49	25	0.12	(<i>R</i>)	80
6 ^f	Pr (1f)	EtOH–water ^{d,e}	(1f) ₂ ·(–)- 11 ^{g,h}	96	18	0.17	(<i>S</i>)	80
7	i-Pr (1g)	EtOH ^{d,e}	(1g) ₂ ·(–)- 11	43	8	0.03	(<i>R</i>)	62
8	Bu (1h)	MeCN–EtOH ^{d,e}	(1h) ₂ ·(–)- 11	76	21	0.16	(<i>S</i>)	63
9	Bu (1h)	EtOAc–EtOH ^{d,e}	(1h) ₂ ·(–)- 11	13	14	0.02	(<i>R</i>)	63
10	i-Bu (1i)	EtOAc–EtOH ^{d,e}	(1i) ₂ ·(–)- 11	29	7	0.02	(<i>R</i>)	64
11	i-Pent (1j)	MeCN–EtOH ^{d,e}	(1j) ₂ ·(–)- 11	99	17	0.17	(<i>S</i>)	65

^a Enantiomeric excess, yield and resolving capability obtained after digestions. ^b Based on the half of the racemate that is regarded to be 100% for each antipode. ^c Resolving capability, also known as the Fogassy parameter [$S = (\text{Yield}/100) \times (\text{ee}/100)$].⁶⁶ ^d Solvent used for crystallization. ^e Solvent used for digestions. ^f 0.5 equiv. of $\text{Ca}(\text{H-DBTA})_2$ **[(-)-11]** was used. ^g Water was incorporated into the diastereomer. ^h The composition of the diastereomer was different from that of the predominant associate shown in Scheme 12.

and 5–11). The underlying reason for this difference may be that the strength and the nature of the non-bonding interactions between the resolving agent [(–)-**11**] and the given 3-phospholene oxide are presumably different in the case of the aryl- and alkyl derivatives (**1a–d** or **1e–j**).

Generally, the enantiomeric excess values obtained with the acidic Ca^{2+} -salt of the *O,O'*-dibenzoyl-(2*R*,3*R*)-tartaric acid [(–)-**11**] were comparable with the ee values obtained with TADDOL derivatives [(–)-**9** or (–)-**10**]. However, $\text{Ca}(\text{H-DBTA})_2$ [(–)-**11**] underperformed TADDOL and spiro-TADDOL [(–)-**9** and (–)-**10**] in terms of the resolving capability values (*S*) (compare Table 6 with Tables 1, 3 and 4).


In this series of resolution experiments, the solvent did not influence in which an antipode of the 3-phospholene oxide (**1a-g**, **1i** and **1j**) was incorporated into the diastereomer. The only exception was the 1-butyl-3-phospholene oxide (**1h**) (Table 6, entries 8 and 9). However, the significance of this solvent dependent enantioselection was not as pronounced, as in the case of spiro-TADDOL [(–)-**10**] (Tables 3 and 4).

The ^1H NMR studies revealed that the predominant composition of the diastereomers may be described by the formula $\text{Ca}(\mathbf{1})_2(\text{H-BDTA})_2$ (Table 6). The only exception was the propyl-3-phospholene oxide (**1f**), where one 3-phospholene oxide

molecule was incorporated into the diastereomer (Table 6, entry 6).

2.2.2. Resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (1a-j) with calcium hydrogen *O,O'*-di-*p*-toluoyl-(2*R*,3*R*)-tartrate [(–)-12]. Beside the acidic calcium salt of *O,O'*-dibenzoyl-(2*R*,3*R*)-tartaric acid [(–)-11], the corresponding *O,O'*-di-*p*-toluoyl derivative [(–)-12] was also applied to prepare optically active 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a-j**) according to the procedure described in section 2.2.1 (Scheme 13). The selected results are summarized in Table 7.

Regarding the aryl-3-phospholene oxides (**1a-d**), lower resolving capability values (*S*) could be obtained with $\text{Ca}(\text{H-DPTTA})_2$ [(-)-**12**] than with $\text{Ca}(\text{H-DBTA})_2$ [(-)-**11**] (compare Table 6, entries 2–4 and Table 7, entries 2–4). The only exception was the 1-phenyl-3-phospholene oxide (**1a**), in which case the results showed parity (compare Table 6, entry 1 and Table 7, entry 1). Interestingly, this tendency reversed for the series of the alkyl-3-phospholene oxides (**1e-j**), where, in 5 out of 6 instances, the application of $\text{Ca}(\text{H-DPTTA})_2$ [(-)-**12**] was more advantageous (compare Table 6, entries 5 and 7–11 with Table 7, entries 5 and 7–10). Both $\text{Ca}(\text{H-DBTA})_2$ and $\text{Ca}(\text{H-DPTTA})_2$ [(-)-**11** and (-)-**12**] were equally applicable for



Scheme 13 General protocol for the resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a–j**) with Ca(H-DPTTA)₂ [(–)-**12**].

Table 7 Resolution of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a–j**) with 0.25 equiv. of Ca(H-DPTTA)₂ [(–)-**12**]

Entry	Y	Solvent	Diastereomeric complex	ee ^a (%)	Yield ^{a,b} (%)	S ^{a,c} (–)	Abs. config.	Ref.
1 ^d	Ph (1a)	EtOH–water	(1a) ₂ ·(–)- 12 ^{e,f}	93	55	0.51	(S)	80
2	2-MeC ₆ H ₄ (1b)	EtOH–water	(1b) ₂ ·(–)- 12	45	13	0.06	(R)	80
3	4-MeC ₆ H ₄ (1c)	EtOH–water	(1c) ₂ ·(–)- 12	32	15	0.05	(S)	80
4	1-Naphthyl (1d)	EtOH–water	(1d) ₂ ·(–)- 12 ^e	69	29	0.20	(R)	80
5	Et (1e)	EtOH–water	(1e) ₂ ·(–)- 12 ^e	73	34	0.25	(S)	80
6	Pr (1f)	EtOH–water	(1f) ₂ ·(–)- 12	41	42	0.17	(S)	80
7	<i>i</i> -Pr (1g)	EtOAc–EtOH–water	(1g) ₂ ·(–)- 12	92	10	0.09	(S)	62
8	Bu (1h)	EtOH–water	(1h) ₂ ·(–)- 12	77	27	0.21	(S)	63
9	<i>i</i> -Bu (1i)	EtOAc–EtOH–water	(1i) ₂ ·(–)- 12	73	23	0.17	(R)	64
10	<i>i</i> -Pent (1j)	EtOH–water	(1j) ₄ ·[(–)- 12] ₅ ^f	95	32	0.30	(S)	81

^a Enantiomeric excess, yield and resolving capability obtained after digestions. ^b Based on the half of the racemate that is regarded to be 100% for each antipode. ^c Resolving capability, also known as the Fogassy parameter [$S = (\text{Yield}/100) \times (\text{ee}/100)$]. ^d 0.5 equiv. of Ca(H-DPTTA)₂ [(–)-**12**] was used. ^e Water was incorporated into the diastereomer. ^f The composition of the diastereomer was different from that of the predominant associate shown in Scheme 13.

the enantiomeric separation of 1-propyl-3-phospholene oxide (**1f**) in terms of resolving capability (S), but a higher enantiomeric excess could be obtained with Ca(H-DBTA)₂ [(–)-**11**] (compare Table 6, entry 6 and Table 7, entry 6). In this series of experiments, the formation of diastereomers with a composition of Ca(**1**)₂(H-DPTTA)₂ was observed in most of the cases (Table 7) similarly to the instances described for the use of Ca(H-DBTA)₂ [(–)-**11**] (Table 6).

The impact of the similar chiral environment of the *O,O'*-dibenzoyl- or *O,O'*-di-*p*-toluoyl-(2*R*,3*R*)-tartaric acid was that the resolving agents (–)-**11** and (–)-**12** preferred diastereomeric complex formation with the same enantiomers of a given 3-phospholene oxide (**1b–d**, **1f** and **1h–1j**) in the majority of the instances (compare Table 6, entries 2–4, 6 and 8–11 with Table 7, entries 2–4, 6 and 8–10). However, in the case of the phenyl-, ethyl or *i*-propyl-derivatives (**1a**, **1e** and **1g**) different antipodes could be prepared with Ca(H-DBTA)₂ and Ca(H-DPTTA)₂ [(–)-**11** or (–)-**12**] (compare Table 6, entries 1, 5 and 7 with Table 7, entries 1, 5 and 7). It is worth mentioning that no solvent dependent enantioselectivity could be observed, when Ca(H-DPTTA)₂ [(–)-**12**] was used as the resolving agent.


2.2.3. Resolution of 1-alkoxy-3-methyl-3-phospholene 1-oxides (4a–c) with calcium hydrogen *O,O'*-dibenzoyl- and di-*p*-toluoyl-(2*R*,3*R*)-tartrate [(–)-11** and (–)-**12**].** After the resolu-

tion of 1-aryl- and 1-alkyl-3-methyl-3-phospholene 1-oxides (**1a–j**), we extended our resolution methods by applying Ca(H-DBTA)₂ (–)-**11** or Ca(H-DPTTA)₂ (–)-**12** for the enantiomeric separation of the corresponding phosphinate analogues, 1-alkoxy-3-methyl-3-phospholene 1-oxides (**4a–c**). The resolutions were carried out as described in section 2.2.1 (Scheme 14). The selected best results are summarised in Table 8.

When Ca(H-DBTA)₂ [(–)-**11**] was the resolving agent, the alkoxy-3-phospholene oxides (**4a–c**) were prepared with an ee of 17–92%, whereas the corresponding resolving capability values fell in the range of 0.06–0.33 (Table 8, entries 1–4). By applying Ca(H-DPTTA)₂ [(–)-**12**], the (*R*)-enantiomer of ethoxy- and propoxy-3-phospholene oxide could be prepared with an ee of 75% and 55%, respectively, and with a resolving capability value of 0.33 and 0.20, respectively (Table 8, entries 5 and 6). It is noteworthy that Ca(H-DPTTA)₂ [(–)-**12**] could not differentiate between the two antipodes of *i*-propoxy-3-phospholene oxide (**4c**) (Table 8, entry 7).

The results in Table 8 indicate that the longer alkoxy chain may influence the non-bonding interactions with the resolving agent in a negative manner (compare Table 8, entry 1 with 2 and 3 and entry 5 with 6).

The 1-propoxy-3-phospholene oxide (**4b**) was the only alkoxy derivative whose (*R*)- and (*S*)-enantiomers could be prepared



Scheme 14 General method for the resolution of 1-alkoxy-3-methyl-3-phospholene 1-oxides (**4a–c**) with $\text{Ca}(\text{H-DBTA})_2$ or $\text{Ca}(\text{H-DPTTA})_2$ [(–)-**11** or (–)-**12**].

Table 8 Resolution of 1-alkoxy-3-methyl-3-phospholene 1-oxides (**4a–c**) with 0.25 equiv. of Ca(H-DBTA)₂ or Ca(H-DPTTA)₂ [(–)-**11** or (–)-**12**]

Entry	R	Resolving agent	Solvent	Diastereomeric complex	ee ^a (%)	Yield ^{a,b} (%)	S ^{a,c} (–)	Abs. config.	Ref.
1	Et (4a)	(–)- 11	MeCN–EtOH	(4a) ₂ ·(–)- 11	91	29	0.26	(<i>R</i>)	80
2	Pr (4b)	(–)- 11	MeCN–EtOH	(4b) ₂ ·(–)- 11	59	18	0.11	(<i>R</i>)	67
3	Pr (4b)	(–)- 11	EtOAc–EtOH	(4b) ₂ ·(–)- 11	17	34	0.06	(<i>S</i>)	67
4	i-Pr (4c)	(–)- 11	MeCN–EtOH	(4c) ₂ ·(–)- 11	92	36	0.33	(<i>S</i>)	80
5	Et (4a)	(–)- 12	EtOH–water	(4a) ₂ ·(–)- 12	75	44	0.33	(<i>R</i>)	80
6	Pr (4b)	(–)- 12	EtOAc–EtOH–water	(4b) ₂ ·(–)- 12	55	36	0.20	(<i>R</i>)	67
7	i-Pr (4c)	(–)- 12	EtOH–water	(4c) ₂ ·(–)- 12	Rac.	—	—	—	80

^a Enantiomeric excess, yield and resolving capability obtained after digestions. ^b Based on the half of the racemate that is regarded to be 100% for each antipode. ^c Resolving capability, also known as the Fogassy parameter [$S = (\text{Yield}/100) \times (\text{ee}/100)$].⁶⁶

with the same resolving agent, $\text{Ca}(\text{H-DBTA})_2$ [(**-**)-**11**] using different solvents (ee for (*R*)-**4b**: 59% and ee for (*S*)-**4b**: 17%) (Table 8, entries 2 and 3).


The diastereomers formed on resolution contained the corresponding 1-alkoxy-3-phospholene oxides (**4a-c**) and $\text{Ca}(\text{H-DBTA})_2$ [(**-**)-**11**] or $\text{Ca}(\text{H-DPTTA})_2$ [(**-**)-**12**] in a ratio of 2 : 1 similarly to the cases described for the resolution of most of the aryl- and alkyl-3-phospholene oxides (compare Table 8 with Tables 6 and 7).

2.2.4. Resolution of six-membered P-heterocyclic phosphine oxides and a phosphonic ester-amide (5, 7 and 8) with calcium hydrogen *O,O'*-dibenzoyl- and di-*p*-toluoyl-(2*R*,3*R*)-tartrate [(–)-11 and (–)-12]. The success with the acidic calcium salts of *O,O'*-dibenzoyl- or *O,O'*-di-*p*-toluoyl-(2*R*,3*R*)-tartaric acid [(–)-11 or (–)-12] in the optical resolution of five-membered phosphine oxides and phosphinates (**1a–j** and **4a–c**) encouraged us to try the enantiomeric separation of six-membered phosphine


oxides and a phosphonic ester-amide. The 6,6-dichloro-1-methyl-3-phenyl-3-phosphabicyclo[3.1.0]hexane 3-oxide (5), 4-chloro-5-methyl-1-phenyl-1,2,3,6-tetrahydrophosphinine 1-oxide (7) and 6-diethylamino-dibenzo[*c,e*][5,6]oxaphosphorine 6-oxide (8) served as model compounds, and the resolutions were carried out as described in section 2.2.1. The results are summarized in Schemes 15–17.

Both $\text{Ca}(\text{H-DBTA})_2$ [(−)-**11**] and $\text{Ca}(\text{H-DPTTA})_2$ [(−)-**12**] were applicable for the partial enantiomeric separation of phenyl-phosphabicyclo[3.1.0]hexane oxide (**5**). Interestingly, resolving agents (−)-**11** or (−)-**12** formed diastereomeric coordination complexes with the different enantiomers of phosphabicyclo[3.1.0]hexane oxide **5**. In this manner, (−)-**5** and (+)-**5** could be prepared with an ee of 67% and 22%, respectively (Scheme 15).⁷¹


Using $\text{Ca}(\text{H-DPTTA})_2$ [(-)-12] as the resolving agent, enantiopure (*R*)-1-phenyl-1,2,3,6-tetrahydrophosphinine 1-oxide



Scheme 15 The resolution of 6,6-dichloro-1-methyl-3-phenyl-3-phosphabicyclo[3.1.0]hexane 3-oxide (5) with $\text{Ca}(\text{H-DBTA})_2$ or $\text{Ca}(\text{H-DPTTA})_2$ [(*-*)-11 or (*-*)-12].



Scheme 16 The resolution of 4-chloro-5-methyl-1-phenyl-1,2,3,6-tetrahydrophosphinine 1-oxide (**7**) with Ca(H-DPTTA)₂ [(-)-**12**].



Scheme 17 The resolution of 6-diethylamino-dibenzo[*c,e*][5,6]oxaphosphorine 6-oxide (**8**) with Ca(H-DBTA)₂ or Ca(H-DPTTA)₂ [(-)-**11** or (-)-**12**].

[(**R**)-**7**] could be prepared (Scheme 16). It is noted that no crystalline diastereomers were formed, when Ca(H-DBTA)₂ [(-)-**11**] was used.⁷⁹

The acidic calcium salts *O,O'*-dibenzoyl- or *O,O'*-di-*p*-toluoyl-(2*R*,3*R*)-tartaric acid [(-)-**11** or (-)-**12**] were also suitable resolving agents for the preparation of the (-)-6-diethylamino-dibenzo[*c,e*][5,6]oxaphosphorine 6-oxide [(-)-**8**]. Among all of the resolving agents [(-)-**9**–(-)-**12**] used for the enantioseparation of dibenzooxaphosphorine oxide **8**, the highest ee and resolving capability values were obtained with Ca(H-DBTA)₂ [(-)-**11**] (Scheme 17).⁷⁰

2.3. X-Ray analysis of the diastereomeric complexes

The crystallization process plays an important role in these investigations not only by forming well defined solid precipitates, but also through the manipulation possibilities through solubility and insolubility properties, as well as affecting *e.g.* the dielectric properties by a designed engineering approach (fine tuning by crystal engineering). Thus, crystal structure determinations^{56–64,67,73,79} of these complex associate molecular systems may be of use in responding to queries regarding the structural background of the physical processes taking place during the experiments.

Crystallographic data shed light not only on the solvent^{56,64} or co-solvent by identifying not only the presence of these species but also their role in the structure of the respective crystals. An interesting example was provided by the crystal structure of the quaternary (*R*)-*i*-butyl-3-phospholene oxide [(**R**)-**1i**]-spiro-TADDOL [(-)-**10**]-isopropanol 1 : 2 : 1 molecular assembly.⁶⁴ In this case, a molecule of the applied solvent was also included in the crystal. The competitive binding of the

solvent to the two host molecules and also to the (*R*)-*i*-butyl-3-phospholene oxide [(**R**)-**1i**] resolution target could be traced back to the slightly stronger binding of the solvent than that of the respective (*R*)-*i*-butyl-3-phospholene oxide guest [(**R**)-**1i**].

In another neutral complex case, the key feature of the resolution process was the formation of the 2 : 1 associate of spiro-TADDOL [(-)-**10**] with the (*S*)-(-)-1-isopropyl-3-methyl-3-phospholene oxide [(*S*)-**1g**] guest.⁶⁵ Apart from the well-defined





Fig. 5 The two largest attractive interactions (in kJ mol⁻¹) indicated also by red broken lines between the respective molecular centres in the crystal of the (*S*)-(-)-1-isopropyl-3-methyl-3-phospholene oxide guest [(*S*)-**1g**] with two independent spiro-TADDOL [(-)-**10**] hosts. Arrows indicate the donor–acceptor directions of the H-bond vectors.





* Tentative assignment of the absolute P-configuration.

Fig. 6 The best results for the resolution of five- and six-membered P-heterocyclic phosphine oxides, phosphinates, a phosphine sulfide and a phosphonic ester-amide (1–8).



As it turned out clearly from this series of crystal structure determinations,^{56-64,67,73,79} the obvious and common binding of the P-oxides is realized through the P=O oxygen atom. As it is described, the binding is primarily established through H-bonding in the neutral type assemblies incorporating TADDOL derivatives [(-)-**9** and (-)-**10**]. The acidic Ca salt matrices involved the binding of the P=O function into the sixfold coordination sphere of the metal ion. These observations may be explained by the enhanced negative charge on the P=O oxygen atom.

In this paper, our versatile resolution methods applying TADDOL derivatives [(–)-**9** and (–)-**10**] or the Ca²⁺-salts of tartaric acid derivatives [(–)-**11** and (–)-**12**] are summarized, and the methods are widely applicable for the preparation of five- and six-membered P-heterocyclic phosphine oxides, phosphinates, a phosphine sulfide and a phosphonic ester-amide (**1–8**) in an optically active form (Fig. 6). The corresponding enantiomers (**1–8**) were obtained with an ee above 95% in the majority of the instances. The P-heterocycles (**1–8**) discussed are unknown in the literature in an optically active form. The absolute configuration of the P-stereogenic centers was also determined in most of the instances. In a few cases, diastereomeric molecular or coordination complexes were characterized by X-ray crystallography and the characteristic non-bonding interactions between the host and the molecules were identified. The optically active P-heterocyclic phosphine oxides are precursors of the corresponding P(III)-compounds which could be applied as ligands in platinum complexes (**14**).^{62,81–85}

- 1 L. D. Quin, *A guide to organophosphorus chemistry*, John Wiley & Sons, New York, 2000.
- 2 T. Imamoto, in *Handbook of Organophosphorus Chemistry*, ed. R. Engel, Marcel Dekker, New York, 1992, ch. 1, pp. 1–54.
- 3 R. Noyori, *Asymmetric Catalysis in Organic Synthesis*, John Wiley & Sons, New York, 1994.
- 4 *Phosphorus Ligands in Asymmetric Catalysis*, ed. A. Börner, Wiley-VCH, Weinheim, 2008.
- 5 *Catalytic Asymmetric Synthesis*, ed. I. Ojima, Wiley, Weinheim, 2010.
- 6 H. Brunner and W. Zettlmeier, *Handbook of enantioselective catalysis with transition metal compounds*, VCH, Weinheim, 1993.
- 7 *Comprehensive Asymmetric Catalysis*, ed. E. N. Jacobsen, A. Pfaltz and H. Yamamoto, Springer-Verlag, Heidelberg, 2000.
- 8 W. Tang and X. Zhang, *Chem. Rev.*, 2003, **103**, 3029–3070.
- 9 A. Grabulosa, *P-Stereogenic Ligands in Enantioselective Catalysis*, The Royal Society of Chemistry, Cambridge, 2010.
- 10 S. Lühr, J. Holz and A. Börner, *ChemCatChem*, 2011, **3**, 1708–1730.
- 11 K. M. Pietrusiewicz and M. Zablocka, *Chem. Rev.*, 1994, **94**, 1375–1411.
- 12 O. Korpiun and K. Mislow, *J. Am. Chem. Soc.*, 1967, **89**, 4784–4786.
- 13 S. Juge, M. Stephan, J. A. Laffitte and J. P. Genet, *Tetrahedron Lett.*, 1990, **31**, 6357–6360.
- 14 J. V. Carey, M. D. Barker, J. M. Brown and M. J. H. Russell, *J. Chem. Soc., Perkin Trans. 1*, 1993, 831–839.
- 15 E. J. Corey, Z. Chen and G. J. Tanoury, *J. Am. Chem. Soc.*, 1993, **115**, 11000–11001.
- 16 M. J. Johansson and N. Kann, *Mini-Rev. Org. Chem.*, 2004, **1**, 233–247.
- 17 A. Grabulosa, J. Granell and G. Muller, *Coord. Chem. Rev.*, 2007, **251**, 25–90.
- 18 D. S. Glueck, *Synlett*, 2007, 2627–2634.
- 19 O. I. Kolodiazhnyi, *Tetrahedron: Asymmetry*, 2012, **23**, 1–46.
- 20 O. I. Kolodiazhnyi, V. P. Kukhar and A. O. Kolodiazhna, *Tetrahedron: Asymmetry*, 2014, **25**, 865–922.
- 21 E. Bergin, C. T. O'Connor, S. B. Robinson, E. M. McGarrigle, C. P. O'Mahony and D. G. Gilheany, *J. Am. Chem. Soc.*, 2007, **129**, 9566–9567.
- 22 K. Nikitin, K. V. Rajendran, H. Müller-Bunz and D. G. Gilheany, *Angew. Chem., Int. Ed.*, 2014, **53**, 1906–1909.
- 23 I. G. M. Campbell and J. K. Way, *J. Chem. Soc.*, 1961, 2133–2141.

- 69 V. Ujj, T. Szuhánszki, J. Schinder, M. Czugler, E. Fogassy and G. Keglevich, *Hung. J. Ind. Chem.*, 2008, **36**, 131–136.
- 70 V. Ujj, Ph.D Thesis, 2010.
- 71 P. Bagi, A. Laki and G. Keglevich, *Heteroatom Chem.*, 2013, **24**, 179–186.
- 72 V. Ujj, A. Kerényi, A. Laki, E. Fogassy and G. Keglevich, *Lett. Org. Chem.*, 2010, **7**, 110–113.
- 73 P. Bagi, K. Karaghiosoff, M. Czugler, D. Hessz, M. Kállay, M. Kubinyi, T. Szilvási, P. Pongrácz, L. Kollár, I. Timári, K. E. Kövér, L. Drahos, E. Fogassy and G. Keglevich, *Heteroatom Chem.*, 2015, submitted.
- 74 G. Keglevich, *Synthesis*, 1993, 931–942.
- 75 G. Keglevich, *Curr. Org. Chem.*, 2006, **10**, 93–111.
- 76 A. Mravik, Z. Böcskei, Z. Katona, I. Markovits and E. Fogassy, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 1534–1536.
- 77 A. Mravik, Z. Böcskei, K. Simon, F. Elekes and Z. Izsaki, *Chem. – Eur. J.*, 1998, **4**, 1621–1627.
- 78 A. Mravik, Z. Böcskei, Z. Katona, I. Markovits, G. Pokol, D. K. Menyhárd and E. Fogassy, *J. Chem. Soc., Chem. Commun.*, 1996, 1983–1984.
- 79 V. Ujj, J. Schindler, T. Novák, M. Czugler, E. Fogassy and G. Keglevich, *Tetrahedron: Asymmetry*, 2008, **19**, 1973–1977.
- 80 V. Ujj, P. Bagi, J. Schindler, J. Madarász, E. Fogassy and G. Keglevich, *Chirality*, 2010, **22**, 699–705.
- 81 P. Bagi, T. Kovács, T. Szilvási, P. Pongrácz, L. Kollár, L. Drahos, E. Fogassy and G. Keglevich, *J. Organomet. Chem.*, 2014, **751**, 306–313.
- 82 A. Kerényi, V. Kovács, T. Körtvélyesi, K. Ludányi, L. Drahos and G. Keglevich, *Heteroatom Chem.*, 2010, **21**, 63–70.
- 83 P. Pongrácz, L. Kollár, A. Kerényi, V. Kovács, V. Ujj and G. Keglevich, *J. Organomet. Chem.*, 2011, **696**, 2234–2237.
- 84 G. Keglevich, P. Bagi, Á. Szöllősy, T. Körtvélyesi, P. Pongrácz, L. Kollár and L. Drahos, *J. Organomet. Chem.*, 2011, **696**, 3557–3563.
- 85 P. Bagi, T. Szilvási, P. Pongrácz, L. Kollár, L. Drahos and G. Keglevich, *Curr. Org. Chem.*, 2014, **18**, 1529–1538.

