

CrossMark
click for updatesCite this: *Chem. Sci.*, 2016, 7, 1245

Targeting antioxidant pathways with ferrocenylated N-heterocyclic carbene supported gold(i) complexes in A549 lung cancer cells†

J. F. Arambula,^{*a} R. McCall,^a K. J. Sidoran,^b D. Magda,^c N. A. Mitchell,^d C. W. Bielawski,^{ef} V. M. Lynch,^g J. L. Sessler^g and K. Arumugam^{*h}

Ferrocene containing N-heterocyclic carbene (NHC) ligated gold(i) complexes of the type $[\text{Au}(\text{NHC})_2]^+$ were prepared and found to be capable of regulating the formation of reactive oxygen species (ROS) via multiple mechanisms. Single crystal X-ray analysis of bis(1-(ferrocenylmethyl)-3-mesitylimidazol-2-ylidene)-gold(i) chloride (5) and bis(1,3-di(ferrocenylmethyl)imidazol-2-ylidene)-gold(i) chloride (6) revealed a quasi-linear geometry around the gold(i) centers (*i.e.*, the C–Au–C bond angle were measured to be $\sim 177^\circ$ and all the Au–C_{carbene} bonds distances were in the range of 2.00 (7)–2.03 (1) Å). A series of cell studies indicated that cell proliferation inhibition and ROS generation were directly proportional to the amount of ferrocene contained within the $[\text{Au}(\text{NHC})_2]^+$ complexes (IC_{50} of **6** < **5** < bis(1-benzyl-3-mesitylimidazol-2-ylidene)-gold(i) chloride (4)). Complexes **4**–**6** were also confirmed to inhibit thioredoxin reductase as inferred from lipoate reduction assays and increased chelatable intracellular zinc concentrations. RNA microarray gene expression assays revealed that **6** induces endoplasmic reticulum stress response pathways as a result of ROS increase.

Received 17th September 2015

Accepted 28th October 2015

DOI: 10.1039/c5sc03519h

www.rsc.org/chemicalscience

1. Introduction

The regulation of reactive oxygen species (ROS) within biological systems plays a vital role in the health and longevity of many organisms.^{1,2} In disease states, such as cancer, the basal levels of ROS are elevated due to increased cellular growth combined with reduced waste elimination.³ Cellular adaptation to ROS in pre-neoplastic cells exposed to inducers (*i.e.*, hypoxia, metabolic defects, ER stress, oncogene activation, *etc.*) often result in increased antioxidant pathway activity.⁴ Biological systems have a host of natural defenses to ROS, including enzymes such as superoxide dismutase (SOD), glutathione peroxidase, and

catalase. Additional cofactors, such as glutathione (GSH), thioredoxin (Trx)/thioredoxin reductase (TrxR), ascorbate (vitamin C), and α -tocopherol (vitamin E), are also able to serve as ROS scavengers.

Although cancer cells can thrive in the oxidative environment that they create, their ability to buffer ROS has limits. Both tumor and normal cells are driven to apoptosis when ROS levels become too high.⁵ This oxidative-stress pathway to apoptosis, if exploited, could be a new cancer treatment option.⁴ While any compound that disrupts redox homeostasis will negatively affect all cells, normal cells are thought to have a greater capacity for adaptation.⁴ Thus, it is expected that an agent that acts to increase oxidative stress will overload the capabilities of neoplastic cells, while being relatively less lethal to normal cells.

The chemotherapeutic development of agents that alter the redox environment within cancer cells have been categorized into (1) ROS generators (*e.g.*, motexafin gadolinium (MGd), β -lapachone, *etc.*) and (2) antioxidant system inhibitors (*e.g.*, buthionine sulphoximine, tetrathiomolybdate).^{6–14} Collectively, data gleaned from this work have provided insight into the cellular antioxidant system and has resulted in the proposal of oxidative stress modulation as an anticancer strategy.^{4,15}

Within the antioxidant system, Trx plays a central role in mediating cellular response to environmental stress making the inhibition of Trx/TrxR an attractive strategy for patients undergoing radiation therapy.^{16,17} Studies with auranofin (see Fig. 1) revealed the ability of Au(i)-based compounds to inhibit TrxR via binding to the selenylsulfide/selenothiol redox center

^aDepartment of Chemistry, Georgia Southern University, Statesboro, Georgia, 30460, USA. E-mail: jarambula@georgiasouthern.edu; Tel: +1-912-478-2346

^bDepartment of Chemistry, St. Bonaventure University, 3261 West State Road, New York, 14778, USA

^cLumiphore, Inc., Berkeley, California, 94710, USA

^dDepartment of Health Sciences, Gettysburg College, Gettysburg, PA 17325-1400, USA

^eCenter for Multidimensional Carbon Materials (CMCM), Institute for Basic Science (IBS), Ulsan 689-798, Republic of Korea

^fDepartment of Chemistry and Department of Energy Engineering, Ulsan National Institute of Science and Technology (UNIST), Ulsan 689-798, Republic of Korea

^gDepartment of Chemistry, University of Texas at Austin, Austin, Texas, 78712, USA

^hDepartment of Chemistry, Wright State University, 3640 Colonel Glenn Hwy, Dayton, Ohio, 45435, USA. E-mail: kuppiswamy.arumugam@wright.edu

† Electronic supplementary information (ESI) available. CCDC 1419940 and 1419941. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5sc03519h



fetal bovine serum, and antibiotics (200 U cm⁻³ penicillin and 200 µg cm⁻³ streptomycin). Stock solutions of complex (10 mM in DMSO or 5 mM in 50/50 v/v water/DMSO) were formulated and then diluted in medium for secondary stocks of 100–200 µM depending on the complex being tested. Secondary stock solutions were serially diluted in medium and immediately added to wells, whereupon plates were incubated at 37 °C under a 5% CO₂/95% air atmosphere. After a total of three days, a 50 µL aliquot of 3 mg mL⁻¹ tetrazolium dye, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Sigma Chemical Co.), was added to each well, followed by a four hour incubation period at 37 °C. The medium was then removed, the resulting formazan was dissolved in 50 µL DMSO and the respective absorbances were measured at 560–650 nm using a microplate reader (Molecular Devices, Sunnyvale, CA). Absorbance values were corrected for background and then normalized to wells containing untreated cells to allow for plate-to-plate comparisons. The data are shown as mean inhibition of proliferation or growth as a percentage of control cells and are from 2–3 replicate experiments.

2.5 ICP-MS determination of Fe and Au

To determine complex uptake in tumor cell lines, ICP-MS studies were undertaken using an A549 lung cancer cell line. Cells were seeded in 150 cm² cell culture flasks and grown to confluence in 30 mL RPMI 1640 medium supplemented with 2 mM L-glutamine, 10% heat inactivated fetal bovine serum, and antibiotics (200 U cm⁻³ penicillin and 200 µg cm⁻³ streptomycin). The media was removed and supplemented with 30 mL of media (with FBS and antibiotics) containing 2.5 µM of complex originating from 2.5 mM stock complex in DMSO. The cells were then allowed to incubate at 37 °C under a 5% CO₂/95% air atmosphere for 6 h. The medium from each sample was then removed, and cells were washed with PBS (made in-house with ICP-MS grade deionized water), treated with trypsin, and pelleted in 15 mL conical tubes. The pellets were washed 2× with 10 mL of PBS and the cells were counted with a hemocytometer. Cell counts consisted of 25–40 million cells per sample. Samples were pelleted, frozen over dry ice, digested with conc. HNO₃, and analyzed by Applied Analytical, Inc. for total Fe and Au content.

2.6 Determination of relative ROS levels through FACS analysis

Tumor cells (2–3 × 10⁶) were plated overnight and then incubated with media containing one of the complexes described above at concentrations of 2.5 µM. Control cells were treated with vehicle only. At defined time-points, the media was collected and the cells were washed with PBS. The PBS washing was collected and the attached cells were treated with trypsin and collected. The loosened cells were passed through a 40 µm cell strainer. All media and washings were collected, pelleted by centrifugation (3 min @ 300 g) and washed twice with cold PBS. The cells were once again pelleted and suspended in PBS at a final concentration of 2 × 10⁶ cells per mL. To each of the 15 mL centrifuge tubes was added 100 µL of the cell suspension

before being incubated in the dark at 37 °C for 15 min at a final concentration of 1 mg mL⁻¹ CM-DCFA. PBS (2 mL) was added to each sample. The cells were then pelleted, washed 2× with PBS, and re-suspended in 5 µg mL⁻¹ of propidium iodide (PI) in PBS. Control samples of unstained cells, cells stained with only CM-DCFA, and cells stained with only PI were also prepared. Each sample was added to one well of a 96-well plate. Samples were then subjected to FACS analysis using a Millipore Guava easy-Cyte 8 and analyzed using the Guava inCyte software.

2.7 Lipote reduction

To 96-well plates containing plateau phase A549 lung cells was added 2.5 µM of each complex described above in RPMI-16 media containing 10% FBS and P/S. The cells were left to incubate for 6 h at 37 °C at 5% CO₂. At this point, the media was removed and replaced with HBSS buffer containing 20 mM lipoate and 1 mM 5,5'-dithiobis-2-nitrobenzoic acid (DTNB). The plates were immediately monitored at 405 nm in a time dependent fashion. Time points were collected every 20 min and the T₀ value was subtracted from each point.

2.8 Determination of intracellular Zn levels through FACS analysis

The concentration of intracellular free zinc was assessed using the ion-specific fluorescent probe, FluoZin-3-AM (FluoZin-3, Molecular Probes, Inc.) Plateau phase cultures grown in T-25 flasks were treated with control vehicle or zinc acetate in the presence or absence of 2.5 µM of a complex as described above for 4 h. After treatment, the cells were washed with PBS and treated with trypsin/EDTA for 5 min. Complete medium was then added and the cells were isolated by centrifugation. Cell pellets were washed and re-suspended in PBS. An aliquot of 1 × 10⁶ cells was removed, centrifuged, and re-suspended in 100 µL of 20 µmol L⁻¹ FluoZin-3 in PBS. After a 25 min incubation period under ambient conditions, the cells were washed twice with PBS, re-suspended in 0.5 mL PBS, and then incubated for 20 min. An aliquot of the cell suspension was supplemented with 2 µg mL⁻¹ propidium iodide (Sigma), incubated for 5 min, and subjected to two-parameter flow cytometric analysis.

2.9 Isolation of RNA for microarray analysis

Ribonucleic acid was isolated for microarray analysis using the QIAGEN RNeasy Plus Universal Mini Total RNA extraction protocol. About 5 000 000 cells of A549 were seeded into each cell culture dish (Corning 430293) with 11 mL RPMI 1640 with 2 mM L-glutamine, 10% heat-inactivated fetal bovine serum, 200 U cm⁻³ penicillin, and 200 µg cm⁻³ streptomycin. Each pair of dishes was seeded from the same T-75 flask. After being incubated overnight at 37 °C with 5% CO₂, another 11 mL of the previously-described media were added to control dishes, while treated dishes received 11 mL of 6 at a concentration of 5 µM in the previously-described media, for a final concentration of 2.5 µM. After incubating at 37 °C with 5% CO₂ for 6 h, the media was removed by aspiration. The cells were washed with warm PBS and then treated with trypsin. The trypsin was quenched using complete medium and the cells were transferred into



3.2 Ability to inhibit cell proliferation

To assess the ability, if any, of the individual complexes of this study to inhibit cell growth, cell proliferation assays were conducted following exposure of A549 lung cells to 2–6 and a control compound, auranofin; key data are summarized in Table 2. Typical dose–response curves were observed with all complexes investigated (*cf.* Fig. 3a). It was observed that potency was directly proportional to amount of ferrocene contained within the complex (*i.e.*, IC_{50} of $6 < 5 < 4$). The potency of **6** ($IC_{50} = 0.14 \pm 0.03 \mu\text{M}$) was found to be >10 fold greater than auranofin ($IC_{50} = 1.67 \pm 0.05 \mu\text{M}$) in this cell line. In addition, it was found that the Au-containing complexes displayed significantly greater potency (>100-fold) than the individual ferrocene subunits (*i.e.*, compounds **2** and **3**). To assess the contribution of each moiety of **6** to the observed cell proliferation inhibitory effects, A549 cells were exposed to variable concentrations of **3** + **4** and auranofin + **3**, both in a 1 : 2 molar ratio, and compared to **6**. A combined dose of $[\text{Au}(\text{NHC})_2]^+$ **4** and ferrocene **3** provided a slight synergistic effect (*i.e.* $IC_{50} = 0.61 \pm 0.05 \mu\text{M}$ vs. $0.71 \pm 0.03 \mu\text{M}$), while mixtures of auranofin and **3** provided no improvement. Regardless, the ability of **6** to inhibit cell proliferation was still significantly greater than the sum of its constituent parts (*c.f.* Fig. 3b).

Previous studies had indicated an increase in the potency of ferrocenium relative to ferrocene.²⁶ To test if this relationship was relevant to the complexes described above, **5** was oxidized




Fig. 3 (a) Cell proliferation profiles and (b) bar graph plot summarizes the potency of all the compounds explored. Note that the potency of **6** is greater than the 2 : 1 mixture of compounds **3** and **4**. Error bars represent one standard deviation. A one-way ANOVA with Dunnet's *post-hoc* test was used to compare each compound with auranofin ($*P < 0.05$; $**P < 0.01$).

Table 2 IC_{50} values for compounds tested on A549 lung cancer cells

Complex	IC_{50} (μM) ^a	Complex	IC_{50} (μM)
2	6.4 ± 1.1	6	0.14 ± 0.03
3	13 ± 1.5	Auranofin (A)	1.67 ± 0.05
4	0.71 ± 0.03	3 + 4	0.61 ± 0.05
5	0.39 ± 0.01	Auranofin (A) + 3	1.61 ± 0.09

^a Standard deviation is noted (3–5 repeat runs).

to $[\mathbf{5}][\text{BF}_4]_2$ and examined for its ability to inhibit cell proliferation of A549 lung cells. The oxidized ferrocenium compound ($[\mathbf{5}][\text{BF}_4]_2$) was tested in conjunction with **5** and no difference was observed in its ability to inhibit A549 cell growth (see ESI†).

Complex **6** was further screened with PC-3 prostate (p53 null), A2780 (wt-p53 platinum sensitive), and 2780CP (wt-p53 isogenic partner to A2780 displaying multidrug resistance (MDR)) (see Table 3).^{66,67} Inspection of the IC_{50} values indicated similar potencies across all cell lines with no observed resistance in 2780CP relative to A2780 cell lines.

3.3 Assessing cellular uptake via ICP-MS

To assess complex integrity and uptake, inductively coupled plasma mass spectrometry (ICP-MS) studies were carried out with the goal of quantifying Fe and Au levels. In brief, A549 cells were independently exposed to 2.5 μM of **6** or **3** for 6 h. Cells were then collected, counted, and quantified for intracellular uptake of Fe and Au *via* ICP-MS (*cf.* Fig. 4). An increase in Fe was evident in both samples treated with **6** as well as **3**. Relative to **3**, Fe was quantified as 11-fold higher in cells after exposure to **6**. This result suggested to us that cellular uptake of **6** was 5.6-fold higher than that of **3** in A549 lung cancer cells and is consistent with the ~100-fold difference observed in the ability of **6** to inhibit cell proliferation. Gold was also detected in cells exposed to **6**. Subsequent analysis resulted in a 4 : 1 ratio of Fe : Au in cells exposed to **6**, indicative that the $[\text{Au}(\text{NHC})_2]^+$ complex is stable and enters the cell as a whole complex.

3.4 Reactive oxygen species disruption

It is proposed that the combination of ferrocene moieties and Au–NHC complexes present in the compounds of this study results in a system that is capable of disrupting ROS regulation *via* multiple mechanisms. To assess the ability of the aforementioned complexes to disrupt and increase ROS levels, fluorescence assisted cell-sorting (FACS) analyses were conducted utilizing 5-(and-6)-chloromethyl-2',7'-dichlorodihydrofluorescein diacetate, acetyl ester (CM-H₂DCFDA), a fluorescein based indicator for

Table 3 IC_{50} values for compound **6** in various cancer cell lines

Cell line	A549 lung	A2780 ovarian	2780CP ovarian	PC-3 prostate
IC_{50} (μM) ^a	0.14 ± 0.03	0.19 ± 0.01^c	0.12 ± 0.01^c	0.48 ± 0.15^b

^a Standard deviation is noted (3–5 repeat runs). ^b A Dunnet's *post-hoc* test revealed that the IC_{50} for compound **6** was only different in PC-3 cells. ^c Tukey's test was used to verify that the potency of compound **6** was equal in the A2780 and A2780CP cell lines.



Acknowledgements

J. F. A. acknowledges support in part by funds from the Office of the Vice President for Research & Economic Development and the Jack N. Averitt College of Graduate Studies. K. A. acknowledges start-up funds provided by St. Bonaventure University and a Keenan-Martine grant. C. W. B. acknowledges the Institute for Basic Science (IBS-R019-D1) as well as the BK21 Plus Program as funded by the Ministry of Education and the National Research Foundation of Korea for their support. J. L. S. acknowledges the National Institutes of Health (CA 68682) for partial support. The authors would also like to thank Prof. Zahid H. Siddik (M.D. Anderson Cancer Center, Houston) for providing A2780 and 2780CP Ovarian cell lines.

Notes and references

- J. M. McCord and I. Fridovich, *J. Biol. Chem.*, 1968, **243**, 5753–5760.
- J. M. McCord and I. Fridovich, *Free Radical Biol. Med.*, 1988, **5**, 363–369.
- D. Trachootham, J. Alexandre and P. Huang, *Nat. Rev. Drug Discovery*, 2009, **8**, 579–591.
- M. Diehn, R. W. Cho, N. A. Lobo, T. Kalisky, M. J. Dorie, A. N. Kulp, D. Qian, J. S. Lam, L. E. Ailles, M. Wong, B. Joshua, M. J. Kaplan, I. Wapnir, F. M. Dirbas, G. Somlo, C. Garberoglio, B. Paz, J. Shen, S. K. Lau, S. R. Quake, J. M. Brown, I. L. Weissman and M. F. Clarke, *Nature*, 2009, **458**, 780–783.
- Y. Zhao, E. B. Butler and M. Tan, *Cell Death Dis.*, 2013, **4**, e532.
- E. A. Bey, M. S. Bentle, K. E. Reinicke, Y. Dong, C. R. Yang, L. Girard, J. D. Minna, W. G. Bornmann, J. Gao and D. A. Boothman, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 11832–11837.
- J. Chen and J. Stubbe, *Curr. Opin. Chem. Biol.*, 2004, **8**, 175–181.
- F. M. de Groot, E. W. Damen and H. W. Scheeren, *Curr. Med. Chem.*, 2001, **8**, 1093–1122.
- U. Galm, M. H. Hager, S. G. van Lanen, J. Ju, J. S. Thorson and B. Shen, *Chem. Rev.*, 2005, **105**, 739–758.
- J. C. Juarez, M. Manuia, M. E. Burnett, O. Betancourt, B. Boivin, D. E. Shaw, N. K. Tonks, A. P. Mazar and F. Doñate, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 7147–7152.
- H. Maeda, S. Hori, H. Ohizumi, T. Segawa, Y. Kakehi, O. Ogawa and A. Kakizuka, *Cell Death Differ.*, 2004, **11**, 737–746.
- D. Magda and R. A. Miller, *Semin. Cancer Biol.*, 2006, **16**, 466–476.
- J. L. Sessler, G. Hemmi, T. D. Mody, T. Murai, A. Burrell and S. W. Young, *Acc. Chem. Res.*, 1994, **27**, 43–50.
- M. Tomasz, *Chem. Biol.*, 1995, **2**, 575–579.
- C. Gorrini, I. S. Harris and T. W. Mak, *Nat. Rev. Drug Discovery*, 2013, **12**, 931–947.
- D. L. Kirkpatrick, G. Ehrmantraut, S. Stettner, M. Kunkel and G. Powis, *Oncol. Res.*, 1997, **9**, 351–356.
- S. J. Wei, A. Botero, K. Hirota, C. M. Bradbury, S. Markovina, A. Laszlo, D. R. Spitz, P. C. Goswami, J. Yodoi and D. Gius, *Cancer Res.*, 2000, **60**, 6688–6695.
- C. Marzano, V. Gandin, A. Folda, G. Scutari, A. Bindoli and M. P. Rigobello, *Free Radical Biol. Med.*, 2007, **42**, 872–881.
- C. K. Mirabelli, R. K. Johnson, C. M. Sung, L. Faucette, K. Muirhead and S. T. Crooke, *Cancer Res.*, 1985, **45**, 32–39.
- M. P. Rigobello, A. Folda, B. Dani, R. Menabò, G. Scutari and A. Bindoli, *Eur. J. Pharmacol.*, 2008, **582**, 26–34.
- M. P. Rigobello, G. Scutari, R. Boscolo and A. Bindoli, *Br. J. Pharmacol.*, 2002, **136**, 1162–1168.
- T. M. Simon, D. H. Kunishima, G. J. Vibert and A. Lorber, *Cancer*, 1979, **44**, 1965–1975.
- (a) J. L. Hickey, R. A. Ruhayel, P. J. Barnard, M. V. Baker, S. J. Berners-Price and A. Filipovska, *J. Am. Chem. Soc.*, 2008, **130**, 12570–12571; (b) M.-L. Teyssot, A.-S. Jarrousse, M. Manin, A. Chevy, S. Roche, F. Norre, C. Beaudoin, L. Morel, D. Boyer, R. Mahiou and A. Gautier, *Dalton Trans.*, 2009, 6894–6902; (c) I. Ott, X. Qian, Y. Xu, D. H. W. Vlecken, I. J. Marques, D. Kubutat, J. Will, W. S. Sheldrick, P. Jesse, A. Prokop and C. P. Bagowski, *J. Med. Chem.*, 2009, **52**, 763–770; (d) T. V. Serebryanskaya, A. A. Zolotarev and I. Ott, *MedChemComm*, 2015, **6**, 1186–1189; (e) J.-J. Zhang, C.-M. Che and I. Ott, *J. Organomet. Chem.*, 2015, **782**, 37–41; (f) S. B. Aher, P. N. Muskawar, K. Thenmozhi and P. R. Bhagat, *Eur. J. Med. Chem.*, 2014, **81**, 408–419; (g) E. Vergara, A. Casini, F. Sorrentino, O. Zava, E. Cerrada, M. P. Rigobello, A. Bindoli, M. Laguna and P. J. Dyson, *ChemMedChem*, 2010, **5**, 96–102; (h) M. V. Baker, P. J. Barnard, S. J. Berners-Price, S. K. Brayshaw, J. L. Hickey, B. W. Skelton and A. H. White, *Dalton Trans.*, 2006, 3708–3715; (i) P. J. Barnard, M. V. Baker, S. J. Berners-Price, B. W. Skelton and A. H. White, *Dalton Trans.*, 2004, 1038–1047; (j) C.-M. Che and R. W.-Y. Sun, *Chem. Commun. (Cambridge, U. K.)*, 2011, **47**, 9554–9560; (k) J. C. Y. Lin, R. T. W. Huang, C. S. Lee, A. Bhattacharyya, W. S. Hwang and I. J. B. Lin, *Chemical Reviews (Washington, DC, United States)*, 2009, **109**, 3561–3598; (l) W. Liu and R. Gust, *Chem. Soc. Rev.*, 2013, **42**, 755–773; (m) E. Schuh, C. Pflüger, A. Citta, A. Folda, M. P. Rigobello, A. Bindoli, A. Casini and F. Mohr, *J. Med. Chem.*, 2012, **55**, 5518–5528; (n) T. J. Siciliano, M. C. Deblock, K. M. Hindi, S. Durmus, M. J. Panzner, C. A. Tessier and W. J. Youngs, *J. Organomet. Chem.*, 2011, **696**, 1066–1071.
- O. Rackham, S. J. Nichols, P. J. Leedman, S. J. Berners-Price and A. Filipovska, *Biochem. Pharmacol.*, 2007, **74**, 992–1002.
- E. Viry, E. Battaglia, V. Deborde, T. Müller, R. Réau, E. Davioud-Charvet and D. Bagrel, *ChemMedChem*, 2008, **3**, 1667–1670.
- R. Rubbiani, I. Kitanovic, H. Alborzinia, S. Can, A. Kitanovic, L. A. Onambele, M. Stefanopoulou, Y. Geldmacher, W. S. Sheldrick, G. Wolber, A. Prokop, S. Wolf and I. Ott, *J. Med. Chem.*, 2010, **53**, 8608–8618.
- S. J. Berners-Price and P. J. Barnard, in *Ligand Design in Medicinal Inorganic Chemistry*, John Wiley & Sons, Ltd, 2014, pp. 227–256 DOI: 10.1002/9781118697191.ch9 and references therein.



- 28 M. A. Cinellu, I. Ott and A. Casini, in *Bioorganometallic Chemistry*, Wiley-VCH Verlag GmbH & Co. KGaA, 2014, pp. 117–140, DOI: 10.1002/9783527673438.ch04 and references therein.
- 29 I. Ott, *Coord. Chem. Rev.*, 2009, **253**, 1670–1681.
- 30 B. Bertrand, L. Stefan, M. Pirrotta, D. Monchaud, E. Bodio, P. Richard, P. le Gendre, E. Warmerdam, M. H. de Jager, G. M. M. Groothuis, M. Picquet and A. Casini, *Inorg. Chem.*, 2014, **53**, 2296–2303.
- 31 E. M. Barranco, M. C. Gimeno, A. Laguna and M. D. Villacampa, *Inorg. Chim. Acta*, 2005, **358**, 4177–4182.
- 32 M. Concepción Gimeno and A. Laguna, *Gold Bull.*, 1999, **32**, 90–95.
- 33 M. C. Gimeno, A. Laguna, C. Sarroca and P. G. Jones, *Inorg. Chem.*, 1993, **32**, 5926–5932.
- 34 D. T. Hill, G. R. Girard, F. L. McCabe, R. K. Johnson, P. D. Stupik, J. H. Zhang, W. M. Reiff and D. S. Eggleston, *Inorg. Chem.*, 1989, **28**, 3529–3533.
- 35 A. Houlton, R. M. G. Roberts, J. Silver and R. V. Parish, *J. Organomet. Chem.*, 1991, **418**, 269–275.
- 36 M. Viotte, B. Gautheron, M. M. Kubicki, Y. Mugnier and R. V. Parish, *Inorg. Chem.*, 1995, **34**, 3465–3473.
- 37 M. Viotte, B. Gautheron, I. Nifant'ev and L. G. Kuz'mina, *Inorg. Chim. Acta*, 1996, **253**, 71–76.
- 38 U. E. I. Horvath, G. Bentivoglio, M. Hummel, H. Schottenberger, K. Wurst, M. J. Nell, C. E. J. van Rensburg, S. Cronje and H. G. Raubenheimer, *New J. Chem.*, 2008, **32**, 533–539.
- 39 C. Y. Acevedo-Morantes, E. Meléndez, S. P. Singh and J. E. Ramírez-Vick, *J. Cancer Sci. Ther.*, 2012, **4**, 271–275.
- 40 M. F. R. Fouda, M. M. Abd-Elzaher, R. A. Abdelsamaia and A. A. Labib, *Appl. Organomet. Chem.*, 2007, **21**, 613–625.
- 41 G. Gasser, I. Ott and N. Metzler-Nolte, *J. Med. Chem.*, 2011, **54**, 3–25.
- 42 M. Gormen, P. Pigeon, S. Top, A. Vessieres, M.-A. Plamont, E. A. Hillard and G. Jaouen, *MedChemComm*, 2010, **1**, 149–151.
- 43 M. Gormen, D. Plažuk, P. Pigeon, E. A. Hillard, M.-A. Plamont, S. Top, A. Vessières and G. Jaouen, *Tetrahedron Lett.*, 2010, **51**, 118–120.
- 44 Á. Mooney, R. Tiedt, T. Maghoub, N. O'Donovan, J. Crown, B. White and P. T. M. Kenny, *J. Med. Chem.*, 2012, **55**, 5455–5466.
- 45 D. Plažuk, A. Vessières, E. A. Hillard, O. Buriez, E. Labbé, P. Pigeon, M.-A. Plamont, C. Amatore, J. Zakrzewski and G. Jaouen, *J. Med. Chem.*, 2009, **52**, 4964–4967.
- 46 G. Tabbi, C. Cassino, G. Cavigliolo, D. Colangelo, A. Ghiglia, I. Viano and D. Osella, *J. Med. Chem.*, 2002, **45**, 5786–5796.
- 47 D. Hamels, P. M. Dansette, E. A. Hillard, S. Top, A. Vessières, P. Herson, G. Jaouen and D. Mansuy, *Angew. Chem., Int. Ed.*, 2009, **48**, 9124–9126.
- 48 G. Jaouen and S. Top, in *Advances in Organometallic Chemistry and Catalysis*, John Wiley & Sons, Inc., 2013, pp. 563–580 DOI: 10.1002/9781118742952.ch42 and references therein.
- 49 D. Osella, M. Ferrali, P. Zanello, F. Laschi, M. Fontani, C. Nervi and G. Cavigliolo, *Inorg. Chim. Acta*, 2000, **306**, 42–48.
- 50 H. Tamura and M. Miwa, *Chem. Lett.*, 1997, **26**, 1177–1178.
- 51 For other ferrocenylated NHC complexes see, U. Siemeling, *Eur. J. Inorg. Chem.*, 2012, **2012**, 3523–3536 and references therein; K. Arumugam, J. Chang, V. M. Lynch and C. W. Bielawski, *Organometallics*, 2013, **32**, 4334–4341; K. Arumugam, C. D. Varnado, S. Sproules, V. M. Lynch and C. W. Bielawski, *Chem.–Eur. J.*, 2013, **19**, 10866–10875; B. Bildstein, M. Malaun, H. Kopacka, K.-H. Ongania and K. Wurst, *J. Organomet. Chem.*, 1998, **552**, 45–61; B. Bildstein, M. Malaun, H. Kopacka, K.-H. Ongania and K. Wurst, *J. Organomet. Chem.*, 1999, **572**, 177–187; B. Bildstein, M. Malaun, H. Kopacka, K. Wurst, M. Mitterböck, K.-H. Ongania, G. Opromolla and P. Zanello, *Organometallics*, 1999, **18**, 4325–4336; N. Debono, A. S. Labande, E. Manoury, J.-C. Daran and R. Poli, *Organometallics*, 2010, **29**, 1879–1882; W. A. Herrmann and C. Köcher, *Angew. Chem., Int. Ed.*, 1997, **36**, 2162–2187; D. M. Khramov, E. L. Rosen, V. M. Lynch and C. W. Bielawski, *Angew. Chem., Int. Ed.*, 2008, **47**, 2267–2270; A. Labande, J.-C. Daran, E. Manoury and R. Poli, *Eur. J. Inorg. Chem.*, 2007, 1205–1209; S. Leuthäuser, D. Schwarz and H. Plenio, *Chem.–Eur. J.*, 2007, **13**, 7195–7203; E. L. Rosen, C. D. Varnado, A. G. Tennyson, D. M. Khramov, J. W. Kamplain, D. H. Sung, P. T. Cresswell, V. M. Lynch and C. W. Bielawski, *Organometallics*, 2009, **28**, 6695–6706; U. Siemeling, *Eur. J. Inorg. Chem.*, 2012, 3523–3536; U. Siemeling, T. C. Auch, O. Kuhnert, M. Malaun, H. Kopacka and B. Bildstein, *Z. Anorg. Allg. Chem.*, 2003, **629**, 1334–1336; U. Siemeling, C. Farber and C. Bruhn, *Chem. Commun.*, 2009, 98–100; U. Siemeling, C. Farber, C. Bruhn, M. Leibold, D. Selent, W. Baumann, M. von Hopffgarten, C. Goedecke and G. Frenking, *Chem. Sci.*, 2010, **7**, 697–704; U. Siemeling, C. Färber, M. Leibold, C. Bruhn, P. Mücke, R. F. Winter, B. Sarkar, M. von Hopffgarten and G. Frenking, *Eur. J. Inorg. Chem.*, 2009, 4607–4612; M. Süßner and H. Plenio, *Angew. Chem., Int. Ed.*, 2005, **44**, 6885–6888; C. D. Varnado Jr, V. M. Lynch and C. W. Bielawski, *Dalton Trans.*, 2009, 7253–7261.
- 52 G. Occhipinti, V. R. Jensen, K. W. Törnroos, N. Å. Frøystein and H.-R. Bjørsvik, *Tetrahedron*, 2009, **65**, 7186–7194.
- 53 A. Flahaut, S. Roland and P. Mangeney, *J. Organomet. Chem.*, 2007, **692**, 5754–5762.
- 54 M. K. Samantaray, C. Dash, M. M. Shaikh, K. Pang, R. J. Butcher and P. Ghosh, *Inorg. Chem.*, 2011, **50**, 1840–1848.
- 55 K. Arumugam, C. D. Varnado, S. Sproules, V. M. Lynch and C. W. Bielawski, *Chem.–Eur. J.*, 2013, **19**, 10866–10875.
- 56 J. Howarth, J. L. Thomas, K. Hanlon and D. McGuirk, *Synth. Commun.*, 2000, **30**, 1865–1878.
- 57 A. S. K. Hashmi, T. Hengst, C. Lothschütz and F. Rominger, *Adv. Synth. Catal.*, 2010, **352**, 1315–1337.
- 58 J. R. Aranzaes, M.-C. Daniel and D. Astruc, *Can. J. Chem.*, 2006, **84**, 288–299.
- 59 I. Noviandri, K. N. Brown, D. S. Fleming, P. T. Gulyas, P. A. Lay, A. F. Masters and L. Phillips, *J. Phys. Chem. B*, 1999, **103**, 6713–6722.



- 60 M. C. Burla, R. Caliendo, M. Camalli, B. Carrozzini, G. L. Cascarano, L. de Caro, C. Giacobozzo, G. Polidori and R. Spagna, *J. Appl. Crystallogr.*, 2005, **38**, 381–388.
- 61 M. C. Burla, M. Camalli, B. Carrozzini, G. L. Cascarano, C. Giacobozzo, G. Polidori and R. Spagna, *J. Appl. Crystallogr.*, 2003, **36**, 1103.
- 62 G. Sheldrick, *Acta Crystallogr., Sect. C: Struct. Chem.*, 2015, **71**, 3–8.
- 63 G. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 2008, **64**, 112–122.
- 64 J. J. Dunsford, K. J. Cavell and B. M. Kariuki, *Organometallics*, 2012, **31**, 4118–4121.
- 65 P. J. Barnard, M. V. Baker, S. J. Berners-Price, B. W. Skelton and A. H. White, *Dalton Trans.*, 2004, 1038–1047, DOI: 10.1039/b316804b.
- 66 J. F. Arambula, J. L. Sessler and Z. H. Siddik, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 1701–1705.
- 67 J. F. Arambula, J. L. Sessler and Z. H. Siddik, *MedChemComm*, 2012, **3**, 1275–1281.
- 68 E. Obrador, J. Navarro, J. Mompo, M. Asensi, J. A. Pellicer and J. M. Estrela, *BioFactors*, 1998, **8**, 23–26.
- 69 S. Tuttle, A. Horan, C. Koch, K. Held, Y. Manevich and J. Biaglow, *Int. J. Radiat. Oncol., Biol., Phys.*, 1998, **42**, 833–838.
- 70 D. Magda, C. Lepp, N. Gerasimchuk, I. Lee, J. L. Sessler, A. Lin, J. E. Biaglow and R. A. Miller, *Int. J. Radiat. Oncol., Biol., Phys.*, 2001, **51**, 1025–1036.
- 71 K. R. Gee, Z. L. Zhou, D. Ton-That, S. L. Sensi and J. H. Weiss, *Cell Calcium*, 2002, **31**, 245–251.
- 72 D. Magda, P. Lecane, R. A. Miller, C. Lepp, D. Miles, M. Mesfin, J. E. Biaglow, V. V. Ho, D. Chawannakul, S. Nagpal, M. W. Karaman and J. G. Hacia, *Cancer Res.*, 2005, **65**, 3837–3845.
- 73 D. Magda, P. Lecane, Z. Wang, W. Hu, P. Thiemann, X. Ma, P. K. Dranchak, X. Wang, V. Lynch, W. Wei, V. Csokai, J. G. Hacia and J. L. Sessler, *Cancer Res.*, 2008, **68**, 5318–5325.
- 74 M. Hirsilä, P. Koivunen, L. Xu, T. Seeley, K. I. Kivirikko and J. Myllyharju, *FASEB J.*, 2005, **19**, 1308–1310.
- 75 P. S. Lecane, M. W. Karaman, M. Sirisawad, L. Naumovski, R. A. Miller, J. G. Hacia and D. Magda, *Cancer Res.*, 2005, **65**, 11676–11688.
- 76 C. J. Schofield and P. J. Ratcliffe, *Nat. Rev. Mol. Cell Biol.*, 2004, **5**, 343–354.
- 77 G. L. Semenza, *Nat. Rev. Cancer*, 2003, **3**, 721–732.
- 78 J. H. Choi, T. N. Kim, S. Kim, S. H. Baek, J. H. Kim, S. R. Lee and J. R. Kim, *Anticancer Res.*, 2002, **22**, 3331–3335.
- 79 D. L. Kirkpatrick, M. Kuperus, M. Dowdeswell, N. Potier, L. J. Donald, M. Kunkel, M. Berggren, M. Angulo and G. Powis, *Biochem. Pharmacol.*, 1998, **55**, 987–994.
- 80 D. K. Smart, K. L. Ortiz, D. Mattson, C. M. Bradbury, K. S. Bisht, L. K. Sieck, M. W. Brechbiel and D. Gius, *Cancer Res.*, 2004, **64**, 6716–6724.

