

# Chemical Science

rsc.li/chemical-science



ISSN 2041-6539



ROYAL SOCIETY  
OF CHEMISTRY

Celebrating  
IYPT 2019

#### EDGE ARTICLE

Werner M. Nau, Oren A. Scherman *et al.*  
Preferential binding of unsaturated hydrocarbons in  
aryl-bisimidazolium-cucurbit[8]uril complexes furthers  
evidence for small-molecule  $\pi$ - $\pi$  interactions

Cite this: *Chem. Sci.*, 2019, 10, 10240

All publication charges for this article have been paid for by the Royal Society of Chemistry

# Preferential binding of unsaturated hydrocarbons in aryl-bisimidazolium·cucurbit[8]uril complexes furberishes evidence for small-molecule $\pi-\pi$ interactions†

Steven J. Barrow, ‡<sup>a</sup> Khaleel I. Assaf, ‡<sup>b</sup> Aniello Palma,<sup>a</sup> Werner M. Nau \*<sup>b</sup> and Oren A. Scherman \*<sup>a</sup>

Whilst cucurbit[*n*]urils (CB*n*) have been utilized in gas encapsulation, only the smaller CB*n* (*n* = 5 and 6) have utility given their small cavity size. In this work, we demonstrate that the large cavity of CB8 can be tailored for gaseous and volatile hydrocarbon encapsulation by restricting its internal cavity size with auxiliary aryl-bisimidazolium (Bis, aryl = phenyl, naphthyl, and biphenyl) guests. The binding constants for light hydrocarbons (*C* ≤ 4) are similar to those measured with CB6, while larger values are obtained with Bis·CB8 for larger guests. A clear propensity for higher affinities of alkenes relative to alkanes is observed, most pronounced with the largest delocalized naphthalene residue in the auxiliary Bis guest, which provides unique evidence for sizable small-molecule  $\pi-\pi$  interactions.

Received 3rd July 2019

Accepted 7th October 2019

DOI: 10.1039/c9sc03282g

rsc.li/chemical-science

## Introduction

The development of refined gas-encapsulation materials is driven by economic (methane binding) and environmental (CO<sub>2</sub> capture) promises and holds additional potential for advanced sensing and photochemical applications.<sup>1,2</sup> Besides metal-organic frameworks,<sup>3-5</sup> porous coordination polymers<sup>6,7</sup> exhibit very high surface areas and have demonstrated selectivity and capacity for adsorbing gases like CO<sub>2</sub>, making them competitive candidates for gas encapsulation, with the common disadvantage of being water sensitive.<sup>8-10</sup> Discrete host-guest chemistry presents an alternative approach largely by-passing stability issues.<sup>11-13</sup> In particular, cucurbit[*n*]urils, CB*n*, present a class of macrocycles that have demonstrated gas uptake capacities comparable to several porous materials.<sup>11,12,14-21</sup> CB*n* are based on glycoluril subunits, which have been shown to encapsulate a variety of cationic and neutral guest species.<sup>20-27</sup> The unique binding capabilities of the CB*n* family arise due to ion-dipole interactions at the carbonyl-lined portals, in addition to the size and hydrophobicity of the inner cavity.<sup>28</sup> The size of the inner cavity of CB*n* is a powerful predictor in terms of the breadth of chemical species that can bind to the macrocycle. CB8 can bind

two small aromatic compounds simultaneously, whereas the smaller CB7 and CB6 can generally bind only one at a time.<sup>23</sup> As a consequence of size complementarity, gas binding tends to be favoured by the smaller CB5 and CB6 homologues.<sup>14,21,29-31</sup>

Herein, we demonstrate that aryl-bisimidazolium (Bis) guests can tailor the interior cavity space of CB8 toward preferential binding of gaseous and volatile hydrocarbons (Fig. 1) with increased selectivity for unsaturated ones. The Bis guests differ based on the hydrophobic linker between the two imidazolium units, specifically, phenyl (Bis1), naphthyl (Bis2), and biphenyl (Bis3). We show that a wide variety of hydrocarbon guests can be encapsulated, *via* <sup>1</sup>H NMR and fluorescence spectroscopy, and that by changing the hydrophobic moiety within the Bis guests, CB8 can be made selective toward particular guest molecules by changing the size and shape of the remaining cavity space within the macrocycle. The three Bis guests<sup>32</sup> (Fig. 1) have been previously used in conjunction with CB8 to enable encapsulation of small solvent molecules in a 1 : 1 : 1 binding stoichiometry and other auxiliary guests without “lids”.<sup>33,34</sup>

Auxiliary aromatic guest-hydrocarbon complexes have been earlier assembled inside capsular assemblies or coordination cages.<sup>35,36</sup> How these auxiliary guests allow for a large macrocycle to encapsulate small molecular species lies in the fact that the auxiliary guests occupy a large portion of the internal cavity, such as that of CB8, effectively altering the packing coefficient (PC) of small molecules inside the cavity.<sup>28,37</sup> Fig. 1d and e show DFT calculations that reveal the extent of the internal cavity volume restriction for a U-shaped conformation of the Bis guests in CB8 that templates the encapsulation of a second

<sup>a</sup>Melville Laboratory for Polymer Synthesis, Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK. E-mail: oas23@cam.ac.uk

<sup>b</sup>Department of Life Sciences and Chemistry, Jacobs University Bremen, Campus Ring 1, D-28759 Bremen, Germany. E-mail: w.nau@jacobs-university.de

† Electronic supplementary information (ESI) available: Materials and methods section, fluorescence titrations, ITC experiments, recognition of diiodine, and additional NMR data. See DOI: 10.1039/c9sc03282g

‡ These authors contributed equally to this work.





Fig. 1 Chemical structures of (a) CB8, (b) the bisimidazolium auxiliary guests, and (c) their schematic complexes with CB8; differently shaped cavities shown in grey. (d) and (e) show the side and top views of the DFT-optimized (wB97XD/3-21G\* level of theory) binary complex structures with the respective accessible cavity volume ( $V_{\text{cavity}}$ ) highlighted in blue.

guest. Vacant CB8 has a cavity volume of  $367 \text{ \AA}^3$ ,<sup>28</sup> however, the formation of Bis1·CB8, Bis2·CB8, and Bis3·CB8 complexes reduces the available cavity volume to 114, 126, and  $146 \text{ \AA}^3$ , respectively (Table 1). In terms of capacity, these complexes fall in between CB5 and CB6 for which gas encapsulation has been documented,<sup>11,16,17,19,20,29,31</sup> driven, among others, by the release of high-energy water (for CB6)<sup>38,39</sup> and cavitation energy (for CB5).<sup>21</sup> Moreover, as reflected by the calculations, the resulting templated cavities differ in shape, from equatorially elongated (for Bis1·CB8) to spherically (for Bis2·CB8) to axially elongated (for Bis3·CB8), which offers an interesting design approach.

Table 1 Calculated cavity volumes of  $\text{CB}_n$  and Bis·CB8 complexes

$\text{CB}_n$	Cavity volume/ $\text{\AA}^3$	Bis·CB8	Cavity volume/ $\text{\AA}^3$
CB5	$68^a$	Bis1·CB8	$114^b$
CB6	$142^a$	Bis2·CB8	$126^b$
CB7	$242^a$	Bis3·CB8	$146^b$
CB8	$367^a$		

<sup>a</sup> From ref. 28. <sup>b</sup> Calculated from optimized structures, see Fig. 1.

Finally, by varying the size of the aryl unit from phenyl to naphthyl and biphenyl, it should not only be possible to vary size and shape of the resulting cavity, but also secondary C–H– $\pi$  and  $\pi$ – $\pi$  guest–guest interactions, the importance of which remains under debate.<sup>33,36,40–51</sup> In addition to playing a role in rational secluded-cavity design, the Bis guests are fluorescent, which enables working at micromolar concentrations and offers the opportunity for direct and real-time optical sensing of gaseous and volatile guests at micromolar concentrations.<sup>19,20</sup>

## Results and discussion

The formation of 1 : 1 host–guest complexes between CB8 and Bis1–Bis3 with high binding constants ( $K_a > 10^6 \text{ M}^{-1}$ ) has been established by using different spectroscopic methods (see ESI,† and ref. 32). As anticipated, the binary Bis·CB8 complexes should act as receptors for small molecules, which would otherwise not complex to free CB8. Visual evidence for the entrapment of small molecules – and for their potential in sensing – can be obtained by their addition to diiodine ( $\text{I}_2$ ) solutions, which affects an immediate color change (Fig. S11†) from yellow-brown (in water) to violet, an iodine color otherwise only observed in nonpolar solvents. In this case, the binding constants were determined by direct UV-visible titrations ( $K_a \sim 2 \times 10^4 \text{ M}^{-1}$ , see Table 2 and Fig. S11†). These binding constants are lower than the previously measured value with CB6 ( $K_a = 1.4 \times 10^6 \text{ M}^{-1}$ )<sup>52</sup> but similar to that obtained for  $\alpha$ -cyclodextrin.<sup>53</sup> Accordingly, Bis·CB8 complexes are competitive binders for small guests.

Table 2 Association constants ( $K_a$ ) of hydrocarbons with Bis·CB8 systems and  $\text{CB}_n$ , measured in neat water

Guest	$K_a^a/(10^3 \text{ M}^{-1})$				
	Bis1·CB8	Bis2·CB8	Bis3·CB8	$\text{CB}_6^b$	$\text{CB}_7^c$
Methane	$0.6 \pm 0.1$	$0.4 \pm 0.1$	0.6	<2	3
Ethane		$0.5 \pm 0.1$		24	3.4
Propane		14		180	6
<i>n</i> -Butane	580	89	$35 \pm 11$	280	170
<i>cis</i> -Butene		430		150	35
<i>trans</i> -Butene		24		21	14
Isobutane	186	31	410	850	265
Isobutene	18	65	66	84	43
Neopentane	14	5600	9.3	<2	1000
Cyclopentane		67		1300	196
Cyclopentene	$290^d$	$260 [480]^d$	$960^d$	$140^e$	$25^e$
Cyclopentanol	$2.0^d$	$6.7^d$	$5.9^d$		
Cyclohexane		66		<2	1500
1,3-Cyclohexadiene	$88^d$	$530^d$	$1900^d$		
Benzene	$85^d$	$170 [520]^d$	$[710]^d$	<2	17
Phenol	$3.0^d$	$32 \pm 5^d$	$18^d$		
$\text{I}_2$	$17 \pm 2^e$	$21 \pm 8^f$	$19 \pm 4^e$	$1400^g$	$100^g$

<sup>a</sup> Error in  $K_a$  values is 15% unless stated differently. <sup>b</sup> From ref. 20. <sup>c</sup> From ref. 19. <sup>d</sup> Values in square brackets measured by ITC, 10% error unless stated differently, see Table S3 in ESI. <sup>e</sup> Measured in this work by indicator displacement. <sup>f</sup> Measured by UV-vis absorption titrations, see ESI. <sup>g</sup> From ref. 52.





For optically inactive guests, such as hydrocarbons,  $^1\text{H}$  NMR spectroscopy was used for structural characterization of the ternary hydrocarbon complexes; very large upfield shifts ( $>2$  ppm) were observed for the encapsulated guests (see Fig. 2 and ESI $^\dagger$ ), which were found to be in slow exchange even for the smallest guests. This was in contrast to the binary Bis·CB8 complexes, in which host and guest were in fast exchange, resulting in sharper NMR bands. For example, in the  $^1\text{H}$  NMR spectra for complex formation of the Bis1·CB8·methane system (see ESI $^\dagger$ ), significant shifts and line broadening were observed. While upfield shifts up to 1 ppm are characteristic for CB $n$  encapsulation itself,<sup>19</sup> the larger values in the Bis1·CB8 guest complexes are due to anisotropic shielding effects from the adjacent aryl groups of the pre-complexed Bis guests.<sup>8</sup> The  $^1\text{H}$  NMR spectra for the formation of the *trans*-butene·Bis1·CB8 complex can be seen in Fig. 2a. An upfield shift of the peaks associated with *trans*-butene of 1.4–3.0 ppm is observed once the guest is encapsulated within the Bis1·CB8 complex. Similarly, the encapsulation of *cis*-butene inside the Bis2·CB8 complex was confirmed by the upfield shifts of the guest peaks (Fig. 2b). Other gases, including CO<sub>2</sub> and SF<sub>6</sub>, also form complexes with the Bis·CB8 systems (see ESI $^\dagger$ ). The complexation of SF<sub>6</sub> with the Bis·CB8 was investigated by using  $^{19}\text{F}$  NMR, in which the fluorine atoms experienced an upfield shift (see Fig. S20 $^\dagger$ ), in accordance with the inclusion of perfluorinated guests within CB $n$ .<sup>26</sup>

The binding affinities of selected hydrocarbons were independently measured by using fluorescence titrations, through monitoring the fluorescence response of the auxiliary Bis guests upon binding of the second guest molecule (Fig. 3 and ESI $^\dagger$ ). For gases, pressure was adjusted to control their concentration




Fig. 3 (a) The encapsulation of a second guest within a Bis·CB8 complex enhances the fluorescence of the first guest; changes in fluorescence can be directly correlated to the binding strength of the second guest. (b) and (c) Changes in fluorescence spectra for Bis2·CB8 complexes, plotted versus wavelength and second guest concentration (inset) for (b) *n*-butane and (c) neopentane.

(see ESI $^\dagger$ ), while for volatile liquid guests stock solutions were used. Although concentration variations in gas titrations are greatly limited compared to conventional titrations with stock solutions,<sup>19,20</sup> aqueous hydrocarbon solubilities are accurately known, which allowed for good reproducibilities. Enhancement of the fluorescence intensity was observed upon formation of the 1 : 1 : 1 ternary complexes, which is attributed to the replacement of the residual water molecules from the cavity by the hydrophobic guests (Fig. 3). As can be seen from Fig. 3 and Table S2, $^\dagger$  the different hydrocarbons showed markedly different fluorescence responses. The measured binding affinities are shown in Table 2, as well as data for CB6 (ref. 20) and CB7 (ref. 19) (the PC analysis for each system is provided in Table S1 $^\dagger$ ).

The binding constants vary with the auxiliary guest. For example, *n*-butane binds more strongly to the Bis1·CB8 complex compared to Bis2·CB8 and Bis3·CB8, while neopentane binds tightly to Bis2·CB8. This might be attributable to the different cavity shapes (Fig. 1), in which the more spherical cavity of the Bis2·CB8 complex prefers globular guests such as neopentane, while those of Bis1·CB8 and Bis3·CB8 preferentially bind elongated guests, along the equatorial and axial voids, respectively. Interestingly, Bis2·CB8 markedly and consistently showed higher binding affinities for alkenes than for the corresponding alkanes. Specifically, isobutene and cyclopentene bind more strongly than isobutane and cyclopentane, respectively. The opposite selectivity applies for CB6 and CB7, to which saturated hydrocarbons bind more strongly than their unsaturated counterparts.<sup>19,20</sup> Strong binding of alkenes is counterintuitive, because they are 3–5 times more water soluble (Table 3) than alkanes and, therefore, less hydrophobic. This hints at another prevailing aspect that contributes to hydrocarbon binding in these Bis·CB8




Fig. 2  $^1\text{H}$  NMR spectra for the binding of (a) *trans*-butene to the Bis1·CB8 complex and (b) *cis*-butene to Bis2·CB8, in D<sub>2</sub>O.



**Table 3** Guest solubility ( $S$ ), guest volume ( $V$ ), polarizability ( $\alpha$ ), hydration free energy ( $\Delta G_{\text{hydr}}$ ), binding free energy ( $\Delta G_{\text{a}}$ ) as measured in neat water, and corrected binding free energy ( $\Delta G'_{\text{a}}$ ) for different CBn host–guest complexes; all energy values in kcal mol<sup>-1</sup>

Guest	$S^a/\text{mM}$	$V^b/\text{\AA}^3$	$\alpha^c/\text{\AA}^3$	$\Delta G_{\text{hydr}}^d$	Bis2·CB8		CB6		CB7	
					$\Delta G_{\text{a}}^e$	$\Delta G'_{\text{a}}^f$	$\Delta G_{\text{a}}^g$	$\Delta G'_{\text{a}}^f$	$\Delta G_{\text{a}}^h$	$\Delta G'_{\text{a}}^f$
Methane	1.40	29	2.59	1.99	$-3.53 \pm 0.17$	-1.54			-4.74	-2.75
Ethane	1.89	45	4.43	1.82	$-3.73 \pm 0.13$	-1.92	-5.98	-4.16	-4.82	-3.00
Propane	1.52	63	6.37	1.94	-5.66	-3.72	-7.17	-5.23	-5.15	-3.21
<i>n</i> -Butane	1.25	80	8.2	2.06	-6.75	-4.69	-7.43	-5.37	-7.14	-5.08
<i>cis</i> -Butene	3.99	74	8.0	1.37	-7.69	-6.31	-7.06	-5.69	-6.20	-4.83
<i>trans</i> -Butene	4.11	74	8.49	1.36	-5.98	-4.62	-5.90	-4.54	-5.66	-4.30
Isobutane	0.92	79	8.14	2.24	-6.13	-3.89	-8.09	-5.85	-7.40	-5.16
Isobutene	4.69	75	8.29	1.28	-6.57	-5.29	-6.72	-5.44	-6.32	-5.04
Neopentane	0.46	96	9.99	2.65	-9.21	-6.56			-8.19	-5.54
Cyclopentane	2.24	86	9.15	1.20	-6.58	-5.38	-8.33	-7.13	-7.22	-6.02
Cyclopentene	7.93	81	8.87 <sup>i</sup>	0.55	-7.39	-6.84	-7.02	-6.47	-6.00	-5.45
Cyclohexane	0.69	102	11.0	1.19	-6.57	-5.38			-8.4	-7.21
Benzene	22.79	89	10.7	-0.89	-7.13	-8.02			-5.77	-6.66
I <sub>2</sub>	0.13 <sup>i</sup>	71	10.3 <sup>j</sup>	-1.20	$-5.90 \pm 0.28$	-7.10	-8.38 <sup>k</sup>	-9.58	-6.82 <sup>k</sup>	-8.02

<sup>a</sup> From ref. 62. <sup>b</sup> Obtained from AM1-optimized structures by using the QSAR module of Hyperchem. <sup>c</sup> From ref. 19. <sup>d</sup> Calculated from the solubility ( $S$ ) and vapor pressure ( $p_{\text{vap}}$ ) according to  $\Delta G_{\text{hydr}} = -RT \ln(Sp^0/p_{\text{vap}}) - 1.90$  kcal mol<sup>-1</sup> with  $p^0 = 101.325$  kPa and  $p_{\text{vap}}$  in kPa. <sup>e</sup> Obtained from binding constants in Table 2; error  $\pm 0.10$  kcal mol<sup>-1</sup>, unless explicitly stated. <sup>f</sup>  $\Delta G'_{\text{a}} = \Delta G_{\text{a}} + \Delta G_{\text{hydr}}$ . <sup>g</sup> From ref. 20. <sup>h</sup> From ref. 19. <sup>i</sup> From ref. 63. <sup>j</sup> From ref. 64. <sup>k</sup> From ref. 52.

complexes, namely,  $\pi$ - $\pi$  interactions between the first guest and the second guest. Although the estimated cavity size of Bis2·CB8 is slightly less than that of CB6 (Table 1), it binds cyclohexane better than CB6 ( $6.6 \times 10^4$  versus  $<0.2 \times 10^4$  M<sup>-1</sup>), presumably because the ternary complex is somewhat more flexible and can adapt its lids to the encapsulated guest. However, it binds cyclohexane less tightly than CB7 ( $6.6 \times 10^4$  versus  $150 \times 10^4$  M<sup>-1</sup>), which can be attributed in this case to a tight packing (PC = 81% versus 72%). In contrast, benzene binds to the Bis2·CB8 system 10-times more strongly than to CB7.

To corroborate the preferential binding of unsaturated over saturated guests with a different method, we conducted isothermal titration calorimetry (ITC) experiments, which afforded additional thermochemical information (see ESI, Table S3 and Fig. S10<sup>†</sup>). This technique requires higher concentrations of the guest, which limited the accessible guest range; to remedy, we included two additional water-soluble derivatives (phenol and cyclopentanol), which set up an additional pair to probe for  $\pi$ - $\pi$  interactions. In general, guest binding to the Bis·CB8 binary complexes was found to be enthalpically driven by 8–12 kcal mol<sup>-1</sup>, with a negative entropic component (Fig. S10<sup>†</sup>), a signature which could be related to the non-classical hydrophobic effect (removal of high-energy water molecules from the cavity).<sup>38,39,54</sup> The binding constants for cyclopentene and benzene with Bis2·CB8 obtained by ITC and fluorescence displacement (see Table 2) showed satisfactory agreement, if one considers the known systematic variations in binding constants when different methods are being employed.<sup>23</sup> The higher binding affinity for phenol than cyclopentanol with the three Bis·CB8 systems confirmed  $\pi$ - $\pi$  interactions: although both guests have identical volume<sup>55</sup> and although the highly water-soluble phenol

should display a lower hydrophobic component to the driving force, it has a higher affinity to the Bis·CB8 receptors than the less water-soluble cyclopentanol (see also Table S4 in ESI<sup>†</sup>). This affinity pattern for the Bis·CB8 receptors (guests with phenyl residue binding more strongly than those with cyclopentyl ones) contrasts that observed early for CB6 (three order of magnitude lower binding of phenyl than of the corresponding cyclopentyl guests)<sup>20,55</sup> and later for CB7.<sup>19</sup>

The hydrocarbons in Table 2 cover a homologous series with a wide range of guest size and hydrophobicity, but without interference from electrostatic interactions (ion–ion, ion–dipole, dipole–dipole, and hydrogen-bonding).<sup>19</sup> In order to decouple the classical hydrophobic effect<sup>38,39</sup> associated with guest desolvation as a driving force for host–guest complexation, the hydration free energy of the guest – experimentally known for hydrocarbons – needs to be added to the experimental binding free energy.<sup>19,33</sup> This affords a guest-desolvation corrected value for the driving force ( $\Delta G'_{\text{a}} = \Delta G_{\text{a}} + \Delta G_{\text{hydr}}$ ) that reflects the intrinsic affinity of a particular hydrocarbon to Bis2·CB8, CB6, and CB7 (Table 3). For each host, the  $\Delta G'_{\text{a}}$  values increase roughly with the size/polarizability of the hydrocarbons (Table 3 and Fig. 4). This trend is reasonable as dispersion interactions<sup>56,57</sup> are expected to increase with guest size<sup>19,21</sup> as long as the PC of a guest does not become too large.<sup>24,28,58,59</sup>

A comparison of guest desolvation-corrected free binding energies ( $\Delta G'_{\text{a}}$ ) of the arene-loaded Bis·CB8 complexes with the equally large CB6 cavity is immediately instructive. This is because the interpretation of the driving force can be reduced to (i) direct host–guest interactions (dispersion,  $\pi$ - $\pi$ , C–H- $\pi$ , and cation- $\pi$  interactions) and (ii) the non-classical hydrophobic effect (removal of high-energy water).<sup>38,39</sup> Considering first the alkanes (and iodine), the  $\Delta G'_{\text{a}}$  values for CB6 are consistently higher (more negative) than the corresponding ones for







effects and direct molecular interactions, including dispersion and  $\pi$ - $\pi$  interactions.<sup>67-69,72</sup> The Bis-CB8 systems provide a solvent-shielded environment through the CB walls, while the imidazolium moieties act as 'lids' that close the CB carbonyl portals. To further expand the usage of Bis-CB8 systems as "intermolecular interaction chambers", see Fig. 3a, we plan to study alkyl, perfluoroalkyl, and substituted aryl linkers between the imidazolium units. It may well be that intermolecular interactions inside the chambers become more pronounced than in the corresponding molecular balances, due to more effective desolvation and forced proximity.

## Conclusions

In summary, we have shown that CB8, one of the largest members of the CB $n$  family, can be tailored towards selective hydrocarbon binding. This was achieved through the formation of aryl-bisimidazolium complexes, the purpose of which was three-fold. Firstly, the aromatic units are fluorescent, such that binding events can be directly monitored. Secondly, the auxiliaries restrict the available cavity space within CB8, which increases the packing coefficient and favours the complexation of small guests, including gases. Thirdly, the incorporation of a naphthyl unit between the imidazolium caps allows for sizable  $\pi$ - $\pi$  interactions with the encapsulated small guest molecules, which tips the selectivity towards alkenes and arenes.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

S. J. B. thanks the European Commission for a Marie Curie Fellowship (NANOSPHERE, 658360), O. A. S. and A. P. acknowledge an ERC Starting Investigator grant (ASPIRe240629) and EPSRC Programme Grant (NOtCH, EP/L027151/1) for the support. W. M. N. and K. I. A. are grateful to the DFG for grant NA-686/8 within the priority program SPP 1807 "Control of London Dispersion Interactions in Molecular Chemistry".

## Notes and references

- J.-R. Li, Y. Ma, M. C. McCarthy, J. Sculley, J. Yu, H.-K. Jeong, P. B. Balbuena and H.-C. Zhou, *Coord. Chem. Rev.*, 2011, **255**, 1791-1823.
- D. J. Wales, J. Grand, V. P. Ting, R. D. Burke, K. J. Edler, C. R. Bowen, S. Mintova and A. D. Burrows, *Chem. Soc. Rev.*, 2015, **44**, 4290-4321.
- D. N. Dybtsev, H. Chun, S. H. Yoon, D. Kim and K. Kim, *J. Am. Chem. Soc.*, 2004, **126**, 32-33.
- N. L. Rosi, J. Eckert, M. Eddaoudi, D. T. Vodak, J. Kim, M. O'Keeffe and O. M. Yaghi, *Science*, 2003, **300**, 1127-1129.
- J.-R. Li, R. J. Kuppler and H.-C. Zhou, *Chem. Soc. Rev.*, 2009, **38**, 1477-1504.
- C. R. Kim, T. Uemura and S. Kitagawa, *Chem. Soc. Rev.*, 2016, **45**, 3828-3845.
- J. Duan, W. Jin and S. Kitagawa, *Coord. Chem. Rev.*, 2017, **332**, 48-74.
- A. V. Leontiev, A. W. Saleh and D. M. Rudkevich, *Org. Lett.*, 2007, **9**, 1753-1755.
- S. Xiang, Y. He, Z. Zhang, H. Wu, W. Zhou, R. Krishna and B. Chen, *Nat. Commun.*, 2012, **3**, 954.
- Z. Zhang, Y. Zhao, Q. Gong, Z. Li and J. Li, *Chem. Commun.*, 2013, **49**, 653-661.
- H. Kim, Y. Kim, M. Yoon, S. Lim, S. M. Park, G. Seo and K. Kim, *J. Am. Chem. Soc.*, 2010, **132**, 12200-12202.
- J. Tian, S. Ma, P. K. Thallapally, D. Fowler, B. P. McGrail and J. L. Atwood, *Chem. Commun.*, 2011, **47**, 7626-7628.
- Y. Ruan, P. W. Peterson, C. M. Hadad and J. D. Badjic, *Chem. Commun.*, 2014, **50**, 9086-9089.
- K. A. Kellersberger, J. D. Anderson, S. M. Ward, K. E. Krakowiak and D. V. Dearden, *J. Am. Chem. Soc.*, 2001, **123**, 11316-11317.
- Y. Miyahara, K. Abe and T. Inazu, *Angew. Chem., Int. Ed.*, 2002, **41**, 3020-3023.
- B. S. Kim, Y. H. Ko, Y. Kim, H. J. Lee, N. Selvapalam, H. C. Lee and K. Kim, *Chem. Commun.*, 2008, 2756-2758.
- G. Huber, F.-X. Legrand, V. Lewin, D. Baumann, M.-P. Heck and P. Berthault, *ChemPhysChem*, 2011, **12**, 1053-1055.
- C. P. Kumar, F. Wu, C. E. Woodward and A. I. Day, *Supramol. Chem.*, 2014, **26**, 670-676.
- K. I. Assaf, M. Florea, J. Antony, N. M. Henriksen, J. Yin, A. Hansen, Z.-w. Qu, R. Sure, D. Klapstein, M. K. Gilson, S. Grimme and W. M. Nau, *J. Phys. Chem. B*, 2017, **121**, 11144-11162.
- M. Florea and W. M. Nau, *Angew. Chem., Int. Ed.*, 2011, **50**, 9338-9342.
- S. He, F. Biedermann, N. Vankova, L. Zhechkov, T. Heine, R. E. Hoffman, A. De Simone, T. T. Duignan and W. M. Nau, *Nat. Chem.*, 2018, **10**, 1252-1257.
- E. Masson, X. Ling, R. Joseph, L. Kyeremeh-Mensah and X. Lu, *RSC Adv.*, 2012, **2**, 1213-1247.
- S. J. Barrow, S. Kasera, M. J. Rowland, J. del Barrio and O. A. Scherman, *Chem. Rev.*, 2015, **115**, 12320-12406.
- K. I. Assaf and W. M. Nau, *Chem. Soc. Rev.*, 2015, **44**, 394-418.
- D. Shetty, J. K. Khedkar, K. M. Park and K. Kim, *Chem. Soc. Rev.*, 2015, **44**, 8747-8761.
- K. I. Assaf and W. M. Nau, *Supramol. Chem.*, 2014, **26**, 657-669.
- X. Lu and L. Isaacs, *Angew. Chem., Int. Ed.*, 2016, **55**, 8076-8080.
- W. M. Nau, M. Florea and K. I. Assaf, *Isr. J. Chem.*, 2011, **51**, 559-577.
- M. El Haouaj, M. Luhmer, Y. H. Ko, K. Kim and K. Bartik, *J. Chem. Soc., Perkin Trans. 2*, 2001, 804-807.
- G. Huber, F. X. Legrand, V. Lewin, D. Baumann, M. P. Heck and P. Berthault, *ChemPhysChem*, 2011, **12**, 1053-1055.
- M. Kunth, C. Witte, A. Hennig and L. Schröder, *Chem. Sci.*, 2015, **6**, 6069-6075.
- D. Jiao, F. Biedermann and O. A. Scherman, *Org. Lett.*, 2011, **13**, 3044-3047.
- R. Rabbani and E. Masson, *Org. Lett.*, 2017, **19**, 4303-4306.



- 34 S. Liu, A. D. Shukla, S. Gadde, B. D. Wagner, A. E. Kaifer and L. Isaacs, *Angew. Chem., Int. Ed.*, 2008, **47**, 2657–2660.
- 35 A. Shivanyuk, A. Scarso and J. Rebek Jr, *Chem. Commun.*, 2003, 1230–1231.
- 36 M. Yoshizawa, M. Tamura and M. Fujita, *J. Am. Chem. Soc.*, 2004, **126**, 6846–6847.
- 37 S. Mecozzi and J. Rebek Jr, *Chem.–Eur. J.*, 1998, **4**, 1016–1022.
- 38 F. Biedermann, V. D. Uzunova, O. A. Scherman, W. M. Nau and A. De Simone, *J. Am. Chem. Soc.*, 2012, **134**, 15318–15323.
- 39 F. Biedermann, W. M. Nau and H. J. Schneider, *Angew. Chem., Int. Ed.*, 2014, **53**, 11158–11171.
- 40 J.-Y. Ortholand, A. M. Z. Slawin, N. Spencer, J. F. Stoddart and D. J. Williams, *Angew. Chem., Int. Ed.*, 1989, **28**, 1394–1395.
- 41 C. A. Hunter and J. K. M. Sanders, *J. Am. Chem. Soc.*, 1990, **112**, 5525–5534.
- 42 S. L. Cockroft, C. A. Hunter, K. R. Lawson, J. Perkins and C. J. Urch, *J. Am. Chem. Soc.*, 2005, **127**, 8594–8595.
- 43 S. E. Wheeler and K. N. Houk, *J. Am. Chem. Soc.*, 2008, **130**, 10854–10855.
- 44 S. Grimme, *Angew. Chem., Int. Ed.*, 2008, **47**, 3430–3434.
- 45 C. R. Martinez and B. L. Iverson, *Chem. Sci.*, 2012, **3**, 2191–2201.
- 46 F. Biedermann and O. A. Scherman, *J. Phys. Chem. B*, 2012, **116**, 2842–2849.
- 47 L.-J. Riwar, N. Trapp, B. Kuhn and F. Diederich, *Angew. Chem., Int. Ed.*, 2017, **56**, 11252–11257.
- 48 L. Yang, J. B. Brazier, T. A. Hubbard, D. M. Rogers and S. L. Cockroft, *Angew. Chem., Int. Ed.*, 2016, **55**, 912–916.
- 49 C. Bravin, G. Licini, C. A. Hunter and C. Zonta, *Chem. Sci.*, 2019, **10**, 1466–1471.
- 50 V. Martí-Centelles, A. L. Lawrence and P. J. Lusby, *J. Am. Chem. Soc.*, 2018, **140**, 2862–2868.
- 51 H.-J. Schneider, *Acc. Chem. Res.*, 2015, **48**, 1815–1822.
- 52 H. S. El-Sheshtawy, B. S. Bassil, K. I. Assaf, U. Kortz and W. M. Nau, *J. Am. Chem. Soc.*, 2012, **134**, 19935–19941.
- 53 T.-L. Neoh, Y. Noda, H. Yoshii and T. Furuta, *J. Inclusion Phenom. Macrocyclic Chem.*, 2006, **56**, 117–123.
- 54 F. Biedermann, M. Vendruscolo, O. A. Scherman, A. De Simone and W. M. Nau, *J. Am. Chem. Soc.*, 2013, **135**, 14879–14888.
- 55 C. Márquez, R. R. Hudgins and W. M. Nau, *J. Am. Chem. Soc.*, 2004, **126**, 5806–5816.
- 56 T. Liu and H. J. Schneider, *Angew. Chem., Int. Ed.*, 2002, **41**, 1368–1370.
- 57 G. Haberhauer, S. Woitschetzki and H. Bandmann, *Nat. Commun.*, 2014, **5**, 3542.
- 58 I<sub>2</sub> is an expected outlier in Fig. 4a, because of additional stabilizing interactions, namely halogen bonding, see ref. 52. The reason for *trans*-butene being an outlier for Bis2·CB8 in Fig. 4c is presumably due to specific steric interactions; note that *trans*-butene is the most elongated, rigid guest.
- 59 S. Löffler, A. Wuttke, B. Zhang, J. J. Holstein, R. A. Mata and G. H. Clever, *Chem. Commun.*, 2017, **53**, 11933–11936.
- 60 K. Fucke, K. M. Anderson, M. H. Filby, M. Henry, J. Wright, S. A. Mason, M. J. Gutmann, L. J. Barbour, C. Oliver, A. W. Coleman, J. L. Atwood, J. A. K. Howard and J. W. Steed, *Chem.–Eur. J.*, 2011, **17**, 10259–10271.
- 61 V. D. Uzunova, K. I. Assaf, A. I. Lazar, Y. Liu and W. M. Nau, *Supramol. Chem.*, 2016, **28**, 384–395.
- 62 W. M. Haynes, *CRC Handbook of Chemistry and Physics*, CRC Press, Boca Raton, FL, 2012.
- 63 R. W. Ramette and R. W. Sandford, *J. Am. Chem. Soc.*, 1965, **87**, 5001–5005.
- 64 G. Maroulis, C. Makris, U. Hohm and D. Goebel, *J. Phys. Chem. A*, 1997, **101**, 953–956.
- 65 While auxiliary intermolecular interactions (dispersion, C–H– $\pi$ ) with the imidazolium units will contribute to the overall complex stabilities, we do not invoke dominant  $\pi$ – $\pi$  or cation– $\pi$  interactions with the imidazolium rings, because the  $\pi$  systems of the unsaturated guests prefer, according to DFT calculations (see Table S5 and Fig. S50 in ESI<sup>†</sup>), to form  $\pi$ – $\pi$  stacking with the naphthalene ring in the Bis2·CB8 complexes.
- 66 V. Corne, A. M. Sarotti, C. Ramirez de Arellano, R. A. Spanevello and A. G. Suárez, *Beilstein J. Org. Chem.*, 2016, **12**, 1616–1623.
- 67 S. L. Cockroft and C. A. Hunter, *Chem. Commun.*, 2006, 3806–3808.
- 68 S. L. Cockroft and C. A. Hunter, *Chem. Soc. Rev.*, 2007, **36**, 172–188.
- 69 L. Yang, C. Adam, G. S. Nichol and S. L. Cockroft, *Nat. Chem.*, 2013, **5**, 1006.
- 70 S. Paliwal, S. Geib and C. S. Wilcox, *J. Am. Chem. Soc.*, 1994, **116**, 4497–4498.
- 71 J. Hwang, B. E. Dial, P. Li, M. E. Kozik, M. D. Smith and K. D. Shimizu, *Chem. Sci.*, 2015, **6**, 4358–4364.
- 72 F. R. Fischer, W. B. Schweizer and F. Diederich, *Chem. Commun.*, 2008, 4031–4033.
- 73 A. E. Aliev and W. B. Motherwell, *Chem.–Eur. J.*, 2019, **25**, 10516–10530.

