

Cite this: *Chem. Sci.*, 2019, 10, 10733

All publication charges for this article have been paid for by the Royal Society of Chemistry

The butterfly effect in bisfluorenylidene-based dihydroacenes: aggregation induced emission and spin switching†

Xiaodong Yin,^{ab} Jonathan Z. Low,^b Kealan J. Fallon,^b Daniel W. Paley^b and Luis M. Campos^{*b}

Linear acenes are a well-studied class of polycyclic aromatic hydrocarbons and their established physical properties have led to their widespread application across the field of organic electronics. However, their quinoidal forms – dihydroacenes – are much less explored and exhibit vastly different photophysical and electronic properties due to their non-planar, cross-conjugated nature. In this work, we present a series of difluorenylidene dihydroacenes which exhibit a butterfly-like structure with a quinoidal skeleton, resulting in comparatively higher optical gaps and lower redox activities than those of their planar analogs. We found that these compounds exhibit aggregation induced emission (AIE), activated through restriction of the “flapping” vibrational mode of the molecules in the solid state. Furthermore, anthracene-containing dihydroacenes exhibit thermally activated ground-state spin switching as evidenced by planarization of the acene core and diradical activity recorded by EPR. These two characteristics in this relatively unexplored class of materials provide new insights for the design of multifunctional materials.

Received 16th August 2019
Accepted 4th October 2019

DOI: 10.1039/c9sc04096j

rsc.li/chemical-science

Introduction

Polycyclic aromatic hydrocarbons (PAHs) are ubiquitous materials – they have been found in space¹ and have made their way into a variety of optoelectronic devices^{2,3} and strong composite materials.^{4,5} Oligoacenes are a subclass of PAHs that have sparked interest in organic electronics^{6,7} due to the realization of their potential as active materials in organic field effect transistors (OFETs),^{8,9} organic light-emitting diodes (OLEDs),^{10,11} and third-generation photovoltaics.^{12–16} In most of these systems, fine-tuning the chemistry of the core structure and intermolecular interactions is essential for stability and the overall performance of the materials. These molecules are generally flat and exhibit strong pi-stacking interactions that can be beneficial or detrimental for particular types of applications. For example, aggregation can impair optical emission, but it can be beneficial for directional charge transport.¹⁷ In this vein, contorted PAHs can be tuned for practical applications. Additionally, introducing substituents with steric hindrance and conformational degrees of freedom can drastically alter

their physical properties. For example, acenes with a butterfly structure were originally studied a couple of decades ago, and the butterfly structure can be changed to a flat structure by electrochemical oxidation.^{18–21}

Through precise molecular engineering of the core structure of oligoacenes, Tang and co-workers have shown that quinoidal derivatives of oligoacenes can exhibit aggregation induced restricted intramolecular vibration (RIV).^{22,23} AIE is counterintuitive since aggregation tends to quench photoluminescence of organic chromophores.^{24–26} For example, bis(diphenylmethylene)dihydroacenes are non-emissive in solution, but upon aggregation, they fluoresce in the solid state (Fig. 1A). This property does not translate to the fluoreno-quinoidal oligomers. In fact, fluoreno adducts are prone to stabilizing the radical character, as elegantly demonstrated by Wudl,²⁷ Wu,²⁸ Haley,²⁹ and others.^{30,31} The fluoreno-derivative of Chichibabin's radical (Fig. 1B) is non-emissive; however, in its most stable structure, the diradical is readily characterized. This diradical character is not surprising, given that extending anthracene oligomers along the 9,10-positions disrupts pi-orbital overlap, thus forcing aromatization of the central ring and yielding the stable open-shell fluorenyl groups shown in Fig. 1B as the most favorable resonance structure. Such phenomena motivated our interest in investigating the structure–property relationship, extending the oligoacene core with bisfluoreno adducts. Interestingly, we found that the bisfluorenoanthracene in Fig. 1C exhibits both AIE in the solid state and the thermally accessible open-shell diradicaloid character.

^aBeijing Key Laboratory of Photoelectronic/Electrophotonic Conversion Materials, School of Chemistry and Chemical Engineering, Beijing Institute of Technology, Beijing 102488, P. R. China

^bDepartment of Chemistry, Columbia University, New York, New York 10027, USA. E-mail: lcampos@columbia.edu

† Electronic supplementary information (ESI) available. CCDC 1947366–1947368. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc04096j



The main focus of this study lies in understanding the fine interplay between the optical and magnetic properties by varying the aromaticity of the bisfluoreno-quinoidal core through the number of fused aromatic rings (Fig. 1C), **bFA** (anthracene core), **bFT** (tetracene) and **bFP** (pentacene). Surprisingly, we found that these compounds exhibit properties of both systems described above – AIE and switchable diradical character. The AIE fluorescence quantum yield was found to be up to 180 times higher for **bFP** in the solid-state than in solution. While AIE typically occurs due to RIR around the connecting bond of the flanking phenyl rings and the polyaromatic hydrocarbon core in aggregates or in the solid,^{24,32} the fluoreno-quinoidal molecules shown have no such degrees of freedom. These materials provide blueprints for a new class of luminogens where AIE arises solely from RIV of the strained butterfly-like structures. Moreover, the aromaticity of the quinoidal core is essential to drive the switching between low-spin (quinoid) and high-spin (planar oligoacene diradical) states. Notably, whilst spin switching is widely studied in transition metal complexes for applications in memory devices,^{33–38} pure hydrocarbon organic compounds with such properties are less common.³⁹

Results and discussion

Bisfluoreno-oligoacenes were synthesized from their corresponding acene-diones using the Corey–Fuchs reaction to generate *para*-bis(dibrominated) species,⁴⁰ which were then reacted with 9,9-dimethyl-9-stannafluorene *via* Stille coupling to yield **bFA** (anthracene), **bFT** (tetracene) and **bFP** (pentacene) as colorless or light yellow crystals (see the ESI† for details). Single crystals of the three compounds were obtained by slow evaporation in a solution of chloroform/hexane (1 : 4 v/v), and the structures obtained by XRD are shown in Fig. 1C. The topology of all the molecules resembles butterfly shapes, with both fluorenes puckered on the same side.

The planes of the two fluorene moieties form a dihedral angle of 76.7° for **bFA**, 73.5° for **bFT** and 86.4° for **bFP**, respectively. The acene backbones of the molecules are bent at angles of 113.9°, 113.0° and 112.1° at the central ring for **bFA**, **bFT** and **bFP**, respectively. The bending angles of the acenes in this series are all more acute than those of the tetraphenyl derivatives, such as PDHA (Fig. 1), which bend at ~132°. This is attributed to the increase in steric bulk of the fluorene moiety.²³ As shown in Fig. S2,† the acene backbones also display a typical quinoidal structure – for example in **bFA**, C1–C15 and C8–C28 have double bond character (~1.35 Å) while C1–C2, C1–C14, C8–C7, and C8–C9 have single bond character (~1.49 Å). The molecules pack as dimers in the solid state where one fluorene group from each molecule engages in π - π stacking with the adjacent molecule at an average interplanar distance of ~4.0 Å. The central fluorene groups in the dimer form a herringbone configuration, with a close CH- π interaction, ~2.6 Å. The optimized gas phase structure of **bFA** determined by DFT (B3LYP/6-31G**) is similar to the crystal structure, except for a widening of the dihedral angle between the two fluorenes to 97.9°, which can be assigned to the intermolecular interactions




Fig. 1 Schematic illustration of (A) Tang's work on aggregation induced emission properties of tetraphenyl dihydroanthracene (PDHA); (B) Wu's work on the diradical character of fluorene-capped Chichibabin's hydrocarbon; (C) AIE/spin dual mode switching properties of bisfluorenylidene dihydroanthracene (**bFA**) in this work.

in closely packed crystals. However, the bending angle of the central anthracene ring calculated to be 111.1° matches well with the crystallographic data.

During the synthesis and characterization of the molecules, we noticed remarkably bright fluorescent spots by thin-layer chromatography. Thus, we sought to probe the optical properties of **bFA**, **bFT** and **bFP** in solution (CH₂Cl₂). The results are summarized in Table 1 and Fig. S3.† All three compounds exhibit similar UV-Vis absorption spectra with an onset of ~400 nm and maxima located at ~350 nm. Unexpectedly, there is a slight blue shift in the absorption onset as the acene skeleton is extended from anthracene to pentacene, which was further confirmed by DFT calculations (B3LYP/6-31G**, see the ESI†). The electronic transition of the absorption maxima can be assigned to HOMO → LUMO (S₀ → S₁) for **bFA**, HOMO–1 → LUMO (S₀ → S₂) for **bFT** and HOMO–1 → LUMO and HOMO–1 → LUMO+1 (S₀ → S₂ and S₀ → S₃) transitions for **bFP**, respectively. Thus, we deduce that the blue shift in the spectra results from transitions to higher excited states in **bFP**.

In solutions of dichloromethane and THF, all three compounds exhibit weak fluorescence, with quantum yields below 1% (Table 1). However, when an antisolvent (H₂O) is





Fig. 2 (A) Photographic image of quartz vials containing difluorenylidene acenequinoids displaying fluorescence in THF/H₂O solutions with different water fractions irradiated using a 365 nm UV lamp; (B) UV-Vis and photoluminescence spectra of bFT in THF and THF/water = 1/9; (C) plots of quantum yield as a function of water fraction with excitation at 380 nm; the inset is the aggregation induced emission efficiency ($\Phi_{\text{aggr}}/\Phi_{\text{sol}}$); absolute quantum yields of these samples were determined with an integrating sphere (Horiba Fluorolog-3).

of the rotation (ΔG^\ddagger) in bFA-t is 17.3 kcal mol⁻¹ (Fig. S11[†]). In contrast to bFA, neither bFT nor bFP show line broadening of the proton signals as the temperature is increased; rather, the



Fig. 3 (a) Optimized structures of the ground state (S_0 , red) and excited state (S_1 , blue) conformations of bFP in the gas and crystal state with the B3LYP/6-31G** and ONIOM (B3LYP/6-31G** : UFF) methods, respectively; (b) the ONIOM model, with the high layer presented as ball-and-stick molecules and the low layer presented as wireframe molecules.



Fig. 4 (a) VT-¹H NMR of bFA in tetrachloroethane-d₂; (b) EPR data of bFA in the solid state at 300 K (black) and 450 K (red) and in dichlorobenzene (4 × 10⁻² M) at 430 K (blue); inset: spin density illustration of the T₁ state of bFA_{twist} (twisted configuration) at the UM06L/6-311G** level of theory; (c) VT-¹H NMR of bFA-t and the molecular structure of bFA-t.

signals are sharpened, and no signal of EPR was observed (see Fig. S12[†]), indicating that these molecules do not have a thermally accessible open-shell ground-state configuration. Besides, the barrier to rotation (ΔG^\ddagger) for bFT is estimated to be ca. 21.7 kcal mol⁻¹ by the NMR data (Fig. S13[†]), which is much higher than that of bFA-t.

It is noteworthy to report our observation of a diprotonated derivative of bFA-t (bFAH-t) with strong blue fluorescence as the main product of the microwave-assisted Stille coupling reaction at ~180 °C (see Fig. S14[†]). This is strong evidence of H-abstraction of a diradical intermediate of bFA-t at high temperature.^{46,47} The UV-Vis absorbance of bFAH-t indicates that the core of this molecule has an aromatic anthracene structure with a finely split peak at ca. 400 nm. Moreover, this





Fig. 5 (a) Energy diagram of **bFA** with rotation of the fluorene moiety; (b) HOMOs of the fluorene twisted structure of **bFA** ($\mathbf{bFA}_{\text{twist}}$), **bFT** ($\mathbf{bFT}_{\text{twist}}$), and **bFP** ($\mathbf{bFP}_{\text{twist}}$), with the y_0 value and bond length (*) of the C–C bond between fluorene and acenes.

diprotonated compound can be converted back to the quinoid structure by oxidation with DDQ, as shown in Fig. S15.†

In order to understand the spin switching process, we computed the stationary points and transition state for **bFA**, as it changed into the open-shell structure from the closed-shell ground state (DFT, UB3LYP/6-31G**, Fig. 5A). As the dihedral angle joining fluorene to anthracene was planarized, an energy saddle point was observed in the conformation where one fluorene is twisted relative to anthracene whilst the other one is not. The theoretical energy barrier of 18 kcal mol⁻¹ compares well with the experimental result of deuterium-NMR of **bFA-t** with the same aromatic core (see Fig. S11†). When both fluorenes are twisted, a relatively stable state is formed with a comparable energy level (+3 kcal mol⁻¹) to that of the original state. Single point energy calculations of both the single twisted structure and double twisted structure were conducted at the UM06L/6-311G** level of theory.

The biradical character of organic materials can be assessed by the biradical character index (y_0) and indicates the

contribution of the open-shell resonance form to the overall ground state structure, ranging from a pure biradical ($y_0 = 1$) to a purely closed shell ($y_0 = 0$).^{48–51} For **bFA**, y_0 values of 0 and 0.16 were obtained for the single twisted structure and double twisted structure, respectively, indicating the open-shell character of the double twisted structure ($\mathbf{bFA}_{\text{twist}}$) as shown in Fig. 5b. In addition, the theoretical C–C double bond length in $\mathbf{bFA}_{\text{twist}}$ between the fluorene and anthracene moieties shows a remarkable extension (1.44 Å) compared to that in the butterfly configuration (1.36 Å), indicating the open shell character from a structural point of view. Crucially, the triplet state of $\mathbf{bFA}_{\text{twist}}$ is calculated to be lower in energy than the diradical singlet state (as shown in Fig. 5), indicating that intersystem crossing is thermodynamically favorable in this compound. The spin density of the T₁ state of $\mathbf{bFA}_{\text{twist}}$ is mainly distributed on the fluorene moiety, especially on carbons a and b, which match very well with the signal broadening observed by VT-¹H NMR (see Fig. 4b). Theoretical analysis was also conducted for the twisted structures of **bFT** ($\mathbf{bFT}_{\text{twist}}$) and **bFP** ($\mathbf{bFP}_{\text{twist}}$), and both of them exhibit a closed shell ground state electronic structure ($y_0 = 0$, see Fig. 5b). These calculations match well with our experimental observations. Furthermore, the carbon–carbon bond length between fluorene and acenes of $\mathbf{bFT}_{\text{twist}}$ and $\mathbf{bFP}_{\text{twist}}$ is ca. 1.41 Å which is more comparable to a typical double bond, indicating closed-shell quinoidal structures.

From this analysis we conclude that, at elevated temperatures, aromatization is a strong enough driving force to access the open shell ground state in the anthraquinoid **bFA**, whereas the longer acenequinoid structures in **bFT** and **bFP** do not possess sufficient aromatic character in their aromatic conformation to exhibit this trait. This finding is supported by Clar's theory of aromatic sextets, where higher order acene molecules exhibit greater stability as two small acenes *versus* one large acene system (e.g. 2 × naphthalene *vs.* 1 × pentacene for **bFP**).^{52,53} Besides, since the open-shell character of **bFA** is strongly related to the rotation of fluorene moieties, the weak EPR signal in the solid state even at a high temperature can be assigned to the constriction of intramolecular motion.

Conclusions

In summary, we synthesized a series of difluorenylidene dihydroacene compounds using a facile and efficient method. These compounds exhibit wider optical gaps than expected due to their bent, butterfly-like quinoid structures. All compounds exhibit aggregation induced emission, which is rarely seen in such a kind of structure without any freely rotatable functionalities. A flapping intramolecular vibration was identified as the mechanism based on theoretical analysis (QM/MM calculations). Furthermore, a transformation between the bent quinoid butterfly configuration and the aromatic planar configuration is thermally accessible in the anthracene-containing material. As a result, the electronic structure is also switchable between the closed shell at a low temperature and open shell at a high temperature, which is confirmed by VT-¹H NMR and EPR. This phenomenon can be assigned to



rotation of the fluorene moieties at a high temperature, leading to an intermediate structure with a flattened anthracene skeleton and moderate diradical character. Conversely, this transition is not thermally accessible in longer acenequinoids, due to the lower driving force as a function of the lower aromaticity of the aromatic conformation of these acenes. We envisage that the structure–property relationship reported herein will create new pathways for designing novel multi-functional materials.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was funded by the National Science Foundation (DMR-1807580). K. J. F. acknowledges the funding by the US-UK Fulbright Commission and Lloyd's Tercentenary Research Foundation. Single crystal X-ray diffraction was performed at the Shared Materials Characterization Laboratory at Columbia University. Use of the SMCL was made possible by funding from Columbia University.

Notes and references

- 1 F. J. Lovas, R. J. McMahon, J. U. Grabow, M. Schnell, J. Mack, L. T. Scott and R. L. Kuczkowski, *J. Am. Chem. Soc.*, 2005, **127**, 4345–4349.
- 2 J. Wu, W. Pisula and K. Müllen, *Chem. Rev.*, 2007, **107**, 718–747.
- 3 A. Narita, X. Y. Wang, X. Feng and K. Müllen, *Chem. Soc. Rev.*, 2015, **44**, 6616–6643.
- 4 Z. Qin, G. S. Jung, M. J. Kang and M. J. Buehler, *Sci. Adv.*, 2017, **3**, e1601536.
- 5 C. Lee, X. Wei, J. W. Kysar and J. Hone, *Science*, 2008, **321**, 385–388.
- 6 J. E. Anthony, *Chem. Rev.*, 2006, **106**, 5028–5048.
- 7 J. E. Anthony, *Angew. Chem., Int. Ed.*, 2008, **47**, 452–483.
- 8 K. Ito, T. Suzuki, Y. Sakamoto, D. Kubota, Y. Inoue, F. Sato and S. Tokito, *Angew. Chem., Int. Ed.*, 2003, **115**, 1191–1194.
- 9 A. Naibi Lakshminarayana, A. Ong and C. Chi, *J. Mater. Chem. C*, 2018, **6**, 3551–3563.
- 10 L. C. Picciolo, H. Murata and Z. H. Kafafi, *Appl. Phys. Lett.*, 2001, **78**, 2378–2380.
- 11 S. A. Odom, S. R. Parkin and J. E. Anthony, *Org. Lett.*, 2003, **5**, 4245–4248.
- 12 M. B. Smith and J. Michl, *Chem. Rev.*, 2010, **110**, 6891–6936.
- 13 S. N. Sanders, E. Kumarasamy, A. B. Pun, M. T. Trinh, B. Choi, J. Xia, E. J. Taffet, J. Z. Low, J. R. Miller, X. Roy, X. Y. Zhu, M. L. Steigerwald, M. Y. Sfeir and L. M. Campos, *J. Am. Chem. Soc.*, 2015, **137**, 8965–8972.
- 14 E. Kumarasamy, S. N. Sanders, A. B. Pun, S. A. Vaselabadi, J. Z. Low, M. Y. Sfeir, M. L. Steigerwald, G. E. Stein and L. M. Campos, *Macromolecules*, 2016, **49**, 1279–1285.
- 15 S. N. Sanders, E. Kumarasamy, A. B. Pun, K. Appavoo, M. L. Steigerwald, L. M. Campos and M. Y. Sfeir, *J. Am. Chem. Soc.*, 2016, **138**, 7289–7297.
- 16 M. Einzinger, T. Wu, J. F. Kompalla, H. L. Smith, C. F. Perkinson, L. Nienhaus, S. Wieghold, D. N. Congreve, A. Kahn, M. G. Bawendi and M. A. Baldo, *Nature*, 2019, **571**, 90–94.
- 17 V. C. Sundar, J. Zaumseil, V. Podzorov, E. Menard, R. L. Willett, T. Someya, M. E. Gershenson and J. A. Rogers, *Science*, 2004, **303**, 1644–1646.
- 18 N. Martin, L. Sánchez, C. Seoane, E. Ortí, P. M. Viruela and R. Viruela, *J. Org. Chem.*, 1998, **63**, 1268–1279.
- 19 M. C. Díaz, B. M. Illescas, N. Martín, R. Viruela, P. M. Viruela, E. Ortí, O. Brede, I. Zilbermann and D. M. Guldi, *Chem.–Eur. J.*, 2004, **10**, 2067–2077.
- 20 N. E. Gruhn, N. A. Macías-Ruvalcaba and D. H. Evans, *Langmuir*, 2006, **22**, 10683–10688.
- 21 S. L. Broman, C. L. Andersen, T. Jousselein-Oba, M. Mansø, O. Hammerich, M. Frigoli and M. B. Nielsen, *Org. Biomol. Chem.*, 2017, **15**, 807–811.
- 22 Z. He, L. Shan, J. Mei, H. Wang, J. W. Y. Lam, H. H.-Y. Sung, I. D. Williams, X. Gu, Q. Miao and B. Z. Tang, *Chem. Sci.*, 2015, **6**, 3538–3543.
- 23 Z. He, L. Zhang, J. Mei, T. Zhang, J. W. Y. Lam, Z. Shuai, Y. Q. Dong and B. Z. Tang, *Chem. Mater.*, 2015, **27**, 6601–6607.
- 24 J. Mei, N. L. C. Leung, R. T. K. Kwok, J. W. Y. Lam and B. Z. Tang, *Chem. Rev.*, 2015, **115**, 11718–11940.
- 25 Z. Zhao, H. Nie, C. Ge, Y. Cai, Y. Xiong, J. Qi, W. Wu, R. T. K. Kwok, X. Gao, A. Qin, J. W. Y. Lam and B. Z. Tang, *Adv. Sci.*, 2017, **4**, 1700005.
- 26 D. Wang, M. M. S. Lee, G. Shan, R. T. K. Kwok, J. W. Y. Lam, H. Su, Y. Cai and B. Z. Tang, *Adv. Mater.*, 2018, **30**, 1802105.
- 27 X. Gong, M. Tong, F. G. Brunetti, J. Seo, Y. Sun, D. Moses, F. Wudl and A. J. Heeger, *Adv. Mater.*, 2011, **23**, 2272–2277.
- 28 Z. Zeng, Y. M. Sung, N. Bao, D. Tan, R. Lee, J. L. Zafra, B. S. Lee, M. Ishida, J. Ding, J. T. López Navarrete, Y. Li, W. Zeng, D. Kim, K. W. Huang, R. D. Webster, J. Casado and J. Wu, *J. Am. Chem. Soc.*, 2012, **134**, 14513–14525.
- 29 G. E. Rudebusch, J. L. Zafra, K. Jorner, K. Fukuda, J. L. Marshall, I. Arrechea-Marcos, G. L. Espejo, R. Ponce Ortiz, C. J. Gómez-García, L. N. Zakharov, M. Nakano, H. Ottosson, J. Casado and M. M. Haley, *Nat. Chem.*, 2016, **8**, 753–759.
- 30 D. Beaudoin, O. Levasseur-Grenon, T. Maris and J. D. Wuest, *Angew. Chem., Int. Ed.*, 2016, **55**, 894–898.
- 31 K. Cocq, V. Maraval, N. Saffon-Merceron, A. Saquet, C. Poidevin, C. Lepetit and R. Chauvin, *Angew. Chem., Int. Ed.*, 2015, **54**, 2703–2706.
- 32 M. E. Howe and M. A. Garcia-Garibay, *J. Org. Chem.*, 2019, **84**, 9570–9576.
- 33 F. Schwarz, G. Kastlunger, F. Lissel, C. Egler-Lucas, S. N. Semenov, K. Venkatesan, H. Berke, R. Stadler and E. Lörtscher, *Nat. Nanotechnol.*, 2015, **11**, 170–176.
- 34 O. Kahn and C. Jay Martinez, *Science*, 1998, **279**, 44–48.
- 35 M. Mannini, F. Pineider, P. Sainctavit, C. Danieli, E. Otero, C. Sciancalepore, A. M. Talarico, M. Arrio, A. Cornia, D. Gatteschi and R. Sessoli, *Nat. Mater.*, 2009, **8**, 194–197.
- 36 T. Matsumoto, G. N. Newton, T. Shiga, S. Hayami, Y. Matsui, H. Okamoto, R. Kumai, Y. Murakami and H. Oshio, *Nat. Commun.*, 2014, **5**, 3865.



- 37 A. Bouseksou, G. Molnár and G. Matouzenko, *Eur. J. Inorg. Chem.*, 2004, 4353–4369.
- 38 P. Gütllich, A. Hauser and H. Spiering, *Angew. Chem., Int. Ed.*, 1994, **33**, 2024–2054.
- 39 M. R. Geraskina, A. T. Buck and A. H. Winter, *J. Org. Chem.*, 2014, **79**, 7723–7727.
- 40 C. Y. Chiu, B. Kim, A. A. Gorodetsky, W. Sattler, S. Wei, A. Sattler, M. Steigerwald and C. Nuckolls, *Chem. Sci.*, 2011, **2**, 1480–1486.
- 41 N. L. C. Leung, N. Xie, W. Yuan, Y. Liu, Q. Wu, Q. Peng, Q. Miao, J. W. Y. Lam and B. Z. Tang, *Chem.–Eur. J.*, 2014, **20**, 15349–15353.
- 42 Z. Zhao, X. Zheng, L. Du, Y. Xiong, W. He, X. Gao, C. Li, Y. Liu, B. Xu, J. Zhang, F. Song, Y. Yu, X. Zhao, Y. Cai, X. He, R. T. K. Kwok, J. W. Y. Lam, X. Huang, D. Lee Phillips, H. Wang and B. Z. Tang, *Nat. Commun.*, 2019, **10**, 2952.
- 43 Z. Zhao, S. Chen, X. Shen, F. Mahtab, Y. Yu, P. Lu, J. W. Y. Lam, H. S. Kwok and B. Z. Tang, *Chem. Commun.*, 2010, **46**, 686–688.
- 44 M. Desroches, P. M. Burrezo, J. Boismenu-lavoie, M. Peça, J. G. Carlos, J. M. Matxain, D. Casanova, J. Morin and J. Casado, *Angew. Chem., Int. Ed.*, 2017, **56**, 16212–16217.
- 45 X. Lu, S. Lee, J. O. Kim, T. Y. Gopalakrishna, H. Phan, T. S. Heng, Z. Lim, Z. Zeng, J. Ding, D. Kim and J. Wu, *J. Am. Chem. Soc.*, 2016, **138**, 13048–13058.
- 46 H. Sakurai, H. Tobita, M. Kira and Y. Nakadaira, *Angew. Chem., Int. Ed.*, 1980, **19**, 620.
- 47 C. Wentrup, M. J. Regimbald-Krnel, D. Müller and P. Comba, *Angew. Chem., Int. Ed.*, 2016, **55**, 14600–14605.
- 48 G. E. Rudebusch, J. L. Zafra, K. Jorner, K. Fukuda, J. L. Marshall, I. Arrechea-Marcos, G. L. Espejo, R. Ponce Ortiz, C. J. Gómez-García, L. N. Zakharov, M. Nakano, H. Ottosson, J. Casado and M. M. Haley, *Nat. Chem.*, 2016, **8**, 753–759.
- 49 L. K. Montgomery, J. C. Huffman, E. A. Jurczak and M. P. Grendze, *J. Am. Chem. Soc.*, 1986, **108**, 6004–6011.
- 50 M. Abe, *Chem. Rev.*, 2013, **113**, 7011–7088.
- 51 W. Zeng, Z. Sun, T. S. Heng, T. P. Gonçalves, T. Y. Gopalakrishna, K.-W. Huang, J. Ding and J. Wu, *Angew. Chem., Int. Ed.*, 2016, **55**, 8615–8619.
- 52 E. Clar, in *Mobile Source Emissions Including Polycyclic Organic Species*, ed. D. Rondia, M. Cooke and R. K. Haroz, Springer Netherlands, Dordrecht, 1983, pp. 49–58.
- 53 M. Solà, *Front. Chem.*, 2013, **1**, 22.

