

PAPER

[View Article Online](#)
[View Journal](#) | [View Issue](#)

Cite this: *Dalton Trans.*, 2023, **52**, 5496

Received 9th February 2023,
Accepted 28th March 2023

DOI: 10.1039/d3dt00425b

rsc.li/dalton

Substitution lability of the perfluorinated Cp* ligand in Rh(I) complexes†

Joshua Parche, Susanne M. Rupf,  Robin Sievers and Moritz Malischewski *

Several cationic rhodium(I) complexes $[\text{Rh}(\text{COD})\text{L}_2][\text{C}_5(\text{CF}_3)_5]^-$ have been synthesized through substitution of the weakly bound $[\text{C}_5(\text{CF}_3)_5]^-$ ligand from $[\text{Rh}(\text{COD})(\text{C}_5(\text{CF}_3)_5)]$, further emphasizing its unique reactivity. Besides acetonitrile, pyridine derivatives with varying degrees of fluorination have been employed as ligands in order to investigate the influence of fluorination upon the binding affinity towards the resulting $[\text{Rh}(\text{COD})]^+$ fragment and the limit as to which the $[\text{C}_5(\text{CF}_3)_5]^-$ ligand can be displaced. Furthermore, the newly synthesized compounds represent rare examples of rhodium complexes containing fluorinated pyridines as ligands.

Introduction

Fluorinated ligands, in particular F- or CF_3 -substituted aromatics, have been gaining increasing attention in organometallic chemistry over the last years. The electron withdrawing effect of fluorine may lead to a decreased bonding strength of fluorinated ligands towards metal centers in comparison to their electron rich counterparts. For example, η^6 -fluorobenzenes coordinated at rhodium(I) centers have been shown to have lower binding affinity with increasing degree of fluorination.¹ These comparably weak binding interactions can be of great advantage in catalysis, for systems proceeding *via* an inner-sphere mechanism that requires vacant coordination sites for substrate binding. Due to the high reactivity and inherent instability of coordinatively unsaturated transition metal complexes, their vacant sites can be masked by labile ligands, which stabilize the respective complex while being easily displaced and not interfering with the catalytic process.^{1,2} Furthermore, the strong carbon–fluorine bond is relatively inert towards potential side-reactions such as oxidative addition, while also decreasing the probability of coordination through the electron-withdrawing group itself (*e.g.* cyano groups).^{3,4} In this sense, Weller *et al.* have demonstrated the synthesis of catalytically active rhodium(I) complexes containing a η^6 -monofluorobenzene ligand, prepared by using a simple one-pot procedure starting from $[\text{Rh}(\text{COD})_2][\text{BAR}^{\text{F}}_4]$ (COD = cycloocta-1,5-diene, $\text{Ar}^{\text{F}} = \text{C}_6\text{H}_3-(3,5-\text{CF}_3)_2$).⁵ The

respective bench-stable complexes $[\text{Rh}(\text{C}_6\text{H}_5\text{F})(\text{R}_2\text{PCH}_2\text{PR}'_2)][\text{BAR}^{\text{F}}_4]$ ($\text{R}, \text{R}' = \text{'Bu}$ or Cy) have been proven as efficient precatalysts for various reactions, including intermolecular hydroacylations and dehydrogenative couplings of substituted amine–boranes.^{5–10} In addition, similar rhodium(I) complexes containing weaker coordinating 1,2-difluorobenzene or 1,2,3-trifluorobenzene ligands allowed the isolation and structural characterization of previously inaccessible amine–borane σ -complexes, *via* displacement of the weakly coordinating ligands.^{11,12} Interestingly, the further reduced coordinating ability of 1,2,3,4-tetra- and pentafluorobenzene led to the π -complexation of the weakly coordinating anion (WCA) $[\text{BAR}^{\text{F}}_4]^-$ in solution, forming zwitterionic complexes $[\text{Rh}(\text{R}_2\text{PCH}_2\text{PR}'_2)_2(\eta^6-(3,5-(\text{CF}_3)_2\text{C}_6\text{H}_3)\text{BAR}^{\text{F}}_3)]$ (Fig. 1, top).¹² In order to access these complexes containing highly fluorinated benzenes, the use of $[\text{Al}(\text{OR}^{\text{F}})_4]^-$, popularized by the group of Krossing, proved to be crucial to avoid displacement of the respective fluorobenzene.^{13,14} Complexes containing highly fluorinated benzenes were shown to effectively catalyze the Tishchenko reaction of cyclohexanecarboxaldehyde ($\text{C}_6\text{H}_{11}\text{CHO}$), while leading to shorter reaction times for




Fig. 1 Examples of rhodium(I) complexes with fluorinated aromatic ligands.

Freie Universität Berlin, Institute of Inorganic Chemistry, Fabeckstr. 34-36, 14195 Berlin, Germany. E-mail: moritz.malischewski@fu-berlin.de

†Electronic supplementary information (ESI) available: General conditions. Copies of NMR spectra. DFT calculations. CCDC 2240960 and 2240962. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3dt00425b>

Dalton Trans., 2023, 52, 5496–5502 | 5497




Fig. 2 Investigated substitutions of $[C_5(CF_3)_5]^-$ in $[Rh(COD)(C_5(CF_3)_5)]$ with different ligands.

signals, in analogy to the substitution by toluene.²⁹ The coordinated $[C_5(CF_3)_5]^-$ ligand shows a singlet at $\delta = -51.1$ ppm in $CDCl_3$ (or $\delta = -51.46$ ppm in CD_2Cl_2), while the uncoordinated $[C_5(CF_3)_5]^-$ possesses a downfield shifted signal at $\delta = -50.18$ ppm in $CDCl_3$ (or $\delta = -50.52$ ppm in CD_2Cl_2). Integration of the respective signals allows to observe the ratio of coordinated to non-coordinated $[C_5(CF_3)_5]^-$. Any unreacted $[Rh(COD)(C_5(CF_3)_5)]$ can be removed from cationic complexes by washing the respective mixture with *n*-pentane.

First, the substitution of the perfluorinated Cp* from $[Rh(COD)(C_5(CF_3)_5)]$ in MeCN was investigated, in order to assess the displacement of $[C_5(CF_3)_5]^-$ by two N-donor ligands. It was found that stirring the reaction for 1 h at room temperature, led to the complete substitution of the Cp ligand, turning it into a WCA and quantitatively forming the isolatable cationic 16-electron complex $[Rh(COD)(MeCN)_2][C_5(CF_3)_5]$ (Fig. 3). The ^{19}F NMR spectrum revealed a singlet at $\delta = -51.1$ ppm in $CDCl_3$, which is in agreement with the signals of $[Rh(COD)$

$(PhMe)][C_5(CF_3)_5]$.²⁹ In addition, a downfield shifted singlet was observed in the 1H NMR spectrum for the MeCN ligands ($\delta = 2.20$ ppm in CD_2Cl_2) in comparison to free MeCN ($\delta = 1.97$ ppm in CD_2Cl_2). Integration of the MeCN signals revealed a twofold coordination, when referencing to the COD signals.

Furthermore, it was found that the backreaction to $[Rh(COD)(C_5(CF_3)_5)]$ proceeds rapidly, in comparison to the back reaction of $[Rh(COD)(PhMe)][C_5(CF_3)_5]$. Within 30 min after dissolving $[Rh(COD)(MeCN)_2][C_5(CF_3)_5]$ in $CDCl_3$ (15 mg mL^{-1}), both signals of substituted as well as coordinated $[C_5(CF_3)_5]^-$ are observed with a ratio of 4:1 respectively (Fig. 3). The equilibrium of the backreaction is strongly dependent on the concentration and is reached within 1 h. For a concentration of 2 mg mL^{-1} of $[Rh(COD)(MeCN)_2][C_5(CF_3)_5]$ in $CDCl_3$, an equilibrium of cationic to neutral complex of 1:2, respectively, is observed. Under strongly diluted conditions (0.5 mg mL^{-1}) $[Rh(COD)(MeCN)_2][C_5(CF_3)_5]$ can be fully reacted to $[Rh(COD)(C_5(CF_3)_5)]$. Dissolving the substituted complex in an increasingly polar solvent like CD_2Cl_2 led to a significantly less pronounced backreaction, reaching an equilibrium of 10:1 of cationic to neutral complex at a comparable concentration as in Fig. 3 (15 mg mL^{-1}).

Single crystals of $[Rh(COD)(MeCN)_2][C_5(CF_3)_5]$ in the monoclinic $P2_1/n$ space group were obtained from solvent mixtures of CH_2Cl_2 and *n*-pentane by slowly cooling to -75 °C (Fig. 4). The crystal structure revealed an averaged Rh–N bond length of 2.057(7) Å and an average Rh–C distance to the COD ligand of 2.121(7) Å, comparable to the bond lengths of $[Rh(COD)(MeCN)_2][BF_4]$, with average distances of 2.080(9) Å and 2.125(11) Å respectively.³³

The substitution of $[C_5(CF_3)_5]^-$ was further carried out with pyridine and several fluorine substituted derivatives (Scheme 2). Hereby, pyridine derivatives were employed with further decreasing basicity, depending on the position of the fluorine substituent as well as the degree of fluorination, in order to discover the substitution limit of $[C_5(CF_3)_5]^-$. The reactions could be performed by addition of the respective pyridine into a solution of $[Rh(COD)(C_5(CF_3)_5)]$ in *n*-pentane, which led to a rapid precipitation of a colorless solid. For all substitutions shown in Scheme 2 (except 2,6-difluoropyridine),




Fig. 3 Substitution of $[C_5(CF_3)_5]^-$ in $[Rh(COD)(C_5(CF_3)_5)]$ by MeCN and backreaction in $CHCl_3$. ^{19}F NMR spectrum (377 MHz, $CDCl_3$, rt) measured 30 min after dissolving $[Rh(COD)(MeCN)_2][C_5(CF_3)_5]$ in $CDCl_3$, showing the presence of both MeCN substituted complex (left) and $[Rh(COD)(C_5(CF_3)_5)]$ (right).





Fig. 4 Molecular structure in solid state of $[Rh(COD)(MeCN)_2][C_5(CF_3)_5]$. Disorders are omitted for clarity. Ellipsoids are depicted with 50% probability level. Color code: white-hydrogen; grey-carbon; blue-nitrogen; green-fluorine; light-blue-rhodium.





Scheme 2 Substitution of $[C_5(CF_3)_5]^-$ in $[Rh(COD)(C_5(CF_3)_5)]$ by different (fluorinated) pyridines with their corresponding proton affinities (B3LYP-D3/def2tzvp). The substitution by 2,6-difluoropyridine was carried out overnight.

1H NMR studies revealed a twofold coordination of the respective pyridine ligand, similar to the substitution with MeCN. However, no noteworthy shift of the pyridine signals was observed. ^{19}F NMR spectra clearly confirmed the conversion of $[C_5(CF_3)_5]^-$ into a WCA and furthermore showed a downfield shifted signal for the respective fluorine substituent(s), e.g. $\delta = -68.3$ ppm for uncoordinated 2-fluoropyridine vs. $\delta = -61.2$ ppm for 2-fluoropyridine coordinated complex $[Rh(COD)(2-C_5H_4FN)_2][C_5(CF_3)_5]$ in CD_2Cl_2 .

Single crystals of $[Rh(COD)(3-C_5H_4FN)_2][C_5(CF_3)_5]$ were obtained from solvent mixtures of CH_2Cl_2 and *n*-pentane by slowly cooling to -75 °C. The compound crystallizes in the triclinic space group $P\bar{1}$. The asymmetric unit contains two $[Rh(COD)(3-C_5H_4FN)_2][C_5(CF_3)_5]$ fragments, further confirming the twofold coordination of the pyridines (Fig. 5). Additionally, the solid-state structure shows an average Rh–N bond length of 2.111(3) Å and an average of 2.133(5) Å for the Rh–C distance to the COD ligand. A database survey in the Cambridge Crystallographic Database (CSD) revealed no results for rhodium complexes with fluorinated pyridine ligands, thus $[Rh(COD)(3-C_5H_4FN)_2][C_5(CF_3)_5]$ represents a rare example thereof.




Fig. 5 Molecular structure in solid state of $[Rh(COD)(3-C_5H_4FN)_2][C_5(CF_3)_5]$. Disorders are omitted for clarity. Ellipsoids are depicted with 50% probability level. Color code: white-hydrogen; grey-carbon; blue-nitrogen; green-fluorine; light-blue-rhodium.




Fig. 6 ^{19}F NMR spectrum (377 MHz, $CDCl_3$, rt) of $[Rh(COD)(3-C_5H_4FN)_2][C_5(CF_3)_5]$ after 3 d in $CDCl_3$ at rt.

Surprisingly in contrast to the MeCN and toluene substitutions, no backreaction was observed for the pyridine substituted complexes in solution with CD_2Cl_2 , except for the 2,6-difluoropyridine substituted complex. When dissolving $[Rh(COD)(C_5H_5N)_2][C_5(CF_3)_5]$ in $CHCl_3$, no backreaction was observed either, despite prolonged reaction times (up to 4 d). Dissolving $[Rh(COD)(C_5H_5N)_2][C_5(CF_3)_5]$ in coordinating solvents such as MeCN did not lead to a substitution of the pyridine ligands, which was confirmed by 1H NMR. The backreaction of $[Rh(COD)(3-C_5H_4FN)_2][C_5(CF_3)_5]$ to $[Rh(COD)(C_5(CF_3)_5)]$ proved to be minimal in $CHCl_3$, despite low concentrations and prolonged reaction times (Fig. 6). On the other hand, substitution of complexes $[Rh(COD)(2-C_5H_4FN)_2][C_5(CF_3)_5]$ and $[Rh(COD)(3,5-C_5H_3F_2N)_2][C_5(CF_3)_5]$ showed to be reversible in $CDCl_3$, so they could be fully converted back into $[Rh(COD)(C_5(CF_3)_5)]$.

In strong contrast, the substitution with 2,6-difluoropyridine behaved differently in comparison to the other conducted pyridine substitutions. Performing the reaction in *n*-pentane did not lead to a fast precipitation of a solid, instead a yellow emulsion was formed after stirring the reaction overnight, discoloring the yellow *n*-pentane solution. After removing the remaining solvent the residue was washed with *n*-pentane, revealing an insoluble solid, strongly indicating the formation of a cationic complex under the given conditions. While 1H NMR spectra revealed the presence of two equivalents of 2,6-difluoropyridine when referenced to the COD signals, no signals of substituted $[C_5(CF_3)_5]^-$ could be observed in the ^{19}F NMR spectrum in either CD_2Cl_2 or $CDCl_3$, indicating a rapid backreaction to $[Rh(COD)(C_5(CF_3)_5)]$ (see ESI, Fig. 20 and 21†). Elemental analysis studies however strongly indicate that the substitution with 2,6-difluoropyridine did proceed, as well as IR spectra of the newly formed species, which show new bands with strong similarities compared to IR spectra of the other (fluoro)pyridine substituted complexes (see Experimental section). It is therefore likely, that 2,6-difluoropyridine is able to substitute $[C_5(CF_3)_5]^-$, however the resulting complex $[Rh(COD)(2,6-C_5H_3F_2N)_2][C_5(CF_3)_5]$ is rather unstable, due to the backreaction to $[Rh(COD)(C_5(CF_3)_5)]$ in solution.

Conclusion

In conclusion, substitution of the $[C_5(CF_3)_5]^-$ ligand of $[Rh(COD)(C_5(CF_3)_5)]$ was investigated with different N-donor



ligands, emphasizing its extraordinary electron deficiency and the weak bonding interactions towards the metal center. Hereby the displacement of the $[\text{C}_5(\text{CF}_3)_5]^-$ ligand was first investigated with MeCN, leading to the formation of isolatable 16-electron complex $[\text{Rh}(\text{COD})(\text{MeCN})_2][\text{C}_5(\text{CF}_3)_5]$. Different pyridines with decreasing basicities were subsequently employed, to explore the limits of the substitution for the $[\text{C}_5(\text{CF}_3)_5]^-$ ligand. While all pyridines were shown to quantitatively substitute the perfluorinated Cp^* ligand, the stability of the resulting cationic complexes was found to be strongly dependent on the basicity of the competing ligand. Cationic complexes containing relatively basic ligands, such as unsubstituted pyridine, were found to be stable in weakly coordinating solvents (e.g., CHCl_3 or CH_2Cl_2), whereas for pyridines with weaker basicities, e.g. 2,6-difluoropyridine, the substitution proved to be fully reversible in solution, favouring the reformation of $[\text{Rh}(\text{COD})(\text{C}_5(\text{CF}_3)_5)]$. In general, the ionic products are less stable in CHCl_3 than in CH_2Cl_2 . Furthermore, newly synthesized complexes $[\text{Rh}(\text{COD})(\text{C}_5\text{H}_{5-x}\text{F}_x\text{N})_2][\text{C}_5(\text{CF}_3)_5]$ represent rare examples of rhodium(i) complexes containing (fluorinated) pyridine ligands.

Experimental

General procedure

The substitutions with different pyridine derivatives were carried out by dissolving $[\text{Rh}(\text{COD})(\text{C}_5(\text{CF}_3)_5)]$ (15.0 mg, 24.0 μmol , 1.0 eq.) with anhydrous *n*-pentane (1 mL) in a 10 mL Schlenk tube, followed by addition of the respective pyridine (0.2 mL, excess). The solution was then stirred at rt for 5 min, after which a colorless precipitate formed. The solvent was removed under high vacuum and the residue was washed with *n*-pentane (2×1 mL). The remaining solvent was removed under high vacuum to afford the respective cationic complex.

Synthetic procedures

Preparation of $[\text{Rh}(\text{COD})(\text{MeCN})_2][\text{C}_5(\text{CF}_3)_5]$. $[\text{Rh}(\text{COD})(\text{C}_5(\text{CF}_3)_5)]$ (15.0 mg, 24.0 μmol , 1.0 eq.) was placed in a 10 mL Schlenk tube and dissolved in anhydrous MeCN (2 mL, excess). The solution was stirred at rt for 1 h and the solvent was removed under high vacuum. The residue was washed with *n*-pentane (2×1 mL) and the remaining solvent was removed under high vacuum, to afford $[\text{Rh}(\text{COD})(\text{MeCN})_2][\text{C}_5(\text{CF}_3)_5]$ (16.8 mg, 24.0 μmol , quant.) as a yellow solid.

^1H NMR (600 MHz, CD_2Cl_2) δ [ppm]: 4.43 (s, 4H), 2.45 (m, 4H), 2.20 (s, 6H), 1.94 (q, $^3J = 8.1$ Hz, 4H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (151 MHz, CD_2Cl_2) δ [ppm]: 125.1 (s), 123.3 (s), 122.2 (s), 85.8 (d, $^1J_{\text{Rh}} = 12.5$ Hz), 30.9 (s). **^{19}F NMR** (377 MHz, CD_2Cl_2) δ [ppm]: -50.5 (s, 15F). **FT-IR** (ATR) ν [cm^{-1}]: 2963 (w), 2322 (w), 2293 (w), 1648 (w), 1497 (m), 1205 (s), 1108 (s), 972 (m), 875 (w), 802 (m), 633 (m). **HRMS** (ESI-TOF, positive) m/z for $[\text{C}_{10}\text{H}_{15}\text{NRh}]^+$ calculated: 252.0260; measured: 252.0260. **HRMS** (ESI-TOF, negative) m/z for $[\text{C}_{10}\text{F}_{15}]^-$ calculated: 404.9766; measured: 404.9486. **EA** ($\text{C}_{22}\text{H}_{18}\text{F}_{15}\text{N}_2\text{Rh}$) calculated:

C: 37.84%, H: 2.60%, N: 4.01%; measured: C: 37.86%, H: 2.60%, N: 4.02%.

Preparation of $[\text{Rh}(\text{COD})(\text{C}_5\text{H}_5\text{N})_2][\text{C}_5(\text{CF}_3)_5]$. The substitution was performed according to the general procedure with anhydrous pyridine, to afford $[\text{Rh}(\text{COD})(\text{C}_5\text{H}_5\text{N})_2][\text{C}_5(\text{CF}_3)_5]$ (18.6 mg, 24.0 μmol , quant.) as a pale-yellow solid.

^1H NMR (600 MHz, CD_2Cl_2) δ [ppm]: 8.62–8.56 (m, 4H), 7.74 (tt, $^3J = 7.7$ Hz, $^4J = 1.6$ Hz, 2H), 7.37–7.31 (m, 4H), 4.09 (s, 4H), 2.67–2.57 (m, 4H), 2.04 (q, $^3J = 7.9$ Hz, 4H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (151 MHz, CD_2Cl_2) δ [ppm]: 149.9 (s), 138.9 (s), 126.2 (s), 124.7 (s), 122.9 (s), 85.7 (d, $^1J_{\text{Rh}} = 12.1$ Hz), 30.5 (s). **^{19}F NMR** (565 MHz, CD_2Cl_2) δ [ppm]: -50.5 (s, 15F). **FT-IR** (ATR) ν [cm^{-1}]: 3021 (w), 2965 (w), 2925 (w), 2894 (w), 2877 (w), 2844 (w), 2360 (w), 1600 (w), 1496 (m), 1444 (m), 1296 (w), 1201 (vs), 1111 (vs), 997 (m), 971 (m), 874 (w), 836 (w), 802 (w), 760 (s), 698 (s), 632 (s). **HRMS** (ESI-TOF, negative) m/z for $[\text{C}_{10}\text{F}_{15}]^-$ calculated: 404.9766; measured: 404.9686. **EA** ($\text{C}_{28}\text{H}_{22}\text{F}_{15}\text{N}_2\text{Rh}$) calculated: C: 43.43%, H: 2.86%, N: 3.62%; measured: C: 44.60%, H: 3.13%, N: 3.57%.

Preparation of $[\text{Rh}(\text{COD})(3\text{-C}_5\text{H}_4\text{FN})_2][\text{C}_5(\text{CF}_3)_5]$. The substitution was performed according to the general procedure with anhydrous 3-fluoropyridine, to afford $[\text{Rh}(\text{COD})(3\text{-C}_5\text{H}_4\text{FN})_2][\text{C}_5(\text{CF}_3)_5]$ (19.5 mg, 24.0 μmol , quant.) as a yellow solid.

^1H NMR (600 MHz, CD_2Cl_2) δ [ppm]: 8.54 (t, $^3J = 2.5$ Hz, 2H), 8.46 (dd, $^3J = 5.4$ Hz, 2H), 7.57–7.51 (m, 2H), 7.39 (dt, $^3J = 8.6$, $^3J = 5.3$ Hz, 2H), 4.12 (s, 4H), 2.68–2.59 (m, 4H), 2.06 (q, $^3J = 7.7$ Hz, 4H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (151 MHz, CD_2Cl_2) δ [ppm]: 161.2 (s), 159.5 (s), 146.7 (d, $J = 4.4$ Hz), 139.3 (d, $J = 29.6$ Hz), 127.7 (d, $J = 6.2$ Hz), 126.8 (d, $J = 17.5$ Hz), 86.9 (d, $^1J_{\text{Rh}} = 12.0$ Hz), 30.9 (s). **^{19}F NMR** (565 MHz, CD_2Cl_2) δ [ppm]: -50.5 (s, 15F), -119.4 (s, 2F). **FT-IR** (ATR) ν [cm^{-1}]: 3011 (w), 2957 (w), 2894 (w), 2847 (w), 1586 (w), 1481 (m), 1435 (m), 1258 (m), 1208 (vs), 1117 (vs), 976 (w), 845 (m), 799 (s), 696 (s), 633 (s), 537 (m). **HRMS** (ESI-TOF, negative) m/z for $[\text{C}_{10}\text{F}_{15}]^-$ calculated: 404.9766; measured: 404.9486. **EA** ($\text{C}_{28}\text{H}_{20}\text{F}_{17}\text{N}_2\text{Rh}$) calculated: C: 41.50%, H: 2.49%, N: 3.46%; measured: C: 41.88%, H: 2.71%, N: 3.59%.

Preparation of $[\text{Rh}(\text{COD})(2\text{-C}_5\text{H}_4\text{FN})_2][\text{C}_5(\text{CF}_3)_5]$. The substitution was performed according to the general procedure with anhydrous 2-fluoropyridine, to afford $[\text{Rh}(\text{COD})(2\text{-C}_5\text{H}_4\text{FN})_2][\text{C}_5(\text{CF}_3)_5]$ (19.5 mg, 24.0 μmol , quant.) as an orange solid.

^1H NMR (600 MHz, CD_2Cl_2) δ [ppm]: 8.49 (s, 2H), 7.85 (q, $^3J = 7.6$ Hz, 2H), 7.29 (s, 2H), 7.06–6.98 (d, $^3J = 8.3$ Hz, 2H), 4.19 (s, 4H), 2.69–2.60 (m, 4H), 2.02 (q, $^3J = 7.8$ Hz, 4H). **$^{13}\text{C}\{^1\text{H}\}$ NMR** (151 MHz, CD_2Cl_2) δ [ppm]: 148.3 (d, $J = 6.5$ Hz), 144.4 (s), 125.0 (s), 123.7 (s), 123.2 (s), 112.4 (s), 85.1 (d, $^1J_{\text{Rh}} = 12.5$ Hz), 30.9 (s). **^{19}F NMR** (565 MHz, CD_2Cl_2) δ [ppm]: -50.5 (s, 15F), -61.2 (s, 2F). **FT-IR** (ATR) ν [cm^{-1}]: 3012 (w), 2962 (w), 2922 (w), 2894 (w), 2845 (w), 1615 (m), 1576 (w), 1533 (m), 1479 (s), 1446 (s), 1337 (m), 1260 (m), 1205 (vs), 1100 (vs), 865 (s), 800 (s), 771 (vs), 734 (m), 696 (m), 633 (vs), 560 (m). **HRMS** (ESI-TOF, negative) m/z for $[\text{C}_{10}\text{F}_{15}]^-$ calculated: 404.9766; measured: 404.9668. **EA** ($\text{C}_{28}\text{H}_{20}\text{F}_{17}\text{N}_2\text{Rh}$) calculated: C:



41.50%, H: 2.49%, N: 3.46%; measured: C: 41.78%, H: 2.93%, N: 3.54%.

Preparation of [Rh(COD)(3,5-C₅H₃F₂N)₂][C₅(CF₃)₅]. The substitution was performed according to the general procedure with anhydrous 3,5-difluoropyridine, to afford [Rh(COD)(3,5-C₅H₃F₂N)₂][C₅(CF₃)₅] (20.3 mg, 24.0 μmol, quant.) as a pale-yellow solid.

¹H NMR (600 MHz, CD₂Cl₂) δ [ppm]: 8.46 (d, ³J = 2.3 Hz, 4H), 7.41 (tt, ³J = 7.9 Hz, ⁴J = 2.3 Hz, 2H), 4.13 (s, 4H), 2.68–2.60 (m, 4H), 2.06 (q, ³J = 7.6 Hz, 4H). **¹³C{¹H} NMR** (151 MHz, CD₂Cl₂) δ [ppm]: 161.3 (s), 136.0 (d, J = 27.3 Hz), 125.1 (s), 115.5 (s), 87.5 (s), 30.8 (s). **¹⁹F NMR** (565 MHz, CD₂Cl₂) δ [ppm]: −50.6 (s, 15F), −116.3 (s, 4F). **FT-IR** (ATR) ν [cm^{−1}]: 3088 (w), 3013 (w), 2961 (w), 2925 (w), 2897 (w), 2361 (w), 1599 (m), 1494 (m), 1438 (m), 1317 (m), 1210 (vs), 1116 (vs), 977 (m), 871 (m), 801 (w), 688 (m), 633 (m), 533 (m). **HRMS** (ESI-TOF, negative) *m/z* for [C₁₀F₁₅][−] calculated: 404.9766; measured: 404.9684. **EA** (C₂₈H₁₈F₁₉N₂Rh) calculated: C: 39.74%, H: 2.14%, N: 3.31%; measured: C: 37.68%, H: 3.15%, N: 3.44%.

Preparation of [Rh(COD)(2,6-C₅H₃F₂N)₂][C₅(CF₃)₅]. The substitution was performed according to the general procedure with anhydrous 2,6-difluoropyridine but with a prolonged reaction time of 17 h, to afford [Rh(COD)(2,6-C₅H₃F₂N)₂][C₅(CF₃)₅] (20.3 mg, 24.0 μmol, quant.) as a yellow solid.

FT-IR (ATR) ν [cm^{−1}]: 2961 (w), 2927 (w), 2893 (w), 2844 (w), 1629 (s), 1567 (w), 1529 (m), 1493 (s), 1467 (w), 1329 (w), 1207 (vs), 1112 (vs), 1007 (s), 876 (w), 792 (vs), 741 (m), 697 (w), 632 (s), 569 (m). **EA** (C₂₈H₁₈F₁₉N₂Rh) calculated: C: 39.74%, H: 2.14%, N: 3.31%; measured: C: 39.76%, H: 2.18%, N: 3.32%.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Gefördert durch die Deutsche Forschungsgemeinschaft (DFG) —Projekt Nummer 387284271—SFB 1349. Computing time was made available by High-Performance Computing ZEDAT/FU Berlin. The authors acknowledge the assistance of the Core Facility BioSupraMol supported by the DFG. The authors acknowledge generous support by the Fonds of the Chemical Industry (Kekulé fellowship for Robin Sievers).

References

- 1 S. D. Pike, I. Pernik, R. Theron, J. S. McIndoe and A. S. Weller, *J. Organomet. Chem.*, 2015, **784**, 75–83.
- 2 S. T. H. Willems, P. H. M. Budelaar, N. N. P. Moonen, R. de Gelder, J. M. M. Smits and A. W. Gal, *Chem. – Eur. J.*, 2002, **8**, 1310–1320.
- 3 V. V. Grushin and H. Alper, *Chem. Rev.*, 1994, **94**, 1047–1062.
- 4 R. J. Less, T. C. Wilson, M. McPartlin, P. T. Wood and D. S. Wright, *Chem. Commun.*, 2011, **47**, 10007–10009.
- 5 A. B. Chaplin, J. F. Hooper, A. S. Weller and M. C. Willis, *J. Am. Chem. Soc.*, 2012, **134**, 4885–4897.
- 6 A. Prades, M. Fernández, S. D. Pike, M. C. Willis and A. S. Weller, *Angew. Chem., Int. Ed.*, 2015, **54**, 8520–8524.
- 7 I. Pernik, J. F. Hooper, A. B. Chaplin, A. S. Weller and M. C. Willis, *ACS Catal.*, 2012, **2**, 2779–2786.
- 8 T. M. Douglas, J. L. Nôtre, S. K. Brayshaw, C. G. Frost and A. S. Weller, *Chem. Commun.*, 2006, 3408–3410.
- 9 R. Dallanegra, A. P. M. Robertson, A. B. Chaplin, I. Manners and A. S. Weller, *Chem. Commun.*, 2011, **47**, 3763–3765.
- 10 J. F. Hooper, R. D. Young, I. Pernik, A. S. Weller and M. C. Willis, *Chem. Sci.*, 2013, **4**, 1568–1572.
- 11 T. M. Douglas, A. B. Chaplin, A. S. Weller, X. Yang and M. B. Hall, *J. Am. Chem. Soc.*, 2009, **131**, 15440–15456.
- 12 A. L. Colebatch, A. I. McKay, N. A. Beattie, S. A. Macgregor and A. S. Weller, *Eur. J. Inorg. Chem.*, 2017, **2017**, 4533–4540.
- 13 I. Krossing, *Chem. – Eur. J.*, 2001, **7**, 490–502.
- 14 A. I. McKay, J. Barwick-Silk, M. Savage, M. C. Willis and A. S. Weller, *Chem. – Eur. J.*, 2020, **26**, 2883–2889.
- 15 S. T. Belt, S. B. Duckett, M. Helliwell and R. N. Perutz, *J. Chem. Soc., Chem. Commun.*, 1989, 928–930.
- 16 C. L. Higgitt, A. H. Klahn, M. H. Moore, B. Oelckers, M. G. Partridge and R. N. Perutz, *J. Chem. Soc., Dalton Trans.*, 1997, 1269–1280.
- 17 L. Lefort, T. W. Crane, M. D. Farwell, D. M. Baruch, J. A. Kaeuper, R. J. Lachicotte and W. D. Jones, *Organometallics*, 1998, **17**, 3889–3899.
- 18 R. Sievers, J. Parche, N. G. Kub and M. Malischewski, *Synlett*, 2023, DOI: [10.1055/s-0042-1751426](https://doi.org/10.1055/s-0042-1751426).
- 19 T. J. Kealy and P. L. Pauson, *Nature*, 1951, **168**, 1039–1040.
- 20 C. Janiak and H. Schumann, in *Adv. Organomet. Chem.*, ed. F. G. A. Stone and R. West, Academic Press, 1991, vol. 33, pp. 291–393.
- 21 D. W. Macomber, W. P. Hart and M. D. Rausch, in *Adv. Organomet. Chem.*, Academic Press, 1982, vol. 21, pp. 1–55.
- 22 S. Lauk and A. Schäfer, *Eur. J. Inorg. Chem.*, 2021, **2021**, 5026–5036.
- 23 F. Mazzotta, G. Zitzer, B. Speiser and D. Kunz, *Chem. – Eur. J.*, 2020, **26**, 16291–16305.
- 24 G. Paprott and K. Seppelt, *J. Am. Chem. Soc.*, 1984, **106**, 4060–4061.
- 25 O. J. Curnow and R. P. Hughes, *J. Am. Chem. Soc.*, 1992, **114**, 5895–5897.
- 26 K. Sünkel, S. Weigand, A. Hoffmann, S. Blomeyer, C. G. Reuter, Y. V. Vishnevskiy and N. W. Mitzel, *J. Am. Chem. Soc.*, 2015, **137**, 126–129.
- 27 E. D. Laganis and D. M. Lemal, *J. Am. Chem. Soc.*, 1980, **102**, 6633–6634.
- 28 R. D. Chambers, S. J. Mullins, A. J. Roche and J. F. S. Vaughan, *J. Chem. Soc., Chem. Commun.*, 1995, 841–842.



- 29 R. Sievers, M. Sellin, S. M. Rupf, J. Parche and M. Malischewski, *Angew. Chem., Int. Ed.*, 2022, e202211147.
- 30 D. W. Slocum, T. R. Engelmann, R. L. Fellows, M. Moronski and S. Duraj, *J. Organomet. Chem.*, 1984, **260**, C21–C25.
- 31 S. G. Shore, W. L. Hsu, M. R. Churchill and C. Bueno, *J. Am. Chem. Soc.*, 1983, **105**, 655–656.
- 32 R. Alberto, personal communication.
- 33 S. O. Kang, V. M. Lynch, V. W. Day and E. V. Anslyn, *Organometallics*, 2011, **30**, 6233–6240.

