



Cite this: *Phys. Chem. Chem. Phys.*,
2025, 27, 14985

Surface hopping simulations reveal deactivation pathways of a charge transfer system with planarizing and twisting motion†

Julia Haberhauer, ^{*a} Sebastian Mai, ^b Leticia González ^b and
Christof Hättig ^{*a}

Nonadiabatic surface hopping simulations are used to investigate the relaxation pathways after photoexcitation of 4-(indol-1-ylamino)benzonitrile, a prototypical example of *N*-aryl-substituted 1-aminoindoles. This molecule combines functional groups that can undergo in the excited state twisted as well as planarized intramolecular charge transfer and are potential building blocks for molecular motors. The results of the nonadiabatic dynamics simulation show that after excitation with light in the range of 4.2–4.8 eV, the *N*-aryl-substituted 1-aminoindole decays rapidly from S_3 and S_2 into the S_1 state. On the S_1 potential energy surface, the system relaxes then to one of the charge-transfer minima with twisted amino-phenyl and planarized amino-indole torsional angles. The twist and the planarization occur nearly synchronously, with a slight advance of the twist. The geometric relaxation is accompanied by a change of the electronic structure from a $\pi\pi^*$ excitation localized on indole to a charge transfer excitation from the indole and amino groups into the benzonitrile moiety. The simulated time-resolved fluorescence spectrum exhibits a bright band from a locally excited state and a weaker charge-transfer band. The transition energies suggest that the experimentally observed emission happens from the S_1 minima. After transition to the ground state, half of the trajectories relax back to the original minimum and half of the trajectories to its enantiomer. The twist and planarization occur nearly synchronously, with planarization slightly leading. The combination of preference to rotate over twisted intermediate structure in the excited state and over more twisted structures in the ground state indicates that some kinetic work is done during a full cycle. Even though the kinetic work done per cycle is small, this might be increased by e.g. introducing substituents.

Received 6th March 2025,
Accepted 19th June 2025

DOI: 10.1039/d5cp00889a

rs.c.li/pccp

1 Introduction

While molecular motors are widespread in nature, their synthetic realization is a young field of research.¹ A system in which controlled movement is triggered by an external influence is referred to as a molecular machine. A subtype thereof are molecular switches,¹ which are systems that can be switched reversibly between two or more states or geometries that differ in their properties. All mechanical work is undone when a switch returns to its original state. A molecular motor, on the other hand, is a subtype of molecular machines in which the mechanic work performed is not canceled when the motor

returns to its original state.¹ According to Feynman, a molecular motor requires an energy source, a periodic potential energy, and structural asymmetry.²

Molecular motors can be categorized according to the type of movement and drive type. The most common molecular motors are rotary motors – or rotors – in which a repetitive, unidirectional 360° movement is performed under energy supply.¹ Another type of molecular motor is the push motor, inspired by the movement of cilia.^{3–7} Currently available synthetic molecular motors usually require several energetic stimuli (often including thermal stimuli as rate determining step) to run through an entire cycle.^{1,8–15} An example of this is Feringa's pioneering rotor, which has also been subject of intensive theoretical simulations.^{16–19}

Even though most synthetic motors use thermal stimuli, some exclusively light-driven motors that use multiple light impulses have also been investigated.^{8,9} A theoretical study by Wang *et al.*¹⁰ concluded that a weak asymmetry as introduced by isotopic chirality is already sufficient to achieve the unidirectionality required for molecular motors.

^a Institute of Theoretical Chemistry I, Faculty of Chemistry, Ruhr University Bochum, Universitätsstraße 150, 44780 Bochum, Germany.

E-mail: julia.haberhauer@rub.de, christof.haettig@rub.de

^b Institute of Theoretical Chemistry, Faculty of Chemistry, University of Vienna, Währingerstrasse 17, 1090 Vienna, Austria

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d5cp00889a>



In 2017, a system was proposed in which a single light pulse should be sufficient to achieve an entire cycle for a molecular motor.²⁰ The motion sequence of the motor is most similar to a push motor. It is a combination of two excited-state relaxation phenomena that are combined in one molecule: one relaxation motion which is similar to the relaxation pathway of a TICT (twisted intramolecular charge transfer) system^{21,22} and a second motion that is similar to the relaxation pathway of a PLICT (planar intramolecular charge transfer) system.²³ In a TICT system, two functional groups are coplanar in the ground state but twisted relative to each other in the excited state.^{24,25} The prime example of this is 4-(dimethylamino)benzonitrile (DMABN).^{26–28} Experimentally, the twist in the S_1 state is recognized by a large Stokes shift of about 9800 cm^{-1} (in polar solvents like 1-chlorobutane)^{29,30} and an almost forbidden emission, *i.e.* a very low quantum yield for the TICT band.^{29,30} PLICT systems, on the other hand, are twisted in the ground state and rotate to a planar structure in the excited state.²³ They also exhibit large Stokes shifts for the emission (*e.g.* 9300 cm^{-1} for 1-aminoindole in cyclohexane),²³ but the transition to the ground state is usually dipole-allowed and has a high quantum yield.²³

The *N*-aryl-substituted 1-aminoindoles proposed in ref. 20 combine TICT and PLICT chromophores to achieve two independent rotations in the excited state, as sketched in Fig. 1. Molecules that can undergo TICT and PLICT rotations in the excited state are referred to as PLATICT systems.²⁰ Some

N-aryl-substituted 1-aminoindoles are such systems: after electronic excitation they undergo an intramolecular charge transfer (CT) from the aminoindole to the aryl chromophore, which is accompanied by a planarization of the amino-indole and a twist of the amino-aryl bond (relaxation from I^* to IV^* in Fig. 1a). After electronic relaxation to the ground state, the PLATICT-like geometry IV rotates back to the ground-state equilibrium structure I (see Fig. 1a). The pathways for the geometric relaxations in the excited and in the ground state are not yet known. They might proceed either *via* TICT-like (pathways A and D) or PLICT-like (pathways C and F) intermediate structures, or both torsions could occur simultaneously (B and E). One of the PLATICT compounds studied in ref. 20 experimentally and with static quantum chemical calculations is 4-(indol-1-ylamino)benzonitrile (IYABN), a combination of 1-aminoindole and aminobenzonitrile (ABN). The reported results comprise experimental absorption and fluorescence spectra and computed equilibrium structures for S_0 and S_1 with the respective vertical $S_0 \rightarrow S_1$ and $S_1 \rightarrow S_0$ transition energies at the CC2/def2-TZVP//((TD)-CAM-B3LYP/def2-TZVP level. This molecule shows experimentally very large Stokes shifts between 15700 cm^{-1} (in *n*-hexane) and 18200 cm^{-1} (in 1-chlorobutane) and very low quantum yields.²⁰ The strong Stokes shifts suggest the presence of two rotations in the excited state, which is also supported by the computed S_1 equilibrium structure. However, the relaxation pathways after photoexcitation, both in the excited state and, after fluorescence, in the ground state, remain elusive. Thus, the extent to which this or other PLATICT systems can perform mechanical work is still unknown. This depends on how different the deactivation on the excited state is from the evolution of the system on the ground state.

The molecule combines three structural units – indole, amino group, and benzonitrile – linked by two single bonds and coupled in the excited states by possible CTs between the three units. It is an open question whether during the relaxation after photoexcitation the torsions around the two linking single bonds occur sequentially in a specific order or in a concerted fashion. In order to answer this question, here we perform nonadiabatic dynamics simulations using trajectory surface hopping,³¹ as implemented in the software package Surface Hopping including Arbitrary Couplings (SHARC),^{32–34} combined with the algebraic diagrammatic construction through second order,^{35,36} ADC(2). We study the relaxation of the molecule after photoexcitation in the UV/Vis regime (4.2 eV to 4.8 eV) and after deexcitation from the excited-state minima and compare the relaxation paths in the excited state with those in the ground state.

2 Computational details

To validate the computational setup for the nonadiabatic dynamics simulations, we first benchmarked the electronic structure methods and basis sets. Then, as starting point for the excited-state dynamics, Wigner distributions are generated




Fig. 1 Suggested pathways for *N*-aryl-substituted 1-aminoindoles. (a) After excitation from the ground state minimum I to I^* the structure relaxes to an excited-state minimum IV^* . After deexcitation to IV it can relax back to the I . Pathways *via* twisted structures (A) and (D), concerted pathways (B) and (E), or pathways *via* planarized structures (C) and (F) are conceivable for the relaxations. (b) Possible structures for the 4-(indol-1-ylamino)benzonitrile studied in the current work.

For the description of nonadiabatic effects between the remaining electronic states, the local diabaticization scheme from ref. 58 was used. The required wave function overlaps were computed with the WFOverlap program,⁵⁹ using the single-excitation amplitudes as though they were CIS coefficients, and truncating the coefficients to 0.999 of the norm.⁶⁰ The kinetic energy was corrected by adjusting the full velocity vectors; frustrated hops were not reflected. For decoherence corrections, the energy-based method of Granucci, Persicio, and Zocante⁶¹ with the default decoherence parameter ($C = 0.1$ a.u.) was used. In the final analysis, trajectories with problems in energy conservation, potential or kinetic energy smoothness, hopping energy restriction, or intruder states were excluded as in the default settings of SHARC’s diagnostic tool (further details are specified in the ESI†).

The surface hopping molecular dynamics simulations for the relaxation after photoexcitation from the ground state minima and after the de-excitation from the excited state minima were carried out with the SHARC package version 3.0.^{32–34} The necessary energies, gradients and overlaps were calculated on-the-fly at the MP2/cc-pVDZ level of theory for the ground and at the ADC(2)/cc-pVDZ level of theory for the excited state.^{35,36,41,43,44} Initial nuclear ensembles for starting the



Experimental spectroscopic studies²⁰ of the fluorescence used an excitation energy of ≈ 4.4 eV. According to these experiments, we included for the excited-state dynamics starting geometries and velocities in the excited states S_1 , S_2 , and S_3 with transition energies between 4.2 and 4.8 eV. From the 563 initial conditions generated for the excited state dynamics, 82 have been stochastically selected⁶² for starting trajectories (36 in S_1 , 31 in S_2 , and 15 in S_3). All stochastically chosen trajectories start from geometries obtained from S_0 -Min-1. 55 of the in total 82 trajectories terminated after 1000 fs without issues, being still in an excited state (see Fig. S11, ESI†).

For the simulations of the relaxation in S_0 after emission, 196 start geometries and velocities were chosen stochastically⁶² from the Wigner distributions for the excited-state minima (44 for S_1 -Min-a, 64 for S_1 -Min-b, 49 for S_1 -Min-c, and 39 for S_1 -Min-d, see Section 3.1 for the structure of the minima). The geometries were vertically projected down onto the S_0 potential energy surface (PES). The setting for the dynamics were as described above (only using MP2 instead of ADC(2)). In the analysis, the contributions from the trajectories starting in different excited state minima have been weighted according to the number of trajectories that ended in the excited-state simulations in the respective basin.

3 Results and discussion

In this section we first present the results of the static calculations and spectra and then provide an overview of the electronic evolution for the excited state dynamics. We then consider the geometric changes that occur in the excited and ground state dynamics, and finally discuss the implications of these geometric changes for the role of IYABN as a potential motor.

3.1 Static calculations: geometry and excitations

On the ground-state PES, we found two conformational minima that differ in the pyramidalization of the amino group (see Fig. 2, for details how the minima on S_0 as well as on S_1 were found see ESI,† Chapter 7). In all ground state minima the indole and the amino group are twisted relative to each other by about 90° whereas the phenyl ring is roughly coplanar with the amino group. The energetically lower minimum is designated as S_0 -Min-1 and the higher one as S_0 -Min-2. They differ by 4 kJ mol^{-1} in energy and are separated by a transition state with a barrier height of 10 kJ mol^{-1} . Both minima are chiral. The mirror image of S_0 -Min-1 is also a rotamer of S_0 -Min-2 and, similarly, the mirror image of S_0 -Min-2 is a rotamer of S_0 -Min-1.

In the excited state, four minima were found at the ADC(2)/cc-pVDZ level used for the simulations (*vide infra*). The four minima are referred to as S_1 -Min-a, S_1 -Min-b, S_1 -Min-c and S_1 -Min-d (see Fig. 2). In the larger cc-pVTZ and aug-cc-pVTZ basis sets, geometry optimizations for S_1 -Min-a collapse to S_1 -Min-b. In contrast to the ground state, the indole and the amino group are in all S_1 minima coplanar while the phenyl ring is twisted by about 90° relative to the amino group, *i.e.* these minima show the structural signatures of a PLATICT state.²⁰ S_1 -Min-a and S_1 -

Min-c as well as the minima S_1 -Min-b and S_1 -Min-d are each other's rotamer. S_1 -Min-a and S_1 -Min-b differ through inversion of the pyramidalization at the phenyl ring. The same holds for S_1 -Min-c and S_1 -Min-d. Due to C_s symmetry of the S_1 equilibrium structures they are not chiral. The structures S_1 -Min-c and S_1 -Min-d are, apart from the pyramidalization at atom C_4 similar to the S_1 equilibrium structure obtained in ref. 20 with TD-CAM-B3LYP/def2-TZVP. Fig. 2 summarizes the relative energies of the equilibrium structures. The minima in S_1 differ by about 340 kJ mol^{-1} or 3.5 eV from the ground state minima. S_1 -Min-b and S_1 -Min-d are slightly lower in energy than S_1 -Min-c. The rotamers S_1 -Min-b and S_1 -Min-d differ only by 3 kJ mol^{-1} .

When comparing the ground- and excited-state structures, in addition to the torsions further changes in the structures can be observed: upon excitation to S_1 , the bond between C_4 and N_3 as well as the bond between C_{12} and N_2 are elongated while the bond length of the N–N bond decreases. Furthermore, in the S_1 state the carbon atom C_4 is no longer purely sp^2 -hybridized. If we compare the structure of the ABN subsystem of IYABN in the S_1 state with results²⁷ for isolated DMABN, similar changes are found in the TICT state of the latter: the elongated C–N bond, the pyramidalization at C_4 , and, most prominently, the twist of the amino group by 90° relative to the phenyl ring. Both, the TICT minimum in DMABN and the PLATICT minima of IYABN exhibit (almost) C_s symmetry. In contrast to IYABN, DMABN has on the S_1 PES a locally excited (LE) minimum where the amino group is almost coplanar with the phenyl ring.²⁷ Such a minimum could not be located on the S_1 PES of IYABN with ADC(2). This is consistent with the TD-CAM-B3LYP/def2-TZVP results of ref. 20 where a minimum without twist between amino group and phenyl ring was only found for the parent compound without cyano group, *N*-(1*H*-indol-1-yl)aniline, and the fluor derivative, 4-(indol-1-ylamino)fluorobenzene.

Also, when comparing the aminoindole subsystem of IYABN with the ground-state and PLICT excited-state minima²³ of 1-aminoindole, we find similar structural changes upon excitation: a decrease of the N–N and an increase of the C–N bond length, and, most prominently, the indole and the amino group are coplanar in the PLICT state while they are twisted in the ground state. In the PLICT minimum of 1-aminoindole, C_{12} is not sp^2 -hybridized.²³ This pyramidalization is not observed in the PLATICT state of IYABN. Similar as DMABN, also 1-aminoindole has on the S_1 PES in addition to the ICT minimum a LE minimum,²³ which is not found for IYABN.²³ The current results for the S_1 and S_2 structures of IYABN are consistent with those obtained in ref. 20 with (TD)-CAM-B3LYP/def2-TZVP. In particular, the changes between the ground and the excited state minima are similar.²⁰ The more exhaustive search for conformers in the current work revealed on the ground state PES a second minimum (S_0 -Min-2) and two additional minima on the S_1 PES with ADC(2)/cc-pVDZ (S_1 -Min-a and S_1 -Min-c) of which, however, only S_1 -Min-c remains stable when reoptimized with a larger basis set. In the previous study the minima S_1 -Min-b and S_1 -Min-d were called *syn* and *anti*.²⁰

Fig. 3 depicts the most important natural transition orbitals⁶³ (NTOs) at the ground-state minimum S_0 -Min-1 (NTOs






Fig. 3 Vertical transitions obtained with ADC(2)/aug-cc-pVTZ in vacuum at the ground-state minimum S_0 -Min-1 (structure optimized at the MP2/aug-cc-pVTZ level). For each state the most important occupied and virtual NTO is shown.

for the other ground-state minimum given in Fig. S1, ESI†). At the ground-state structure, the lowest first three singlet transitions are $\pi\pi^*$ excitations localized in one of the two aromatic subsystems: S_1 is localized on the aminobenzonitrile, and S_2 and S_3 on the indole fragment. The NTOs of first excitation are

similar to those of the $S_1(\pi\pi^*)$ transition at the ground-state structure in aminobenzonitrile, which relaxes for DMABN to the LE state (see Fig. S7 and S8, ESI†). The second excitation corresponds to the $S_1(\pi\pi^*)$ transition of aminoindole, relaxing to a PLICT state (see Fig. S5 and S6, ESI†).²³ The first two excited states of IYABN are very close in energy, separated at the Franck–Condon (FC) point at the ADC(2)/aug-cc-pVTZ level by only 0.02 eV, with S_1 having a small oscillator strength. The NTOs of the third excitation at the FC point are similar those of S_2 in aminoindole, which relaxes to a LE state (see Fig. S5 and S6, ESI†).²³ The fourth and the fifth transition are plus and minus combinations of a CT transition from the indole to the benzonitrile group and a second $\pi\pi^*$ transition localized in the benzonitrile group. The latter contribution is similar to the second transition in aminobenzonitrile, relaxing in DMABN to the TICT state (see Fig. S7 and S8, ESI†).²⁷

The excitation energies for the third, forth, and fifth transition differ at the FC point in the aug-cc-pVTZ basis by only ≈ 0.1 eV. They have higher oscillator strengths than the first two transitions. For the cc-pVDZ basis the same trend is observed.

The NTOs for the deexcitation from S_1 to S_0 are very similar at all excited state minima. Fig. 4 shows them at the S_1 -Min-b structure (NTOs for S_1 -Min-c and S_1 -Min-d are given in Fig. S2–S4, ESI†). In all cases, it represents a CT transfer transition from the aminoindole to benzonitrile unit. Since the π -systems of the two subunits are almost perpendicular to each other, the transition is (almost) dipole-forbidden, which can be seen from an oscillator strength of 0.000. The occupied NTO corresponds to the occupied NTO of the PLICT state of 1-aminoindole while the virtual NTO is similar to the virtual NTO of DMABN's TICT state.^{23,27}

The results for the lowest transitions agree qualitatively with ref. 20 although the emission energies differ slightly: in the previous study the lowest vertical excitation at the ground-state structure was 0.07 eV higher with an oscillator strength of 0.022 while the emission energy was 0.48 eV higher but had also an oscillator strength of 0.000. The higher emission energy in ref. 20 is related to the DFT geometry: the emission energy of ADC(2)/aug-cc-pVTZ computed at this geometry is only 0.11 eV higher than in ref. 20. Tables 1 and 2 compare the ADC(2) results for the lowest vertical excitation and emission energies and the respective oscillator strengths with results from the




Fig. 4 Electronic transitions of the S_1 -Min-b, calculated using ADC(2)/aug-cc-pVTZ in vacuum. Previous geometry optimization using ADC(2)/aug-cc-pVTZ in vacuum. The most important occupied and virtual NTO of the transition is shown.



Table 1 Vertical excitation energies and oscillator strengths at the ground state minimum S_0 -Min-1 for different basis sets and methods. The optimized MP2/cc-pVDZ was used for these calculations

Basis set	Method	S_1		S_2	
		eV	Osc	eV	Osc
aug-cc-pVTZ	ADC(2)	4.61	0.001	4.62	0.065
	CC2	4.62	0.002	4.64	0.054
cc-pVDZ	ADC(2)	4.78	0.005	4.80	0.058
	CC2	4.79	0.006	4.82	0.047

approximate coupled cluster singles and doubles model⁶⁴ CC2. At the ground-state structure, the ADC(2) and CC2 results for the excitation energies to S_1 and S_2 agree within 0.02 eV, for the vertical emission at the S_1 equilibrium structure they still differ only by 0.13 eV.

As the experimental spectroscopic studies²⁰ were carried out in solution (*n*-hexane, 1-chlorobutane, and dioxane), solvent effects should be included when comparing with them. Since the focus of the current work is on the relaxation pathways after photoexcitation and after the emission, we only considered solvent effects for comparison with the absorption spectrum in 1-chlorobutane with COSMO. The solvent effects lead mainly to a red shift in particular of the bright $\pi\pi^*$ excitations S_3 , S_4 , and S_5 (*vide infra*). Depending on whether CC2 or ADC(2) is used, the ordering of the (at the ground state structure) energetically close-lying states S_1 and S_2 as well as the order within the pair S_4 and S_5 switches (see Fig. S9 and S10, ESI†). Geometry optimizations for the excited state with COSMO-ADC(2) in 1-chlorobutane showed only minor changes in the S_1 equilibrium structures and the NTOs of the $S_0 \rightarrow S_1$ transition are almost the same as in the vacuum calculation.

Since the aug-cc-pVTZ basis used above would have been extremely costly for surface hopping molecular dynamics calculations, the more cost-efficient alternative cc-pVDZ basis was chosen. The corresponding excitation energies and oscillator strengths are compared with those from the aug-cc-pVTZ basis in Tables 1 and 2. At the ground-state structure, the excitation energies obtained with the smaller basis set are both blue-shifted by ≈ 0.2 eV. A similar blue shift of ≈ 0.25 eV is found for the third, fourth and fifth excited state (see Table S1, ESI†). The de-excitation energy at the S_1 structure is blue shifted by 0.05 eV. Since the characters of the lowest transitions remains in the smaller basis set qualitatively the same as in the reference aug-cc-pVTZ basis, we safely concluded that the cc-pVDZ basis is sufficient for a qualitatively correct description of

the photophysics of IYABN. Accordingly, we used ADC(2)/cc-pVDZ for the nonadiabatic molecular dynamics simulations and the corresponding geometry optimizations and frequency calculations needed for the Wigner distributions. Test calculations showed very small spin-orbit couplings, suggesting that intersystem crossing is not expected.^{65,66} Therefore, triplet states and spin-orbit couplings have been excluded in the surface hopping dynamics.

3.2 Spectrum

Fig. 5 shows a comparison between the experimental absorption spectrum recorded in 1-chlorobutane and computed spectra. For the simulated absorption spectrum shown in black, the transition energies and oscillator strengths were calculated for the 5 lowest singlet transitions in gas phase for the 563 structures obtained from the Wigner distribution of the ground-state minima. The obtained stick spectra have been broadened with Gaussians (full width at half maximum, FWHM = 0.1 eV) and accumulated. Regarding the band shape and position, the spectrum simulated for the gas phase agrees qualitatively with the experimental spectrum recorded in chlorobutane as much as can be expected considering that the calculations were done without diffuse functions and the neglect of solvent effects which are expected to red-shift the excitation energies of the brighter $\pi\pi^*$ states S_3 – S_5 slightly more than the lower states with small oscillator strengths.

To investigate the origin of the difference between the vacuum absorption spectrum simulated with ADC(2)/cc-pVDZ and the experimental spectrum recorded in 1-chlorobutane, we estimated how the computed spectrum changes if (a) solvation




Fig. 5 Comparison of the ground-state absorption spectrum. The experimental absorption spectrum²⁰ recorded in 1-chlorobutane is shown in red (note that the increase beyond 5.3 eV originates from a solvent band). In black the spectrum computed from 563 structures from the Wigner distribution for S_0 -Min-1 and S_0 -Min-2 at the ADC(2)/cc-pVDZ level is shown. The contributions of the different states are given as shaded areas. The purple line shows the spectrum obtained by shifting the vacuum results for each state by the difference in the vertical excitation energy between ADC(2)/cc-pVDZ and COSMO-ADC(2)/cc-pVDZ for 1-chlorobutane. The blue spectrum is obtained using the differences between ADC(2)/cc-pVDZ and COSMO-ADC(2)/aug-cc-pVTZ for shifting.

Table 2 Vertical transition energies and oscillator strengths at the S_1 minimum S_1 -Min-b for different basis sets and methods. The excited state structure was optimized with ADC(2)/cc-pVDZ

Basis set	Method	S_1	
		eV	Osc
aug-cc-pVTZ	ADC(2)	1.81	0.000
	CC2	1.93	0.000
cc-pVDZ	ADC(2)	1.85	0.000
	CC2	1.98	0.000



effects are included with COSMO and (b) the larger basis set is used. For this, we evaluated at the MP2/cc-pVDZ optimized structure for S_0 -Min-1 for each state the difference between the ADC(2)/cc-pVDZ excitation energies and those obtained with COSMO-ADC(2)/cc-pVDZ and, to address (b), those obtained with COSMO-ADC(2)/aug-cc-pVTZ in 1-chlorobutane. We then corrected the excitation energies for the Wigner ensemble by adding for each state the respective energy shifts. For different geometries, the adiabatic characters for the same state may not match, but we are only considering a qualitative comparison. The solvation effects shift S_3 – S_5 closer to S_1 and S_2 which leads to a compression of the overall band shape from these transitions. The additional inclusion of diffuse functions red-shifts the whole band by about 0.25 eV. The final band position and band shape are in good agreement with the experimental band shape, confirming the accurate description of this system with the applied electronic structure methods. As absorption to S_4 and S_5 is minor for the energy range of 4.2–4.8 eV and higher states are found to relax rapidly to lower states, considering S_1 to S_3 (for trajectories and therefore also for starting points) in the excited state dynamics is a valid compromise between accuracy and computational effort.

Fig. 6 compares simulated emission spectra for the four minima on the ADC(2)/cc-pVDZ PES for S_1 , each obtained from a Wigner ensemble with 500 structures (*vide supra*) with the experimental spectrum measured in 1-chlorobutane. The spectra computed separately for the four S_1 minima are all similar. Compared to the experimental spectrum in solution, they are red-shifted by about 0.5–0.6 eV. This red shift is most likely partly due to the known underestimation of the vertical excitation energy for the TICT state in the ABN fragment by ADC(2) and CC2.²⁷ A second reason might be that experimentally the fluorescence can take place before a vibrational ground-state on the S_1 PES is reached (*vide infra*).




Fig. 6 Comparison of the fluorescence spectrum. The experimental spectrum²⁰ in 1-chlorobutane is shown in red, the computed spectra for the four different excited-state minima are shown in orange (S_1 -Min-a), purple (S_1 -Min-b), green (S_1 -Min-c) and blue (S_1 -Min-d based). Computed on ADC(2)/cc-pVDZ level of theory.

3.3 Electronic evolution

To analyze the population of the electronic states, we counted for each state how many trajectories populate it, *i.e.* have this state set as active state. The adiabatic populations (including those from trajectories terminated prematurely) are shown in Fig. 7. The net number of hops provides evidence that trajectories initially excited to S_3 sequentially decay *via* S_2 to S_1 , and that trajectories initially excited to S_2 decay to S_1 , *i.e.* the dominant electronic relaxation pathway seems to be $S_3 \rightarrow S_2 \rightarrow S_1$. From the 15 trajectories that hopped during the simulation to the ground state, 11 were forced to hop due to an energy difference lower than 0.1 eV between the active state and S_0 . To obtain time constants for the population transfer, the kinetic model $S_{32} \rightarrow S_1 \rightarrow S_0$ was chosen where S_{32} denotes the joint population of S_3 and S_2 . The population fit implemented in SHARC was used for setting up the differential equations for the rate laws, solving them with a fifth-order Runge–Kutta scheme, and fitting the kinetic parameters using a nonlinear least squares fit with bootstrapping.⁶⁷ Time constants of 47 ± 7.4 fs for $S_{32} \rightarrow S_1$ and 6.4 ± 2.4 ps for $S_1 \rightarrow S_0$ were obtained.

Fig. 8 shows a simulated transient fluorescence spectrum computed from the 55 successfully completed trajectories that remained in an excited state using for each trajectory and time step the excitation energy and oscillator strength of the current active state. The spectrum was smoothened using Gaussian convolution in energy (FWHM of 0.5 eV) and time (FWHM of 25 fs). Two emission bands can be identified from the simulated spectrum in vacuum: a bright band that appears immediately after excitation (0–300 fs) at around 4 eV which is attributed to locally excited states and a second emission band which builds up with the decay of the first band at energies around 2.1 eV and a low oscillator strength which is assigned to the charge transfer state. The latter transition energy is in rather good agreement with the experimental value of 2.41 eV, considering the expected underestimation of the transition energies for TICT states by ADC(2) (*vide supra*).




Fig. 7 Classic electronic population of singlet S_1 (green), S_2 (blue), S_3 (purple) and S_0 (red) over all converged trajectories.






Fig. 8 Left: Simulated transient fluorescence spectrum at the ADC(2)/cc-pVDZ level from the 55 completed trajectories that remained in the excited state for 1000 fs. Right: Time evolution of the fluorescence signals from the locally excited and the charge-transfer band, dashed lines from the exponential fit for the relaxation from locally excited to charge transfer state (see text).

To analyze the time evolution of the two emission bands, we determined the integrated relative population of the LE and the CT bands as a function of time. For this purpose, we defined two energy windows (above and below 3.5 eV) and integrated the time-dependent spectrum in these ranges. The integrated fluorescence intensities (without including the Gaussian convolution in time) have then been fitted to an exponential model for the decay of the LE and the rise of the CT fluorescence signal as, respectively $A \cdot \exp(-t/\tau_1) + B \cdot \exp(-t/\tau_2)$ and $(C(1 - \exp(-t/\tau_1)))$, using the nonlinear least-squares (NLLS) Marquardt–Levenberg algorithm of gnuplot⁶⁸ via the parameters A , B , C , τ_1 , and τ_2 . The time constant τ_1 was obtained as 190 fs, τ_2 as 5 fs. The full time evolutions of the LE and CT bands and the fitted exponential model are shown in Fig. 8.

To verify the interpretation of the two bands as originating from locally excited and charge-transfer configurations and to further analyse them, we computed charge transfer numbers⁶⁹ as partial summations over squared transition density matrix elements, using the TheoDORÉ program.^{70,71} For this, we partitioned IYABN into the indole, the amino, and the benzonitrile (phenyl plus cyano) groups. The analysis of the transitions and the assignment to the defined groups in TheoDORÉ is based on a Löwdin population analysis. The time evolution of the electronic character of the excitations obtained from this analysis is shown in Fig. 9.

Initially, the molecule is predominantly in an excited state localized in the indole moiety. With increasing time, the population of the local excitation on indole decays while the population of different CT contributions increases. After 1000 fs the excitation is dominated by CT transitions from the indole and amino groups into the benzonitrile group. For comparison with the time evolution for the decay of the LE and rise of the CT band in the spectrum, we fitted the contribution from the local excitation on indole to $A' \exp(-t/\tau')$, the rise of the CT excitation between indole and phenyl to $b + B'(1 - \exp(-t/\tau'))$, and the rise of the CT excitation between amine and phenyl to $c + C(1 - \exp(-t/\tau'))$ with a common time constant τ' using the




Fig. 9 Charge transfer character over time of the electronic wave function in terms of local excitations of the indole, amine, phenyl, CT excitation from indole to phenyl, CT excitation from amine to phenyl and other CT excitations.

same NLLS approach as above. The time constant for the decay of the local excitation on indole and the rise of the CT contributions obtained from this fit is 232 fs, which fits qualitatively to the time constant fitted to the time evolution of the two bands in the simulated fluorescence spectrum and corroborates the above interpretation of the spectrum.

3.4 Nuclear motion

For the relaxation pathways after photoexcitation from the S_0 to the S_1 minima and the return pathways after deexcitation, the most important coordinates are the torsion angles τ_{E_1,E_2} between the indole and the amino group and τ_{E_2,E_3} between the amino and the phenyl group (see Fig. 10). For the following, we use the shorthand notation (X,Y) for $\tau_{E_2,E_3} = X$ and $\tau_{E_1,E_2} = Y$. The dihedrals are mapped into the interval $[0^\circ, 180^\circ]$. The S_0 minima are at $\approx(0^\circ, 90^\circ)$ and $\approx(180^\circ, 90^\circ)$ (after rotation of the



Fig. 10 Definition of the planes for the indol, amino, and phenyl groups that have been used to measure the torsion angles τ_{E_1, E_2} between the indol and the amino and τ_{E_2, E_3} between the amino and the phenyl groups. The planes and their normal vectors have been determined using the recipe from ref. 72 for indol (E_1) from the positions of the atoms C_1 , C_{12} , C_{13} , and C_{14} , for the amino group (E_2) from the positions of the atoms N_2 , N_3 , C_4 , and H, and for the phenyl ring (E_3) from the positions of the atoms C_5 , C_6 , C_{10} , and C_{11} .




Fig. 11 Left: Time evolution of the relaxation pathways after photoexcitation. Shown are the angle between the indol and the amino group, τ_{E_1, E_2} , on the y axes and the angle between the amino and the phenol group, τ_{E_2, E_3} , on the x axes. The time evolution (in fs) is indicated by the color. Indicated are the possible pathways over the planar (dashed, C), twisted (dashed, A) or synchron (dashed, B) structure (as in Fig. 1). Shown are the 55 trajectories that were successfully completed in an excited state. Right: Analysis of the asynchronicity of the indol-amino and amino-phenyl torsions for the relaxation after photoexcitation. Shown are mean value (black line) and the mean value plus/minus one standard deviation (blue shaded area) of δ_{Asy} for the 55 successfully completed trajectories that remained in an excited state along the coordinate s (see text).

conversion from S_3 and S_2 to S_1 is only ≈ 47 fs and after about 300–400 fs the PLATICT structure is reached. On the other hand, the simulated emission spectrum in Fig. 9 shows a small oscillator strength, meaning that the emission and deexcitation to the ground state is slow. Thus, in solution, the hop to the ground state should occur mostly after some of the kinetic excess energy has been dissipated to the environment. Therefore, we started the simulation of the dynamics for the relaxation after deexcitation to the ground state from Wigner distributions for the four excited state minima.

Fig. 12 shows the time evolution of the torsion angles τ_{E_1, E_2} and τ_{E_2, E_3} during the simulation of the relaxation on the

ground-state PES after fluorescence for the selected 164 trajectories. The angle between the phenyl ring and the amino group, τ_{E_2, E_3} , starts at around 90° and reaches values around the ground-state minima of $\approx 0^\circ$ or $\approx 180^\circ$ after about 200 fs. (Note, that because of the symmetry of the phenyl ring the structures at 0° and 180° are symmetry-equivalent.) The angle between the amino and the indol group, τ_{E_1, E_2} , has values around 0° or 180° at the start of the trajectories. It reaches $\approx 90^\circ$ after about 150 fs and then oscillates between $\approx 50^\circ$ and $\approx 130^\circ$ (around 90°) every 200–300 fs. The simulation shows that after emission the molecule returns to one of the ground-state minima in Fig. 2, *i.e.* τ_{E_1, E_2} relaxes back to a twisted and




Fig. 12 Left: Time evolution of the relaxation pathways on the ground-state PES after emission. Shown are the angle between the indol and the amino group, τ_{E_1, E_2} , on the y axes and the angle between the amino and the phenol group, τ_{E_2, E_3} , on the x axes. The time evolution (in fs) is indicated by the color. Initial conditions were taken from separate Wigner distributions for the four excited-state minima S_1 -Min-a, S_1 -Min-b, S_1 -Min-c and S_1 -Min-d. Indicated are the possible pathways over the planar (dashed, F), twisted (dashed, D) or synchron (dashed, E) structure (as in Fig. 1). Right: Analysis of the asynchronicity of the indol-amino and amino-phenyl torsions for the relaxation after photoemission. Shown are the mean value (black line) and the mean value plus/minus one standard deviation (blue shaded area) of δ_{Asy} for all successfully completed trajectories, weighted by the relative populations of the S_1 minima found after the end of the excited state trajectories along the coordinate s (see text).



τ_{E_2, E_3} to a planar orientation. Apart from a 180° rotation of the phenyl ring, we observe a bifurcation of the pathways into groups relaxing either to S_0 -Min-1 or S_0 -Min-2 with final geometries around $(0^\circ, 90^\circ)$ and into their enantiomers with final geometries around $(180^\circ, 90^\circ)$.

To assign the structures assumed at the end of trajectories at 1000 fs to the different S_0 minima, we started again for each of them from the final structure a geometry optimization. Depending on the excited state minimum from which the trajectories were started from, we found different distributions of the end structures to the ground-state minima (for more information about the disentangled instead of the cumulative behavior of the ground state simulations see Table S2, ESI†). If the number of trajectories are weighted according to the ratio with which they are reached in the excited state dynamics, $28\% \pm 6\%$ of the trajectories arrive in S_0 -Min-1 or its enantiomer and $72\% \pm 6\%$ in S_0 -Min-2 or its enantiomer.

To analyze the asynchronicity between the rotations around the indol-amino and the amino-phenyl bonds, we determined again δ_{Asy} as function of the coordinate s . Fig. 12 depicts its mean value and its standard deviation that have been evaluated by weighting the ensembles generated for the four S_1 minima by the respective populations found at the end of the excited-state dynamics.

Similar as during the excited-state dynamics, the two torsion angles change also for the relaxation after photoemission on the ground-state PES to a large extend but not fully synchronously. Over the whole pathway, the trajectories exhibit on average a small negative asynchronicity, *i.e.* a slight preference towards twisted intermediate structures. When analyzing the trajectories started from the different excited state minima separately, different trends are observed: for the first fifth of the pathway, the trajectories show on average negative asynchronicity, regardless of the minimum from which they have been started. Thereafter, the trajectories started from S_1 -Min-a have on average mostly a twisted character only showing a small planar character around $s = 0.5$. The trajectories started from S_1 -Min-b exhibit on average exclusively a twisted character. During the latter phase, the asynchronicity increases up to 20% of a fully twisted structure. Overall, the trajectories started from S_1 -Min-b show on average the largest asynchronicity. The trajectories started from S_1 -Min-c exhibit on average a twisted character, with a small planar character at around $s = 0.2$. The trajectories started from S_1 -Min-d oscillate between a planarized (on average up to 5% of the asynchronicity of a completely planarized structure) and a twisted asynchronicity (up to 5%). Across all time steps, the trajectories started from all minima have a preferentially twisted character (while S_1 -Min-d has the smallest asynchronicity character).

3.5 Pathway of ground and excited state compared

As pointed out in the introduction, the pathways for the rotation around the indol-amino and the amino-phenyl bonds are important to assess the potential of *N*-aryl-substituted 1-aminoindoles for applications as molecular motors. For this, the full cycle of movements in the excited and in the ground

state should form an unidirectional 360° rotation or there has to be at least a difference in the pathways for the forward and backward motions. Comparing Fig. 11 and 12, we can observe that the relaxation pathways in both the ground and the excited state run on average over intermediate structures with some degree of asynchronicity towards fully twisted structures. For the whole simulation pathway, this is more pronounced in the ground state.

Integrating the asynchronicity over the pathways gives an idea about the effectiveness as motor. The difference of the integrals for the forward motion in the excited state and the backward motion on the ground state PES is a diagnostic that shows whether some mechanical work is done during a full cycle. The maximal mechanical work would be done along a hypothetical pathway which in the excited state would run through totally planarized structure and in the ground state through totally twisted structure or *vice versa*, *i.e.* a pathway with full asynchronicity ($\delta_{\text{Asy}} = \pm 1$) in one direction in the excited state and the opposite asynchronicity in the ground state.

For IYABN we obtained for the trajectories on average a result from the Riemann summation that is 1% of the hypothetical maximum (with a standard deviation of 13%). Even if this number should be taken with caution, it indicates that although the pathways in the ground and excited states are very similar, they are not identical and that a small amount of kinetic work could be achieved during a full cycle.

4 Conclusions

The photophysical behaviour of IYABN after excitation with UV/VIS light has been investigated with static calculations as well as with surface hopping molecular dynamics simulations. The ground-state absorption spectrum has been computed by sampling vertical spectra calculated with ADC(2)/cc-pVDZ for a Wigner distribution for the ground-state equilibrium structures. Band shape and band position agree for the absorption within about 0.1 eV with the experimental absorption spectrum recorded in 1-chlorobutane once solvent and remaining basis set effects have been accounted for based on corrections from a single-point COSMO-ADC(2)/aug-cc-pVTZ calculation.

The relaxation dynamics surface hopping simulations including the first 3 excited singlet states were carried out at the MP2/cc-pVDZ and ADC(2)/cc-pVDZ levels of theory, for the ground and the excited states, respectively. The results show that after the photoexcitation the system decays fast, with a time constant of about 47 fs to the lowest excited singlet state S_1 and on the S_1 PES within a few hundred fs to an intermolecular charge-transfer state with planarized amino-indol and twisted amino-phenyl torsion angles. No other minima in the excited state are reached during the relaxation. The time constant for the decay to the ground state found from the surface hopping dynamics is about 6.4 ps.

In the PLATICT state, one electron has been moved from the amino-indole donor group to the benzonitrile acceptor group. As a consequence, the oscillator strength for emission is very low.



The simulated transient emission spectrum shows two bands: a bright band around 4 eV which appears immediately after excitation and decays rapidly with a time constant of 190 fs and a second, almost dark band at ≈ 2.4 eV which appears with the decay of the first band and persists until the end of the trajectories. The transition energy of the latter band is in good agreement with the experimental fluorescence spectrum recorded in 1-chlorobutane.²⁰

The geometric analysis of the relaxation pathways after photoexcitation and after emission show that the rotations around the indol-amino and the amino-phenyl bonds happen both, in the excited state and on the ground state PES, almost synchronously. We observe only small asynchronicities of the rotations which are in the excited and in the ground state directed such that the pathways are on average a bit closer to fully twisted than to fully planarized intermediate structures. Comparison of the ground and excited state analysis reveals that the ground state dynamics occurs on average over more twisted structures than the relaxation in the excited states, depending on which of the energetically close-lying PLATICT minima is reached during relaxation in the excited state.

Because of the similarity of the relaxation pathways in the excited and the ground state, overall, only a small amount of mechanical work is done during a full relaxation cycle of IYABN. Still, the results indicate that the combination of PLICT and TICT chromophores to a PLATICT functionality can in principle be used to build molecular motors. The amount of accessible mechanical work per relaxation cycle can probably be increased by introducing higher structural asymmetry and sterical guidance by adding substituents with steric demands to achieve a higher degree of directionality for the relaxation pathways. Alternatively, the nitrile group can be changed or other substituents with electronic effects be introduced either in the phenyl or in the indole fragment of the *N*-aryl-1-aminoindole.

In this regard, it is already known from previous work²⁰ that removing the nitrile group or replacing it by a carboxylic ester (MeOOC-) group, or a fluor atom can change the excited-state potential energy surfaces significantly.²⁰ In computational studies of the fluoro and the unsubstituted counterparts of IYABN in vacuum not only a PLATICT but also a PLICT minimum was found. The PLATICT minima showed again large Stokes shifts of approx. 18 000 cm⁻¹ (for the unsubstituted *N*-(1*H*-indole-1-yl)aniline), and 16 000 cm⁻¹ (for the F-substituted counterpart *N*-(4-fluorophenyl)indol-1-amine), for the PLICT minima the Stokes shifts are only 11 000 cm⁻¹ (for the unsubstituted *N*-aryl-1-aminoindole) and 10 000 cm⁻¹ (for the F-substituted-*N*-aryl-1-aminoindole). Experimentally, only one fluorescence signal with a Stokes shift of 10 000 cm⁻¹ to 11 000 cm⁻¹ was found for both molecules in different solvents, which was assigned to a PLICT minimum. This shows that the potential energy surface for S₁ can be altered such that the PLICT structure becomes a minimum on the S₁ PES by changing the substituents. Our calculations have shown that for the CN-substituted derivative the PLICT structure is not a minimum and also not accessible as intermediate structure for the relaxation after photoexcitation. Tuning the S₁ PES by substitution at the

phenol and the indole subsystems such that the PLICT structure is low enough in energy to be accessible along the relaxation pathways, but not a stable minimum could open up relaxation pathways running over planarized structures in S₁. If the planar structure on the S₀ potential surface is not strongly stabilized by the same substitution so that the relaxation pathways for the backward motion on the S₀ PES doesn't change in the same way, the difference in forward and backward motions could be significantly enlarged and more kinetic work could become accessible.

Conflicts of interest

The authors declare no conflict of interest.

Data availability

The data that support the findings of this study are available on Zenodo at <https://doi.org/10.5281/zenodo.15267389>, reference number 15267389 and in the ESI.†

Acknowledgements

Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy-EXC 2033-390677874-RESOLV. We further acknowledge Vienna Scientific Cluster (VSC4 and VSC5) for allocation of computational resources and the University of Vienna for continuous support. We thank Prof. Gebhard Haberhauer for inspiring discussions and access to original data from ref. 20 for visualization.

Notes and references

- 1 S. Kassem, T. van Leeuwen, A. S. Lubbe, M. R. Wilson, B. L. Feringa and D. A. Leigh, *Chem. Soc. Rev.*, 2017, **46**, 2592–2621.
- 2 R. Feynman, R. Leighton, M. Sands and E. Hafner, *The Feynman Lectures on Physics*; Vol. I, AAPT, 1965, vol. 33, p. 750.
- 3 K. Verhey, J. Dishinger and H. Kee, *Biochem. Soc. Trans.*, 2011, **39**, 1120–1125.
- 4 M. J. Lynch, R. Levenson, E. A. Kim, R. Sircar, D. F. Blair, F. W. Dahlquist and B. R. Crane, *Structure*, 2017, **25**, 317–328.
- 5 C. Kallweit, G. Haberhauer and S. Woitschetzki, *Chem. – Eur. J.*, 2014, **20**, 6358–6365.
- 6 E. R. Kay, D. A. Leigh and F. Zerbetto, *Angew. Chem., Int. Ed.*, 2007, **46**, 72–191.
- 7 G. Haberhauer, *Angew. Chem., Int. Ed.*, 2011, **50**, 6415–6418.
- 8 C. García-Iriepa, M. Marazzi, F. Zapata, A. Valentini, D. Sampedro and L. M. Frutos, *J. Phys. Chem. Lett.*, 2013, **4**, 1389–1396.
- 9 J. Wang and B. Durbeej, *ChemistryOpen*, 2018, **7**, 583–589.
- 10 J. Wang, B. Oruganti and B. Durbeej, *Org. Lett.*, 2020, **22**, 7113–7117.



- 11 J. Michl and E. C. H. Sykes, *ACS Nano*, 2009, **3**, 1042–1048.
- 12 D. T. Payne, J. Labuta, Z. Futera, V. Březina, L. Hanyková, M. K. Chahal and J. P. Hill, *Org. Chem. Front.*, 2022, **9**, 39–50.
- 13 R. Eelkema, M. Pollard, J. Vicario, N. Katsonis, B. Serrano Ramon, C. Bastiaansen, D. Broer and B. Feringa, *Nature*, 2006, **440**, 163.
- 14 S. Stolz, O. Gröning, J. Prinz, H. Brune and R. Widmer, *Proc. Natl. Acad. Sci. U. S. A.*, 2020, **117**, 14838–14842.
- 15 S. P. Fletcher, F. Dumur, M. M. Pollard and B. L. Feringa, *Science*, 2005, **310**, 80–82.
- 16 F. Romeo-Gella, I. Corral and S. Faraji, *J. Chem. Phys.*, 2021, **154**, 064111.
- 17 N. Koumura, R. Zijlstra, R. van Delden, N. Harada and B. Feringa, *Nature*, 1999, **401**, 152–155.
- 18 G. Pérez-Hernández and L. González, *Phys. Chem. Chem. Phys.*, 2010, **12**, 12279–12289.
- 19 J. Wen, S. Mai and L. González, *J. Phys. Chem. A*, 2023, **127**, 9520–9529.
- 20 G. Haberhauer, *Chem. – Eur. J.*, 2017, **23**, 9288–9296.
- 21 S. Sasaki, G. P. C. Drummen and G.-i Konishi, *J. Mater. Chem. C*, 2016, **4**, 2731–2743.
- 22 C. Wang, W. Chi, Q. Qiao, D. Tan, Z. Xu and X. Liu, *Chem. Soc. Rev.*, 2021, **50**, 12656–12678.
- 23 G. Haberhauer, R. Gleiter and C. Burkhardt, *Chem. – Eur. J.*, 2016, **22**, 971–978.
- 24 W. Rettig, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 971–988.
- 25 R. Misra and S. Bhattacharyya, *Intramolecular Charge Transfer: Theory and Applications*, Wiley-VCH Verlag GmbH Co. KGaA, London, 2018.
- 26 W. Rettig and B. Zietz, *Chem. Phys. Lett.*, 2000, **317**, 187–196.
- 27 A. Köhn and C. Hättig, *J. Am. Chem. Soc.*, 2004, **126**, 7399–7410.
- 28 C. Zhong, *Phys. Chem. Chem. Phys.*, 2015, **17**, 9248–9257.
- 29 W. Rettig, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 971–988.
- 30 Z. R. Grabowski, K. Rotkiewicz and W. Rettig, *Chem. Rev.*, 2003, **103**, 3899–4032.
- 31 J. C. Tully, *J. Chem. Phys.*, 1990, **93**, 1061–1071.
- 32 M. Richter, P. Marquetand, J. González-Vázquez, I. Sola and L. González, *J. Chem. Theory Comput.*, 2011, **7**, 1253–1258.
- 33 S. Mai, P. Marquetand and L. González, *Wiley Interdiscip. Rev.:Comput. Mol. Sci.*, 2018, **8**, e1370.
- 34 S. Mai, D. Avagliano, M. Heindl, P. Marquetand, M. F. S. J. Menger, M. Oppel, F. Plasser, S. Polonius, M. Ruckebauer, Y. Shu, D. G. Truhlar, L. Zhang, P. Zobel and L. González, *SHARC3.0: Surface Hopping Including Arbitrary Couplings – Program Package for Non-Adiabatic Dynamics*, 2023, DOI: [10.5281/zenodo.7828641](https://zenodo.org/record/7828641).
- 35 A. B. Trofimov and J. Schirmer, *J. Phys. B:At., Mol. Opt. Phys.*, 1995, **28**, 2299.
- 36 A. Trofimov, I. Krivdina, J. Weller and J. Schirmer, *Chem. Phys.*, 2006, **329**, 1–10.
- 37 TURBOMOLE V7.7 2022, a development of University of Karlsruhe and Forschungszentrum Karlsruhe GmbH, 1989–2007, TURBOMOLE GmbH, since 2007; available from <https://www.turbomole.org>.
- 38 S. G. Balasubramani, G. P. Chen, S. Coriani, M. Diedenhofen, M. S. Frank, Y. J. Franzke, F. Furche, R. Grotjahn, M. E. Harding, C. Hättig, A. Hellweg, B. Helmich-Paris, C. Holzer, U. Huniar, M. Kaupp, A. Marefat Khah, S. Karbalaei Khani, T. Müller, F. Mack, B. D. Nguyen, S. M. Parker, E. Perlt, D. Rappoport, K. Reiter, S. Roy, M. Rückert, G. Schmitz, M. Sierka, E. Tapavicza, D. P. Tew, C. van Wüllen, V. K. Voora, F. Weigend, A. Wodyski and J. M. Yu, *J. Chem. Phys.*, 2020, **152**, 184107.
- 39 M. Häser and R. Ahlrichs, *J. Comput. Chem.*, 1989, **10**, 104–111.
- 40 C. Hättig and F. Weigend, *J. Chem. Phys.*, 2000, **113**, 5154–5161.
- 41 C. Hättig, A. Hellweg and A. Köhn, *Phys. Chem. Chem. Phys.*, 2006, **8**, 1159–1169.
- 42 C. Hättig and A. Köhn, *J. Chem. Phys.*, 2002, **117**, 6939–6951.
- 43 C. Hättig, *Adv. Quantum Chem.*, 2005, **50**, 37–60.
- 44 J. Schirmer, *Phys. Rev. A*, 1982, **26**, 2395–2416.
- 45 F. Weigend, A. Köhn and C. Hättig, *J. Chem. Phys.*, 2001, **116**, 3175–3183.
- 46 T. H. Dunning, *J. Chem. Phys.*, 1989, **90**, 1007–1023.
- 47 R. A. Kendall, T. H. Dunning and R. J. Harrison, *J. Chem. Phys.*, 1992, **96**, 6796–6806.
- 48 D. E. Woon and T. H. Dunning, *J. Chem. Phys.*, 1994, **100**, 2975–2988.
- 49 A. Klamt and G. Schüürmann, *J. Chem. Soc., Perkin Trans. 2*, 1993, 799–805.
- 50 S. Karbalaei Khani, A. Marefat Khah and C. Hättig, *Phys. Chem. Chem. Phys.*, 2018, **20**, 16354–16363.
- 51 A. A. Maryott, E. R. Smith, *Table of Dielectric Constants of Pure Liquids*, U. S. Govt. Print. Off., 1951, p. 13, <https://apps.dtic.mil/dtic/tr/fulltext/u2/a278956.pdf>.
- 52 S. Budavari, *The Merck index: an encyclopedia of chemicals, drugs, and biologicals*, Merck, Rahway, N.J., U.S.A., 11th edn, 1989, p. 239.
- 53 R. Schinke, *Photodissociation Dynamics: Spectroscopy and Fragmentation of Small Polyatomic Molecules*, Cambridge University Press, 1993.
- 54 J. P. Dahl and M. Springborg, *J. Chem. Phys.*, 1988, **88**, 4535–4547.
- 55 S. Mai, P. Marquetand and L. González, *Int. J. Quantum Chem.*, 2015, **115**, 1215–1231.
- 56 D. Tuna, D. Lefrançois, Ł. Wolański, S. Gozem, I. Schapiro, T. Andruniów, A. Dreuw and M. Olivucci, *J. Chem. Theory Comput.*, 2015, **11**, 5758–5781.
- 57 F. Plasser, R. Crespo-Otero, M. Pederzoli, J. Pittner, H. Lischka and M. Barbatti, *J. Chem. Theory Comput.*, 2014, **10**, 1395–1405.
- 58 G. Granucci, M. Persico and A. Toniolo, *J. Chem. Phys.*, 2001, **114**, 10608–10615.
- 59 F. Plasser, M. Ruckebauer, S. Mai, M. Oppel, P. Marquetand and L. González, *J. Chem. Theory Comput.*, 2016, **12**, 1207–1219.
- 60 S. Mai, F. Plasser, M. Pabst, F. Neese, A. Köhn and L. González, *J. Chem. Phys.*, 2017, **147**, 184109.
- 61 G. Granucci, M. Persico and A. Zocante, *J. Chem. Phys.*, 2010, **133**, 134111.
- 62 M. Barbatti, G. Granucci, M. Persico, M. Ruckebauer, M. Vazdar, M. Eckert-Maksić and H. Lischka, *J. Photochem. Photobiol., A*, 2007, **190**, 228–240.



- 63 R. L. Martin, *J. Chem. Phys.*, 2003, **118**, 4775–4777.
- 64 O. Christiansen, H. Koch and P. Jørgensen, *Chem. Phys. Lett.*, 1995, **243**, 409–418.
- 65 S. Mai, F. Plasser, M. Pabst, F. Neese, A. Köhn and L. González, *J. Chem. Phys.*, 2017, **147**, 184109.
- 66 F. Neese, *Wiley Interdiscip. Rev.:Comput. Mol. Sci.*, 2012, **2**, 73–78.
- 67 S. Nangia, A. W. Jasper, T. F. Miller and D. G. Truhlar, *J. Chem. Phys.*, 2004, **120**, 3586–3597.
- 68 T. Williams, C. Kelley *et al.* *Gnuplot 5.2.8: an interactive plotting program*, <https://gnuplot.sourceforge.net/>, 2019.
- 69 S. Mai, F. Plasser, J. Dorn, M. Fumanal, C. Daniel and L. González, *Coord. Chem. Rev.*, 2018, **361**, 74–97.
- 70 F. Plasser and H. Lischka, *J. Chem. Theory Comput.*, 2012, **8**, 2777–2789.
- 71 F. Plasser, *J. Chem. Phys.*, 2020, **152**, 084108.
- 72 D. Cremer and J. A. Pople, *J. Am. Chem. Soc.*, 1975, **97**, 1354–1358.

