


Cite this: *RSC Adv.*, 2020, 10, 9309

The effect of solvents on the thermal degradation products of two Amadori derivatives†

Shiyi Zhang, Rui Li, * Yudan Zhang and Mingqin Zhao *

To enrich the flavor additives of the Maillard reaction, two Amadori analogs, *N*-(1-deoxy-*D*-fructosyl-1-yl)-*L*-phenylalanine ester (Derivative 1) and di-*O*-isopropylidene-2,3:4,5- β -*D*-fructopyranosyl phenylalanine ester (Derivative 2), were chemically synthesized starting from *D*-fructose. The samples were reacted at 120 and 180 °C for 2 h, and the effects of solvents (water and ethanol) on their degradation products were studied. The analyses of thermogravimetry (TG), derivative thermogravimetry (DTG), differential scanning calorimetry (DSC), and gas chromatography-mass spectrometry (GC/MS) were used to investigate the thermal behavior and degradation products of the samples. TG–DTG curves show that the T_p values of the samples corresponding to the largest mass-loss rates are 132 and 275 °C, respectively. The degradation products of Derivative 1 are mainly phenyl acetaldehyde and phenylalanine ethyl ester in water and ethyl benzoate and benzaldehyde diethyl acetal in ethanol. For Derivative 2, the major degradation products both in water and ethanol are phenylalanine ethyl ester and diacetonefructose, but the products have different relative contents affected by solvent media. The products of the pyrolysis of the samples at 350 °C were analyzed and compared with the degradation compounds obtained in solvent. These results show that organic solvents can greatly influence the degradation pathway and products. Finally, possible mechanisms of the degradation processes are proposed.

Received 25th November 2019
Accepted 26th February 2020

DOI: 10.1039/c9ra09854b

rsc.li/rsc-advances

1. Introduction

The Amadori compounds, produced in the initial stage of the Maillard reaction, are ubiquitous during the manufacture and storage of food such as bread, cakes, potatoes, meat, coffee, and tobacco.^{1–4} The degradation products of Amadori compounds include volatile aromas, which directly influence the organoleptic and nutritional qualities of foodstuffs.^{5–7} Mills *et al.*⁸ and Huyghues-Despointes *et al.*⁹ studied the pyrolysis of 1-*L*-proline-1-deoxy-*D*-fructose at various temperatures and reported the generation of compounds such as dihydrofuran, dihydropyrones, and pyrrolidines with bread aromas. Additionally, Shigematsu *et al.*,¹⁰ Vernin *et al.*¹¹ and Mao *et al.*^{12–14} reported the formation of volatile aroma compounds *via* the thermal degradation of 1-*L*-alanine-1-deoxy-*D*-fructose, 1-*L*-valine-1-deoxy-*D*-fructose, 1-*L*-methionine-1-deoxy-*D*-fructose, 1-*L*-glutamic acid-1-deoxy-*D*-fructose, 1-*L*-phenylalanine-1-deoxy-*D*-

fructose, 1-*L*-leucine-1-deoxy-*D*-fructose, *etc.* Since the degradation products of Amadori compounds play a major role in the flavors of food, factors influencing the degradation processes were investigated. Davidek *et al.*¹⁵ studied the effects of phosphate ions on the degradation products of 1-*L*-glycine-1-deoxy-*D*-fructose, and the results showed that the addition of phosphate ions could accelerate the degradation processes. Shu *et al.*¹⁶ reported that at a pH of 8.5, the major products formed from the degradation of glucosamine were pyrazine and methylpyrazine. Hofmann *et al.*¹⁷ reported that the formation of phenylacetaldehyde was favored during the degradation of 1-*L*-phenylalanine-1-deoxy-*D*-fructose catalyzed by ambient oxygen. However, the effect of organic solvents such as ethanol on the generated degradation products from Amadori derivatives has not yet been studied.

In the present paper, two compounds were chemically synthesized, *N*-(1-deoxy-*D*-fructosyl-1-yl)-*L*-phenylalanine ester (Derivative 1) and di-*O*-isopropylidene-2,3:4,5- β -*D*-fructopyranosyl phenylalanine ester (Derivative 2) (Fig. 1), starting from *D*-fructose.^{18,19} In Derivative 2, the hydroxyl groups on the pyranose ring are protected by acetal groups to disfavor the isomerization of the sugar moiety. The two derivatives were analyzed by TG, DTG, and DSC, and they were pyrolyzed. The effects of the solvent media on the thermal degradation products were then assessed by GC/MS. The results show that the solvent can influence the degradation of products to a certain extent.

College of Tobacco Science, Henan Agricultural University, Zhengzhou 450002, China.
E-mail: ganguo1989@henau.edu.cn; zhaomingqin@126.com

† Electronic supplementary information (ESI) available: Including ¹H NMR and ¹³C NMR spectra and TLC results of Derivative 1 and Derivative 2 (Fig. S1–S5, Scheme S1), all the total ion chromatograms (TICs) from the solvent-phase and pyrolytic degradations of Derivative 1 and Derivative 2 at different temperatures (Fig. S6–S15), mass spectra of degradation compounds extracted from the TICs (Fig. S16–S38), and the proposed degradation mechanisms of Derivative 2 (Scheme S2). See DOI: 10.1039/c9ra09854b






Fig. 1 Chemical structures of Derivative 1 and Derivative 2.




Fig. 2 TG and DTG curves of Derivative 1.

2. Materials and methods

2.1 Materials

Methanol, dimethylformamide, dichloromethane, acetone, ethyl acetate, trifluoroacetic acid, and petroleum ether were purchased from Tianjin Kermel Chemical Reagent Co., Ltd (China). The above reagents were of analytical grade. Deionized water (filtered through a 0.22 μm membrane filter, $>18.2 \text{ M}\Omega \text{ cm}$) was purified using a Milli-Q system from Millipore. Silica gel was purchased from Qingdao Haiyang Chemical Reagent Co., Ltd (China). Phenylalanine ethyl ester, di-*O*-isopropylidene-2,3,4,5- β -D-fructose and di-*O*-isopropylidene-2,3,4,5- β -D-fructopyranosyl trifluoromethanesulfonate were synthesized according to ref. 18 and 19, respectively.

2.2 Structure identification

^1H NMR and ^{13}C NMR spectra data were recorded on a Bruker 400 MHz spectrometer. Each sample was dissolved in CDCl_3 or CD_3OD containing tetramethylsilane (TMS) as the internal

reference. HRMS data were obtained using an Agilent 1290LC-6540 Accurate Q-TOF mass spectrometer.

2.3 Preparation of Derivative 2

Di-*O*-isopropylidene-2,3,4,5- β -D-fructopyranosyl tri-fluoromethanesulfonate (736.70 mg, 1.88 mmol) and phenylalanine ethyl ester (544.27 mg, 2.82 mmol) were mixed in dimethylformamide (30 mL) and then heated at 120 $^\circ\text{C}$ for 6 h. The progress of the reaction was monitored by thin-layer chromatography (TLC) with petroleum ether–ethyl acetate (3 : 1, v : v). When the starting materials were consumed, the solvent was evaporated under reduced pressure. The crude residue was purified on a silica gel column with petroleum ether–ethyl acetate (3 : 1, v : v) to give the target compound as a yellow oil (342.20 mg, 41.85%). ^1H NMR (400 MHz, CDCl_3 containing 0.03% TMS, 25 $^\circ\text{C}$): δ : 7.28–7.15 (m, 5H, Ar), 4.58 (dd, $J = 7.9 \text{ Hz}$, $J = 2.6 \text{ Hz}$, 1H, CH), 4.38 (d, $J = 2.6 \text{ Hz}$, 1H, CH), 4.22 (dd, $J = 7.9 \text{ Hz}$, $J = 1.0 \text{ Hz}$, 1H, CH), 4.06 (q, $J = 7.2 \text{ Hz}$, 2H, CH_2), 3.81 (dd, $^2J = 64.6 \text{ Hz}$, $^3J = 12.9 \text{ Hz}$, 2H, CH_2), 3.66 (t, $J = 6.8 \text{ Hz}$, 1H, CH), 2.97 (m, 2H, CH_2), 2.84 (s, 2H, CH_2), 1.51 (s, 3H, CH_3), 1.46 (s, 3H, CH_3), 1.38 (s, 3H, CH_3), 1.34 (s, 3H, CH_3), 1.13 (t, $J = 7.2 \text{ Hz}$, 3H, CH_3). ^{13}C NMR (100 MHz, CDCl_3 , 25 $^\circ\text{C}$): δ : 174.21 (CO), 137.19 (C, Ar), 129.27 (C, Ar), 128.36 (C, Ar), 126.64 (C, Ar), 108.91 (C), 108.15 (C), 103.48 (C), 71.12 (CH), 71.07 (CH), 70.40 (CH), 63.23 (CH_2), 61.24 (CH_2), 60.46 (CH), 52.94 (CH_2), 39.26 (CH_2), 26.55 (CH_3), 25.89 (CH_3), 25.56 (CH_3), 24.06 (CH_3), 14.11 (CH_3). HRMS ESI-TOF: $m/z = 458.2154$ [$\text{M} + \text{Na}$] $^+$ (calcd for $\text{C}_{23}\text{H}_{33}\text{NO}_7\text{Na}$, 458.2155).

2.4 Preparation of Derivative 1

Derivative 1 was prepared through the hydrolysis of Derivative 2. The reaction procedure is described below. Derivative 2 (119.10 mg, 0.27 mmol) was stirred in 3.97 mL of trifluoroacetic acid–water (v : v 9 : 1) for 1 h at room temperature. Then, the solvent was removed on a rotary evaporator at 45 $^\circ\text{C}$. The crude residue was purified on a silica gel column with ethyl acetate to give the target compound as a white oil (65.67 mg, 68.44%). ^1H NMR (400 MHz, MeOD, 25 $^\circ\text{C}$): δ : 7.33–7.30 (m, 2H, Ar), 7.28–7.22 (m, 3H, Ar), 4.20–4.15 (m, 1H, CH), 4.11 (q, $J = 7.1 \text{ Hz}$, 2H, CH_2), 3.98–3.95 (m, 1H, CH), 3.87 (m, 1H, CH), 3.81–3.77 (m, 1H, CH), 3.69–3.63 (m, 2H, CH_2), 3.28–3.08 (m, 2H, CH_2), 3.15 (s, 2H, CH_2), 1.10 (t, $J = 7.1 \text{ Hz}$, 3H, CH_3). ^{13}C NMR (100 MHz, MeOD, 25 $^\circ\text{C}$): pyranose form δ : 171.47 (CO), 136.36 (C, Ar), 130.49 (C, Ar), 130.06 (C, Ar), 128.83 (C, Ar), 97.26 (C), 71.46 (CH), 71.25 (CH), 70.81 (CH), 64.93 (CH), 63.23 (CH_2), 63.14 (CH_2), 53.56 (CH_2), 37.81 (CH_2), 14.22 (CH_3). Furanose form δ :

Table 1 Thermal analysis data for Derivative 1 and Derivative 2

Sample name	DSC				TG-DTG	
	$T_{\text{onset}}/^\circ\text{C}$	$T_{\text{peak}}/^\circ\text{C}$	$T_{\text{end}}/^\circ\text{C}$	$\Delta H/\text{kJ mol}^{-1}$	$T_{\text{p}}/^\circ\text{C}$	$T_{\text{range}}/^\circ\text{C}$
Derivative 1 (peak 1)	110	132	146	25	132	103–193
Derivative 1 (peak 2)	247	255	272	–13	254	217–279
Derivative 2	81	226	351	–852	275	160–341



170.21 (CO), 135.66 (C, Ar), 130.44 (C, Ar), 129.87 (C, Ar), 128.54 (C, Ar), 103.20 (C), 84.54 (CH), 84.32 (CH), 84.10 (CH), 77.62 (CH), 62.61 (CH₂), 52.82 (CH₂), 51.79 (CH₂), 37.18 (CH₂), 14.22 (CH₃). HRMS ESI-TOF: $m/z = 356.1708$ [$M + H$]⁺ (calcd for C₁₇H₂₆NO₇, 356.1709).

2.5 Thermal degradation process²⁰

The thermal degradation of each sample (5 mg) was conducted at 120 °C or 180 °C for 120 min in water (5 mL) or ethanol (5 mL) utilizing sealed 48 mL Synthware glass vials (bought from Beijing Synthware Glass Instrument Co., Ltd.) in oil baths. The reactions were immediately stopped by cooling under a stream of cold water. Solvents were evaporated from the reaction mixtures, and the residues were redissolved in methanol for GC/MS analysis.

2.6 Thermal analysis²¹

TG, DTG, and DSC analyses were conducted using a simultaneous thermal analyzer (NETZSCH STA 449 F3, Germany). The mass of each sample loaded into the alumina crucible for each run was approximately 5 mg. The tests were carried out under an argon atmosphere at a flow rate of 60 mL min⁻¹. Thermal degradation was monitored from 30 to 900 °C at a constant heating rate of 10 °C min⁻¹. The TG, DTG and DSC curves of the samples were simultaneously determined.

2.7 Pyrolysis-GC/MS and GC/MS analyses²²

Pyrolysis-GC/MS analysis was conducted on a combined CDS Pyroprobe 5000 and Agilent 7890/5975 GC/MS system. The pyroprobe was heated at a rate of 5 °C ms⁻¹ from an initial temperature of 50 °C to a final temperature of 350 °C, which was then held for 15 s. The sample mass was approximately 1 mg. The pyrolysis was conducted in air. The corresponding pyrolysis pressure was 1.03×10^6 Pa, and it was higher than the pressure of the GC column. The final pyrolysis products were directly carried into the GC/MS column. Details of the GC/MS analysis conditions are shown below. A DB-5MS fused silica capillary column (50 m × 250 μm id × 0.25 μm df) was used. The injector temperature was 250 °C. The initial oven temperature was 50 °C, and this was held for 2 min; the oven was then heated to 250 °C at a rate of 5 °C min⁻¹, and held at 250 °C for 5 min. The helium carrier gas flow rate remained at 1 mL min⁻¹, and the split ratio was 50 : 1. The EI energy was 70 eV, and the transfer line temperature was 280 °C. The ion source temperature was 230 °C. The quadrupole temperature was 150 °C. The EI-MS scan range was 30–400 amu, and the solvent delay time was 3.6 min. Qualitative analysis of the compound was achieved with the NIST mass spectral search program for the NIST/EPA/NIH mass spectral library version 2.2 (NIST2014, Gaithersburg, USA). Quantitative analysis was based on the area normalization method, which was used to calculate the relative content of each component. All experiments were performed in duplicate, and the quantitative data were expressed by the average values.




Fig. 3 TG and DTG curves of Derivative 2.




Fig. 4 DSC curve of Derivative 1.

3. Results and discussion

3.1 TG-DTG analysis

The TG curve (%) and the DTG curve (DTG, % min⁻¹) of Derivative 1 are presented in Fig. 2. Three stages of thermal degradation were observed. The initial stage, occurring between 103 and 193 °C (Table 1), was the main degradation stage, and approximately 31.3% of the total mass was lost. In this stage,




Fig. 5 DSC curve of Derivative 2.

the highest mass-loss rate appeared at a T_p of 132 °C, and at this point, approximately 11.6% of the initial mass of the sample had been lost. The second stage occurred between 217 and 279 °C (Table 1), and the maximum decomposition rate was at 254 °C. Approximately 14.1% of the initial mass of the sample was lost during this stage. From 284 to 900 °C, the sample exhibited a slow mass-loss process with a mass loss of 30.3%. According to the molecular structure of Derivative 1, the benzyl and carbonyl groups are relatively stable, while the C–OH and C–COO–CH₂CH₃ moieties are unstable groups. The two major mass-loss processes are likely induced by dehydration, deesterification, and decarboxylation.¹³ Upon temperature elevation, C–N bonds are broken as are other chemical bonds.

Fig. 3 shows the mass loss and mass-loss rates of Derivative 2 with increasing temperature. The curve reveals that the main

degradation stage occurred between 160 and 341 °C (Table 1), and in this stage, approximately 98.3% of the total mass was lost. In this stage, the highest mass-loss rate appeared at a T_p of 275 °C, and at this point, approximately 53.0% of the initial mass had been lost. TG–DTG analysis indicates that Derivative 1 and Derivative 2 have good stability at room temperature, and the temperature corresponding to the highest mass-loss rate of Derivative 1 is lower than that of Derivative 2 (Table 1). These phenomena suggest that Derivative 1, with unprotective hydroxyl groups, is more susceptible to thermal degradation.

3.2 DSC analysis

The DSC curves of Derivative 1 and Derivative 2 are shown in Fig. 4 and 5, respectively. In Fig. 4, the T_{peak} values of Derivative 1 are at 132 and 255 °C. The former peak with the onset

Table 2 Pyrolysis products of Derivative 1 and Derivative 2 at 350 °C^a

No.	RT min ^{−1}	Products	Structure	Match (%)	Relative content (%)	
					Derivative 1	Derivative 2
1	8.33	Tetrahydrofuran-3,4-diol		85	2.60	—
2	10.45	Toluene		80	0.29	—
3	12.25	2-Formyl furan		87	0.69	—
4	13.53	Benzaldehyde		87	0.49	0.21
5	14.09	Styrene		89	0.70	—
6	15.69	Phenylmethanol		87	—	0.14
7	19.91	Ethyl benzoate		85	—	0.18
8	21.32	Phenethyl acetate		95	1.32	55.84
9	24.94	Ethyl-3-phenylpropionate		90	38.84	0.84
10	28.04	Ethyl cinnamate		80	4.84	0.83
11	28.33	1-(2-Furyl)-1,2-butanediol		85	—	0.64
12	29.13	Phenylalanine ethyl ester		91	—	2.86
13	31.30	Diacetonefructose		92	—	4.62

^a “—” means that the compound was not detected. The table shows only the main degradation products.



temperature (T_{onset}) of 110 °C corresponds to the first mass-loss process displayed in TG-DTG curves, showing an endothermic degradation. The latter peak with T_{onset} of 247 °C illustrates an exothermic degradation consistent with the second stage shown in TG-DTG curves. On the other hand, the DSC curve of Derivative 2 displays one main exothermic peak with a T_{peak} of 226 °C (Fig. 5). These results suggest that the degradation processes in Derivative 1 are more complex than that in Derivative 2. The onset temperature (T_{onset}), end temperatures (T_{end}) and thermal enthalpy change (ΔH) of the samples in the DSC curves are detected by the apparatus and summarized in Table 1. In general, the DSC data are corresponding to that of TG-DTG results.

3.3 Pyrolysis-GC/MS analysis (degradation without solvents)

TG-DTG and DSC analyses helped elucidate the relationship between temperature change and mass loss (or enthalpy change) in the samples, but these data are not sufficient to identify the molecular structure of the degradation products. To further understand the influence of solvent on the decomposition process, online pyrolysis-GC/MS experiments (degradation without solvents) were conducted at 350 °C in air for Derivative 1 and Derivative 2. The pyrolysis conditions were set based on the degradation temperatures of Derivative 1 and Derivative 2 shown in the TG-DTG curves.

Table 2 lists the main components released from Derivative 1 and Derivative 2 under the pyrolysis conditions. The main pyrolysis products of Derivative 1 are ethyl-3-phenylpropionate, ethyl cinnamate and tetrahydrofuran-3,4-diol. Heterocyclic

aroma-active compounds such as 2-formyl furan are dehydration and oxidation products of pyranoses, while phenyl derivatives are obtained from phenylalanine ester. For the pyrolysis of Derivative 2, the products are mainly phenethyl acetate, phenylalanine ethyl ester, and diacetonefructose. Flavor compounds such as phenylmethanol, benzaldehyde, ethyl benzoate, ethyl-3-phenylpropionate and ethyl cinnamate are also formed. Comparing the products between Derivative 1 and Derivative 2, the number of furan or hydrofuran compounds from Derivative 2 is lower than that from Derivative 1, indicating that the protection of the hydroxyl groups notably restricted the degradation of the pyranose ring.

3.4 Thermal degradation in water and ethanol

To investigate the thermal degradation processes and the distribution of the degradation products obtained in the solvent, Derivative 1 and Derivative 2 were heated at 120 °C and 180 °C in water and ethanol. After evaporating the solvents, the thermal degradation products were analyzed by GC/MS.

Table 3 shows the distribution of degradation products for Derivative 1 and Derivative 2 when the samples were heated at 120 °C and 180 °C in aqueous medium. Derivative 1 releases three compounds, 5-(dimethoxymethyl)-2-furanmethanol (0.71%), ethyl-3-phenylpropionate (0.95%) and phenylalanine ethyl ester (93.38%), at 120 °C. Upon heating to 180 °C, the relative content of phenylalanine ethyl ester drastically decreased to 38.88%, and this was accompanied by the

Table 3 Degradation products of Derivative 1 and Derivative 2 in water at 120 and 180 °C^a

No.	RT min ⁻¹	Products	Structure	Match (%)	Relative content (%)			
					Derivative 1 (120 °C)	Derivative 1 (180 °C)	Derivative 2 (120 °C)	Derivative 2 (180 °C)
1	16.52	Phenylacetaldehyde		88	—	3.85	—	—
2	23.93	5-(Dimethoxymethyl)-2-furanmethanol		87	0.71	2.79	—	—
3	24.94	Ethyl-3-phenylpropionate		90	0.95	—	—	—
4	27.42	Methyl-2-amino-3-phenylpropanoate		89	—	2.53	—	—
5	28.65	2,3:4,5-Bis-O-(1-methylethylidene)-β-D-arabinohexos-2-ulo-2,6-pyranose		88	—	—	—	11.03
6	29.13	Phenylalanine ethyl ester		89	93.38	38.88	5.10	4.23
7	31.30	Diacetonefructose		91	—	—	84.80	70.56














^a “—” means that the compound was not detected. The table shows only the main degradation products.



formation of phenylacetaldehyde (3.85%). As expected, the main degradation products of Derivative 2 in water are phenylalanine ethyl ester (5.10%) and diacetonefructose (84.80%). The relative contents of these two compounds at 180 °C are lower than that at 120 °C because of the formation of a compound derived from diacetonefructose. Besides, fewer

degradation products (3 and 4 compounds from Derivative 1, 2 and 3 compounds from Derivative 2 at 120 and 180 °C respectively) are detected relative to what is seen after pyrolysis (8 compounds from Derivative 1 and 9 compounds from Derivative 2). The lower degradation temperature and the solvent medium are responsible for these results.

Table 4 Products of the degradation of Derivative 1 and Derivative 2 in ethanol at 120 and 180 °C^a

No.	RT min ⁻¹	Products	Structure	Match (%)	Relative content (%)			
					Derivative 1 (120 °C)	Derivative 1 (180 °C)	Derivative 2 (120 °C)	Derivative 2 (180 °C)
1	13.53	Benzaldehyde		90	1.99	—	—	—
2	14.19	Diethyl oxalate		91	0.31	—	—	—
3	15.69	Phenylmethanol		87	0.44	—	—	—
4	16.97	1,1,3-Triethoxypropane		89	0.36	0.99	—	—
5	19.91	Ethyl benzoate		90	6.70	4.52	—	—
6	20.08	2-Methoxy-2-phenylacetic acid		87	10.02	—	—	—
7	21.71	Benzaldehyde diethyl acetal		92	36.53	37.67	—	—
8	22.16	Ethyl phenylacetate		84	1.14	2.13	—	—
9	23.93	5-(Dimethoxymethyl)-2-furanmethanol		85	0.43	—	—	—
10	24.39	(2,2-Diethoxyethyl)benzene		82	2.25	1.76	—	—
11	24.94	Ethyl-3-phenylpropionate		90	5.72	5.11	—	—
12	28.04	Ethyl cinnamate		88	1.84	3.62	—	—
13	28.65	2,3:4,5-Bis-O-(1-methylethylidene)-β-D-arabino-hexos-2-ulo-2,6-pyranose		87	—	—	—	4.30
14	29.13	Phenylalanine ethyl ester		89	5.86	—	6.75	14.10
15	31.30	Diacetonefructose		91	—	—	76.86	69.98

^a “—” means that the compound was not detected. The table shows only the main degradation products.



degradation processes. For Derivative 2, the degradation products are phenylalanine ethyl ester and diacetonefructose with relative contents of 6.75% and 76.86% at 120 °C and 14.10% and 69.98% at 180 °C, respectively. The relative content of phenylalanine ethyl ester from ethanol is higher while the amount of diacetonefructose is lower relative to their levels from water, suggesting that changing the degradation medium is one method to control the degradation products.

Based on the degradation products formed in water and ethanol, possible degradation pathways of the two samples are proposed (Scheme 1). For Derivative 1 in water, the degradation processes essentially follow the rules of 1-L-phenylalanine-1-deoxy-D-fructose. The acyclic form of Derivative 1 undergoes



cyclization and dehydration followed by the elimination of the amino groups, producing 5-hydroxymethyl-2-furfural (HMF) and phenylalanine ethyl ester. The research of Yaylayan *et al.*²³ and Gottschalk *et al.*²⁴ has proven this. 5-(Dimethoxymethyl)-2-furanmethanol is likely formed through the condensation of HMF with methanol (Scheme 1). For the degradation of phenylalanine ethyl ester, ethyl cinnamate is produced by the deamination of phenylalanine ethyl ester, and the hydrogenation of this compound generates ethyl-3-phenylpropionate.²⁵ Additionally, the formation of phenylacetaldehyde occurs through the Strecker degradation of phenylalanine interacting with α -dicarbonyl compounds (Scheme 1).²⁶ The degradation product with largest relative contents in water is phenylalanine ethyl ester (93.38% at 120 °C and 38.88% at 180 °C), which is one of the reactants of Derivative 1.

For the degradation process in ethanol, more degradation products, especially the compounds containing ethyl group, are generated than that produced in water. The content of phenylalanine ethyl ester is greatly reduced. Instead, the aldol condensation reactions between aldehyde compounds and ethanol generate the acetol compounds, that is, benzaldehyde for benzaldehyde diethyl acetal and phenylacetaldehyde for (2,2-diethoxyethyl)benzene (Scheme 1). Here, benzaldehyde could be formed by the free radical-initiated oxidative cleavage of the carbon-carbon double bond of the enolized phenylacetaldehyde.^{27,28} Other ethyl group-containing compounds are also formed. For instance, ethyl phenylacetate and ethyl benzoate are produced through the esterification of ethanol and corresponding acids (Scheme 1).²⁹

In the pyrolysis process in air, the degradation mechanism is of complexity. For example, furfural is generated, which is consistent with previous reports.^{30,31} A possible mechanism is initiated by the oxidation of the C1 or C6 carbon atoms of the glucose moieties, producing aldopentose.³² Other compounds such as toluene and styrene are likely caused by the initial benzylic cleavage followed by the generation of reactive radicals.²⁵

For Derivative 2, the hydrolysis of the C-N bonds in the junction generates the final products, phenylalanine ethyl ester, and diacetonefructose (Scheme S2†). Both elevating the temperature and changing the reaction media could influence the decomposition processes. Specifically, the degradation products in water and ethanol are similar while the relative contents of each compound are different. When pyrolysed in air at 350 °C, abundant phenyl group-containing compounds are generated from phenylalanine ethyl ester (Scheme S2†). Among them, phenethyl acetate with the largest relative content (55.84%) is likely formed from the esterification of acetic acid and phenyl ethanol. Here, acetic acid is reported to generate *via* the hydrolytic β -dicarbonyl cleavages from 1-deoxy-2,4-hexodiulose,³³ one possible intermediate during the pyrolysis process. For phenyl ethanol, it is formed from the hydrogenation of phenylacetaldehyde³⁴ which is a Strecker degradation product of phenylalanine.

4. Conclusions

The thermal behavior and degradation products of Derivative 1 and Derivative 2 were investigated by TG-DTG, DSC, pyrolysis-

GC/MS, and GC/MS analysis. TG analysis indicated that Derivative 1 and Derivative 2 have good stability at room temperature, and the T_p values corresponding to the largest mass losses are 132 and 275 °C, respectively. DSC curves demonstrated that the T_{peak} values are 132 and 255 °C for Derivative 1 and 226 °C for Derivative 2. Pyrolysis-GC/MS analysis showed that characteristic aroma compounds are produced during the degradation of these samples. Further degradation processes of Derivative 1 in water generate new flavor compounds such as phenylacetaldehyde, whereas the reactions in ethanol produce compounds such as phenylacetaldehyde diethyl acetal through the aldol condensation between phenylacetaldehyde and ethanol. The number and content of degradation products from Derivative 2 are also affected by solvent media. These results show that the degradation media can influence the degradation processes by reacting with the Amadori intermediates. The synthesis of other Amadori derivatives and the effects of organic solvents on their degradation products are under investigation.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This research was funded by the Key Science and Technology Program of Science and the Technology Department of Henan Province (No. 132102210042) and the program of Henan Agricultural University (No. 30500569). We would also like to thank Dr Zhiyong Wu from Henan Agricultural University and Dr Yuntao Wang from the Zhengzhou University of Light Industry for data analysis and language modification.

References

- 1 M. N. Lund and C. A. Ray, *J. Agric. Food Chem.*, 2017, **65**, 4537–4552.
- 2 C. Rannou, D. Laroque, E. Renault, C. Prost and T. Sérot, *Food Res. Int.*, 2016, **90**, 154–176.
- 3 A. D. Troise and V. Fogliano, *Trends Food Sci. Technol.*, 2013, **33**, 63–74.
- 4 V. V. Mossine and T. P. Mawhinney, *Adv. Carbohydr. Chem. Biochem.*, 2010, **64**, 291–402.
- 5 K. Aalaei, M. Rayner and I. Sjöholm, *Crit. Rev. Food Sci. Nutr.*, 2019, **59**, 1829–1839.
- 6 R. Jeantet, T. Croguennec, P. Schuck and G. Brulé, *Handbook of Food Science and Technology 1: Food Alteration and Food Quality*, Wiley, Hoboken, 2016, pp. 133–157.
- 7 D. A. Baines, S. Bishara, J. K. Parker and D. S. Mottram, *ACS Symp. Ser.*, 2010, **1024**, 63–69.
- 8 F. D. Mills and J. E. Hodge, *Carbohydr. Res.*, 1976, **51**, 9–21.
- 9 A. Huyghues-Despointes, V. A. Yaylayan and A. Keyhani, *J. Agric. Food Chem.*, 1994, **42**, 2519–2524.
- 10 H. Shigematsu, S. Shibata, T. Kurata, H. Kato and M. Fujimaki, *Agric. Biol. Chem.*, 1977, **41**, 2377–2385.



- 11 G. Vernin, J. Metzger, C. Boniface, M.-H. Murello, A. Siouffi, J.-L. Larice and C. Párkányi, *Carbohydr. Res.*, 1992, **230**, 15–29.
- 12 D. Mao, S. Li, D. Mou, G. Peng, X. Bai, C. Jia and J. Yang, *Acta Tab. Sin.*, 2014, **20**, 18–29.
- 13 Y. Ou, H. Ju, C. Jia and D. Mao, *Tob. Sci. Technol.*, 2011, **4**, 41–46.
- 14 D. Mao, H. Ju, D. Mou and X. Bai, *Acta Tab. Sin.*, 2010, **16**, 1–9.
- 15 T. Davidek, N. Clety, S. Aubin and I. Blank, *J. Agric. Food Chem.*, 2002, **50**, 5472–5479.
- 16 C.-K. Shu, *J. Agric. Food Chem.*, 1998, **46**, 1129–1131.
- 17 T. Hofmann and P. Schieberle, *J. Agric. Food Chem.*, 2000, **48**, 4301–4305.
- 18 D. Wang, Y. Liang, P. Gong and Y. Zhao, *Chin. J. Med. Chem.*, 2002, **12**, 94–96.
- 19 D. Xenakis, N. Moll and B. Gross, *Synthesis*, 1983, **7**, 541–543.
- 20 G. Lu, T.-H. Yu and C.-T. Ho, *J. Agric. Food Chem.*, 1997, **45**, 233–236.
- 21 M. Lai, X. Ji, T. Tao, Y. Shan, P. Liu and M. Zhao, *J. Therm. Anal. Calorim.*, 2017, **128**, 1627–1638.
- 22 S. Zeng, S. Sun, S. Liu, J. Hu and B. He, *J. Therm. Anal. Calorim.*, 2014, **115**, 1049–1056.
- 23 V. A. Yaylayan and P. Sporns, *Food Chem.*, 1987, **26**, 283–305.
- 24 A. Gottschalk, *Biochem. J.*, 1952, **52**, 455–460.
- 25 S. Wang, B. Liu and Q. Su, *J. Anal. Appl. Pyrolysis*, 2004, **71**, 393–403.
- 26 V. A. Yaylayan, *Food Sci. Technol. Res.*, 2003, **9**, 1–6.
- 27 F. L. Chu and V. A. Yaylayan, *J. Agric. Food Chem.*, 2008, **56**, 10697–10704.
- 28 P. C. Wietstock, T. Kunz and F.-J. Methner, *J. Agric. Food Chem.*, 2016, **64**, 8035–8044.
- 29 T. Hofmann, P. Münch and P. Schieberle, *J. Agric. Food Chem.*, 2000, **48**, 434–440.
- 30 Y. Zhang and C.-T. Ho, *J. Agric. Food Chem.*, 1991, **39**, 760–763.
- 31 J. Chen and C.-T. Ho, *J. Agric. Food Chem.*, 1998, **46**, 1518–1522.
- 32 V. A. Yaylayan and A. Keyhani, *J. Agric. Food Chem.*, 2000, **48**, 2415–2419.
- 33 T. Davidek, S. Devaud, F. Robert and I. Blank, *J. Agric. Food Chem.*, 2006, **54**, 6667–6676.
- 34 S. Zöchling and M. Murkovic, *Food Chem.*, 2002, **79**, 125–134.

