



Cite this: *Phys. Chem. Chem. Phys.*,
2016, 18, 11932

New insights into aromatic pathways of carbachlorins and carbaporphyrins based on calculations of magnetically induced current densities†

Isaac Benkyi,^{*a} Heike Fliegl,^{*b} Rashid R. Valiev^{*cd} and Dage Sundholm^{*a}

Magnetically induced current densities have been calculated and analyzed for a number of synthesized carbachlorins and carbaporphyrins using density functional theory and the gauge including magnetically induced current (GIMIC) method. Aromatic properties have been determined by using accurate numerical integration of the current flow yielding reliable current strengths and pathways that are related to the degree of aromaticity and the aromatic character of the studied molecules. All investigated compounds are found to be aromatic. However, the obtained aromatic pathways differ from those previously deduced from spectroscopic data and magnetic shielding calculations. For all studied compounds, the ring current divides into an outer and an inner branch at each pyrrolic subring, showing that all π -electrons of the pyrrolic rings take part in the delocalization pathway. The calculations do not support the common notion that the main share of the current takes the inner route at the pyrrolic rings without an inner hydrogen and follows an 18π aromatic pathway. The aromatic pathways of the investigated carbaporphyrins and carbachlorins are very similar, since the current strength via the $C_\beta=C_{\beta'}$ bond of the cyclopentadienyl ring of the carbaporphyrins is almost as weak as the current density passing the corresponding saturated $C_\beta-C_{\beta'}$ bond of the carbachlorins.

Received 14th November 2015,
Accepted 25th February 2016

DOI: 10.1039/c5cp06987d

www.rsc.org/pccp

1 Introduction

Carbaporphyrinoids are porphyrin analogues in which one of the nitrogen atoms of the porphyrin macroring is replaced by an isoelectronic CH moiety.^{1–6} Carbachlorins are similar molecules with a saturated $C_\beta-C_{\beta'}$ bond of the cyclopentadienyl ring. This class of organic compounds is interesting for chemists due to their potential use in catalysis as they are able to bind metals in unusual oxidation states.^{1,6} In particular carbaporphyrinoid compounds such as carbachlorins with strong absorption bands at around 650 nm or even at longer wavelengths have the potential to be used as photosensitizers in photodynamic therapy.⁷

A clear picture of the full utilization of these compounds cannot be obtained without elucidating their aromatic properties.

In spite of their unusual structure and chemistry, carbaporphyrinoids can be expected to be aromatic like most planar organic (heterocyclic) compounds that exhibit conjugation pathways fulfilling the Hückel $4n + 2\pi$ rule for aromaticity. However, very little is known about their aromatic character and electron delocalization pathways,^{8,9} mainly because it is challenging to experimentally quantify the electron delocalization of complicated multiring molecules, and computational studies of the aromatic character of porphyrinoids are also demanding. By using the gauge-including magnetically induced current (GIMIC) method,^{10–12} it is though possible to determine current strength susceptibilities and current pathways by explicitly calculating the susceptibility of the magnetically induced current density passing selected chemical bonds. A careful analysis of the current density provides information about how electronic charge can be transported around molecular rings of fused multiring molecules.^{10,12–19} Comprehensive and detailed current-density studies have proven to be very helpful in elucidating the aromatic properties of porphyrinoids.^{20–28} Current density calculations carried out using the GIMIC program are an invaluable computational means for designing carbaporphyrinoids or other compounds with distinct aromaticity and electron delocalization features, since they can accurately predict aromatic pathways and ring-current strengths along different routes in the molecule.^{10–12,29} The GIMIC program, which is a stand-alone

^a University of Helsinki, Department of Chemistry, P. O. Box 55 (A.I. Virtanens plats 1), FIN-00014 University of Helsinki, Finland. E-mail: Dage.Sundholm@helsinki.fi

^b Centre for Theoretical and Computational Chemistry (CTCC), Department of Chemistry, University of Oslo, P. O. Box 1033 Blindern, 0315 Oslo, Norway. E-mail: Heike.Fliegl@kjemi.uio.no

^c Tomsk Polytechnic University, 43a Lenin Avenue, Building 2, Tomsk 634050, Russian Federation

^d Tomsk State University, Lenina Avenue 36, Tomsk, Russian Federation. E-mail: valievrashid@mail.ru

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c5cp06987d



code, has been employed in a number of aromaticity studies of porphyrinoids.^{10,22–28} The studies have shown that numerical integration of the current strength susceptibilities passing selected chemical bonds is a reliable tool for quantifying molecular aromaticity according to the magnetic criterion.^{30–32} By calculating the current strength susceptibilities of selected chemical bonds one obtains information about the electron-delocalization pathways. The approach has proven to be very useful for determining current pathways in multiring molecules, where many other approaches are prone to fail.^{22,33–47}

In this work, we have employed the GIMIC method at the density functional theory (DFT) level to investigate the aromatic character of a number of traditional carbaporphyrinoids and modified carbaporphyrinoids such as oxybenzporphyrin,^{48–50} benzocarbaporphyrin,^{51,52} azuliporphyrin^{51,53,54} and tropiporphyrin.^{55,56} In addition, we have also studied a number of carbachlorins that have been synthesized and characterized.⁹ The proton nuclear magnetic resonance (¹H NMR) and ultraviolet-visible (UV-Vis) spectra have been recorded to assess the aromatic properties of the synthesized carbachlorins and carbaporphyrins.⁹ The calculated values for nucleus independent chemical shifts (NICS)⁵⁷ have been previously used for assessing the aromatic character and delocalization pathways of carbaporphyrinoids,⁹ concluding that all the investigated carbaporphyrinoids are aromatic with a similar 18 π aromatic pathway as the classical aromatic pathway of free-base porphyrin.⁹ Nowadays it is well known that the NICS approach has difficulties in accurately determining the degree of aromaticity of single molecular rings.^{33–43} Furthermore, NICS has even larger problems to provide reliable current pathways in multiring molecules such as free-base porphyrin.^{22,58–62} Here we aim at providing novel insights regarding the aromatic pathways and electron delocalization pathways of the investigated compounds using the reliable current-density integration technique for analyzing the current flow.

The computational methods are described in Section 2. The molecular structures of the carbaporphyrinoids are discussed in Section 3, whereas the results of the current density calculations are presented in detail for all studied molecules in Section 4. The results of the study are summarized in Section 5 where the main conclusions are also drawn.

2 Computational methods

The optimization of the molecular structures as well as the calculations of the NMR shieldings were performed at the density functional theory (DFT) level using the Becke-three-parameter functional combined with the Lee–Yang–Parr exchange–correlation functional (B3LYP)^{63,64} as implemented in Turbomole 6.6.^{65,66} The Karlsruhe triple- ζ quality basis set (def2-TZVP) was used for all atoms.^{67,68} The NMR shielding calculations were performed using the mpshift module of Turbomole.^{69,70} Benchmark calculations have shown that B3LYP/def2-TZVP calculations of ¹H NMR and ¹³C NMR chemical shifts are close to the basis-set limit and agree qualitatively with experimental data for organic molecules.⁷¹

The magnetically induced current densities were calculated at the B3LYP/def2-TZVP level using the GIMIC program.^{10–12} GIMIC is an independent program that uses the atomic orbital density matrix as well as the corresponding first-order magnetically perturbed density matrices from the NMR shielding calculations and basis-set information as input data.^{10,11} GIMIC employs gauge-including atomic orbitals (GIAOs), which imply that the basis set convergence is faster than with magnetic-field independent basis functions.^{10,72,73} When GIAOs are employed, the obtained gauge-origin independent current densities were close to the complete basis-set limit already when standard basis sets are used.^{72,73} The current densities can be analyzed by determining the current pathways, which are obtained by the numerical integration of the current-strength susceptibilities (in nA T^{–1}) flowing along the chosen chemical bonds. The current pathways are visualized using the XMakeMol⁷⁴ and paint.net⁷⁵ programs. The pictures of the molecules have been drawn using the Marvin program.⁷⁶ The streamline representations of the current density have been obtained using PyNgl⁷⁷ and the schematic ring labeling displayed in Fig. 15 has been done using ChemBioDraw⁷⁸ and Gimp.⁷⁹

The effects of dispersion corrections⁸⁰ on the geometry and the current strength were investigated for oxybenzporphyrin. No significant changes in the molecular structure or the ring-current strength were observed. See ESI.†

3 Molecular structures and nomenclature

A number of recently synthesized carbaporphyrins and carbachlorins⁹ have been investigated computationally. The alkyl substituents have been omitted to save computational costs, since previous current density studies have shown that alkyl substituents do not significantly influence the current pathways and current strengths of aromatic porphyrinoids,²⁵ whereas for antiaromatic porphyrinoids substituents such as ethylformate or pentafluorophenyl may significantly reduce the strength of the ring current.^{24–26} The labeling of the investigated molecules follows the one in ref. 9, which inspired us to perform the present study. The common notation enables easier comparison of the present results with previously published ones.^{9,50}

Carbaporphyrin **20** (Fig. 1) is the most simple carbaporphyrin without any substituents. The carbaporphyrin cation **20H⁺** (Fig. 2) is the protonated form of **20** with one extra inner hydrogen and a positive charge. The doubly protonated carbaporphyrin dication **20H₂²⁺** (Fig. 3) has five inner hydrogens of which two saturate the inner carbon of the cyclopentadienyl ring. Carbaporphyrin **14** (Fig. 4) has an aldehyde group substituted in one of the C _{β} positions of the cyclopentadienyl ring of the carbaporphyrin.

Carbachlorin **19** is the most simple carbachlorin with a saturated C _{β} –C _{β'} bond of the cyclopentadienyl ring and without any substituents. Carbachlorin **19** (Fig. 5) lacks the inner hydrogen in the *trans* position to the cyclopentadienyl ring, whereas carbachlorin **19'** (Fig. 6) is the corresponding *cis*






Fig. 1 (a) A streamline plot of the current density 1 bohr above the carbaporphyrin **20** molecule that sustains a net ring-current strength of 26.8 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for carbaporphyrin **20**.




Fig. 4 (a) A streamline plot of the current density 1 bohr above the carbaporphyrin **14** molecule that sustains a net ring-current strength of 24.6 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for carbaporphyrin **14**.




Fig. 2 (a) A streamline plot of the current density 1 bohr above the carbaporphyrin **20H⁺** molecule that sustains a net ring-current strength of 25.7 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for carbaporphyrin **20H⁺**.




Fig. 5 (a) A streamline plot of the current density 1 bohr above the carbachlorin **19** molecule that sustains a net ring-current strength of 24.6 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for carbachlorin **19**.




Fig. 3 (a) A streamline plot of the current density 1 bohr above the carbaporphyrin **20H₂²⁺** molecule that sustains a net ring-current strength of 27.8 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for carbaporphyrin **20H₂²⁺**.




Fig. 6 (a) A streamline plot of the current density 1 bohr above the carbachlorin **19'** molecule that sustains a net ring-current strength of 26.1 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for carbachlorin **19'**.

tautomer. The carbachlorin cation **19H⁺** (Fig. 7) is the protonated form of **19** and **19'** with four inner hydrogens.

In compound **6b**, a propene moiety is fused to the cyclopentadienyl ring forming a fused nonaromatic cyclopentene ring.






Fig. 7 (a) A streamline plot of the current density 1 bohr above the carbachlorin **19H⁺** molecule that sustains a net ring-current strength of 26.5 nA T⁻¹ around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T⁻¹) passing selected bonds are given for carbachlorin **19H⁺**.




Fig. 10 (a) A streamline plot of the current density 1 bohr above the carbaporphyrin **9** molecule that sustains a net ring-current strength of 25.9 nA T⁻¹ around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T⁻¹) passing selected bonds are given for carbaporphyrin **9**.




Fig. 8 (a) A streamline plot of the current density 1 bohr above the carbachlorin **6b** (*cis*) molecule that sustains a net ring-current strength of 25.5 nA T⁻¹ around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T⁻¹) passing selected bonds are given for carbachlorin **6b** (*cis*).




Fig. 11 (a) A streamline plot of the current density 1 bohr above the oxybenzoporphyrin **1** molecule that sustains a net ring-current strength of 24.4 nA T⁻¹ around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T⁻¹) passing selected bonds are given for oxybenzoporphyrin **1**.




Fig. 9 (a) A streamline plot of the current density 1 bohr above the carbachlorin **6b** (*trans*) molecule that sustains a net ring-current strength of 25.7 nA T⁻¹ around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T⁻¹) passing selected bonds are given for carbachlorin **6b** (*trans*).

The two ends of the propene moiety can bind to the cyclopentadienyl ring in the *cis* **6b** (Fig. 8) or *trans* **6b** (Fig. 9) position.

Compounds **6b** are propencarbachlorins, because the C_β–C_{β'} bond of the cyclopentadienyl ring becomes saturated when binding the propene moiety to the cyclopentadienyl. Carbaporphyrins like compound **9** (Fig. 10) could in principle be obtained by oxidizing the corresponding carbachlorin. However, experimentally this is not straightforward since Li and Lash reported that they were not able to oxidize **6b**,⁹ which would have yielded the corresponding carbaporphyrin **9**.

Compounds **1–4** are modified carbaporphyrinoids. In oxybenzoporphyrin **1** (Fig. 11), the cyclopentadienyl ring is replaced by a cyclohexadienone moiety. In benzocarbaporphyrin **2** (Fig. 12), a benzoic ring is fused to the cyclopentadienyl ring. In azuliporphyrin **3** (Fig. 13), a cycloheptatriene ring is fused to the cyclopentadienyl ring, and in tropiporphyrin **4** (Fig. 14), the cyclopentadienyl ring is replaced by a cycloheptatrienyl ring.

4 Current-density calculations

All the investigated compounds are found to be aromatic according to the magnetic criterion. Streamline representations






Fig. 12 (a) A streamline plot of the current density 1 bohr above the benzocarporphyrin **2** molecule that sustains a net ring-current strength of 26.5 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for benzocarporphyrin **2**.




Fig. 13 (a) A streamline plot of the current density 1 bohr above the azuliporphyrin **3** molecule that sustains a net ring-current strength of 15.1 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for azuliporphyrin **3**.




Fig. 14 (a) A streamline plot of the current density 1 bohr above the tropiporphyrin **4** molecule that sustains a net ring-current strength of 21.9 nA T^{-1} around the porphyrinoid macroring. (b) The calculated net current strengths (in nA T^{-1}) passing selected bonds are given for tropiporphyrin **4**.

of the current densities obtained in a plane placed 1 bohr above the molecular plane and the calculated current density pathways

Table 1 The total ring-current strength (in nA T^{-1}) and the current strengths (in nA T^{-1}) of the outer and inner pathways of the studied carbaporphyrinoids. The numbering of the currents is shown in Fig. 15

Ring → molecule	Total	A		B		C		D	
		1	2	3	4	5	6	7	8
20	26.8	6.3	20.5	17.7	7.3	13.0	13.8	17.7	7.3
20H⁺	25.7	1.8	23.9	19.2	6.5	19.8	6.0	19.2	6.5
20H₂²⁺	27.8	27.0	0.9	19.6	8.1	21.1	6.4	19.7	8.2
14	24.8	4.8	20.3	17.8	6.8	12.8	12.3	17.8	6.8
19	24.6	1.5	23.8	18.5	5.9	13.5	10.8	18.5	5.9
19'	26.1	1.6	24.3	13.6	12.7	20.0	6.2	20.4	5.6
19H⁺	26.5	1.4	24.7	20.3	6.4	20.9	5.6	20.3	6.4
6b (cis)	25.5	1.4	24.5	19.0	6.5	13.7	11.5	19.0	6.5
6b (trans)	25.7	1.3	24.3	19.5	6.3	14.1	11.3	19.5	6.3
9	25.9	3.4	23.0	18.4	7.2	13.1	12.8	18.4	7.2
1	24.4	0.7	23.4	18.0	6.5	12.4	12.0	18.7	5.8
2	26.5	3.8	22.5	19.1	6.9	13.9	12.6	19.1	6.9
3	15.1	18.7	−3.2	8.9	5.9	12.0	3.2	8.9	5.9
4	22.1	−15.9	38.0	17.8	4.2	12.8	9.6	17.8	4.2

that are obtained through explicit integration of the current flow across several chemical bonds are given in Fig. 1–14. An overview of the calculated results is given in Table 1 and the respective labeling of the pyrrolic rings is shown in Fig. 15.

The current density calculations show that the aromatic pathway of the studied carbaporphyrinoids does not follow the classical 18π aromaticity route of porphyrinoids as suggested by Li and Lash.⁹ This is not surprising, since the classical 18π aromatic pathway of porphyrins and chlorins might not even be correct.^{10,22,59,81} The aromatic pathways of the studied carbaporphyrins and carbachlorins are indeed very reminiscent of the aromatic pathway of porphyrins and chlorins as previously obtained in current density calculations.²² Similar current pathways as obtained for the carbaporphyrinoids have also been obtained in a number of current density studies on other porphyrinoids.^{20–28}

For all studied compounds, the ring current divides into an outer and an inner pathway at each pyrrolic subring. Thus, all π -electrons of the pyrrolic rings take part in the delocalization pathways. The calculations show that the resistance of the inner NH group is generally larger than for the inner nitrogen without a hydrogen. Thus, a stronger current passes the inner N than the inner NH moiety of the pyrrolic rings. For the pyrrolic rings without an inner hydrogen, the current strengths *via* the outer and inner pathways are almost the same, whereas for the




Fig. 15 The numbering of the current pathways of the carbaporphyrinoids. Odd numbers indicate outer routes.





carbachlorin **19**. The calculated current pattern at the pyrrolic rings of compound **6b** (*trans*) is very similar to the one obtained for **6b** (*cis*). The net current strength is 25.7 nA T^{-1} . The current flow at the propene substituted all-carbon five-membered ring of 1.3 nA T^{-1} takes the outermost route. In addition the fused cyclopentene ring due to the propene substitution sustains a tiny local ring current of 0.5 nA T^{-1} .

4.5 Carbaporphyrin 9

Carbaporphyrin **9** can in principle be obtained by oxidizing carbachlorin **6**. However, that reaction step was unsuccessful.⁹ Current-density calculations show that carbaporphyrin **9** is expected to have similar ring-current pathways as obtained for the unsubstituted carbaporphyrin **20**. The net ring-current strength is 25.9 nA T^{-1} as compared to 26.8 nA T^{-1} for **20** and $25.5\text{--}25.7 \text{ nA T}^{-1}$ for **6b**. A small current of 1.0 nA T^{-1} passes the saturated CH_2 group of the cyclopentene ring fused to the cyclopentadienyl ring of the carbaporphyrin, whereas a current strength of 2.4 nA T^{-1} passes the common bond of the two five-membered rings.

4.6 Oxybenzporphyrin 1

Oxybenzporphyrin **1** is a carbaporphyrinoid where the cyclopentadienyl ring is replaced by a cyclohexadienone ring.⁵² **1** sustains a net ring-current strength of 24.4 nA T^{-1} around the macrocycle, which is 3 nA T^{-1} (10%) smaller than for porphyrin.²² The current pattern at the pyrrolic rings is the same as for porphyrin and the other carbaporphyrinoids studied in this work. The current strengths along the different routes are shown in Fig. 11. The carbonyl group of the cyclohexadienone ring prevents the ring-current to take the outer route. Thus, only 0.7 nA T^{-1} flows on the outside of the cyclohexadienone ring and 23.4 nA T^{-1} takes the inner pathway. The net ring-current strength calculated for the molecular structure of **1** optimized at the same level of theory using also the D3 correction differs by only 0.3 nA T^{-1} from the value of 24.4 nA T^{-1} as obtained without the D3 correction. Thus, the use of the D3 correction has almost no effect on the ring current strengths of the investigated class of molecules.

4.7 Benzocarbachlorin 2

Benzocarbachlorin **2** is a carbaporphyrin with a benzoic ring fused to the C_β bond of the cyclopentadienyl ring.⁴⁹ The current pattern at the pyrrolic rings is very similar to that of the other carbaporphyrinoids of this work. The ring-current strength around the macrocycle of 26.5 nA T^{-1} is almost as large as for unsubstituted carbaporphyrin **20**. At the cyclopentadienyl ring, the current prefers the inner route whose strength is 22.5 nA T^{-1} , whereas a current of 3.8 nA T^{-1} flows outwards passing on the outside of the benzoic ring. The benzoic ring sustains a local ring current of 5.6 nA T^{-1} . The current pathway and current strengths are shown in Fig. 12. The current density calculations show that the proposed 18π aromatic pathway is not completely correct.^{49,52}

4.8 Azuliporphyrin 3

Azuliporphyrin **3** consists of a cycloheptatriene ring fused to the cyclopentadienyl ring of carbaporphyrin **20**. Lash *et al.* proposed

that it has some aromatic character, because the structure can formally be described by two resonance structures.^{49,52} The zwitterionic form is thought to sustain a ring current around the carbaporphyrin macrocycle, whereas in the other form the ring current circles only around the azulene moiety.⁴⁹ However, the current density calculations yield a somewhat different picture of the aromatic character. The strength of the ring current circling around the carbaporphyrin macrocycle is 15.1 nA T^{-1} , which is about 55% of the ring-current strength of porphyrin. At the azulene moiety the main current of 18.7 nA T^{-1} takes the outer route around the cyclopentadienyl ring, whereas a weaker current of 4.6 nA T^{-1} passes on the outside of the cycloheptatrienyl ring. The cyclopentadienyl ring sustains a weak local ring current of 3.2 nA T^{-1} . At the pyrrolic rings, the main current flow passes the C_β carbons. At the pyrrolic rings without an inner hydrogen, 40% of the current takes the inner route, whereas at the pyrrolic ring with an inner hydrogen only 20% of the current passes the NH moiety. The current pathways and current strengths are shown in Fig. 13.

4.9 Tropiporphyrin 4

The calculated structure of tropiporphyrin **4**, which is obtained from carbaporphyrin by replacing the cyclopentadienyl ring with a cycloheptatrienyl ring,⁵⁵ is found to be almost planar with the largest out-of-plane torsional angle of 3° at the cycloheptatrienyl ring. The current strength of tropiporphyrin is 22.1 nA T^{-1} , which is 82% of the ring-current strength of carbaporphyrin **20**. The ring current around the macrocycle mainly passes along the outer routes at the pyrrolic rings. The pattern of the current flow around the carbaporphyrin ring is similar to the one for the other carbaporphyrins with about 20% of the ring current passing the NH moiety of the two pyrrolic rings with an inner hydrogen, whereas 43% of the ring current passes the nitrogen of the pyrrolic ring without an inner hydrogen. The cycloheptatrienyl ring is antiaromatic sustaining a strong local paratropic ring current of -15.9 nA T^{-1} , thus forming a strongly antiaromatic ring fused to the aromatic one. Fused rings with opposite tropicity have previously been found for thienopyrrole modified 20π -electron porphyrinoids and thienobridged porphyrins.^{23,25} The current pathways and current strengths in Fig. 14 show that the calculations yield a different aromatic character as compared to the one deduced from the measured NMR chemical shifts.⁵⁵

5 Summary and conclusions

Magnetically induced current densities of a number of synthesized as well as spectroscopically and theoretically characterized carbachlorins and carbaporphyrins⁹ have been studied computationally at the DFT level and analyzed using the gauge including magnetically induced current (GIMIC) method. The investigated compounds are found to be aromatic with calculated ring-current strength susceptibilities ranging from 15 nA T^{-1} to 27 nA T^{-1} . Thus, they can be considered aromatic according to the magnetic criterion. Tropiporphyrin **3** has the weakest aromaticity among the



calculations when aiming at information about the aromatic character of porphyrinoids, because the approaches do not provide very accurate information about molecular aromaticity for more complex molecules. Instead it is recommended to use current-density calculations in combination with numerical integration of current strengths, because this yields electron-delocalization pathways that show how electrons move around the molecular rings, when they are exposed to an external magnetic field.

Acknowledgements

References

- 1 A. Berlicka, P. Dutka, L. Szterenberg and L. Latos-Grażyński, *Angew. Chem., Int. Ed.*, 2014, **53**, 4885–4889.
- 2 S. Aronoff and M. Calvin, *J. Org. Chem.*, 1943, **8**, 205–223.
- 3 M. O. Senge, *Angew. Chem., Int. Ed.*, 2011, **50**, 4272–4277.
- 4 K. Berlin, *Angew. Chem., Int. Ed.*, 1996, **35**, 1820–1822.
- 5 T. D. Lash and S. T. Chaney, *Chem. – Eur. J.*, 1996, **2**, 944–948.
- 6 M. Pawlicki and L. Latos-Grażyński, *Chem. Rec.*, 2006, **6**, 64–78.
- 7 R. K. Pandey and G. Zheng, in *Porphyrin Handbook*, ed. K. M. S. K. M. Kadish and R. Guilard, Academic Press, San Diego, CA, 2000, vol. 6, pp. 157–230.
- 8 T. D. Lash, *Eur. J. Org. Chem.*, 2007, 5461–5481.
- 9 D. Li and T. D. Lash, *J. Org. Chem.*, 2014, **79**, 7112–7121.
- 10 J. Jusélius, D. Sundholm and J. Gauss, *J. Chem. Phys.*, 2004, **121**, 3952–3963.
- 11 S. Taubert, D. Sundholm and J. Jusélius, *J. Chem. Phys.*, 2011, **134**, 054123.
- 12 H. Fliegl, S. Taubert, O. Lehtonen and D. Sundholm, *Phys. Chem. Chem. Phys.*, 2011, **13**, 20500–20518.
- 13 M. P. Johansson, J. Jusélius and D. Sundholm, *Angew. Chem., Int. Ed.*, 2005, **44**, 1843–1846.
- 14 J. Jusélius and D. Sundholm, *Phys. Chem. Chem. Phys.*, 2008, **10**, 6630–6634.
- 15 S. Taubert, J. Jusélius, D. Sundholm, W. Kloppe and H. Fliegl, *J. Phys. Chem. A*, 2008, **112**, 13584–13592.
- 16 S. Taubert, D. Sundholm and F. Pichierri, *J. Org. Chem.*, 2010, **75**, 5867–5874.
- 17 H. Fliegl, O. Lehtonen, D. Sundholm and V. R. I. Kaila, *Phys. Chem. Chem. Phys.*, 2011, **13**, 434–437.

Phys. Chem. Chem. Phys., 2016, 18, 11932–11941 | 11939

- 18 M. Kaipio, M. Patzschke, H. Fliegl, F. Pichierri and D. Sundholm, *J. Phys. Chem. A*, 2012, **116**, 10257–10268.
- 19 D. Sundholm, *Phys. Chem. Chem. Phys.*, 2013, **15**, 9025–9028.
- 20 H. Fliegl, D. Sundholm, S. Taubert and F. Pichierri, *J. Phys. Chem. A*, 2010, **114**, 7153–7161.
- 21 H. Fliegl, D. Sundholm and F. Pichierri, *Phys. Chem. Chem. Phys.*, 2011, **13**, 20659–20665.
- 22 H. Fliegl and D. Sundholm, *J. Org. Chem.*, 2012, **77**, 3408–3414.
- 23 H. Fliegl, N. Özcan, R. Mera-Adasme, F. Pichierri, J. Jusélius and D. Sundholm, *Mol. Phys.*, 2013, **111**, 1364–1372.
- 24 R. R. Valiev, H. Fliegl and D. Sundholm, *J. Phys. Chem. A*, 2013, **117**, 9062–9068.
- 25 R. R. Valiev, H. Fliegl and D. Sundholm, *Phys. Chem. Chem. Phys.*, 2014, **16**, 11010–11016.
- 26 H. Fliegl, F. Pichierri and D. Sundholm, *J. Phys. Chem. A*, 2015, **119**, 2344–2350.
- 27 R. R. Valiev, H. Fliegl and D. Sundholm, *J. Phys. Chem. A*, 2015, **119**, 1201–1207.
- 28 R. R. Valiev, H. Fliegl and D. Sundholm, *Phys. Chem. Chem. Phys.*, 2015, **17**, 14215–14222.
- 29 H. Fliegl, D. Sundholm, S. Taubert, J. Jusélius and W. Klopper, *J. Phys. Chem. A*, 2009, **113**, 8668–8676.
- 30 A. Balaban, P. von Ragué Schleyer and H. S. Rzepa, *Chem. Rev.*, 2005, **105**, 3436–3447.
- 31 T. Heine, C. Corminboeuf and G. Seifert, *Chem. Rev.*, 2005, **105**, 3889–3910.
- 32 Z. Chen, C. S. Wannere, C. Corminboeuf, R. Puchta and P. von Ragué Schleyer, *Chem. Rev.*, 2005, **105**, 3842–3888.
- 33 P. Lazzeretti, *Phys. Chem. Chem. Phys.*, 2004, **6**, 217–223.
- 34 S. Pelloni, G. Monaco, P. Lazzeretti and R. Zanasi, *Phys. Chem. Chem. Phys.*, 2011, **13**, 20666–20672.
- 35 P. Lazzeretti, *Prog. Nucl. Magn. Reson. Spectrosc.*, 2000, **36**, 1–88.
- 36 I. Morao, B. Lecea and F. P. Cossío, *J. Org. Chem.*, 1997, **62**, 7033–7036.
- 37 J. Jusélius and D. Sundholm, *Phys. Chem. Chem. Phys.*, 1999, **1**, 3429–3435.
- 38 J. O. C. Jiménez-Halla, E. Matito, J. Robles and M. Solá, *J. Organomet. Chem.*, 2006, **691**, 4359–4366.
- 39 S. Pelloni and P. Lazzeretti, *J. Phys. Chem. A*, 2013, **117**, 9083–9092.
- 40 Z. Badri, S. Pathak, H. Fliegl, P. Rashidi-Ranjbar, R. Bast, R. Marek, C. Foroutan-Nejad and K. Ruud, *J. Chem. Theory Comput.*, 2013, **9**, 4789–4796.
- 41 G. Monaco and R. Zanasi, *J. Phys. Chem. A*, 2014, **118**, 1673–1683.
- 42 D. Du, H. Fliegl and D. Sundholm, *J. Chin. Chem. Soc.*, 2016, **63**, 93–100.
- 43 Y. C. Lin, D. Sundholm and J. Jusélius, *J. Chem. Theory Comput.*, 2006, **2**, 761–764.
- 44 S. Taubert, D. Sundholm and F. Pichierri, *J. Org. Chem.*, 2009, **74**, 6495–6502.
- 45 R. Islas, G. Martínez-Guajardo, J. O. C. Jiménez-Halla, M. Solá and G. Merino, *J. Chem. Theory Comput.*, 2010, **6**, 1131–1135.
- 46 A. C. Castro, E. Osorio, J. O. Jiménez-Halla, E. Matito, W. Tiznado and G. Merino, *J. Chem. Theory Comput.*, 2010, **6**, 2701–2750.
- 47 J. J. Torres, R. Islas, E. Osorio, J. G. Harrison, W. Tiznado and G. Merino, *J. Phys. Chem. A*, 2013, **117**, 5529–5533.
- 48 T. D. Lash, *Angew. Chem., Int. Ed.*, 1995, **34**, 2533–2535.
- 49 T. D. Lash, M. J. Hayes, J. D. Spence, M. A. Muckey, G. M. Ferrence and L. F. Szczepura, *J. Org. Chem.*, 2002, **67**, 4860–4874.
- 50 D. I. AbuSalim and T. D. Lash, *Org. Biomol. Chem.*, 2014, **12**, 8719–8736.
- 51 T. D. Lash and M. J. J. Hayes, *Angew. Chem., Int. Ed.*, 1997, **36**, 840–842.
- 52 T. D. Lash, S. T. Chaney and D. T. Richter, *J. Org. Chem.*, 1998, **63**, 9076–9088.
- 53 S. R. Graham, D. A. Colby and T. D. Lash, *Angew. Chem., Int. Ed.*, 2002, **41**, 1371–1374.
- 54 T. D. Lash, D. A. Colby, S. R. Graham and S. T. Chaney, *J. Org. Chem.*, 2004, **69**, 8851–8864.
- 55 K. M. Bergman, G. M. Ferrence and T. D. Lash, *J. Org. Chem.*, 2004, **69**, 7888–7897.
- 56 T. D. Lash and S. T. Chaney, *Tetrahedron Lett.*, 1996, **37**, 8825–8828.
- 57 P. von Ragué Schleyer, C. Maerker, A. Dransfeld, H. Jiao and N. J. R. van Eikema Hommes, *J. Am. Chem. Soc.*, 1996, **118**, 6317–6318.
- 58 M. K. Cyrański, T. M. Krygowski, M. Wisiorowski, N. J. R. van Eikema Hommes and P. von Ragué Schleyer, *Angew. Chem., Int. Ed.*, 1998, **37**, 177–180.
- 59 J. Jusélius and D. Sundholm, *Phys. Chem. Chem. Phys.*, 2000, **2**, 2145–2151.
- 60 E. Steiner and P. W. Fowler, *ChemPhysChem*, 2002, **3**, 114–116.
- 61 E. Steiner, A. Soncini and P. W. Fowler, *Org. Biomol. Chem.*, 2005, **4**, 4053–4059.
- 62 J. I. Wu, I. Fernández and P. von Ragué Schleyer, *J. Am. Chem. Soc.*, 2013, **135**, 315–321.
- 63 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- 64 C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1988, **37**, 785–789.
- 65 R. Ahlrichs, M. Bär, M. Häser, H. Horn and C. Kölmel, *Chem. Phys. Lett.*, 1989, **162**, 165–169.
- 66 F. Furche, R. Ahlrichs, C. Hättig, W. Klopper, M. Sierka and F. Weigend, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2014, **4**, 91–100.
- 67 A. Schäfer, H. Horn and R. Ahlrichs, *J. Chem. Phys.*, 1992, **97**, 2571–2577.
- 68 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 69 M. Häser, R. Ahlrichs, H. P. Baron, P. Weis and H. Horn, *Theor. Chim. Acta*, 1992, **83**, 455–470.
- 70 M. Kollwitz, M. Häser and J. Gauss, *J. Chem. Phys.*, 1998, **108**, 8295–8301.
- 71 S. Taubert, H. Konschin and D. Sundholm, *Phys. Chem. Chem. Phys.*, 2005, **7**, 2561–2569.
- 72 R. Ditchfield, *Mol. Phys.*, 1974, **27**, 789–807.



- 73 K. Wolinski, J. F. Hinton and P. Pulay, *J. Am. Chem. Soc.*, 1990, **112**, 8251–8260.
- 74 XMakemol, <http://www.nongnu.org/xmakemol/>.
- 75 paint.net 4.0.6, <http://www.getpaint.net/>.
- 76 Marvin 15.9.28, 2015, ChemAxon, <http://www.chemaxon.com>.
- 77 PyNgl: Scientific visualization Python program package, <http://www.pyngl.ucar.edu>.
- 78 ChemBioDraw Ultra 14.0 by PerkinElmer, <http://scistore.cambridgesoft.com>.
- 79 gimp: GNU Image Manipulation Program, <http://www.gimp.org>.
- 80 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 81 J. Jusélius and D. Sundholm, *J. Org. Chem.*, 2000, **65**, 5233–5237.

