

Cite this: *Chem. Sci.*, 2018, 9, 4152

# An artificial photosynthetic system for photoaccumulation of two electrons on a fused dipyridophenazine (dppz)–pyridoquinolinone ligand†

Jean-François Lefebvre,<sup>ab</sup> Julian Schindler,<sup>id cd</sup> Philipp Traber,<sup>id c</sup> Ying Zhang,<sup>cd</sup> Stephan Kupfer,<sup>id c</sup> Stefanie Gräfe,<sup>id c</sup> Isabelle Baussanne,<sup>b</sup> Martine Demeunynck,<sup>id b</sup> Jean-Marie Mouesca,<sup>f</sup> Serge Gambarelli,<sup>f</sup> Vincent Artero,<sup>id a</sup> Benjamin Dietzek<sup>id \*cde</sup> and Murielle Chavarot-Kerlidou<sup>id \*a</sup>

Increasing the efficiency of molecular artificial photosynthetic systems is mandatory for the construction of functional devices for solar fuel production. Decoupling the light-induced charge separation steps from the catalytic process is a promising strategy, which can be achieved thanks to the introduction of suitable electron relay units performing charge accumulation. We report here on a novel ruthenium tris-diimine complex able to temporarily store two electrons on a fused dipyridophenazine–pyridoquinolinone  $\pi$ -extended ligand upon visible-light irradiation in the presence of a sacrificial electron donor. Full characterization of this compound and of its singly and doubly reduced derivatives thanks to resonance Raman, EPR and (TD)DFT studies allowed us to localize the two electron-storage sites and to relate charge photoaccumulation with proton-coupled electron transfer processes.

Received 7th October 2017  
Accepted 31st March 2018

DOI: 10.1039/c7sc04348a

rsc.li/chemical-science

## Introduction

The development of solar-driven chemistry exploiting sunlight to produce fuels and commodity chemicals (ammonia, hydrogen peroxide...) from readily available feedstocks (water, carbon dioxide, atmospheric nitrogen and oxygen...) is a top research priority for the 21<sup>st</sup> century.<sup>1–4</sup> Natural photosynthesis is an inspiration in that field,<sup>5–7</sup> and the first functional molecular-based photocathodes for hydrogen production and CO<sub>2</sub> reduction recently appeared in the literature,<sup>8–12</sup> combining molecular photosensitizers as a surrogate for photosystems and catalytic centers mimicking enzymes involved in bioenergetic processes. These artificial photosynthetic systems however display low energy conversion efficiencies because the light-

induced charge separation process is a single-electron event that occurs on the fs–ns timescale, *i.e.* several orders of magnitude faster than multielectron catalysis. Consequently, charge recombination and/or degradation processes most of the time take place before catalysis can proceed. A solution to overcome this limitation therefore requires decoupling these two processes and here again, inspiration can be found in nature. For example, reversible quinone/hydroquinone couples are employed as electron relays in the photosynthetic machinery:<sup>13</sup> after two consecutive photon absorption – charge separation steps at photosystem II (PS II), two electrons are transported by plastoquinone Q<sub>B</sub> (among other electron transport cofactors), thanks to proton-coupled electron transfer (PCET) processes. Moreover, the photosynthetic process generates NAD(P)H as a key biochemical intermediate *via* the addition of two electrons and one proton to NAD(P)<sup>+</sup>. NAD(P)H further acts as a versatile reductant to drive reductive enzymatic reactions in the dark, such as CO<sub>2</sub> fixation (in the Calvin cycle).<sup>13</sup> Hence, the integration of charge accumulation sites in the molecular design of the photocatalytic system and the implementation of PCET processes is an appealing strategy towards the temporary storage of photogenerated electrons before delivery to the catalyst.

Since the initial report by Wasielewski and co-workers in 1992,<sup>14</sup> a few molecular systems have been reported that photoaccumulate reducing equivalents.<sup>15–17</sup> The electron storage sites range from organic structures such as naphthalene or

<sup>a</sup>Laboratoire de Chimie et Biologie des Métaux, Univ. Grenoble Alpes, CNRS, CEA, 38000 Grenoble, France. E-mail: murielle.chavarot-kerlidou@cea.fr

<sup>b</sup>Univ. Grenoble Alpes, CNRS, DPM, 38000 Grenoble, France

<sup>c</sup>Institute of Physical Chemistry, Abbe Center of Photonics, Friedrich Schiller University Jena, Helmholtzweg 4, 07743 Jena, Germany


<sup>d</sup>Department Functional Interfaces, Leibniz Institute of Photonic Technology Jena (IPHT), Albert-Einstein-Straße 9, 07745 Jena, Germany. E-mail: benjamin.dietzek@leibniz-ipht.de

<sup>e</sup>Center for Energy and Environmental Chemistry, Friedrich Schiller University Jena, Philosophenweg 8, 07743 Jena, Germany

<sup>f</sup>Univ. Grenoble Alpes, CEA, CNRS, INAC-SyMMES, 38000 Grenoble, France

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c7sc04348a





**Scheme 1** Synthesis of  $[\text{Ru}(\text{bpy})_2(\text{oxo-dppqp})](\text{PF}_6)_2$  (**1**)( $\text{PF}_6$ )<sub>2</sub> from  $[\text{Ru}(\text{bpy})_2(\text{dppqp})](\text{PF}_6)_2$  and chemical structures of reference complexes  $[\text{Ru}(\text{bpy})_3](\text{PF}_6)_2$  and  $[\text{Ru}(\text{bpy})_2(\text{dppz})](\text{PF}_6)_2$ .

perylene bisimides,<sup>14,18</sup> anthraquinone,<sup>19</sup> and viologen derivatives,<sup>20,21</sup> to iridium(III)<sup>22</sup> or rhodium(III)<sup>23</sup> complexes and polyoxometalates.<sup>24,25</sup> Ruthenium complexes containing NAD(P)<sup>+</sup>/NAD(P)H mimics were also elegantly developed by the group of Tanaka.<sup>26–28</sup> Of particular interest is the work of MacDonnell and co-workers on dinuclear ruthenium complexes bearing extended poly-*N*-heterocyclic bridging ligands<sup>29–32</sup> with, in particular, a quinone-containing bridging ligand undergoing a light-driven proton-assisted four-electron reduction process.<sup>29</sup> These authors also clearly showed that a bent bridging ligand can accumulate two electrons at a potential suitable to drive proton reduction.<sup>32</sup>

Implementing such a strategy in fuel-producing photocathodes however puts strong topological constraints on the molecular design. Indeed, unidirectional electron transfer from the electrode substrate to the accumulation site and through the photosensitizer is required. In addition, the photoaccumulation site should be remote from the electrode surface in order to limit the recombination processes.<sup>33</sup> This precludes the use of previously reported systems in which two photosensitizing units straddle the accumulation site.<sup>14,18,19,24,29,30,32</sup> Conversely, systems based on a single light-harvesting center such as iminoimidazole-containing ruthenium photosensitizers, whose properties have been studied by two of us,<sup>34,35</sup> and disubstituted dipyrido-[3,2-*a*:2',3'-*c*]phenazine (dppz)-based ruthenium complexes<sup>36</sup> are instrumental to the aim of designing molecular photoelectrodes with enhanced efficiency. We therefore designed a novel ruthenium tris-diimine complex, **1**( $\text{PF}_6$ )<sub>2</sub> (Scheme 1), that features an original bent-shaped polyazaaromatic ligand, namely a pyridoquinolinone moiety fused to a dppz-chelating unit. We report herein its full steady state spectroscopic and electrochemical characterization and show how the pyridoquinolinone subunit drastically modulates the electronic properties of the complex toward charge accumulation. Photolysis experiments unambiguously established the ability of **1**( $\text{PF}_6$ )<sub>2</sub> to accumulate two electrons upon visible light irradiation in the presence of a sacrificial electron donor,

and to further stabilize the photogenerated two-electron reduced state thanks to a PCET process. The electron storage sites on the  $\pi$ -extended ligand were identified at the (time-dependent) density functional level of theory ((TD)DFT) and by EPR and resonance Raman studies on independently prepared singly and doubly reduced derivatives.


## Results and discussion

We previously reported an efficient “chemistry on the complex” strategy to synthesize variously substituted ruthenium tris-diimine complexes containing  $\pi$ -extended ligands, such as  $[\text{Ru}(\text{bpy})_2(\text{dppqp})](\text{PF}_6)_2$  (dppqp = dipyrido[3,2-*a*:2',3'-*c*]pyrido[2'',3''-4,5]quinolino[2,3-*h*]phenazine, Scheme 1).<sup>37</sup> Following previously published methods,<sup>38,39</sup> deethylation by cerium ammonium nitrate (CAN,  $\text{Ce}(\text{IV})(\text{NH}_4)_2(\text{NO}_3)_6$ ) treatment, followed by spontaneous air oxidation of the latter afforded  $[\text{Ru}(\text{bpy})_2(\text{oxo-dppqp})](\text{PF}_6)_2$  (**1**( $\text{PF}_6$ )<sub>2</sub>), featuring a fused dipyridophenazine (dppz)-pyridoquinolinone ligand (oxo-dppqp = dipyrido[3,2-*a*:2',3'-*c*]pyrido[2'',3''-4,5]quinolino[2,3-*h*]phenazin-15-one) in 80% yield after purification.

Aggregation through  $\pi$ -stacking is commonly observed for planar polyazaaromatic ruthenium complexes.<sup>37,40</sup> Such a  $\pi$ -stacking behavior is supported by quantum chemical simulations for **1**( $\text{PF}_6$ )<sub>2</sub> (Fig. S3 and Table S1†) and was evaluated by a concentration-dependent <sup>1</sup>H NMR study in CD<sub>3</sub>CN (Fig. S1 & S2†). The threshold concentration below which the monomer is predominant in solution was determined to be  $5 \times 10^{-4}$  M (see details in the ESI†), in line with a previous study on related Ru complexes;<sup>37</sup> accordingly, the following spectroscopic and electrochemical measurements were performed at lower concentrations of **1**( $\text{PF}_6$ )<sub>2</sub>.

The UV/vis spectrum of **1**( $\text{PF}_6$ )<sub>2</sub> displays typical features of ruthenium tris-diimine complexes (Fig. 2 and S4†), with an intense ligand-centered (LC) transition at 290 nm, characteristic of the bipyridine ligands, and metal-to-ligand charge-transfer (MLCT) bands observed between 400 and 500 nm. The high





**Fig. 1** UV/vis (A) and RR spectra (B) of  $[1]^{2+}$  (black) in comparison with the spectra collected during controlled potential electrolysis: first (blue; electrolysis: 1 min at  $-0.5$  V vs. Ag/AgCl) and second reduction (red; electrolysis: 1 min at  $-0.85$  V vs. Ag/AgCl) in 0.1 M tetrabutylammonium tetrafluoroborate/acetonitrile. The RR spectrum was obtained upon excitation at 458 nm (solvent bands subtracted). The RR bands characteristic of vibrations of the bpy ligand (symbol B) as well as of the dppz moiety (symbol D) and the iminobenzoquinone moiety (symbol #) of the oxo-dppqp ligand are indicated. (C) MO diagram for  $[1]^{2+}$  including occupied (black) and virtual (red) frontier orbitals; orbital energies are given with respect to the energy of the HOMO.

molar absorptivity in the visible range compared to the reference complex  $[\text{Ru}(\text{bpy})_2(\text{dppz})](\text{PF}_6)_2$  (Fig. S4†) suggests some contribution from the  $\pi$ -extended oxo-dppqp ligand. This is in agreement with TDDFT calculations predicting a strongly absorbing ligand-centered  $^1\pi\pi^*$  state on the  $\pi$ -extended ligand (see  $S_8$  in Fig. S5†).<sup>41</sup>

The redox properties of  $[1]^{2+}$  were determined by cyclic voltammetry (CV) and differential pulse voltammetry (DPV) (Fig. S6†) and are reported in Table 1. Oxidation of the Ru center is observed at  $+0.86$  V vs.  $\text{Fc}^+/\text{Fc}$ , which is slightly more anodic than the  $\text{Ru}^{\text{III}}/\text{Ru}^{\text{II}}$  processes in  $[\text{Ru}(\text{bpy})_3](\text{PF}_6)_2$  and  $[\text{Ru}(\text{bpy})_2(\text{dppz})](\text{PF}_6)_2$  and most likely reflects an increase of the electron-accepting properties of the  $\pi$ -extended ligand. In addition,  $[1]^{2+}$  displays five successive reduction processes; the first one at  $-0.87$  V vs.  $\text{Fc}^+/\text{Fc}$  is not observed for any of the reference complexes (Fig. S7 & S8†) and is tentatively assigned to a reduction on the pyridoquinolinone moiety. The second cathodic process occurs at a potential very close to the one obtained for the parent  $[\text{Ru}(\text{bpy})_2(\text{dppz})](\text{PF}_6)_2$  complex (Fig. S7†) which indicates a reduction located on the dppz moiety of the ligand. The following reductions are attributed to the three diimine ligands, by comparison to the reference complexes (Fig. S7 & S8†).

To gain more insights into the structures of the reduced compounds, the (electro)chemically reduced forms of  $[1]^{2+}$  (first and second reduction, generating  $[1]^+$  and  $[1]^0$ , respectively) were investigated by UV/vis absorption spectroscopy (Fig. 2, S9 and S10†) and UV/vis and resonance Raman (RR) spectroelectrochemistry (SEC; Fig. 1 and Fig. S11†), in combination with TDDFT calculations (see the ESI† for details).

A 15  $\mu\text{M}$  solution of  $[1]^{2+}$  in  $\text{CH}_3\text{CN}$  was chemically reduced with cobaltocene<sup>32</sup> under rigorously anaerobic conditions. The formation of the singly reduced derivative  $[1]^+$  is accompanied by significant changes in the UV/vis spectrum, with the growth of a very characteristic absorption feature at 653 nm (Fig. 2). In addition, three isosbestic points are observed at 326, 394 and 445 nm, associated with the disappearance of the absorption band at 315 nm and the growth of a new one at 348 nm (Fig. S9†). In contrast, the absorption of the doubly reduced species  $[1]^0$ , generated by the addition of a second reducing equivalent to the solution, extends above 700 nm and is almost featureless (Fig. 2 and S10†). Such broad absorption features above 500 nm typically originate from ligand-centered states of  $\pi\pi^*$  nature of the reduced ligand.<sup>42</sup> Similar spectral changes were also observed for electrochemically generated  $[1]^+$  and  $[1]^0$  (Fig. 1A).<sup>43</sup> In the case of  $[\text{Ru}(\text{bpy})_2(\text{dppz})]^{2+}$ ,<sup>42,44</sup> the first reduction is centered on the phenazine moiety and is characterized by a broad absorption band from 500 to above 800 nm with notable absorption contributions beyond 700 nm (Fig. S11†), thus resembling the spectral features of doubly reduced  $[1]^0$ . These data support the above assignment from cyclic voltammetry with the first reduction of  $[1]^{2+}$  centered at the pyridoquinolinone fragment and the second reduction localized on the phenazine fragment.

The RR spectrum of  $[1]^{2+}$  (collected upon excitation at 458 nm) consists of bands characteristic of vibrations of the bpy ligand (marked with B, Fig. 1B) as well as of the dppz moiety






Fig. 2 From top to bottom: summary of the chemical reduction process, EPR spectra, calculated structures and frontier molecular orbitals carrying excess charge, and UV/vis absorption spectra of  $[1]^{2+}$ ,  $[1]^+$ ,  $[1]^0$  and  $[1H_2]^{2+}$  (see the ESI† for experimental details).

(marked with D) and the iminobenzoquinone moiety of the oxo-dppqp ligand (marked with #). These bands were assigned by comparison with the RR spectra of  $[Ru(bpy)_2(dppz)]^{2+}$ ,  $[Ru(bpy)_3]^{2+}$  (Fig. S12†) and  $[Ru(dppz)_3]^{2+}$ .<sup>45</sup> In the first reduction process, a decrease of the bands attributed to the iminobenzoquinone moiety is observed, whereas MLCT transitions to the bpy ligands and the dppz moiety are sparsely affected. This again indicates that the additional charge is likely to be localized at the distal pyridoquinolinone moiety. In the second reduction process, the bands characteristic of the dppz moiety (symbol D) vanish, while the bands arising from bpy vibrations (symbol B) are still observed. Such an effect is also observed during the first reduction of the reference  $[Ru(bpy)_2(dppz)]^{2+}$  (Fig. S13†). Thus, for  $[1]^0$ , MLCT transitions

to the dppz moiety do not take place anymore, in agreement with a second reduction localized on this part of the ligand.

To support these UV/vis and RR studies, quantum chemical calculations were performed at the DFT and TDDFT level of theory on  $[1]^{2+}$ , as well as on the singly and doubly reduced species  $[1]^+$  and  $[1]^0$ .  $[1]^+$  is clearly of doublet multiplicity; TDDFT allows us to assign the absorption feature at 653 nm to the bright intraligand states (localized on the oxo-dppqp ligand)  $D_7$  and  $D_8$ , calculated at 605 and 576 nm respectively (Fig. S5†), of the singly reduced doublet  $[1]^+$ . The doubly reduced complex,  $[1]^0$ , may be either of singlet or triplet character (Fig. S14†). The DFT calculations reveal a near degeneracy of  $^1[1]^0$  and  $^3[1]^0$ , while the triplet is slightly favored by 0.1 eV. Such an energy difference value is below the margin of error of DFT and thus no unambiguous assignment of the spin state of  $[1]^0$  is feasible. However, the simulated UV/vis absorption spectrum of the doubly reduced triplet species can be clearly associated with the experimental absorption spectra of  $[1]^0$  (Fig. S15†). In particular, the broad absorption band between 550 and 800 nm is unambiguously assigned to three bright oxo-dppqp-centered intraligand states ( $T_{12}$ ,  $T_{15}$  and  $T_{17}$ ) calculated at 759, 588 and 574 nm. On the other hand, no agreement is evident for the simulated  $^1[1]^0$  (Fig. S15†). Thus, the TDDFT calculations

Table 1 Electrochemical data for  $[1]^{2+}$  and reference complexes  $[Ru(bpy)_3](PF_6)_2$  and  $[Ru(bpy)_2(dppz)](PF_6)_2$  ( $E_{1/2}$  in V vs.  $Fc^+/Fc$ )

|                        | $Ru^{III/II}$ | Red <sub>1</sub> | Red <sub>2</sub> | Red <sub>3</sub> | Red <sub>4</sub> | Red <sub>5</sub> |
|------------------------|---------------|------------------|------------------|------------------|------------------|------------------|
| $Ru(bpy)_3^{2+}$       | +0.82         | —                | —                | −1.73            | −1.91            | −2.18            |
| $Ru(bpy)_2(dppz)^{2+}$ | +0.84         | —                | −1.36            | −1.79            | −2.01            | −2.49            |
| $[1]^{2+}$             | +0.86         | −0.87            | −1.32            | −1.81            | −2.02            | −2.43            |

suggest that  $[1]^0$  exists in the triplet state. This is further supported below by electronic paramagnetic resonance experiments (EPR, *vide infra*). In addition, from the performed quantum chemical simulations, the first one-electron reduction is found to be localized on the iminobenzoquinone moiety ( $\alpha$ -HOMO of  $[1]^+$ ), while the excess charges in doubly reduced  $^3[1]^0$  (triplet state)—compared to non-reduced  $[1]^{2+}$ —are localized on the iminobenzoquinone moiety ( $\alpha$ -HOMO-1) and on the dppz moiety ( $\alpha$ -HOMO) as illustrated in Fig. 2. Further details with respect to the quantum chemical simulations are presented in the ESI.†

EPR is the spectroscopic technique of choice to probe the chemical environment of radical species. The chemical reduction procedure was thus adapted to record the X-band EPR spectra of  $[1]^+$  and  $[1]^0$  (200  $\mu$ M in  $\text{CH}_3\text{CN}$ /propionitrile; see UV/vis monitoring in Fig. S16†). The X-band EPR spectrum of the former, recorded in deoxygenated solution at room temperature, is characteristic of an organic radical species ( $g = 2.003$ ), with a very slow relaxation rate (Fig. 2). It displays numerous, easily saturated, very narrow overlapping lines (0.3 G peak to peak) and could be simulated as a monoradical coupled to 3 nitrogen atoms (spin 1 nucleus; hyperfine coupling constants of 7.3, 4.8, and 3.9 MHz) and 6 hydrogen atoms (hyperfine coupling constants of 6.3, 5.4, 4.2, 3.7, 2.0, and 1.8 MHz). This finding is in agreement with the spin density of  $[1]^+$ , which is

predominantly localized on the pyridoquinolinone moiety (see Fig. S17†).  $[1]^0$  is also EPR active, which unambiguously confirms that this species exists in a triplet spin state. Its spectrum displays five equidistant broad lines (2.2 G peak to peak) with relative intensity 1–2–3–2–1 (Fig. 2), attributed to a hyperfine coupling with two equivalent N atoms with a coupling constant of 13.8 MHz (4.9 G for  $g = 2$ ). It appeared to be strongly similar to the spectrum of singly reduced  $[\text{Ru}(\text{bpy})_2(\text{dppz})](\text{PF}_6)_2$  (Fig. S18†), which displayed a similar broad quintet with a hyperfine coupling constant of 14.2 MHz (5.05 G for  $g = 2$ ). The only significant difference is a smaller hydrogen hyperfine coupling observed for the latter.<sup>42</sup> Due to the similarity between the EPR spectra of the doubly reduced species  $[1]^0$  and the singly reduced species of  $[\text{Ru}(\text{bpy})_2(\text{dppz})](\text{PF}_6)_2$ , we propose the formation of the dimer  $\pi\text{-}\{[1]_2\}^0$ , through  $\pi$ -stacking of two  $[1]^0$  complexes, occurring under the EPR analysis conditions. In such a structure, pairing of the electrons located on each pyridoquinolinone moiety would take place (thus precluding their EPR detection); on the other side, the electrons located on each pyrazine ring are too far from each other to interact, leading to the observed signature, identical to the one from a single dppz radical species. Indeed, an increase of the  $\pi$ -stacking propensity of these complexes is likely to occur upon ligand reduction,<sup>46</sup> in agreement with the higher binding energy values calculated for the




Fig. 3 Top: UV/vis monitoring of a continuous photolysis experiment ( $\lambda_{\text{exc}} = 450$  nm; 15  $\mu$ M in  $\text{CH}_3\text{CN}$  in the presence of 0.15 M TEA) (inset: zoom-in of the 600–700 nm area where the formation of  $[1]^+$  is observed). Bottom: comparison of the photolysis experiment ( $t = 5$  min; dotted line) with  $[1\text{H}_2]^{2+}$  (green line). Right: summary scheme of the reversible light-driven charge accumulation process.





## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This research was supported by the Fonds der Chemischen Industrie (B. D. and J. S.), the French National Research Agency (Labex Program, ARCANÉ, ANR-11-LABX-0003-01) and the CEA DRF Impulsion 2017 Program (Project “Light-Acc-Cat”). All calculations have been performed at the Universitätsrechenzentrum of the Friedrich Schiller University Jena. We acknowledge the COST Action CM1202, PERSPECT-H<sub>2</sub>O: Supramolecular Photocatalytic Water Splitting.

## Notes and references

- 1 <http://www.euchems.eu/solar-driven-chemistry/>.
- 2 *Solar Fuels and Artificial Photosynthesis: Science and innovation to change our future energy options*, RSC report, January 2012.
- 3 T. A. Faunce, W. Lubitz, A. W. Rutherford, D. MacFarlane, G. F. Moore, P. Yang, D. G. Nocera, T. A. Moore, D. H. Gregory, S. Fukuzumi, K. B. Yoon, F. A. Armstrong, M. R. Wasielewski and S. Styring, *Energy Environ. Sci.*, 2013, **6**, 695.
- 4 A. Thapper, S. Styring, G. Saracco, A. W. Rutherford, B. Robert, A. Magnuson, W. Lubitz, A. Llobet, P. Kurz, A. Holzwarth, S. Fiechter, H. de Groot, S. Campagna, A. Braun, H. Bercegol and V. Artero, *Green*, 2013, **3**, 43.
- 5 S. Berardi, S. Drouet, L. Francas, C. Gimbert-Surinach, M. Guttentag, C. Richmond, T. Stoll and A. Llobet, *Chem. Soc. Rev.*, 2014, **43**, 7501.
- 6 W. Song, Z. Chen, M. K. Brennaman, J. J. Concepcion, A. O. T. Patrocinio, N. Y. M. Iha and T. J. Meyer, *Pure Appl. Chem.*, 2011, **83**, 749.
- 7 E. S. Andreiadis, M. Chavarot-Kerlidou, M. Fontecave and V. Artero, *Photochem. Photobiol.*, 2011, **87**, 946.
- 8 Z. Yu, F. Li and L. Sun, *Energy Environ. Sci.*, 2015, **8**, 760.
- 9 M. K. Brennaman, R. J. Dillon, L. Alibabaei, M. K. Gish, C. J. Dares, D. L. Ashford, R. L. House, G. J. Meyer, J. M. Papanikolas and T. J. Meyer, *J. Am. Chem. Soc.*, 2016, **138**, 13085.
- 10 N. Queyriaux, N. Kaeffer, A. Morozan, M. Chavarot-Kerlidou and V. Artero, *J. Photochem. Photobiol., C*, 2015, **25**, 90.
- 11 N. Kaeffer, J. Massin, C. Lebrun, O. Renault, M. Chavarot-Kerlidou and V. Artero, *J. Am. Chem. Soc.*, 2016, **138**, 12308.
- 12 G. Sahara, H. Kumagai, K. Maeda, N. Kaeffer, V. Artero, M. Higashi, R. Abe and O. Ishitani, *J. Am. Chem. Soc.*, 2016, **138**, 14152.
- 13 J. Barber, *Chem. Soc. Rev.*, 2009, **38**, 185.
- 14 M. P. O'Neil, M. P. Niemczyk, W. A. Svec, D. Gosztola, G. L. Gaines and M. R. Wasielewski, *Science*, 1992, **257**, 63.
- 15 Y. Pellegrin and F. Odobel, *Coord. Chem. Rev.*, 2011, **255**, 2578.
- 16 L. Hammarström, *Acc. Chem. Res.*, 2015, **48**, 840.
- 17 A. G. Bonn and O. S. Wenger, *Chimia*, 2015, **69**, 17.
- 18 G. Izzet and A. Proust, *Energy Environ. Sci.*, 2013, **6**, 1504.
- 19 T.-a. Koizumi and K. Tanaka, *Angew. Chem., Int. Ed.*, 2005, **44**, 5891.
- 20 T. Fukushima, T. Wada, H. Ohtsu and K. Tanaka, *Dalton Trans.*, 2010, **39**, 11526.
- 21 K. Kobayashi, H. Ohtsu, K. Nozaki, S. Kitagawa and K. Tanaka, *Inorg. Chem.*, 2016, **55**, 2076.
- 22 R. Konduri, H. Ye, F. M. MacDonnell, S. Serroni, S. Campagna and K. Rajeshwar, *Angew. Chem., Int. Ed.*, 2002, **41**, 3185.
- 23 R. Konduri, N. R. de Tacconi, K. Rajeshwar and F. M. MacDonnell, *J. Am. Chem. Soc.*, 2004, **126**, 11621.
- 24 N. R. de Tacconi, R. O. Lezna, R. Konduri, F. Ongeri, K. Rajeshwar and F. M. MacDonnell, *Chem.-Eur. J.*, 2005, **11**, 4327.
- 25 S. Singh, N. R. de Tacconi, N. R. G. Diaz, R. O. Lezna, J. Munoz Zuniga, K. Abayan and F. M. MacDonnell, *Inorg. Chem.*, 2011, **50**, 9318.
- 26 N. Queyriaux, R. A. Wahyuono, J. Fize, C. Gablin, M. Wächtler, E. Martinez, D. Léonard, B. Dietzek, V. Artero and M. Chavarot-Kerlidou, *J. Phys. Chem. C*, 2017, **121**, 5891.
- 27 L. Zedler, S. Kupfer, I. R. de Moraes, M. Wächtler, R. Beckert, M. Schmitt, J. Popp, S. Rau and B. Dietzek, *Chem.-Eur. J.*, 2014, **20**, 3793.
- 28 S. Kupfer, *Phys. Chem. Chem. Phys.*, 2016, **18**, 13357.
- 29 J. M. Aslan, D. J. Boston and F. M. MacDonnell, *Chem.-Eur. J.*, 2015, **21**, 17314.
- 30 J.-F. Lefebvre, D. Saadallah, P. Traber, S. Kupfer, S. Grafe, B. Dietzek, I. Baussanne, J. De Winter, P. Gerbaux, C. Moucheron, M. Chavarot-Kerlidou and M. Demeunynck, *Dalton Trans.*, 2016, **45**, 16298.
- 31 L. Bouffier, M. Demeunynck, A. Milet and P. Dumy, *J. Org. Chem.*, 2004, **69**, 8144.
- 32 L. Bouffier, B. Baldeyrou, M.-P. Hildebrand, A. Lansiaux, M.-H. David-Cordonnier, D. Carrez, A. Croisy, O. Renaudet, P. Dumy and M. Demeunynck, *Bioorg. Med. Chem.*, 2006, **14**, 7520.
- 33 K. Ritter, C. Pehlken, D. Sorsche and S. Rau, *Dalton Trans.*, 2015, **44**, 8889.

- 41 J. Schindler, Y. Zhang, P. Traber, J.-F. Lefebvre, S. Kupfer, M. Demeunynck, S. Gräfe, M. Chavarot-Kerlidou and B. Dietzek, *J. Phys. Chem. C*, 2018, **122**, 83.
- 42 J. Fees, W. Kaim, M. Moscherosch, W. Matheis, J. Klima, M. Krejcik and S. Zalis, *Inorg. Chem.*, 1993, **32**, 166.
- 43 A much more pronounced absorbance was measured above 500 nm for electrochemically generated  $[1]^0$  (Fig. 1). The higher complex concentration employed for this measurement, and additionally the presence of an electrolyte, most probably modify the pi-stacking equilibrium for  $[1]^0$  and could account for this observation.
- 44 L. Zedler, J. Guthmuller, I. Rabelo de Moraes, S. Kupfer, S. Kriek, M. Schmitt, J. Popp, S. Rau and B. Dietzek, *Chem. Commun.*, 2014, **50**, 5227.
- 45 C. G. Coates, L. Jacquet, J. J. McGarvey, S. E. J. Bell, A. H. R. Al-Obaidi and J. M. Kelly, *J. Am. Chem. Soc.*, 1997, **119**, 7130.
- 46 N. R. de Tacconi, R. Chitakunye, F. M. MacDonnell and R. O. Lezna, *J. Phys. Chem. A*, 2008, **112**, 497.
- 47 Y. Pellegrin and F. Odobel, *C. R. Chim.*, 2017, **20**, 283.
- 48 S. Fukuzumi, Y.-M. Lee and W. Nam, *Chem.-Eur. J.*, DOI: 10.1002/chem.201704512.

