

RESEARCH ARTICLE

[View Article Online](#)
[View Journal](#) | [View Issue](#)Cite this: *RSC Med. Chem.*, 2025, 16, 3049**Niclosamide: CRL4^{AMBRA1} mediated degradation of cyclin D1 following mitochondrial membrane depolarization†**Seemon Coomar, ^a Jessica A. Gasser,^b Mikotaj Stabicki,^{bcd} Katherine A. Donovan, ^{ef} Eric S. Fischer, ^{ef} Benjamin L. Ebert,^g Dennis Gillingham ^h and Nicolas H. Thomä^{*ai}

Targeted protein degradation has emerged as a promising approach in drug discovery, utilizing small molecules like molecular glue degraders to harness the ubiquitin-proteasome pathway for selective degradation of disease-driving proteins. Based on results from proteomics screens we investigated the potential of niclosamide, an FDA-approved anthelmintic drug with a 50 year history in treating tapeworm infections, as a molecular glue degrader targeting the proto-oncogene cyclin D1. Proteomics screens in HCT116 colon carcinoma and KELLY neuroblastoma cells, found that niclosamide induces rapid cyclin D1 degradation through a mechanism involving the ubiquitin-proteasome pathway. A genetic CRISPR screen identified the E3 ligase CRL4^{AMBRA1} as a key player in this process. Structure–activity relationship studies highlighted critical features of niclosamide necessary for cyclin D1 degradation, demonstrating a correlation between mitochondrial membrane potential (MMP) disruption and cyclin D1 downregulation. Notably, various mitochondrial uncouplers and other compounds with similar drug sensitivity profiles share this correlation suggesting that MMP disruption can trigger cyclin D1 degradation, and that the cellular signal driving the degradation differs from previously described mechanism involving CRL4^{AMBRA1}. Our findings underscore the complexities of proteostatic mechanisms and the multitude of mechanisms that contribute to degrader drug action.

Received 20th January 2025,
Accepted 19th April 2025

DOI: 10.1039/d5md00054h

rsc.li/medchem**Introduction**

Small molecules that induce proximity to yield therapeutically beneficial ternary complexes of disease-relevant proteins are an emerging topic in drug discovery.^{1,2} The most frequent use of the concept has been in the field of targeted protein degradation (TPD), which has generated considerable clinical


and commercial interest.^{3,4} The field of TPD gained considerable momentum after the mechanistic understanding of immunomodulatory drugs (IMiDs). They degrade the transcription factors (TFs) Ikaros and Aiolos and drive anti-myeloma activity in ubiquitin-dependent manner, which catalysed research into the development of molecules that can degrade disease-driving proteins.⁵ In the case of IMiDs such as lenalidomide (Revlimid), small molecules induce the complex formation between the E3 ligase adapter protein CRBN and the TFs leading to the ubiquitination and consequential degradation of the latter.^{6,7} Structural studies suggested that the molecules could be used as anchors to design bifunctional molecules, known as proteolysis targeting chimeras (PROTACs) that could engage CRBN and a protein of interest that would be degraded by the ubiquitin proteasome system (UPS).⁸ Whereas a rational design is possible for PROTACs, the discovery of small molecules which induce a novel interaction between an E3 ligase and a protein, so-called molecular glue degraders (MGDs) is more challenging as the binding partners are unknown *a priori*.^{9–11}

A typical workflow in the discovery process of a MGD begins with treating cells for a short duration with a compound, followed by global proteomics experiments. If a

^a Friedrich Miescher Institute for Biomedical Research, Fabrikstrasse 24, Basel 4056, Switzerland^b Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA^c Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA 02115, USA^d Krantz Family Center for Cancer Research, Massachusetts General Hospital Cancer Center, Charlestown, MA 02129, USA^e Department of Cancer Biology, Dana-Farber Cancer Institute, 450 Brookline Ave., Boston, MA 02215, USA^f Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, 240 Longwood Ave., Boston, MA 02215, USA^g Dana-Farber Cancer Institute, Boston, MA 02115, USA^h Department of Chemistry, University of Basel, Basel 4056, Switzerlandⁱ Swiss Institute for Experimental Cancer Research (ISREC), EPFL, Station 19, Lausanne 1015, Switzerland. E-mail: nicolas.thoma@epfl.ch† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d5md00054h>

On the basis of global quantitative proteomics screening results, we investigated niclosamide as a potential molecular

Our proteomics screens in HCT116 colon carcinoma and KELLY neuroblastoma cells identified niclosamide as a drug that rapidly induced changes to the proteome (Fig. 1A and S1A†). Particularly, we became interested in its potential to degrade cyclin D1 *via* an apparent MGD mechanism. We treated HCT116 with niclosamide and checked the protein and transcript levels of cyclin D1 *via* immunoblotting and qPCR (Fig. 1B and C). In line with the proteomics results, we saw rapid protein degradation while the transcript levels remained unchanged. Co-treatment with E1 inhibitor TAK243, NEDD8 inhibitor MLN4924, and proteasome inhibitor bortezomib rescued protein levels, indicating the involvement of a cullin E3 ligase (Fig. 1B). These findings link niclosamide to cyclin D1 degradation through a UPS-dependent mechanism, potentially acting as an MGD.



Cyclin D1 degradation is driven by CRL4^{AMBRA1}

Intrigued by these results, we sought to identify the E3 ligase necessary for niclosamide-induced degradation of cyclin D1. We employed a FACS-based genetic CRISPR screen, previously successful in our studies.^{19,20} We established a stable cyclin D1 two-color reporter system in Z-138 acute lymphoblastic leukemia cells, using a construct with a cyclin D1-GFP fusion and an independently translated mCherry for comparative quantification (Fig. 2A). After confirming that the reporter system response mirrored that of endogenous protein (Fig. 2B and C), we proceeded with genetic screens.

Due to low Cas9 activity in Z-138 cells, we switched to Cas9-bearing Hep3B hepatoma cells. We transduced the Hep3B reporter cells with the Bison library (a sgRNA pool targeting 713 UPS-related genes) and analyzed cells showing no cyclin D1 degradation upon compound treatment (Fig. 2D). We saw an enrichment for cells lacking AMBRA1, the substrate adaptor protein in the E3 ligase CRL4^{AMBRA1}, which controls endogenous cyclin D1 levels^{21–23} or one of the members of the dimeric UBA3-NAE1 NEDD8 E1 enzyme (Fig. 2E). The importance of AMBRA1 in niclosamide-mediated degradation of cyclin D1 was validated in HCT-116^{AMBRA1} cells where no cyclin D1 degradation was observed within the short treatment time (Fig. S1B†). This suggested that niclosamide either acts as a molecular glue enhancing the AMBRA1–cyclin D1 interaction or that cyclin D1 degradation is part of a cellular response to niclosamide treatment.

SAR studies reveal distinct features of niclosamide required for cyclin D1 degradation

To identify any induced neomorphic interface involving niclosamide and leading to cyclin D1 degradation, we conducted a structure activity relationship (SAR) study. We aimed to find a site on niclosamide that wouldn't affect the interaction interface, enabling probe design (*e.g.*, for photoaffinity labeling) to elucidate any formed complex. We investigated electronic and steric demands of the two phenyl ring cores by introducing various functional groups and using cyclin D1 degradation as a phenotypic readout.

We found that the nitroarene ring was relatively tolerant to changes, provided the ring remained electron-withdrawing to some extent. Even the inductive *tert*-butyl group was tolerated, but increasing electron-donating properties by changing to an aniline or a pyridine abolished all degradation activity (Fig. 3A and S2A†). The loss of degradation of the arylsulfonamide was attributed to possible steric clashes or off-target effects, which were not further investigated. Altering the type and position of the halide on the salicylamide ring revealed that only the acidic^{24,25} hydroxy group *ortho* to the amide bond was necessary to induce cyclin D1 degradation (Fig. 3B and S2A†).

We also examined the importance of the connectivity between the two arene rings. Inverting the amide bonds in

one molecule and increasing flexibility while reducing electronic conjugation by introducing an extra CH₂ in another molecule (Fig. S2B and D†) resulted in no induced degradation, highlighting the crucial role of the amide bond. To mask the charge of a deprotonated hydroxy group on the salicylamide ring, we bridged it *via* a carbamate to the amide bond. Although this molecule was active, we could not rule out the possibility of it being a prodrug hydrolyzed to niclosamide. Taken together these results emphasized the importance of the acidic proton on the salicylamide ring, which has been previously associated with mitochondrial depolarization.²⁶ This led us to investigate a potential link between this proton shuttling to the mitochondria and the observed effects on cyclin D1 degradation.

Mitochondrial depolarization correlates with cyclin D1 degradation

To obtain a quantitative comparison of degradation and MMP, we employed JC-1, a cationic dye that exhibits green fluorescence as a monomer but red fluorescence upon mitochondrial accumulation, to assess MMP (Fig. 4A). Hence the ratio of the observed red/green fluorescence is used to evaluate mitochondrial health, where for instance a depolarization reduces the red signal relative to the green signal.

We measured MMP in HCT116^{AID-AMBRA1} cells (Fig. 4B), where an auxin-inducible degron was knocked into the endogenous AMBRA1. This allowed us to induce rapid AMBRA1 degradation and observe its impact on the observed phenotype (Fig. S3A and B†). Together with our measurements in cyclin D1 reporter cells we could use the ratiometric green/red fluorescence readouts to simultaneously evaluate the relationship between cyclin D1 degradation, MMP changes, and the role of AMBRA1 in these processes.

Among the molecules used in the SAR study, we observed a striking correlation between the extent of cyclin D1 degradation and mitochondrial depolarization (Fig. 4C) when cells were treated with a concentration of 5 μ M for 3 hours. Notably, induced degradation of AMBRA1 had no effect on depolarization but rescued niclosamide-induced cyclin D1 degradation, as expected.

Mitochondria depolarizers and other drugs induce CCND1 degradation

To test whether the observed correlation extended to other reported mitochondrial depolarizers with different modes of action, we examined the protonophores CCCP, FCCP, BAM15, the mitochondrial membrane-targeting C12 TPP, SR4,²⁷ the anaesthetic sevoflurane^{28,29} and the ionophore valinomycin. To our surprise, all these molecules induced potent cyclin D1 degradation and aligned with the trend seen with the previous series of molecules (Fig. 5A and B). Inspired by our findings, we explored whether other molecules correlating






Fig. 2 CCND1 reporter cell lines capture degradation activity. A) Set-up towards establishing the fluorescence-based reporter system for quantifying CCND1 protein levels in Z138 cells. B) The population of Z138 reporter cells (Z138 CCND1-GFP.IRES.mCherry) gated by flow cytometry and used for quantification of CCND1. C) Dose response curve of CCND1 levels relative to vehicle (DMSO) treatment in the reporter cells treated with niclosamide for 3 h. Values represent the ratio of the geometric mean of GFP and mCherry values in the gated population. D) Workflow of setting up a CCND1 reporter in Hep3B cell with Cas9 activity and performing a FACS based genetic screen of members of the UPS which can rescue drug induced CCND1 degradation. E) Genes enriched in a FACS based genetic CRISPR screen (Bison library) rescuing niclosamide induced CCND1 degradation in Hep3B.Cas9.CCND1-GFP.IRES.mCherry reporter cells. The gates used are shown in D) and each contain 5% of the population shown. Gate D contains cells with gene KO's that stabilize CCND1, whereas gate A contains cells with gene KO's that destabilize CCND1.






Fig. 3 Structure activity study of niclosamide towards CCND1 degradation. A) Molecules probing the nitroarene ring of niclosamide. Change in CCND1 levels relative to vehicle (DMSO) treatment in the reporter cells treated at a concentration of 5 μ M for 3 h. Values represent the ratio of the geometric mean of GFP and mCherry values in the gated population. B) As for A) but for molecules probing the salicylamide ring.




Fig. 4 Niclosamide induced CCND1 degradation correlates with mitochondria depolarization. A) Mitochondrial membrane potential (MMP) quantified using flow cytometry in HCT116^{AID-AMBRA1} cells stained with JC-1. B) Change in MMP relative to vehicle (DMSO) treatment in HCT116^{AID-AMBRA1} cells treated at a concentration of 5 μ M for 3 h. Values represent the ratio of the geometric mean of GFP and DsRed values of all the measures cells. C) Change in CCND1 levels relative to vehicle (DMSO) quantified in reporter cells (Z138.CCND1-GFP.IRES.mCherry) treated with the compounds listed in B) and plotted against the change in MMP measured in HCT116^{AID-AMBRA1} cells treated with the same compounds at a concentration of 5 μ M for 3 h.

with niclosamide in publicly available drug sensitivity datasets (PRISM, CTRP, accessible on <http://depmap.org>) exhibited a similar phenotype. Among the top correlates were inhibitors of the anti-apoptotic proteins Bcl-2 and Mcl-1 located on mitochondria, as well as a telomerase inhibitor (BIBR15322) and molecules like KHS101, which targets the cell growth-promoting TACC3, and tyrphostin A9, a tyrosine kinase receptor PDGFR inhibitor. Notably, the correlation between cyclin D1 degradation and mitochondrial depolarization was maintained among the active molecules (Fig. 5B).

While some molecules arguably contained acidic protons, this was not a common structural feature across all compounds (Fig. 5C). Notably, compounds like tyrphostin A9 and KHS-101, whose reported targets do not suggest disruption of mitochondrial function, have been shown to induce cytotoxicity through this mechanism.^{30,31} Interestingly, aside from the lack of niclosamide induced cyclin D1 degradation in HCT116^{AMBRA1-/-} cells, there was no increase in phosphorylation of threonine 286,^{23,32-34} a key PTM for the nuclear export and ubiquitin-mediated degradation (Fig. S3D†).




Fig. 5 Mitochondria depolarizers and other molecules show same correlation between CCND1 and mitochondrial membrane potential. A) Western blot of anti-CCND1 and anti-tubulin of HCT116 and AMBRA1 KO HCT116^{AMBRA1-/-} cells treated with vehicle (DMSO), niclosamide, CCCP, SR4, BAM15, C12TPP, FCCP and sevoflurane at 5 μ M for 2 h. B) Change in CCND1 levels relative to vehicle (DMSO) quantified in reporter cells (Z138.CCND1-GFP.IRES.mCherry) treated with the compounds listed in B) and plotted against the change in mitochondrial membrane potential measured in HCT116^{AID-AMBRA1} cells treated with the same compounds at a concentration of 5 μ M for 3 h. C) Structure of the protonophore CCCP and molecules that correlated with niclosamide drug sensitivity according to the AUC secondary screen in PRISM dataset. D) Structures of niclosamide, tizoxanide and other closely structurally related molecules.

Even though we aimed to find a molecule that induced cyclin D1 degradation *via* a MGD type mechanism, the chemical diversity of the compounds (Fig. S3E[†]), which all serve as mitochondrial depolarizers, render it unlikely that these serve as direct MGDs to CRL4^{AMBRA1}. Instead, this suggests an indirect mechanism, whereby disruption of the MMP results in the downregulation of cyclin D1 with the help of AMBRA1 and the UPS *via* an interaction that appears

distinct from the threonine 286 site used for canonical proteostasis.²¹

Disrupting mitochondria membrane potential is necessary but not sufficient for cyclin D1 downregulation

It is worth noting that the ionophore nanchangmycin, an antibiotic known to disrupt endocytic uptake of viruses and





N. H. T. is a founder and shareholder of Zenith Therapeutics as well a consultant to Ridgeline Discovery and Red Ridge Bio. B. L. E. has received research funding from Novartis and Calico. He has received consulting fees from Abbvie. He is a

 CC BY-SA

- Inhibits Proliferation of HepG2 Hepatocarcinoma Cells, *J. Biol. Chem.*, 2015, **290**(51), 30321–30341.
- 28 P. Zhang, *et al.*, Mitochondria-Related Ferroptosis Drives Cognitive Deficits in Neonatal Mice Following Sevoflurane Administration, *Front. Med.*, 2022, **9**, 887062.
- 29 R. Bains, *et al.*, Sevoflurane and propofol depolarize mitochondria in rat and human cerebrocortical synaptosomes by different mechanisms, *Acta Anaesthesiol. Scand.*, 2009, **53**(10), 1354–1360.
- 30 S. J. Park, *et al.*, A receptor tyrosine kinase inhibitor, Tyrphostin A9 induces cancer cell death through Drp1 dependent mitochondria fragmentation, *Biochem. Biophys. Res. Commun.*, 2011, **408**(3), 465–470.
- 31 E. S. Polson, *et al.*, KHS101 disrupts energy metabolism in human glioblastoma cells and reduces tumor growth in mice, *Sci. Transl. Med.*, 2018, **10**(454), eaar2718.
- 32 J. P. Alao, The regulation of cyclin D1 degradation: roles in cancer development and the potential for therapeutic invention, *Mol. Cancer*, 2007, **6**, 24.
- 33 H. C. Seftel and H. J. Heinz, Treatment of Human Tapeworm Infections with Yomesan: Single Dose Treatment in Non-Fasting Subjects, *S. Afr. Med. J.*, 1964, **38**, 263.
- 34 PATIENT INFORMATION LEAFLET – YOMESAN CHEWABLE TABLETS, https://www.bayer.com/sites/default/files/YOMESAN_EN_PIL.pdf, (accessed April 2025).
- 35 W. Li, *et al.*, Nanchangmycin regulates FYN, PTK2, and MAPK1/3 to control the fibrotic activity of human hepatic stellate cells, *Elife*, 2022, **11**, e74513.
- 36 K. Rausch, *et al.*, Screening Bioactives Reveals Nanchangmycin as a Broad Spectrum Antiviral Active against Zika Virus, *Cell Rep.*, 2017, **18**(3), 804–815.
- 37 B. S. Saller, *et al.*, Acute suppression of mitochondrial ATP production prevents apoptosis and provides an essential signal for NLRP3 inflammasome activation, *Immunity*, 2024, 90–107.e11.
- 38 F. Ek, *et al.*, Sorafenib and nitazoxanide disrupt mitochondrial function and inhibit regrowth capacity in three-dimensional models of hepatocellular and colorectal carcinoma, *Sci. Rep.*, 2022, **12**(1), 8943.
- 39 H. Sun, *et al.*, Nitazoxanide impairs mitophagy flux through ROS-mediated mitophagy initiation and lysosomal dysfunction in bladder cancer, *Biochem. Pharmacol.*, 2021, **190**, 114588.
- 40 Y. Lee, *et al.*, Cyclin D1-Cdk4 controls glucose metabolism independently of cell cycle progression, *Nature*, 2014, **510**(7506), 547–551.
- 41 T. Sakamaki, *et al.*, Cyclin D1 determines mitochondrial function in vivo, *Mol. Cell. Biol.*, 2006, **26**(14), 5449–5469.
- 42 K. Bhalla, *et al.*, Cyclin D1 represses gluconeogenesis via inhibition of the transcriptional coactivator PGC1alpha, *Diabetes*, 2014, **63**(10), 3266–3278.
- 43 C. Van Humbeeck, *et al.*, Parkin interacts with Ambra1 to induce mitophagy, *J. Neurosci.*, 2011, **31**(28), 10249–10261.
- 44 F. Strappazon, *et al.*, AMBRA1 is able to induce mitophagy via LC3 binding, regardless of PARKIN and p62/SQSTM1, *Cell Death Differ.*, 2015, **22**(3), 419–432.
- 45 M. Di Rienzo, *et al.*, Role of AMBRA1 in mitophagy regulation: emerging evidence in aging-related diseases, *Autophagy*, 2024, **20**(12), 2602–2615.
- 46 N. S. Scholes, *et al.*, Inhibitor-induced supercharging of kinase turnover via endogenous proteolytic circuits, *bioRxiv*, 2024, preprint, DOI: [10.1101/2024.07.10.602881](https://doi.org/10.1101/2024.07.10.602881).