



Cite this: *Chem. Commun.*, 2017, 53, 284

Computational tools for the evaluation of laboratory-engineered biocatalysts

Adrian Romero-Rivera,^a Marc Garcia-Borràs^b and Silvia Osuna*^a

Biocatalysis is based on the application of natural catalysts for new purposes, for which enzymes were not designed. Although the first examples of biocatalysis were reported more than a century ago, biocatalysis was revolutionized after the discovery of an *in vitro* version of Darwinian evolution called Directed Evolution (DE). Despite the recent advances in the field, major challenges remain to be addressed. Currently, the best experimental approach consists of creating multiple mutations simultaneously while limiting the choices using statistical methods. Still, tens of thousands of variants need to be tested experimentally, and little information is available on how these mutations lead to enhanced enzyme proficiency. This review aims to provide a brief description of the available computational techniques to unveil the molecular basis of improved catalysis achieved by DE. An overview of the strengths and weaknesses of current computational strategies is explored with some recent representative examples. The understanding of how this powerful technique is able to obtain highly active variants is important for the future development of more robust computational methods to predict amino-acid changes needed for activity.

Received 22nd July 2016,
Accepted 6th September 2016

DOI: 10.1039/c6cc06055b

www.rsc.org/chemcomm

1. Introduction

Billions of years of evolution have made enzymes superb catalysts capable of accelerating reactions by many orders of magnitude while also being compatible with life. This rate acceleration is

achieved by decreasing the activation barriers of reactions, making them possible at lower temperatures and pressures. Among all known catalysts, enzymes (*i.e.*, biocatalysts) are often the most efficient, specific and selective. Enzymes operate under mild aqueous conditions, are biodegradable and non-toxic, and their high selectivities and efficiencies reduce the number of work-up steps and provide product in higher yields. Many enzymes catalyse complex reactions in a few steps, which is in contrast to traditional catalysis that often requires many protection and deprotection steps. These advantageous characteristics of biocatalysis make enzyme-catalysed routes attractive

^a Institut de Química Computacional i Catàlisi and Departament de Química Universitat de Girona, Campus Montilivi, 17071 Girona, Catalonia, Spain.
E-mail: silvia.osuna@udg.edu

^b Department of Chemistry and Biochemistry, University of California, 607 Charles E. Young Drive, Los Angeles, California, 90095, USA



Adrian Romero-Rivera

Adrian Romero Rivera received his bachelor in Chemistry from the University of Girona (UdG) in 2013. He completed his Masters in Advanced Catalysis and Molecular Modeling (MACMOM) in 2014. He is currently a PhD student in the Institut de Química Computacional i Catàlisi (IQCC) working on computational exploration of (bio)inorganic reaction mechanisms under the supervision of Dr Marcel Swart and Dr Silvia Osuna.



Marc Garcia-Borràs

Marc Garcia-Borràs obtained his PhD (University of Girona, 2015) and received the special award for his thesis from the University of Girona. The same year, he joined Dr Osuna's group at the Institut de Química Computacional i Catàlisi (IQCC) as a postdoctoral researcher working on computational rational enzyme design for six months. He is currently a postdoctoral scholar in Prof. K. N. Houk's group at the University of California, Los Angeles (UCLA), where he is working on the application of computational tools for the evaluation and design of new useful biocatalysts.



alternatives for chemical manufacturing due to the minimization of costs and environmental advantages. The vast potential of biocatalysis is exemplified with the enzyme-catalysed synthesis of some blockbuster drugs.^{1–4} Indeed, biocatalysis has won 18 Presidential Green Chemistry awards since 2000 (5 awards given per year). These awards emphasize the 12 Principles of *Green Chemistry*, which consider environmental factors and energy efficiency among other points.

The current use of biocatalysts in industry is still limited, as enzymes need to be modified to be stable for the desired pHs, temperatures, and solvents. Many chemical transformations of industrial interest do not have a natural enzyme capable of catalysing the reaction, and the biocatalyst active site is often too small for proper binding of the desired substrate. In addition, the lack of a precise understanding of enzyme catalysis makes the alteration of the natural activity of enzymes for synthetically relevant targets a tremendous challenge, even though some key factors have been identified.^{5–9}

All of the available strategies for enzyme engineering consist of the following steps: (i) selection of mutation points, (ii) making the mutations, and (iii) evaluation of the new variants for activity. These can be targeted using computational and/or experimental approaches leading to rational, semi-rational or non-rational enzyme design strategies. Rational design limits the screening effort to a small number of mutations. Semi-rational strategies are based on exploiting initial desired enzymatic activities obtained by rational computational design *via* subsequent rounds of laboratory evolution, which is similar to the way that promiscuous side-reactions of natural enzymes are enhanced.^{10–12} In recent years, successful designs for a broad scope of challenging chemical transformations have been produced using semi-rational design approaches.^{13–22} At the other extreme, in non-rational enzyme evolution, powerful screening methods identify active variants from a large random library of mutants.

The enzyme-engineering field was revolutionized after the discovery of molecular biology methods that modify enzymes using an *in vitro* version of Darwinian evolution. This strategy is

now commonly called Directed Evolution (DE).^{23–26} Initially, iterative cycles of random amino-acid changes were introduced and were followed by selection of the variants with improved thermostability, substrate specificity and enantioselectivity. Since then, many subsequent improvements have been introduced, which include protein engineering,^{23,27,28} gene synthesis,²⁹ sequence analysis,^{30,31} bioinformatics tools,^{13,14,32–35} and high-throughput screening techniques, such as an ultrahigh-throughput droplet-based microfluidic screening platform.^{36,37} Indeed, the success of DE experiments depends on genetic diversity and on high-throughput screening or selection methods.^{27,36}

One powerful strategy consists of combining random mutagenesis with statistics to construct mathematical models of protein sequence and function.² This ensures the accumulation of beneficial mutations leading to the desired activity in a stepwise fashion. DE has become a powerful method to produce novel enzymes with enhanced activity, and it has the advantage that mutations can be introduced at the enzyme active site and at distal positions. The latter are found to be relevant for increasing the enzyme catalytic activity (k_{cat}), as highlighted recently by Kell and coworkers.³⁸ Recent examples of DE-engineered enzymes include the use of ketoreductases for the manufacture of chiral intermediates for pharmaceuticals such as atorvastatin, the active ingredient in Lipitor[®], and a transaminase for the manufacture of sitagliptin, the active ingredient in Januvia[®].^{1,3,4,39} The main drawback of DE is that little information is available as to how these mutations lead to enhanced enzyme proficiency. Many efforts are being made to rationalize how DE introduces new mutations to alter the enzyme catalytic activities.⁴⁰

In this feature article, a short review of the different computational strategies that are being applied to rationally design enzymes is first presented and is followed by a more detailed overview of the available computational approaches currently used to evaluate laboratory-generated enzyme variants. Some recent representative examples are discussed to illustrate the *pros* and *cons* of the different methodologies.



Sílvia Osuna

Sílvia Osuna is currently a Ramón y Cajal researcher at the Institut de Química Computacional i Catàlisi (IQCC) at the University of Girona (Spain). She received her PhD in 2010 from the University of Girona (UdG). She was awarded a Marie Curie IOF Fellowship for a postdoctoral position at the group of Prof. Houk at the University of California, Los Angeles (UCLA). Her research is focused on the study of biochemical processes related to enzyme catalysis and

the development of a computational protocol for the design of new enzymes of pharmaceutical interest.

2. Computational tools for the rational design of enzymes

Computational methods provide an attractive alternative to understand, model and rationally construct novel enzyme catalysis at a reduced cost. The development of robust computational strategies capable of improving and enhancing enzymatic catalysis as DE currently does is one of the most challenging and exciting roads in the biocatalysis field.⁴¹

Many computational strategies have been used, and they range from the enhancement of promiscuous activities of natural enzymes employing multiple sequence and structure alignments,^{42,43} the simultaneous design of the entire protein backbone structure and sequence,^{44,45} and the (re)design of the active site of natural enzymes by mutating a subset of the active site residues while maintaining a rigid backbone.^{46–49} The




with the final goal of improving current rational enzyme evolution protocols. Computational techniques offer the possibility to analyse, at the atomic level, how a particular mutation modifies enzymatic chemo-, regio- or stereo-selectivities, or how, for example, the substrate binding is affected due to changes in the protein conformational dynamics induced by these amino acid substitutions.

In this section, a general description of the most important and applied computational strategies is reported and illustrated with different examples. All of the techniques discussed here differ in the level of resolution used to describe the protein interactions and in how they sample the enzyme conformational space (see Fig. 2).

a. Quantum mechanics calculations and quantum chemical cluster approach

Quantum Mechanics (QM) calculations include a wide range of approximations whose main objective is to solve the Schrödinger equation, providing useful information about the molecular properties and energetics of a particular chemical system. QM calculations include all the *ab initio* methodologies based on the Hartree-Fock approximation, semiempirical methods, and Density Functional Theory (DFT), which is currently the most extensively used QM methodology.⁸³ Although DFT provides an accurate description of the chemical system, the associated computational cost increases exponentially when the size of the system grows. This limitation makes the treatment of an entire



Computational strategy	Accounting for:				Useful for studying:							Computational cost	References:	
	QM	MM	Active site	Whole enzyme	Protein flexibility	Enantioselectivity/ Stereoselectivity	Substrate scope	Reaction mechanism	Conformational dynamics	Effect of distal mutations	Computational strategy		Key examples	
Cluster model	✓	✗	✓	✗	✗	✓	✓	✓	✗	✗	+	89, 90	91-93	
Theozyme	✓	✗	✓	✗	✗	✓	✓	✓	✗	✗	+	84-87	88	
QM/MM	✓	✓	✓	✓	✗	✓	✓	✓	✗	✗	++	94-96	97, 98	
QM/MM-MD	✓	✓	✓	✓	✓	✓	✓	✓	✗	✗	++++	99	99	
EVB	✓	✓	✓	✓	✗	✓	✓	✓	✗	✗	++	100-102	103, 104	
MD	✗	✓	✓	✓	✓	✓	✓	✗	✓	✓	+	105-108	109, 110	
MC	✗	✓	✓	✓	✓	✓	✓	✗	✓	✓	+	111, 112	113	

Fig. 2 Schematic of the available computational tools for evaluating and rationalizing the effect of mutations on the catalytic activity of laboratory-engineered enzymes. The strengths (represented with ✓) and weaknesses (with ✗) are highlighted with the associated computational costs (+). Key references for computational details and representative examples are provided.



protein at the DFT level expensive, which explains why only a few active site residues are considered in QM-based methodologies.

QM *theozyme* calculations have been extensively used in the framework of the *inside-out* protocol to computationally design new enzymes. A *theozyme* is a DFT-optimized, three-dimensional arrangement of amino acid side chains that are optimally disposed to stabilize the TS of the targeted reaction.⁴⁹ The *theozyme* strategy can also be used to study enzymatic mechanisms and to unravel new biological pathways.^{84–87} For example, in a very recent study, this methodology was applied to elucidate the unprecedented biosynthetic pathway of penigequinolone, and a cationic epoxide rearrangement under physiological conditions was observed for the first time.⁸⁸ *Theozyme* calculations were used to analyse and evaluate different possible reaction mechanisms catalysed by key active site residues for a new isolated enzyme (PenF), providing a clear explanation for the product formation experimentally observed. Moreover, *theozyme* calculations together with MD simulations have also been used by the Houk group to evaluate the catalytic performance of different DE-engineered variants as described in Section 3.d.

A popular QM strategy is the Cluster Model (CM) approach, which focuses on a well-chosen shell of amino acids from the active site of the enzyme in consideration (see Fig. 3). This methodology was developed and used more than thirty years ago by Siegbahn, despite the fact that the first application to an enzyme reaction mechanism was only achieved in 1997.⁹¹ Only those residues playing a critical role in the enzymatic mechanism are included in the cluster model. If we take a look at uses in the past, the first systems only included 20–30 atoms without imposing any constraints, which is similar to the *theozyme* approach described above. However, thanks to the boost in computational power, more complex systems can now be handled. Current CM calculations contain more than 200 atoms and include some atomic constraints to better

mimic the protein backbone and the enzyme active site cavity. More specific information about the CM approach and details about the size of the systems and some applications can be found in ref. 89 and 90.

Since the first application of CM, a variety of studies using this methodology have been published, and most of them are related to the enzymatic reaction mechanisms.^{114–117} Some recent applications of CM calculations that target different mutated enzymes with the goal of rationalizing the effect of the introduced mutations (some of them *via* DE) will be discussed. The *pros* and *cons* that this strategy offers will be highlighted.

Limonene epoxide hydrolases (LEH) naturally catalyse the hydrolysis of limonene-1,2-epoxide to limonene-1,2-diol. However, some LEH enzymes can also accept other epoxides as substrates to yield their corresponding diols *albeit* with lower enantioselectivities.¹¹⁸ Zheng and Reetz applied DE to produce LEH variants capable of catalysing non-natural epoxide substrates (*meso*-cyclopentene oxide) with high enantioselectivities.¹¹⁸ Himo and coworkers⁹² studied the enantioselectivity and mutational effects using the CM approach (see Fig. 3). Some residues were truncated, and some atoms were frozen during the DFT geometry optimizations performed at B3LYP/6-31G(d,p) level of theory. Single point calculations with larger basis sets including solvent effects through the CPCM model, zero-point correction, and dispersion effects were applied afterwards. This procedure is typically used for the CM approach and yields accurate predictions of enzymatic reaction mechanisms.^{89,90} Their results showed in the case of the wild-type (WT) enzyme similar energy barriers for the opening of the oxirane ring, which is in agreement with the experiments that showed a small 14% ee for the *R,R*-product with an energy difference of 0.2 kcal mol⁻¹. The calculations indicated that the active site cavity of the WT enzyme is quite spacious, so the cyclopentene oxide substrate can be oriented to expose both faces for the nucleophilic attack. Based on these observations, different mutations were proposed. The optimized structures of the transition states (TSS) indicated that the mutations have a direct effect on the substrate-binding pose. The proposed Leu74Ile and Ile80Cys mutations create some additional space on one side of the active site cavity, thus favoring the attack on the less hindered C2 position. This double mutant presents a lower activation barrier for the addition to C2 and, therefore, exhibits a higher selectivity towards the (*R,R*)-product formation. The combination of Leu114Cys and Ile116Val mutations located on the other side of the catalytic pocket make the other active site side less hindered, leading to a preference for the (*S,S*)-product formation (*pro-R,R* TS +1.4 kcal mol⁻¹ higher in energy).

The mechanism and stereoselectivity of AMDase enzymes were also explored with CM, and a variety of substrates were employed.⁹³ AMDase catalyses the asymmetric decarboxylation of α -aryl- α -methyl malonates. In this study, Himo and coworkers applied two different models (I and II) to analyse the substrate preferences and stereoselectivities of AMDase. The substrate used for model I was α -methyl- α -phenylmalonate (methylphenylmalonate), and methylphenylmalonate and α -methyl- α -vinylmalonate (methylvinylmalonate) for model II. The latter has a




Fig. 3 259-atom cluster model structure optimized by Himo and coworkers for LEH in ref. 92. The cluster model consisted of: Asp132–Arg99–Asp101 catalytic triad, the nucleophilic water, two hydrogen-bonded residues (Tyr53–Asn55), and different groups defining the active-site cavity, Met32, Leu35, Leu74, Met78, Ile80, Val83, Leu103, Leu114 and Ile116. The active site residues that are mutated are represented in pink (Leu74 and Ile80) and orange (Leu114 and Ile116). Atoms in orange were fixed in their X-ray coordinates. Hydrogen atoms are omitted for clarity.



smaller size and may influence the stereoselectivity of the reaction. Model I (composed of 81 atoms) consists of Gly74, Thr75, Ser76, Tyr126, Gly189 (dioxanion hole) and Cys188 (responsible for the protonation step of the reaction mechanism). This model lacks important residues involved in the substrate binding and is too small to accurately reproduce the experimentally observed enantioselectivities. In contrast, model II (225 atoms) also includes Pro14, Pro15, Leu40, Val43, Tyr48, Val56, Met159, and Gly190, which mimic the small and big cavities of the active site, and was able to reveal the differences in enantioselectivity for methylphenylmalonate as a substrate. The *S*-product has a small methyl group pointing to the more solvent-accessible pocket and a much bulkier phenyl group for the hydrophobic pocket, which is formed by Leu40, Val43, and Val156 through their side chain steric repulsion. The *S*-product is less stable (+14.1 kcal mol⁻¹) than the *R*-product. The reaction leading to the *R*-product presents an activation energy of 16.2 kcal mol⁻¹, which is in line with the experimental measurements (14–16 kcal mol⁻¹) and the observed ee of > 99%.¹¹⁹ The good agreement observed between the computations and experiments is due to the contribution of the extra residues included in the large model, which account for an extra hydrogen bond between the backbone amide and the carboxylate group of the Thr75 and Ser76, respectively. This study exemplifies the importance of properly selecting the cluster model size for correctly modeling the enzymatic enantioselectivities. Once the best cluster model for reproducing the stereoselectivities observed for the WT enzyme is built, then it can be applied to evaluate some variants and analyse the effect of the new, introduced mutations. Some reported studies^{120–122} showed an enhanced enantioselectivity preference for the *S*-product over the natural enzyme when Gly74Cys, Cys188Ser mutations were introduced. The position of the new Cys74 residue, located at the *Re* face of the enediolate intermediate, was found to determine the stereochemistry of the product yielding the *S*-enantiomer.^{120–122} In the case of the smaller methylvinylmalonate substrate, the energy difference between the *S*/*R*-products was underestimated compared to the experiments,¹²³ demonstrating that a larger model with a more flexible binding pocket is needed to explain the enantioselectivity of the smaller substrate.

The two examples described above demonstrate that the CM approach is a powerful tool for rationalizing the effect of active site mutations on the enantioselectivity of a particular enzymatic reaction. However, CM is limited because this approach is highly dependent on the initial amino acid selection to build the CM and because the flexibility of some loops close to the active site cannot be properly considered.

b. Hybrid quantum mechanics/molecular mechanics calculations

The aggregation of quantum mechanics/molecular mechanics (QM/MM) techniques reported by Warshel and Levitt⁹⁴ is an extensively used technique for studying biomolecular systems and for drug design.^{95,96} The impact and importance of this computational strategy was meritoriously recognized in 2013

with the Nobel Prize in Chemistry.¹²⁴ QM/MM methods were initially developed to allow for the study of those chemical processes that require quantum mechanical treatment, *e.g.*, bond cleavage or bond forming reactions, but that are too large to be fully studied at a high level of theory. Thus, QM/MM calculations consist of the treatment of a small portion of the chemical system, usually the enzyme active site, using QM, and the rest of the system is described using a less rigorous but computationally more efficient level of theory, such as Molecular Mechanics (MM). MM methods use classical mechanics to model atom–atom interactions, and the energy of the system is computed using simple potential energy functions called force fields.¹²⁵ In the particular case of biocatalysis, QM/MM methods allow the study of chemical reactions (*i.e.*, bond forming/cleavage processes) involving systems of millions of atoms at an atomic resolution, explicitly considering the effects of the protein environment and solvent molecules and their influence on the reaction profile. Two of the most successful QM/MM approaches are (i) the Empirical Valence Bond (EVB) theory (see next section)¹⁰⁰ and (ii) the molecular orbital self-consistent field (MO-SCF).^{126–128} The MO-SCF QM/MM approximation is based on solving the time independent Schrödinger equation using a global Hamiltonian that can be divided in three parts, as shown in Fig. 4: (i) the QM part (H_{QM}) for the small portion of interest in the system where the chemistry takes place (fully represented at the QM level), (ii) a classical part (H_{MM}) to treat the rest of the system and environment, and (iii) a coupling term ($H_{QM/MM}$) that describes the interactions between the QM part and the MM part, which includes the electrostatic perturbation that the QM wave-function suffers due to the presence of the polarized MM environment.^{95,108} In this section, we will review some recent applications of the QM/MM MO-SCF methods for the study of enzymatic reactions and the effect of introduced mutations. These examples highlight how powerful this technique is for mutation analysis and prediction.

Ryde and coworkers reported a comparative study evaluating the effect on both geometry and energetics of using only QM models or hybrid QM/MM.¹²⁹ This paper collects all the *pros* and *cons* of using these two strategies and concluded that QM/MM




Fig. 4 Schematic representation of the QM/MM treatment of a biocatalytic system.



extracted. Based on their QM/MM calculations, they suggested that the main role of Glu119 during the methylation step is the formation of a hydrogen bond with the substrate. These results cannot, however, explain experiments where the Glu119Gln variant was found to be substantially less active than the WT, even though their active sites are similar.¹³⁶ These observations suggest that the impact of the Glu119 residue on the catalysis should be higher. QM/MM calculations for the β -elimination step showed that Glu119 plays a key role. The proton of Glu119 is transferred to the N3 position of the DNA substrate with an energy barrier of 4.8 ± 0.3 kcal mol⁻¹. The latter induces an increased charge on the proton to be extracted and facilitates deprotonation by the water molecule. This C5 deprotonation mediated by water was more favorable than having a hydroxide anion in the enzyme active site. In the final step of the reaction mechanism, the cleavage of the C–C bond formed between the cytosine substrate and Cys81 takes place, followed by protonation of Glu119. The computed activation energy for the bond cleavage step is 4.8 ± 0.4 kcal mol⁻¹ and 3.5 ± 0.2 kcal mol⁻¹ for Glu119 protonation. Experiments where Glu119Ala, Glu119Gln, and Glu119Asp mutations were introduced showed a drastic decrease in the activity during the methylation step.¹³⁶ This QM/MM study by Tuñón and coworkers demonstrated how important the Glu119 residue is during the whole reaction path and clarifies the effect of the Glu119 mutation on the pre-steady-state and steady-state rate constants. Additionally, they showed that a crystallographic water molecule, instead of a hydroxide anion, is responsible for the substrate C5 deprotonation. The QM/MM calculations clarified two highly debated issues of this enzymatic mechanism.

The provided examples show how QM/MM calculations can help elucidate the role of certain active site residues and give some invaluable insights into the enzyme reaction mechanism. Most of these studies either consider the X-ray structure of the enzyme, as in the first provided example, or perform some classical MD simulations (see Section 3.d) to generate an ensemble of conformations from which to start the QM/MM analysis (as described in the last study provided). It is, however, recognized that enzymes are highly dynamic structures that can adopt different relevant conformational states in the course of the reaction. QM/MM calculations can also be coupled to MD simulations to properly describe the substrate conformational changes and the dynamic nature of the enzyme. In a recent perspective study by Rovira and coworkers, the importance of QM/MM-MD calculations for some carbohydrate-active enzymes was highlighted.⁹⁹ In particular, they used QM/MM with the MD enhanced sampling technique metadynamics to map the conformational Free Energy Landscape (FEL). QM/MM-MD simulations have a high computational cost, which hampers their application in studying enzyme conformational dynamics.

c. Empirical valence bond calculations

Warshel developed the EVB approach.¹⁰⁰ The EVB method is a QM/MM approach that describes bond forming and breaking in a chemical reaction through diabatic states from the classical Valence Bond (VB) description of the reactants, intermediates,




Fig. 6 Qualitative representation of EVB treatment for a prototype S_N2 reaction. Different possible VB configurations of the system and their combination to describe the different diabatic VB states of the reactants and products are depicted. The bold black line describes the EVB adiabatic ground state of the reaction. The G term corresponds to the gap between the ground state and the charge-transfer state, and the B term refers to the mixing between states that occurs at the TS, which is known as the resonance energy in the VB framework.

and products species as, shown in Fig. 6. EVB takes the energy of the different species from QM calculations and through diagonalization of the energy matrix generated obtains the eigenvalues. This procedure also allows the possibility to apply EVB to obtain the free energy profile of the studied reaction, which is a very relevant tool for the study of biocatalytic processes. A detailed description of the theory, capabilities, applications and limitations of the EVB method can be found in ref. 101 and 102. The EVB method can provide information on key residues and mutant effects.^{137–139} In this section, some recent applications of EVB to study enzymatic reactions and the role of new introduced mutations are summarized. The strengths and weaknesses of this technique when applied to the study of enzyme catalytic activity are highlighted and discussed.

Kamerlin and coworkers applied the EVB Free Energy Perturbation/Umbrella Sampling (EVB-FEP/US) method to study the enantio- and regio-selectivity of an epoxide hydrolase (EH),¹⁰³ which catalyses the hydrolysis of *trans*-stilbene oxide (TSO) into the corresponding diol. The system included in the EVB region was the TSO substrate, the side chain of Asp105 and His300, and the hydrolytic water molecule. The reaction mechanism consists of a nucleophilic attack of Asp105 on the epoxide, leading to the formation of an alkyl-enzyme intermediate. At this step, the side chain of His300 and the water molecule are treated as beholders because they are not involved at this stage, but they are crucial to maintaining an unbroken EVB region. The next important step consists of the hydrolysis of the alkyl-enzyme intermediate, and all the Asp105, His300, and water molecule are simultaneously involved. The *R,R*- and *S,S*-enantiomers of TSO were considered in the study as well as the nucleophilic attack on both the C1 and C2 positions of the epoxide ring. The protonation state of some residues was investigated and corroborated with experiments.^{140,141} A detailed





Fig. 7 (A) Homology model of the nitrating cytochrome P450 TxE (PDB code: 4TPO). The F/G loop is highlighted in purple, and haem, L-tryptophan and His176 are represented by balls and sticks. (B) X-ray structures for the wild-type LovD enzyme (PDB code: 3HL9). (C) Optimized quantum mechanics arrangement of the catalytic triad. Lys79 is represented as a blue sphere, Ser76 as yellow, and Tyr188 as pink. Overlay of 10 snapshots from the ANTON MD simulations for wild-type enzyme in the monomeric state; LovD1 and LovD9 together with the computed percentage of time that the catalytic triad stays in the proper arrangement for catalysis along the MD trajectory. The experimental k_{cat} values (in min^{-1}) are also included.

perform MD simulations on the 100 μs timescale, which is 200–2000 times longer than those previously reported for P450s.¹⁴⁶ The included mutation in the F/G loop was found to modulate the loop dynamics and completely shifted the enzyme regioselectivity from the C4 to the C5 position of L-tryptophan (see Fig. 7A). The simulations revealed that the F/G loop can adopt two different conformations, the open state needed for substrate binding and product release and the closed-lid conformation essential for excluding water molecules from the enzyme active site and promote catalysis. By determining the transition-path (TPT)¹⁴⁷ connecting both open- and closed-lid conformations, they could characterize the interactions that gate the transition and identify a key intermediate state. In the latter state featuring attributes from both open and closed conformations, a key His176 was identified that is hydrogen-bonded to Tyr89 and induces a partial opening of the F/G loop. The mutation of this position was hypothesized to shift the loop equilibrium towards the catalytically competent closed-lid conformational state. Site-saturation mutagenesis at position 176 indicated that the mutation of the aromatic phenylalanine, tyrosine, and tryptophan residues improved the binding of L-tryptophan and resulted in nitration at the C5 position. MD simulations, together with X-ray crystallography, indicated that the new mutants presented a 90:10 ratio of closed-open conformations, which was in contrast to the 50:50 ratio observed

for the wild-type enzyme. In addition to that, the bulky residues at position 176 forced the substrate to place the indole C5 position close to the ferric peroxynitrile, explaining the increase in regioselectivity.

In another study,¹⁰⁹ we demonstrated the utility of all-atom unbiased microsecond MD simulations performed on the ANTON machine for rationalizing the improvement on the catalytic proficiency of some DE-engineered enzymes for the synthesis of the cholesterol-lowering drug simvastatin.

The natural enzyme studied, LovD, is an acetyltransferase that was found to catalyse the transfer of an α -methylbutyrate side chain to the C8 position of monacolin J acid (MJA) to yield lovastatin. In the natural pathway, LovD is acylated at position 176 because of its interaction with the acyl carrier protein (ACP) domain of its binding partner protein LovF. Afterwards, the α -S-methylbutyrate side chain is then regioselectively transferred to the C8 hydroxyl of MJA. Envisioning a potential enzymatic manufacturing route for the synthesis of simvastatin, 9 rounds of directed evolution were applied to yield LovD9 that accepted an unnatural acyl donor, α -dimethylbutyryl-S-methylmercapto-propionate, as a substrate and obviated the need for allosteric regulation exerted by LovF. X-ray crystallography and nanosecond-scale MD simulations were unable to provide an explanation for the increase in the catalytic activity of the last round of mutants because the catalytic residues Ser176, Tyr188, and Lys79 displayed an almost identical catalytically competent arrangement as predicted by QM (see Fig. 7B and C).

The ANTON microsecond timescale MD simulations performed on the apo monomeric state of the enzyme indicated that the introduced mutations along the DE pathway progressively stabilized the catalytically competent arrangement of the triad since the ideal QM geometry was observed with increasing frequency. The higher population of the catalytically competent conformational state suggested that the free energy of the latter conformation was gradually lowered in the DE pathway. These findings for the monomeric state of the enzyme in an explicit solvent contrasted with the MD simulations performed on the wild-type dimer X-ray structure and on a model for the LovD–LovF complex. The latter demonstrated that protein–protein interactions stabilized the QM ideal geometry of the catalytic triad for catalysis.

In other studies, short MD simulations have been found to be crucial to evaluate the effect of the included mutations on the enzyme catalytic activity, especially in cases where active site mutations have been introduced.^{148–151} Janssen and coworkers selected four different haloalkane dehalogenases for which experimental data on the enantioselectivity conversion of a variety of substrates was available.¹⁵¹ They modeled the enantioselectivity by evaluating the frequency of occurrence of Near Attack Conformations (NAC) for pairs of enantiomers during the MD simulation. NAC is defined as the conformations that deviate from the QM TS presenting angles between the reactive atoms within 20° and distances between reactive atoms of less than the sum of their van der Waals radii.¹¹⁰ They were able to accurately model the enantioselectivity using a cluster of short (10 ps) independent MD simulations with



different initial velocities. This approach was then used to design highly stereoselective mutants of limonene epoxide hydrolase.⁵⁹ Similarly, Zhou *et al.* created an esterase with enhanced selectivity in hydrolytic kinetic resolutions using DE and analysed the source of the enantioselectivity with X-ray crystallography and short MD simulations.¹⁵⁰ Zhou and coworkers performed 80 nanosecond MD simulations to evaluate the effect of two mutations located at the product-release site in an epoxide hydrolase for the efficient bioresolution of bulky pharmac substrates.¹⁴⁸ The combination of MD simulations with the software CAVER⁷⁹ were used to evaluate the effect of the included mutations on the substrate access tunnels of a dehalogenase enzyme.¹⁴⁹ The percentage of time that the access tunnel was in an open or closed conformation was found to correlate with the catalytic activity of the variant.

These studies demonstrate how classical MD simulations coupled with QM calculations can capture enzyme conformational states key for catalysis, which cannot be elucidated by visual inspection of the X-ray data nor with high-level calculations based on an enzyme conformation taken from the crystallographic structure. MD simulations can therefore elucidate the role of both the active site and distal mutations on the catalytic activity of the enzyme. This is in contrast to the other methodologies discussed so far.

e. Monte-Carlo conformational sampling simulations

Monte-Carlo (MC) simulations are a stochastic approach to the task of generating a set of representative configurations under given thermodynamic conditions, such as temperature and volume. In the metropolis MC simulation scheme, the method randomly generates potential movements on a given structure, and the movements are accepted or rejected based on the energy of the new pose with respect to the previous structure. If the change in energy is negative, the new configuration is accepted; otherwise, depending on a probability given by a Boltzmann factor, the new structure will be discarded or selected. In contrast to MD simulations, MC gives no information about the time evolution of structural events and is inefficient for exploring the configurational space of large biomolecules.

Recently, MC has been coupled with normal mode analysis methods to sample conformational changes along normal modes.^{111,112} PELE (Protein Energy Landscape Exploration) is a MC algorithm developed by Guallar and coworkers that subjects the ligand to random rotations and translations and perturbs the protein based on the Anisotropic Network Model (ANM).¹¹¹ PELE has been successfully applied to evaluate the effect of mutations introduced *via* DE^{113,152} and has shown great promise for use in metalloenzyme designs.¹⁵³ One recent example is the application of PELE combined with QM/MM calculations to unveil how substrate oxidation was improved in a DE-engineered laccase.¹¹³ The enzyme is a copper-based oxidase that reduces oxygen to water *via* a one-electron oxidation of a reducing substrate, 2,2'-azino-bis(3-ethylbenzo-thiazoline-6-sulfonic acid) or 2,6-dimethoxyphenol. The evolved laccase presented 5 mutations, two located in the T1 pocket, two at the substrate entrance, and an additional one on the protein surface.

PELE was used to locate 20 substrate-bound structures within 5 kcal mol⁻¹ of the lowest binding energy pose. Afterwards, QM/MM calculations were performed to evaluate the amount of spin density localized on the substrate to estimate the electron transfer reactivity. Their simulations showed that mutations introduced *via* DE increased the enzyme catalytic activity by enhancing the substrate binding rather than the metal redox potential. The mutations located at the enzyme active site affected the binding mode of the substrate and provided a more favorable electrostatic environment for oxidation. The same strategy was recently used to evaluate a doubly mutated peroxygenase engineered by DE for the synthesis of 1-naphthol.¹⁵²

MC methodologies are, therefore, a cheap alternative to MD for evaluating enzyme conformational dynamics. However, they do not provide the time-dependence of the observed events and cannot be applied to evaluate correlated inter-residue motions.

4. Conclusions

Directed evolution has become one of the most powerful strategies to design new enzymes with high activities towards non-natural substrates or reactions. Even though the catalytic proficiencies of DE-engineered enzymes are high, the strategy is not rational because how the included mutations affect catalysis and enzyme stability has not been elucidated. This clearly limits the process since the efforts invested in optimizing a given biocatalyst cannot be further applied in other (un)related cases. In addition, DE requires an initial residual activity for the target substrate or reaction.

More powerful enzyme design methods could be developed if a comprehensive understanding of the relationship between mutations and enzyme catalytic activity was achieved. In this review, the available computational approaches that can be used to elucidate the basis of DE rules of operation were presented with some recent representative examples. These can be divided into strategies that tackle: (i) the evaluation of the enzyme reaction mechanism in atomistic detail using quantum mechanics, if only certain active site residues are considered, or hybrid quantum mechanics/molecular mechanics approaches when the whole enzyme is considered; or (ii) the accurate evaluation of the conformational dynamics of the enzyme, such as molecular dynamics or Monte-Carlo simulations. Each technique presents its strengths and weaknesses, but the combination of them provides an invaluable tool to shed light on the effect of the included DE mutations on the enzyme reaction mechanism or the conformational dynamics of certain active site residues. The understanding of DE rules of operation is of the utmost importance to reach the final goal of developing more robust computational protocols to predict the amino-acid changes needed for activity. This would reduce the need for experimentally probing randomized sequences, rendering the route to novel biocatalysts much more efficient. Robust computational enzyme evolution protocols based on the discussed methodologies will, in the future, need to be developed and applied if the routine design of enzymes is to be pursued.



Acknowledgements

A. R. R. thanks the Generalitat de Catalunya for PhD fellowship (2015-FI-B-00165), M. G. B. is grateful to the European Community for CIG project (PCIG14-GA-2013-630978), and Spanish MINECO for project CTQ2014-52525-P. S. O. thanks the Spanish MINECO CTQ2014-59212-P, Ramón y Cajal contract (RYC-2014-16846), the European Community for CIG project (PCIG14-GA-2013-630978), and the funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC-2015-StG-679001).

Notes and references

- M. Breuer, K. Dittrich, T. Habicher, B. Hauer, M. Keßeler, R. Stürmer and T. Zelinski, *Angew. Chem., Int. Ed.*, 2004, **43**, 788–824.
- R. J. Fox, S. C. Davis, E. C. Mundorff, L. M. Newman, V. Gavrilovic, S. K. Ma, L. M. Chung, C. Ching, S. Tam, S. Muley, J. Grate, J. Gruber, J. C. Whitman, R. A. Sheldon and G. W. Huisman, *Nat. Biotechnol.*, 2007, **25**, 338–344.
- S. Panke and M. Wubbolts, *Curr. Opin. Chem. Biol.*, 2005, **9**, 188–194.
- C. K. Savile, J. M. Janey, E. C. Mundorff, J. C. Moore, S. Tam, W. R. Jarvis, J. C. Colbeck, A. Krebber, F. J. Fleitz, J. Brands, P. N. Devine, G. W. Huisman and G. J. Hughes, *Science*, 2010, **329**, 305–309.
- S. J. Benkovic and S. Hammes-Schiffer, *Science*, 2003, **301**, 1196–1202.
- M. Garcia-Viloca, J. Gao, M. Karplus and D. G. Truhlar, *Science*, 2004, **303**, 186–195.
- S. Marti, M. Roca, J. Andres, V. Moliner, E. Silla, I. Tuñón and J. Bertran, *Chem. Soc. Rev.*, 2004, **33**, 98–107.
- Z. D. Nagel and J. P. Klinman, *Nat. Chem. Biol.*, 2009, **5**, 543–550.
- A. Warshel, P. K. Sharma, M. Kato, Y. Xiang, H. Liu and M. H. M. Olsson, *Chem. Rev.*, 2006, **106**, 3210–3235.
- P. A. Romero and F. H. Arnold, *Nat. Rev. Mol. Cell Biol.*, 2009, **10**, 866–876.
- C. Jaeckel, P. Kast and D. Hilvert, *Annu. Rev. Biophys.*, 2008, **37**, 153–173.
- H. Renata, Z. J. Wang and F. H. Arnold, *Angew. Chem., Int. Ed.*, 2015, **54**, 3351–3367.
- L. Jiang, E. A. Althoff, F. R. Clemente, L. Doyle, D. Rothlisberger, A. Zanghellini, J. L. Gallaher, J. L. Betker, F. Tanaka, C. F. Barbas, III, D. Hilvert, K. N. Houk, B. L. Stoddard and D. Baker, *Science*, 2008, **319**, 1387–1391.
- D. Rothlisberger, O. Khersonsky, A. M. Wollacott, L. Jiang, J. DeChancie, J. Betker, J. L. Gallaher, E. A. Althoff, A. Zanghellini, O. Dym, S. Albeck, K. N. Houk, D. S. Tawfik and D. Baker, *Nature*, 2008, **453**, U190–U194.
- L. Giger, S. Caner, R. Obexer, P. Kast, D. Baker, N. Ban and D. Hilvert, *Nat. Chem. Biol.*, 2013, **9**, U494–U449.
- O. Khersonsky, G. Kiss, D. Roethlisberger, O. Dym, S. Albeck, K. N. Houk, D. Baker and D. S. Tawfik, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 10358–10363.
- R. Blomberg, H. Kries, D. M. Pinkas, P. R. E. Mittl, M. G. Gruetter, H. K. Privett, S. L. Mayo and D. Hilvert, *Nature*, 2013, **503**, 418–421.
- E. A. Althoff, L. Wang, L. Jiang, L. Giger, J. K. Lassila, Z. Wang, M. Smith, S. Hari, P. Kast, D. Herschlag, D. Hilvert and D. Baker, *Protein Sci.*, 2012, **21**, 717–726.
- J. K. Lassila, D. Baker and D. Herschlag, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 4937–4942.
- R. Obexer, S. Studer, L. Giger, D. M. Pinkas, M. G. Gruetter, D. Baker and D. Hilvert, *ChemCatChem*, 2014, **6**, 1043–1050.
- X. Garrabou, T. Beck and D. Hilvert, *Angew. Chem., Int. Ed.*, 2015, **54**, 5609–5612.
- X. Garrabou, B. I. M. Wicky and D. Hilvert, *J. Am. Chem. Soc.*, 2016, **138**, 6972–6974.
- U. T. Bornscheuer, G. W. Huisman, R. J. Kazlauskas, S. Lutz, J. C. Moore and K. Robins, *Nature*, 2012, **485**, 185–194.
- J. C. Francis and P. E. Hansche, *Genetics*, 1972, **70**, 59–73.
- S. Lutz and U. T. Bornscheuer, *Protein Engineering Handbook*, Wiley-VCH Verlag GmbH & Co. KGaA, 2008.
- M. S. Packer and D. R. Liu, *Nat. Rev. Genet.*, 2015, **16**, 379–394.
- R. J. Kazlauskas and U. T. Bornscheuer, *Nat. Chem. Biol.*, 2009, **5**, 526–529.
- N. J. Turner, *Nat. Chem. Biol.*, 2009, **5**, 567–573.
- A. Currin, N. Swainston, P. J. Day and D. B. Kell, *Protein Eng., Des. Sel.*, 2014, **27**, 273–280.
- T. A. Addington, R. W. Mertz, J. B. Siegel, J. M. Thompson, A. J. Fisher, V. Filkov, N. M. Fleischman, A. A. Suen, C. Zhang and M. D. Toney, *J. Mol. Biol.*, 2013, **425**, 1378–1389.
- A. Pavelka, E. Chovancova and J. Damborsky, *Nucleic Acids Res.*, 2009, **37**, W376–W383.
- J. B. Siegel, A. Zanghellini, H. M. Lovick, G. Kiss, A. R. Lambert, J. L. Clair, J. L. Gallaher, D. Hilvert, M. H. Gelb, B. L. Stoddard, K. N. Houk, F. E. Michael and D. Baker, *Science*, 2010, **329**, 309–313.
- R. Kourist, H. Jochens, S. Bartsch, R. Kuipers, S. K. Padhi, M. Gall, D. Böttcher, H.-J. Joosten and U. T. Bornscheuer, *ChemBioChem*, 2010, **11**, 1635–1643.
- R. K. Kuipers, H.-J. Joosten, W. J. H. van Berkel, N. G. H. Leferink, E. Rooijen, E. Ittmann, F. van Zimmeren, H. Jochens, U. Bornscheuer, G. Vriend, V. A. P. Martins dos Santos and P. J. Schaap, *Proteins: Struct., Funct., Bioinf.*, 2010, **78**, 2101–2113.
- R. K. P. Kuipers, H.-J. Joosten, E. Verwiel, S. Paans, J. Akerboom, J. van der Oost, N. G. H. Leferink, W. J. H. van Berkel, G. Vriend and P. J. Schaap, *Proteins: Struct., Funct., Bioinf.*, 2009, **76**, 608–616.
- H. Xiao, Z. Bao and H. Zhao, *Ind. Eng. Chem. Res.*, 2015, **54**, 4011–4020.
- R. Obexer, A. Godina, X. Garrabou, P. R. E. Mittl, D. Baker, A. D. Griffiths and D. Hilvert, *Nat. Chem.*, 2016, DOI: 10.1038/nchem.2596.
- A. Currin, N. Swainston, P. J. Day and D. B. Kell, *Chem. Soc. Rev.*, 2015, **44**, 1172–1239.
- I. V. Pavlidis, M. S. Weiß, M. Genz, P. Spurr, S. P. Hanlon, B. Wirz, H. Iding and U. T. Bornscheuer, *Nat. Chem.*, 2016, **8**, 1076–1082.
- D. C. Wedge, W. Rowe, D. B. Kell and J. Knowles, *J. Theor. Biol.*, 2009, **257**, 131–141.
- D. Baker, *Protein Sci.*, 2010, **19**, 1817–1819.
- F. Duarte, B. A. Amrein and S. C. L. Kamerlin, *Phys. Chem. Chem. Phys.*, 2013, **15**, 11160–11177.
- J. A. Gerlt and P. C. Babbitt, *Curr. Opin. Chem. Biol.*, 2009, **13**, 10–18.
- B. Kuhlman, G. Dantas, G. C. Ireton, G. Varani, B. L. Stoddard and D. Baker, *Science*, 2003, **302**, 1364–1368.
- S. Cooper, F. Khatib, A. Treuille, J. Barbero, J. Lee, M. Beenen, A. Leaver-Fay, D. Baker, Z. Popovic and F. Players, *Nature*, 2010, **466**, 756–760.
- D. N. Bolon and S. L. Mayo, *Proc. Natl. Acad. Sci. U. S. A.*, 2001, **98**, 14274–14279.
- B. I. Dahiya and S. L. Mayo, *Protein Sci.*, 1996, **5**, 895–903.
- H. W. Hellinga and F. M. Richards, *J. Mol. Biol.*, 1991, **222**, 763–785.
- G. Kiss, N. Çelebi-Ölçüm, R. Moretti, D. Baker and K. N. Houk, *Angew. Chem., Int. Ed.*, 2013, **52**, 5700–5725, and references cited therein.
- B. I. Dahiya and S. L. Mayo, *Science*, 1997, **278**, 82–87.
- A. Zanghellini, L. Jiang, A. M. Wollacott, G. Cheng, J. Meiler, E. A. Althoff, D. Rothlisberger and D. Baker, *Protein Sci.*, 2006, **15**, 2785–2794.
- K. Świderek, I. Tuñón, V. Moliner and J. Bertran, *Arch. Biochem. Biophys.*, 2015, **582**, 68–79.
- G. R. Nosrati and K. N. Houk, *Protein Sci.*, 2012, **21**, 697–706.
- C. Malisi, O. Kohlbacher and B. Höcker, *Proteins: Struct., Funct., Bioinf.*, 2009, **77**, 74–83.
- H. Fazelinia, P. C. Cirino and C. D. Maranas, *Protein Sci.*, 2009, **18**, 180–195.
- X. Huang, K. Han and Y. Zhu, *Protein Sci.*, 2013, **22**, 929–941.
- H. K. Privett, G. Kiss, T. M. Lee, R. Blomberg, R. A. Chica, L. M. Thomas, D. Hilvert, K. N. Houk and S. L. Mayo, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 3790–3795.
- G. Kiss, D. Röthlisberger, D. Baker and K. N. Houk, *Protein Sci.*, 2010, **19**, 1760–1773.
- H. J. Wijma, R. J. Floor, S. Bjelic, S. J. Marrink, D. Baker and D. B. Janssen, *Angew. Chem.*, 2015, **127**, 3797–3801.
- C. A. Smith and T. Kortemme, *J. Mol. Biol.*, 2008, **380**, 742–756.
- P.-S. Huang, Y.-E. A. Ban, F. Richter, I. Andre, R. Vernon, W. R. Schief and D. Baker, *PLoS One*, 2011, **6**, e24109.
- X. Fu, J. R. Apgar and A. E. Keating, *J. Mol. Biol.*, 2007, **371**, 1099–1117.
- F. Ding and N. V. Dokholyan, *PLoS Comput. Biol.*, 2006, **2**, e85.
- S. M. Larson, J. L. England, J. R. Desjarlais and V. S. Pande, *Protein Sci.*, 2002, **11**, 2804–2813.



- 65 J. A. Davey and R. A. Chica, *Proteins: Struct., Funct., Bioinf.*, 2014, **82**, 771–784.
- 66 M. Sparta, D. Shirvanyants, F. Ding, N. V. Dokholyan and A. N. Alexandrova, *Biophys. J.*, 2012, **103**, 767–776.
- 67 D. J. Mandell and T. Kortemme, *Curr. Opin. Biotechnol.*, 2009, **20**, 420–428.
- 68 G. D. Friedland and T. Kortemme, *Curr. Opin. Struct. Biol.*, 2010, **20**, 377–384.
- 69 J. A. Davey and R. A. Chica, *Protein Sci.*, 2012, **21**, 1241–1252.
- 70 S. Lindert, J. Meiler and J. A. McCammon, *J. Chem. Theory Comput.*, 2013, **9**, 3843–3847.
- 71 M. J. Grisewood, N. P. Gifford, R. J. Pantazes, Y. Li, P. C. Cirino, M. J. Janik and C. D. Maranas, *PLoS One*, 2013, **8**, e75358.
- 72 C.-Y. Chen, I. Georgiev, A. C. Anderson and B. R. Donald, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 3764–3769.
- 73 I. Georgiev and B. R. Donald, *Bioinformatics*, 2007, **23**, i185–i194.
- 74 I. Georgiev, R. H. Lilien and B. R. Donald, *J. Comput. Chem.*, 2008, **29**, 1527–1542.
- 75 J.-S. Yang, S. W. Seo, S. Jang, G. Y. Jung and S. Kim, *PLoS Comput. Biol.*, 2012, **8**, e1002612.
- 76 J. Damborsky and J. Brezovsky, *Curr. Opin. Chem. Biol.*, 2014, **19**, 8–16.
- 77 C. Malisi, M. Schumann, N. C. Toussaint, J. Kageyama, O. Kohlbacher and B. Höcker, *PLoS One*, 2012, **7**, e52505.
- 78 D. Suplatov, E. Kirilin, V. Takhveev and V. Švedas, *J. Biomol. Struct. Dyn.*, 2014, **32**, 1752–1758.
- 79 E. Chovancova, A. Pavelka, P. Benes, O. Strnad, J. Brezovsky, B. Kozlikova, A. Gora, V. Sustr, M. Klvana, P. Medek, L. Biedermannova, J. Sochor and J. Damborsky, *PLoS Comput. Biol.*, 2012, **8**, e1002708.
- 80 J. D. Durrant, L. Votapka, J. Sørensen and R. E. Amaro, *J. Chem. Theory Comput.*, 2014, **10**, 5047–5056.
- 81 H. J. Wijma, R. J. Floor, P. A. Jekel, D. Baker, S. J. Marrink and D. B. Janssen, *Protein Eng., Des. Sel.*, 2014, **27**, 49–58.
- 82 H. Kries, R. Blomberg and D. Hilvert, *Curr. Opin. Chem. Biol.*, 2013, **17**, 221–228.
- 83 C. Cramer, *Essentials of computational chemistry: theories and models*, John Wiley & Sons, 2013.
- 84 K. Hotta, X. Chen, R. S. Paton, A. Minami, H. Li, K. Swaminathan, I. I. Mathews, K. Watanabe, H. Oikawa, K. N. Houk and C.-Y. Kim, *Nature*, 2012, **483**, 355–U154.
- 85 D. J. Tantillo, J. G. Chen and K. N. Houk, *Curr. Opin. Chem. Biol.*, 1998, **2**, 743–750.
- 86 E. H. Krenske, A. Patel and K. N. Houk, *J. Am. Chem. Soc.*, 2013, **135**, 17638–17642.
- 87 G. Ujaque, D. J. Tantillo, Y. F. Hu, K. N. Houk, K. Hotta and D. Hilvert, *J. Comput. Chem.*, 2003, **24**, 98–110.
- 88 Y. Zou, M. Garcia-Borràs, M. Tang, Y. Hirayama, D. Li, L. Li, K. Watanabe, K. N. Houk and Y. Tang, 2016, submitted for publication.
- 89 P. E. M. Siegbahn and F. Himo, *J. Biol. Inorg. Chem.*, 2009, **14**, 643–651.
- 90 P. E. M. Siegbahn and F. Himo, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2011, **1**, 323–336.
- 91 P. E. M. Siegbahn and R. H. Crabtree, *J. Am. Chem. Soc.*, 1997, **119**, 3103–3113.
- 92 M. E. S. Lind and F. Himo, *Angew. Chem., Int. Ed.*, 2013, **52**, 4563–4567.
- 93 M. E. S. Lind and F. Himo, *ACS Catal.*, 2014, **4**, 4153–4160.
- 94 A. Warshel and M. Levitt, *J. Mol. Biol.*, 1976, **103**, 227–249.
- 95 H. M. Senn and W. Thiel, *Angew. Chem., Int. Ed.*, 2009, **48**, 1198–1229.
- 96 M. W. van der Kamp and A. J. Mulholland, *Biochemistry*, 2013, **52**, 2708–2728.
- 97 B. Karasulu and W. Thiel, *ACS Catal.*, 2015, **5**, 1227–1239.
- 98 J. Aranda, K. Zinovjev, A. C. Swiderek, M. Roca and I. Tuñón, *ACS Catal.*, 2016, **6**, 3262–3276.
- 99 A. Ardèvol and C. Rovira, *J. Am. Chem. Soc.*, 2015, **137**, 7528–7547.
- 100 A. Warshel and R. M. Weiss, *J. Am. Chem. Soc.*, 1980, **102**, 6218–6226.
- 101 A. Shurki, E. Derat, A. Barrozo and S. C. L. Kamerlin, *Chem. Soc. Rev.*, 2015, **44**, 1037–1052.
- 102 S. C. L. Kamerlin and A. Warshel, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2011, **1**, 30–45.
- 103 B. A. Amrein, P. Bauer, F. Duarte, A. J. Carlsson, A. Naworyta, S. L. Mowbray, M. Widersten and S. C. L. Kamerlin, *ACS Catal.*, 2015, **5**, 5702–5713.
- 104 G. Oanca, M. Purg, J. Mavri, J. C. Shih and J. Stare, *Phys. Chem. Chem. Phys.*, 2016, **18**, 13346–13356.
- 105 R. O. Dror, R. M. Dirks, J. P. Grossman, H. Xu and D. E. Shaw, *Annu. Rev. Biophys.*, 2012, **41**, 429–452.
- 106 A. Laio and M. Parrinello, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 12562–12566.
- 107 D. Hamelberg, J. Mongan and J. A. McCammon, *J. Chem. Phys.*, 2004, **120**, 11919–11929.
- 108 M. Orozco, *Chem. Soc. Rev.*, 2014, **43**, 5051–5066.
- 109 G. Jiménez-Osés, S. Osuna, X. Gao, M. R. Sawaya, L. Gilson, S. J. Collier, G. W. Huisman, T. O. Yeates, Y. Tang and K. N. Houk, *Nat. Chem. Biol.*, 2014, **10**, 431–436.
- 110 T. C. Bruice, *Acc. Chem. Res.*, 2002, **35**, 139–148.
- 111 A. Madadkar-Sobhani and V. Guallar, *Nucleic Acids Res.*, 2013, **41**, W322–W328.
- 112 M. Rueda, P. Chacón and M. Orozco, *Structure*, 2007, **15**, 565–575.
- 113 E. Monza, M. F. Lucas, S. Camarero, L. C. Alejaldre, A. T. Martínez and V. Guallar, *J. Phys. Chem. Lett.*, 2015, **6**, 1447–1453.
- 114 R.-Z. Liao, S.-L. Chen and P. E. M. Siegbahn, *ACS Catal.*, 2015, **5**, 7350–7358.
- 115 X. Li, E. M. Sproviero, U. Ryde, V. S. Batista and G. Chen, *Int. J. Quantum Chem.*, 2013, **113**, 474–478.
- 116 B. Manta, F. M. Raushel and F. Himo, *J. Phys. Chem. B*, 2014, **118**, 5644–5652.
- 117 P. E. M. Siegbahn and M. R. A. Blomberg, *FEBS Lett.*, 2014, **588**, 545–548.
- 118 H. Zheng and M. T. Reetz, *J. Am. Chem. Soc.*, 2010, **132**, 15744–15751.
- 119 K. Miyamoto and H. Ohta, *Eur. J. Biochem.*, 1992, **210**, 475–481.
- 120 Y. Ijima, K. Matoishi, Y. Terao, N. Doi, H. Yanagawa and H. Ohta, *Chem. Commun.*, 2005, 877–879.
- 121 Y. Terao, Y. Ijima, K. Miyamoto and H. Ohta, *J. Mol. Catal. B: Enzym.*, 2007, **45**, 15–20.
- 122 R. Obata and M. Nakasako, *Biochemistry*, 2010, **49**, 1963–1969.
- 123 K. Okrasa, C. Levy, M. Wilding, M. Goodall, N. Baudendistel, B. Hauer, D. Leys and J. Micklefield, *Angew. Chem., Int. Ed.*, 2009, **48**, 7691–7694.
- 124 Nobel Prizes 2013, M. Karplus, M. Levitt and A. Warshel, *Angew. Chem., Int. Ed.*, 2013, **52**, 11972.
- 125 A. D. Mackerell, *J. Comput. Chem.*, 2004, **25**, 1584–1604.
- 126 J. Gao, *Rev. Comput. Chem.*, 1996, **7**, 119–185.
- 127 J. Gao, *Acc. Chem. Res.*, 1994, **29**, 298–307.
- 128 J. Gao and X. Xia, *Science*, 1992, **258**, 631–635.
- 129 S. Sumner, P. Soderhjelm and U. Ryde, *J. Chem. Theory Comput.*, 2013, **9**, 4205–4214.
- 130 R. Lonsdale and M. T. Reetz, *J. Am. Chem. Soc.*, 2015, **137**, 14733–14742.
- 131 R. P. P. Neves, P. A. Fernandes and M. J. Ramos, *ACS Catal.*, 2016, **6**, 357–368.
- 132 A. J. M. Ribeiro, L. Yang, M. J. Ramos, P. A. Fernandes, Z.-X. Liang and H. Hiraó, *ACS Catal.*, 2015, **5**, 3740–3751.
- 133 A. D. Daniels, I. Campeotto, M. W. van der Kamp, A. H. Bolt, C. H. Trinh, S. E. V. Phillips, A. R. Pearson, A. Nelson, A. J. Mulholland and A. Berry, *ACS Chem. Biol.*, 2014, **9**, 1025–1032.
- 134 G. H. Zhao, H. Song, Z. W. Chen, F. S. Mathews and M. S. Jorns, *Biochemistry*, 2002, **41**, 9751–9764.
- 135 A. Ilari, A. Bonamore, S. Franceschini, A. Fiorillo, A. Boffi and G. Colotti, *Proteins: Struct., Funct., Bioinf.*, 2008, **71**, 2065–2075.
- 136 F.-K. Shieh and N. O. Reich, *J. Mol. Biol.*, 2007, **373**, 1157–1168.
- 137 L. Mones, W.-J. Tang and J. Florian, *Biochemistry*, 2013, **52**, 2672–2682.
- 138 A. Sharir-Ivry, R. Varatharaj and A. Shurki, *Chem. – Eur. J.*, 2015, **21**, 7159–7169.
- 139 J. Aqvist and S. C. L. Kamerlin, *ACS Catal.*, 2016, **6**, 1737–1743.
- 140 L. T. Elfstrom and M. Widersten, *Biochem. J.*, 2005, **390**, 633–640.
- 141 A. Thomaus, J. Carlsson, J. Aqvist and M. Widersten, *Biochemistry*, 2007, **46**, 2466–2479.
- 142 R. M. Geha, I. Rebrin, K. Chen and J. C. Shih, *J. Biol. Chem.*, 2001, **276**, 9877–9882.
- 143 R. Car and M. Parrinello, *Phys. Rev. Lett.*, 1985, **55**, 2471–2474.
- 144 G. R. Bowman, V. S. Pande and F. Noé, *An Introduction to Markov State Models and Their Application to Long Timescale Molecular Simulation*, Springer, 2014.
- 145 S. C. Dodani, G. Kiss, J. K. B. Cahn, Y. Su, V. S. Pande and F. H. Arnold, *Nat. Chem.*, 2016, **8**, 419–425.



- 146 H. Zhang, C. Kenaan, D. Hamdane, G. H. B. Hoa and P. F. Hollenberg, *J. Biol. Chem.*, 2009, **284**, 25678–25686.
- 147 E. Weinan and E. Vanden-Eijnden, *J. Stat. Phys.*, 2006, **123**, 503–523.
- 148 X.-D. Kong, S. Yuan, L. Li, S. Chen, J.-H. Xu and J. Zhou, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 15717–15722.
- 149 V. Liskova, D. Bednar, T. Prudnikova, P. Rezacova, T. Koudelakova, E. Sebestova, I. K. Smatanova, J. Brezovsky, R. Chaloupkova and J. Damborsky, *ChemCatChem*, 2015, **7**, 648–659.
- 150 J. Ma, L. Wu, F. Guo, J. Gu, X. Tang, L. Jiang, J. Liu, J. Zhou and H. Yu, *Appl. Microbiol. Biotechnol.*, 2013, **97**, 4897–4906.
- 151 H. J. Wijma, S. J. Marrink and D. B. Janssen, *J. Chem. Inf. Model.*, 2014, **54**, 2079–2092.
- 152 P. Molina-Espeja, M. Cañellas, F. J. Plou, M. Hofrichter, F. Lucas, V. Guallar and M. Alcalde, *ChemBioChem*, 2016, **17**, 341–349.
- 153 V. Sáez-Jiménez, S. Acebes, V. Guallar, A. T. Martínez and F. J. Ruiz-Dueñas, *PLoS One*, 2015, **10**, e0124750.

