



Cite this: *Mater. Horiz.*, 2019, 6, 11

Received 6th August 2018,
Accepted 2nd October 2018

DOI: 10.1039/c8mh00931g

rsc.li/materials-horizons

Physical magnification of objects

Edward S. Boyden 

For hundreds of years, scientists have aimed photons, electrons, and X-rays at objects in order to obtain magnified images of them, revealing their micro- and nanoscale architecture, and revolutionizing every branch of the sciences. Recently a new modality of magnification was discovered – physical magnification of objects. For objects with certain properties, such as biological specimens, one can synthesize throughout them a dense, even network of swellable polymer, so that further chemical processing steps will isotropically expand the polymer, and thus the object, manyfold in physical dimension. We here explore the principles governing this new modality of magnification, such as the ability to use inexpensive, ubiquitous imaging hardware to perform high-resolution imaging of objects, and the ability to decrowd molecules within an object for detailed *in situ* chemical analysis.


Light microscopes obtain images of objects by delivering light to a specimen, and using lenses to magnify the resulting optical images of the specimen. The resolution of the microscope is limited by the wavelength of the light, meaning that features smaller than a few hundred nanometers are not resolvable. Super-resolution methods were developed that can break this so-called diffraction limit, but they are slow, expensive, and/or complex to use. Recently a new modality of magnification was discovered, namely the direct physical magnification of objects, which enables nanoscale features to be resolved with ordinary, ubiquitous light microscopes (Fig. 1).¹ For physical objects that are porous, such as biological specimens, it is possible to synthesize (Fig. 1A) throughout such objects a dense, crosslinked, evenly synthesized network of swellable polymer – for example, a hydrogel of a polyelectrolyte such as sodium polyacrylate – covalently anchoring key molecules of interest within the object to the polymer network. The spacing between the polymer threads (or mesh size) should ideally be comparable to the spacing between the molecules of interest, or at least the resolution desired. Then, one can mechanically disrupt the endogenous interactions between the molecules of interest, or even destroy molecules that are not of interest, *e.g.* through heat, enzymatic, or chemical treatment. Finally, adding water causes the hydrogel to swell, pulling the molecules of interest apart from one another *via* the anchors. After expanding an object by ~ 4 fold in linear dimension (Fig. 1B and C), a light microscope with diffraction limit 300 nm would have an effective resolution of $\sim 300/4 \sim 70$ nm – enabling nanoscopy on ordinary microscopes commonly available in science. This strategy is called

expansion microscopy (ExM).¹ Given the nanoscale spacing between polymer threads in such samples, the fundamental limit of such expansion processes might be as small as 5–10 nm.⁸ Given that the final object is $\sim 99\%$ water, having been swollen in water, it is effectively completely transparent by the end of the process (Fig. 1D), which further facilitates the power of fast, diffraction-limited methods of imaging such as light-sheet microscopy.⁷

ExM owes a debt to two lines of research that go back many decades. In the early 1980s, the phase transition physics underlying changes in volume of swellable polyelectrolyte hydrogels was worked out,² and around the same time, it was discovered by biologists that polymer hydrogel embedding of tissues (by soaking specimens in a solution of acrylamide monomer, which in turn would undergo free radical polymerization, resulting in a polyacrylamide hydrogel permeating throughout a preserved specimen) could be used to facilitate biological imaging.³ To date this strategy has been primarily applied to preserved biological cells and tissues, which are highly porous, with proteins^{4–6} or nucleic acids⁷ being anchored to the polyelectrolyte hydrogel *via* commercially available anchoring molecules. One can expand an object over and over, forming a second polymer in the space opened up by a first round of expansion, for added resolution ($\sim 4 \times 4 = 16\times$ magnification would enable $\sim 300/16 \sim 20$ nm resolution).⁸ Recently expansion has been applied to other kinds of specimens, such as crystals of proteins, which were expanded and contracted, and assessed by X-ray imaging.⁹ Recently an unrelated chemistry platform, relying on DNA origami, was used to expand multimeric protein complexes so that individual protein building blocks could be optically resolved, for structural analysis.¹⁰ This highlights that the concept of expansion is not limited only to the use of polymer hydrogels; the initial discovery that it was possible to isotropically expand a biological specimen

Media Lab, McGovern Institute, Koch Institute, Department of Biological Engineering, and Department of Brain and Cognitive Sciences, MIT, Cambridge, MA 02139, USA.
E-mail: esb@media.mit.edu





Expansion of an object can be used to improve the resolution of any microscope, in principle. For example, the application of

expansion to bacteria was used to make them detectable by inexpensive, modified webcams,¹⁹ which might make diagnosis of disease or detection of pathogens very cheap. The expansion of brain circuitry makes neural wiring easier to trace.⁵ A perhaps nonobvious advantage of expansion is that it decrowds molecules within a specimen from one another, making room for interesting chemical reactions such as amplification, hybridization, sequencing, and other analytical chemistries. Already it has been shown that expansion of a cell, by decrowding messenger RNA (mRNA) molecules from each other, could facilitate their more accurate counting, and furthermore made room for amplification of the fluorescence through the fluorescence-amplifying hybridization chain reaction (HCR).⁷ This decrowding also greatly facilitates coded serial hybridization methods like MERFISH, which can read out an exponential amount of information in a linear number of steps, by applying carefully designed hybridization probes in multiple rounds.²⁰ The ability to decrowd proteins from each other, if achievable in a controlled way, may enable epitopes that cannot be detected by antibody binding, to be tagged because the decrowding makes room around each protein for antibody approach. In summary, the recent discovery that specimens such as biological samples can be physically magnified by a material science process is opening up frontiers in the nanoscale mapping of extended objects, in the inexpensive probing of molecular complexes and other nanoscale features, and in the multiplexed analysis of biomolecules through decrowding-enhanced *in situ* chemistry. In the future, greater degrees of physical magnification and

higher isotropy through new polymeric chemistries, as well as novel and powerful kinds of *in situ* analysis chemistry, may make the expansion microscopy toolset an even more powerful toolbox for understanding the configuration of molecules throughout cells and cells throughout tissues, and perhaps reveal the molecular configuration of other kinds of specimen as well.

Conflicts of interest

E. S. B. is an inventor on multiple patents related to expansion microscopy, and also has co-founded a company to pursue clinical and translational applications of expansion microscopy.

References

- 1 F. Chen, P. W. Tillberg and E. S. Boyden, Expansion Microscopy, *Science*, 2015, **347**, 543–548.
- 2 T. Tanaka, D. Fillmore, S. T. Sun, I. Nishio, G. Swislow and A. Shah, Phase Transitions in Ionic Gels, *Phys. Rev. Lett.*, 1980, **45**, 1636–1639.
- 3 P. Hausen and C. Dreyer, The Use of Polyacrylamide as an Embedding Medium for Immunohistochemical Studies of Embryonic Tissues, *Stain Technol.*, 1981, **56**, 287–293.
- 4 T. Ku, J. Swaney, J.-Y. Park, A. Albanese, E. Murray, J. H. Cho, Y.-G. Park, V. Mangena, J. Chen and K. Chung, Multiplexed and Scalable Super-Resolution Imaging of Three-Dimensional Protein Localization in Size-Adjustable Tissues, *Nat. Biotechnol.*, 2016, **34**, 973–981.
- 5 P. W. Tillberg, F. Chen, K. D. Piatkevich, Y. Zhao, C.-C. Yu (Jay), B. P. English, L. Gao, A. Martorell, H.-J. Suk, F. Yoshida, M. Ellen, D. H. R. DeGennaro, G. Gong, U. Seneviratne, S. R. Tannenbaum, R. Desimone, D. Cai and E. S. Boyden, Protein-Retention Expansion Microscopy of Cells and Tissues Labeled Using Standard Fluorescent Proteins and Antibodies, *Nat. Biotechnol.*, 2016, **34**, 987–992.
- 6 T. J. Chozinski, A. R. Halpern, H. Okawa, H.-J. Kim, G. J. Tremel, R. O. L. Wong and J. C. Vaughan, Expansion Microscopy with Conventional Antibodies and Fluorescent Proteins, *Nat. Methods*, 2016, **13**, 1–7.
- 7 F. Chen, A. T. Wassie, A. J. Cote, A. Sinha, S. Alon, E. R. D. Shoh Asano, J.-B. Chang, A. Marblestone, G. M. Church, A. Raj and E. S. Boyden, Nanoscale Imaging of RNA with Expansion Microscopy, *Nat. Methods*, 2016, **13**, 679–684.
- 8 J.-B. Chang, F. Chen, Y.-G. Yoon, E. E. Jung, H. Babcock, J. S. Kang, S. Asano, H.-J. Suk, N. Pak, P. W. Tillberg, A. T. Wassie, D. Cai and E. S. Boyden, Iterative Expansion Microscopy, *Nat. Methods*, 2017, **14**, 593–599.
- 9 L. Zhang, J. B. Bailey, R. H. Subramanian and F. A. Tezcan, Hyperexpandable, Self-Healing Macromolecular Crystals with Integrated Polymer Networks, *Nature*, 2018, **557**, 86–91.
- 10 J. Molle, L. Jakob, J. Bohlen, M. Raab and P. Tinnefeld, Dina Grohmann Towards Structural Biology with Super-Resolution Microscopy, *Nanoscale*, 2018, **10**, 16416–16424.
- 11 T. C. Murakami, T. Mano, S. Saikawa, S. A. Horiguchi, D. Shigeta, K. Baba, H. Sekiya, Y. Shimizu, K. F. Tanaka, H. Kiyonari, M. Iino, H. Mochizuki, K. Tainaka and H. R. Ueda, A Three-Dimensional Single-Cell-Resolution Whole-Brain Atlas Using CUBIC-X Expansion Microscopy and Tissue Clearing, *Nat. Neurosci.*, 2018, **21**, 625–637.
- 12 L. Freifeld, I. Odstrcil, D. Förster, A. Ramirez, J. A. Gagnon, O. Randlett, E. K. Costa, S. Asano, O. T. Celiker, R. Gao, D. A. Martin-Alarcon, P. Reginato, C. Dick, L. Chen, D. Schoppik, F. Engert and E. S. Boyden, Expansion Microscopy of Zebrafish for Neuroscience and Developmental Biology Studies, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, E10799–E10808.
- 13 Y. Zhao, O. Bucur, H. Irshad, F. Chen, A. Weins, A. L. Stancu, E.-Y. Oh, M. DiStasio, V. Torous, B. Glass, I. E. Stillman, S. J. Schnitt, A. H. Beck and E. S. Boyden, Nanoscale Imaging of Clinical Specimens Using Pathology-Optimized Expansion Microscopy, *Nat. Biotechnol.*, 2017, **35**, 757–764.
- 14 ExpansionMicroscopy.Org: Physical Specimen Expansion Enabling 3-D Large Volume, Nanoscale Imaging <http://expansionmicroscopy.org/>.
- 15 T. J. Mosca, D. J. Luginbuhl, I. E. Wang and L. Luo, Presynaptic LRP4 Promotes Synapse Number and Function of Excitatory CNS Neurons, *eLife*, 2017, **6**, e27347.
- 16 T. Deshpande, T. Li, M. K. Herde, A. Becker, H. Vatter, M. K. Schwarz, C. Henneberger, C. Steinhäuser and P. Bedner, Subcellular Reorganization and Altered Phosphorylation of the Astrocytic Gap Junction Protein Connexin43 in Human and Experimental Temporal Lobe Epilepsy, *Glia*, 2017, **65**, 1809–1820.
- 17 I. E. Wang, S. W. Lapan, M. L. Scimone, T. R. Clandinin and P. W. Reddien, Hedgehog Signaling Regulates Gene Expression in Planarian Glia, *eLife*, 2016, **5**, e16996.
- 18 A. R. Halpern, G. C. M. Alas, T. J. Chozinski, A. R. Paredez and J. C. Vaughan, Hybrid Structured Illumination Expansion Microscopy Reveals Microbial Cytoskeleton Organization, *ACS Nano*, 2017, **11**, 12677–12686.
- 19 Y. S. Zhang, J.-B. Chang, M. M. Alvarez, G. Trujillo-de Santiago, J. Aleman, B. Batzaya, V. Krishnadoss, A. A. Ramanujam, M. Kazemzadeh-Narbat, F. Chen, P. W. Tillberg, M. R. Dokmeci, E. S. Boyden and A. Khademhosseini, Hybrid Microscopy: Enabling Inexpensive High-Performance Imaging through Combined Physical and Optical Magnifications, *Sci. Rep.*, 2016, **6**, 22691.
- 20 G. Wang, J. R. Moffitt and X. Zhuang, Multiplexed Imaging of High-Density Libraries of RNAs with MERFISH and Expansion Microscopy, *Sci. Rep.*, 2018, **8**, 4847.
- 21 S. Alon, G. Huynh and E. S. Boyden, Expansion Microscopy: Enabling Single Cell Analysis in Intact Biological Systems, *FEBS J.*, 2018, DOI: 10.1111/febs.14597.

