


Nanoscale

rsc.li/nanoscale



ISSN 2040-3372

PAPER

Zvi Hayouka, Ling Peng *et al.*
Dynamic self-assembling supramolecular dendrimer
nanosystems as potent antibacterial candidates against
drug-resistant bacteria and biofilms


Cite this: *Nanoscale*, 2022, **14**, 9286

Dynamic self-assembling supramolecular dendrimer nanosystems as potent antibacterial candidates against drug-resistant bacteria and biofilms†

Dinesh Dhumal,^a Bar Maron,^b Einav Malach,^b Zhenbin Lyu,^a Ling Ding,^a Domenico Marson,^c Erik Laurini,^c Aura Tintaru,^a Brigino Ralahy,^a Suzanne Giorgio,^a Sabrina Pricl,^{c,d} Zvi Hayouka^{*b} and Ling Peng^{*a}

The alarming and prevailing antibiotic resistance crisis urgently calls for innovative “outside of the box” antibacterial agents, which can differ substantially from conventional antibiotics. In this context, we have established antibacterial candidates based on dynamic supramolecular dendrimer nanosystems self-assembled with amphiphilic dendrimers composed of a long hydrophobic alkyl chain and a small hydrophilic poly(amidoamine) dendron bearing distinct terminal functionalities. Remarkably, the amphiphilic dendrimer with amine terminals exhibited strong antibacterial activity against both Gram-positive and Gram-negative as well as drug-resistant bacteria, and prevented biofilm formation. Multidisciplinary studies combining experimental approaches and computer modelling together demonstrate that the dendrimer interacts and binds *via* electrostatic interactions with the bacterial membrane, where it becomes enriched and then dynamically self-assembles into supramolecular nanoassemblies for stronger and multivalent interactions. These, in turn, rapidly promote the insertion of the hydrophobic dendrimer tail into the bacterial membrane thereby inducing bacterial cell lysis and constituting powerful antibacterial activity. Our study presents a novel concept for creating nanotechnology-based antibacterial candidates *via* dynamic self-assembly and offers a new perspective for combatting recalcitrant bacterial infection.

Received 27th April 2022,
Accepted 14th May 2022

DOI: 10.1039/d2nr02305a

rscl.li/nanoscale

Introduction

Bacterial resistance to antibiotics has become a serious global threat with ever increasing prevalence of infections difficult to treat.^{1–3} This pressing public health crisis has been driving the development of new antibacterial agents to overcome drug resistance. Approaches for developing antibacterial agents include, to name but a few, the modification of existing antibiotics, the identification of active agents on novel targets

against resistant bacteria, and the elaboration of antimicrobial peptides with dual antibacterial and immunomodulatory activities.^{4–7} Among these new strategies, the development of new antibacterial agents which are substantially different from conventional antibiotics is of particular interest. In this context, amphiphilic antibacterial agents that mimic natural antimicrobial peptides and amphiphilic antibacterial detergents constitute appealing candidates.^{8–11} Indeed, such agents are expected to harness the antibacterial features of both antimicrobial peptides and amphiphilic antibacterial detergents, while possessing the self-assembly capacity to form supramolecular nanostructures that further enforce the antibacterial activity *via* cooperative and multivalent interaction.^{7,12} Consequently, a myriad of amphiphilic molecules has been established as antibacterial candidates.^{13–15}

In an effort to meet the urgent need for new antibacterial agents, we steered our continuing efforts to develop amphiphilic dendrimers for use in biomedical applications¹⁶ as antibacterial candidates. Dendrimers are a unique family of synthetic molecules with precisely controlled radial architecture and special multivalent cooperativity confined within a small three-dimensional volume. Different dendrimers have been

^aAix Marseille Univ, CNRS, Centre Interdisciplinaire de Nanoscience de Marseille (CINaM), UMR 7325, Equipe Labelisée Ligue Contre le Cancer, Parc Scientifique et Technologique de Luminy, Marseille, France. E-mail: ling.peng@univ-amu.fr

^bInstitute of Biochemistry, Food Science and Nutrition, The Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot, Israel. E-mail: zvi.hayouka@mail.huji.ac.il

^cMolecular Biology and Nanotechnology Laboratory (MolBNL@UniTS), DEA, University of Trieste, Trieste, Italy

^dDepartment of General Biophysics, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, Poland

†Electronic supplementary information (ESI) available: Fig. S1–S6 as well as all detailed experimental methods and protocols. See DOI: <https://doi.org/10.1039/d2nr02305a>



studied for their antimicrobial activity.^{17,18} Among them, the poly(amidoamine) (PAMAM) dendrimers are of special interest thanks to their excellent biocompatibility, owing to their peptide mimicry, and resistance to enzyme degradation by virtue of their dendritic structure. In addition, they are readily available *via* robust synthesis and commercial sources.^{19,20} Therefore, PAMAM dendrimers are considered as the ideal candidates to be elaborated as mimics of antimicrobial peptides. However, the most active PAMAM dendrimers are often high-generation molecules bearing numerous charged terminals that are unfortunately associated with considerable cytotoxicity at their MIC concentrations.^{21,22}

Recently, small amphiphilic dendrimers composed of hydrophobic entities and hydrophilic dendrons were shown to have antibacterial activity that was frequently related to low dendrimer generation, terminal charge, and hydrophilicity/

hydrophobicity balance.^{23–32} Specifically, various peptide, polylysine and poly(amide) dendrimers were synthesized to improve their antimicrobial activity,^{23–27} whereas polyester dendrimers were explored for imparting biodegradability while reducing toxicity.^{30–32} Also, amphiphilic dendrimers with carbosilane entities²⁸ and poly(aryl ether) backbones²⁹ were studied for antibacterial activity.

Here, we report the design, synthesis and evaluation of amphiphilic dendrimers composed of a long hydrophobic alkyl chain and a small hydrophilic PAMAM dendron carrying one of the following charged terminals: primary amine, tertiary amine, guanidine or carboxylate moieties for antibacterial activity (Fig. 1A). Importantly, we show that not only does the charge, the charge density and steric size associated with the terminal functionalities play crucial roles in the antibacterial activity, but the dynamic self-assembling feature of the amphi-




Fig. 1 Amphiphilic dendrimers studied for antibacterial activity in this work. (A) Chemical structures of amphiphilic PAMAM dendrimers **1a–d** bearing different terminal groups; (B) Cartoon illustration of the antibacterial activity shown by these amphiphilic dendrimers *via* membrane adsorption, self-assembling, interaction, insertion, disintegration and disruption.




dendrimers self-assembled into small and spherical supramolecular nanomicelles ranging from 10 to 20 nm in size, as demonstrated by transmission electron microscopy (TEM) (Fig. 2, Fig. S1†). These data are in line with our previous results obtained for the amphiphilic dendrimers, which in water spontaneously form nanomicelles.^{34,36} In addition, NMR diffusion experiments (DOSY) were performed to estimate the average weights of the formed micelles (Fig. S2†).³⁷ DOSY results confirmed the formation of homogeneous assemblies, which were composed of six molecules of **1a** per micelle. Importantly, the nanomicelles formed by the dendrimers **1a–d** had zeta potentials of +35 mV, +23 mV, +40 mV and –13 mV, respectively, the differences relating to the distinct chemical entities at their terminals. The absolute values of the zeta potentials were all over 10 mV, highlighting the colloidal stability of these nanomicelles.³⁸

To further assess the self-assembly of the amphiphilic dendrimers **1a–d**, we determined their critical micelle concentration (CMC) using a fluorescence spectroscopic method based on the fluorescent dye pyrene.³⁹ CMC is defined as the minimum concentration of an amphiphile above which micelles can form readily. **1d** showed the lowest CMC value of 4.0 μM , whereas **1a** and **1b** had similar CMC values of 15 and 17 μM , and **1c** had the highest CMC value of 35 μM (Table 1). The different CMC values for **1a–1d** can reasonably be ascribed to their structural features: **1c** has the largest hydrophilic entity and the greatest steric hindrance at its terminals which limit its packing and assembling ability and explain the highest CMC value; whereas dendrimer **1d** has the smallest hydrophilic entity, hence the highest packing ability with the lowest CMC value.

To evaluate the safety of these dendrimers, we examined their cytotoxicity on fibroblast (L929) and kidney (HEK293) cell lines using the PrestoBlue assay. No notable adverse effect was observed for dendrimers **1a**, **1b** and **1d** at concentrations up to 200 μ M, whereas dendrimer **1c** showed considerable toxicity

We first synthesized the amphiphilic dendrimers **1a–d** using the ester-terminating dendrimer **1** as the common starting material (Scheme 1). The amidation of **1** with respectively ethylenediamine and *N,N*-dimethylethylenediamine, yielded **1a** and **1b**.^{33,34} Further conjugation of **1a** with the protected arginine derivative followed by subsequent deprotection gave the arginine-terminated dendrimer **1c**,³⁵ whereas hydrolysis of **1** using LiOH furnished the acid-terminated dendrimer **1d**. The structural integrity and purity of all the synthesized dendrimers were examined and confirmed using ¹H- and ¹³C-NMR as well as high-resolution mass spectroscopic analysis (Fig. SA1–SA4, see ESI† Annex for details).

All the dendrimers **1a–d** were soluble in water, with concentrations up to 20 mM. By virtue of their amphiphilicity, these



Scheme 1 Synthesis of dendrimers **1a–d** starting with the common dendrimer **1**.




Fig. 2 Transmission electron microscopy (TEM) images of the supramolecular nanomicelles formed by dendrimers **1a–d**.

Table 1 Self-assembly, toxicity and hemolytic activity of the amphiphilic dendrimers **1a–d**

Dendrimer	CMC ^a (μM)	Particle size ^b (nm)	Zeta potential (mV)	Cytotoxicity IC ₅₀ ^c (μM)		Hemolysis IC ₅₀ ^f (μM) RBC ^g
				L929 ^d	HEK 293 ^e	
1a	15	14 ± 2	+35	>200	196	100
1b	17	17 ± 4	+23	>200	>200	100
1c	35	15 ± 3	+40	177	67	50
1d	4	10 ± 2	−13	>200	>200	>250

^a CMC: critical micelle concentration. ^b Particle size: average values calculated with 300 nanoparticles measured in the TEM images. ^c Cytotoxicity IC₅₀: concentration of a dendrimer required for 50% inhibition of cell growth. ^d L929: fibroblast cells. ^e HEK293: human embryonic kidney cell. ^f Hemolysis IC₅₀: concentration required for lysis of 50% red blood cells. ^g RBC: red blood cells.

towards both cell types (Table 1 and Fig. S3†). We also examined the hemolytic activity of these dendrimers on mouse red blood cells (RBCs). Similar to the trend observed for cytotoxicity, **1c** had the highest hemolytic activity with a IC₅₀ value of 50 μM compared to 100 μM for **1a** and **1b**; **1d** showed no notable hemolysis, even at 250 μM (Table 1 and Fig. S4†). The observed toxicity of **1c** may be ascribed to the arginine terminals, which interact and interfere strongly with the eukaryotic cell membrane *via* both electrostatic interaction and bivalent hydrogen bonds. Conversely, dendrimers **1a**, **1b** and **1d** showed low cytotoxicity and good biocompatibility.

We then assessed the antibacterial activity of the dendrimers **1a–1d** against the Gram-negative bacteria *Escherichia coli* (*E. coli*, K12) and *Pseudomonas aeruginosa* (*P. aeruginosa*, PAO1) as well as the Gram-positive bacteria *Staphylococcus aureus* (*S. aureus*, JLA512) and methicillin-resistant *Staphylococcus aureus* (MRSA, 1206). MRSA was isolated from a patient and is resistant to methicillin (MRSA), whereas PAO1 is an aerobe, meso-

philic human pathogen that was isolated from infected wounds. The minimum inhibitory concentrations (MIC), *i.e.* the lowest concentrations required to inhibit 90% of bacteria, of these dendrimers against all the tested bacteria were determined using broth microdilution assay (Table 2). Remarkably, the amine-terminated dendrimer **1a** exhibited strong antibacterial activity against both Gram-negative and Gram-positive bacteria, as well as the drug-resistant strain MRSA, with MIC values of 3.1 μM for all strains. Interestingly, the tertiary amine-terminated dendrimer **1b** showed similar levels of antibacterial activity towards *E. coli* as **1a**, but differed from **1a** in that it was ineffective against all other tested bacteria; this would imply a possible narrow-spectrum antibacterial activity specific to certain bacteria. To our great surprise, the arginine-terminated dendrimer **1c**, which showed the highest toxicity in eukaryotic cells, was inactive against all tested bacteria strains. The carboxylic acid-terminated dendrimer **1d** was also devoid of notable antibacterial activity.

Table 2 Antibacterial activity of the amphiphilic dendrimers **1a–d**

Dendrimer	Minimum inhibition concentration (MIC)							
	<i>Escherichia coli</i>		<i>Pseudomonas aeruginosa</i>		<i>Staphylococcus aureus</i>		Methicillin resistant <i>Staphylococcus aureus</i>	
	μM	μg mL ^{−1}	μM	μg mL ^{−1}	μM	μg mL ^{−1}	μM	μg mL ^{−1}
1a	3.1	6.0	3.1	6.0	3.1	6.0	3.1	6.0
1b	2.8	6.0	30	65	29	63	19	42
1c	63	200	31	100	63	200	>63	>200
1d	62	100	>124	>200	62	100	47	75

MIC (minimum inhibitory concentration): lowest concentration required to inhibit 90% of bacterial growth, and value given in both μM and μg mL^{−1}.



Noteworthy is the finding that the different antibacterial activities shown by the dendrimers could be ascribed to them having different terminal groups and thereby distinct interactions with bacteria. The inactivity shown by **1d** can be readily explained by the repulsive forces in play between the negatively charged carboxylate terminals of the dendrimer and the negatively charged bacterial membrane. Although the dendrimers **1a**, **1b** and **1c** all have positively charged terminals at neutral pH under physiological conditions, they showed different levels of antibacterial activity. The primary amine groups in **1a**, by virtue of their small size and highly positive charge, could interact strongly with the bacterial membrane surface hence explaining the strong antibacterial activity.

However, the arginine residues in **1c** with the delocalized positive charges on the guanidine scaffolds, offered insufficient interaction with the bacterial cell membrane to allow any antibacterial activity. Similar observation was also reported where replacing ammonium functionalities by guanidiniums abolished antibacterial effect.³² Interestingly, dendrimer **1b** bearing tertiary amine terminals, larger than the primary amine thus offering a lower charge density yet smaller than the guanidine moiety and hence providing higher charge density, generated rather variable activity toward different bacteria. This finding may highlight an adaptability of dendrimer **1b** to be further exploited for developing selective and narrow-spectrum antibacterial agents.




Fig. 3 Studies on the live and dead bacterial cells and their membrane integrity and morphology upon treatment. Fluorescent microscopic imaging of live and dead cells of (A) *E. coli* and (B) *S. aureus* upon treatment with **1a**. Live and dead cells were stained using SYTO9 (green) and propidium iodide (PI) (red), respectively. Scanning electron microscopic (SEM) images of (C) *E. coli* (upper panel) and MRSA (down panel), arrow shows altered bacterial membrane surface upon treatment with **1a**.




Encouraged by the excellent antibacterial activity and safety profile of **1a**, we further evaluated its activity against the biofilm formed by *S. aureus*.^{40,41} Remarkably, **1a** retained its antibacterial activity with a MIC value maintained at 3.1 μM against the *S. aureus* biofilm (Fig. S5†), ultimately confirming the potential of **1a** as a promising candidate in antibacterial treatment against drug-resistant bacteria and bacteria within biofilm matrix.

We next examined the ability of the most potent dendrimer **1a** to disrupt the bacterial membrane and kill bacteria, using SYTO9 and propidium iodide (PI) dyes for staining the live and dead cells, respectively. SYTO9 is a universal dye that crosses intact membranes and stains nucleic acids of all live cells in green, whereas PI can only cross compromised bacterial membranes, and emits red fluorescence when binding to nucleic acids. The PI staining is therefore an indicator of membrane disintegration and cell death. Results from SYTO9 and PI staining (Fig. 3A/B) showed that both Gram-negative *E. coli* and Gram-positive *S. aureus* were damaged after treatment with **1a**, highlighting the potent antibacterial activity of **1a**, may kill bacterial cells by a mechanism most likely involving membrane damage and disruption.

To inspect and visualize the bacterial membrane integrity and morphology upon treatment with **1a**, we performed scanning electron microscopic (SEM) studies. As shown in Fig. 3C and Fig. S6,† treatment with **1a** induced obvious irregularities at the cell surface of both *E. coli* and *S. aureus* cells, with the appearance of blebbing and surface deformations indicating loss of membrane integrity. Also, drastic cell lysis was observed with triggered release of cellular materials. Untreated cells used as a control showed normal and smooth cell surface under SEM. These results highlight that bacterial membrane damaged by **1a** may indeed contribute to its potent antibacterial activity.

To confirm that the antibacterial mechanism of action used by **1a** involved interaction with bacterial membrane, we further assessed the permeability and depolarization of the bacterial outer (OM) and inner membranes (IM). Membrane per-



Minimum Inhibition Concentration (MIC)		
	μM	$\mu\text{g/mL}$
<i>E. coli</i> ^a	>300	>496
<i>P. aeruginosa</i> ^b	>300	>496
<i>S. aureus</i> ^c	>300	>496
MRSA ^d	>300	>496

Fig. 5 Hydrophilic dendrimer **2** without the hydrophobic chain is devoid of any antibacterial activity. MIC (minimum inhibitory concentration) values of **2** to inhibit 90% of bacterial growth such as ^a*Escherichia coli*, ^b*Pseudomonas aeruginosa*, ^c*Staphylococcus aureus*, ^dMethicillin-resistant *Staphylococcus aureus* and values are given in both μM and $\mu\text{g mL}^{-1}$.

meability was measured by monitoring the change in the fluorescent properties of the *N*-phenyl-1-naphthylamine (NPN) dye, a probe which displays increased fluorescence upon binding to hydrophobic membrane regions, whereas membrane depolarization was studied by recording the rapid increase in the fluorescence intensity of the 3,3'-dipropylthiadicarbocyanine iodide (diSC3(5)), a lipophilic potentiometric probe. As shown in Fig. 4, treatment with **1a** led to significant outer membrane




Fig. 4 Assessing the permeability and depolarization of the bacterial outer (OM) and inner membranes (IM). Permeability of the bacterial outer membrane (OM) in *E. coli* (A) as evaluated using the *N*-phenyl-1-naphthylamine (NPN) dye and depolarization of the inner membrane (IM) in *E. coli* (B) and MRSA (C) as evaluated using the 3,3'-dipropylthiadicarbocyanine iodide (diSC3(5)) probe, after treatment with **1a** at the indicated concentrations.






Fig. 6 Simulating the interaction of the most active dendrimer **1a** (A) and the least active dendrimer **1d** (B) with bacterial membrane using molecular dynamics (MD) simulations. (A) Initial (A1) and final (A2) MD trajectory snapshots showing that **1a** can form small aggregates when interacting with the bacterial membrane at the end of the simulation (1 μ s); and initial (A3) and final (A4) MD trajectory snapshots showing that the aggregates of **1a** on the bacterial membrane are able to disaggregate, laterally spread across the upper bilayer leaflet, and fully insert their C_{18} tails within the Gram-negative bacterial-like membrane model at the end of the simulation time (2 μ s). (B) Representative snapshots extracted at the beginning (B1) and after 1 μ s (B2), 2 μ s (B3), and 3 μ s (B4) of MD trajectory showing that **1d** form a stable micelle interacting with the bacterial membrane. In the simulated time, **1d** is not able to disaggregate and induce damage to the Gram-negative bacterial-like membrane model. The charged terminal groups of the dendrimers **1a** and **1d** are represented in blue and red, respectively. The dendritic portions and C_{18} hydrocarbon chains of each amphiphile are represented in light blue and grey, respectively. The lipid portion in the membrane model is shown as light grey sticks, whereas the headgroups are highlighted as light green (POPE) and light orange (POPG) spheres, respectively. Hydrogen atoms, water molecules, ions and counterions have been omitted for clarity.



LP coordinated the project. DD, ZL and BR synthesized the agents; DD, ZL, LD, AT and SG characterized the agents; BM and EM carried out all experiments with bacteria; LD performed the toxicity studies; DM, EL, SP performed molecular

simulation experiments; DD, ZL, SP, ZH and LP analyzed data; and LP wrote the paper with contributions from DD, ZH and SP. All authors proofed the manuscript.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

This work was supported by the French National Research Agency under the framework of the ERA-NET EURONANOMED European Research projects 'TARBRAINFECTION' (LP) and 'antineuropatho' (LP), the Chief Scientist office of the Israeli Ministry of Health grant number 3-17903 in the framework of EuroNanoMed 3 European Research project 'antineuropatho' (ZH), EU H2020 Research and Innovation program NMBP "SAFE-N-MEDTECH" (2019-2023) (grant agreement No. 814607, LP, BR), AMIDEX d'Aix-Marseille Université (LP), the Ligue Nationale Contre le Cancer (LP, ZL) and China Scholarship Council (LD). This article/publication is based upon work from COST Action CA 17140 "Cancer Nanomedicine from the Bench to the Bedside" supported by COST (European Cooperation in Science and Technology).

References

- 1 S. M. Schrader, J. Vaubourgeix and C. Nathan, Biology of antimicrobial resistance and approaches to combat it, *Sci. Transl. Med.*, 2020, 12(549), eaaz6992, DOI: [10.1126/scitranslmed.aaz6992](#).
- 2 J. M. A. Blair, M. A. Webber, A. J. Baylay, D. O. Ogbolu and L. J. V. Piddock, Molecular mechanisms of antibiotic resistance, *Nat. Rev. Microbiol.*, 2015, 13, 42–51, DOI: [10.1038/nrmicro3380](#).
- 3 A. H. Holmes, L. S. P. Moore, A. Sundsfjord, M. Steinbakk, S. Regmi, A. Karkey, *et al.*, Understanding the mechanisms and drivers of antimicrobial resistance, *Lancet*, 2016, 387, 176–187, DOI: [10.1016/S0140-6736\(15\)00473-0](#).
- 4 Z. C. Wu and D. L. Boger, Maxamycins: Durable Antibiotics Derived by Rational Redesign of Vancomycin, *Acc. Chem. Res.*, 2020, 53(11), 2587–2599, DOI: [10.1021/acs.accounts.0c00569](#).
- 5 Z. D. Aron, A. Mehrani, E. D. Hoffer, K. L. Connolly, P. Srinivas, M. C. Torhan, J. N. Alumasa, M. Cabrera, D. Hosangadi, J. S. Barbor, S. C. Cardinale, S. M. Kwasny, L. R. Morin, M. M. Butler, T. J. Opperman, T. L. Bowlin, A. Jerse, S. M. Stagg, C. M. Dunham and K. C. Keiler, Translation inhibitors bind to a novel site on the ribosome and clear *Neisseria gonorrhoeae* in vivo, *Nat. Commun.*, 2021, 12(1), 1799, DOI: [10.1038/s41467-021-22012-7](#).
- 6 Y. Jiang, Y. Chen, Z. Song, Z. Tan and J. Cheng, Recent advances in design of antimicrobial peptides and polypeptides toward clinical translation, *Adv. Drug Delivery Rev.*, 2021, 170, 261–280, DOI: [10.1016/j.addr.2020.12.016](#).
- 7 J. M. V. Makabenta, A. Nabawy, C. H. Li, S. Schmidt-Malan, R. Patel and V. M. Rotello, Nanomaterial-based therapeutics for antibiotic-resistant bacterial infections, *Nat. Rev. Microbiol.*, 2021, 19, 23–36.
- 8 N. Mookherjee, M. A. Anderson, H. P. Haagsman and D. J. Davidson, Antimicrobial host defence peptides: functions and clinical potential, *Nat. Rev. Drug Discovery*, 2020, 19(5), 311–332, DOI: [10.1038/s41573-019-0058-8](#).
- 9 B. Findlay, G. G. Zhanel and F. Schweizer, Cationic amphiphiles, a new generation of antimicrobials inspired by the natural antimicrobial peptide scaffold, *Antimicrob. Agents Chemother.*, 2010, 54(10), 4049–4058, DOI: [10.1128/AAC.00530-10](#).
- 10 M. A. Rahman, M. Bam, E. Luat, M. S. Jui, M. S. Ganewatta, T. Shokfai, M. Nagarkatti, A. W. Decho and C. Tang, Macromolecular-clustered facial amphiphilic antimicrobials, *Nat. Commun.*, 2018, 9(1), 5231, DOI: [10.1038/s41467-018-07651-7](#).
- 11 A. G. Elliott, J. X. Huang, S. Neve, J. Zuegg, I. A. Edwards, A. K. Cain, C. J. Boinett, L. Barquist, C. V. Lundberg, J. Steen, M. S. Butler, M. Mobli, K. M. Porter, M. A. T. Blaskovich, S. Lociuero, M. Strandh and M. A. Cooper, An amphipathic peptide with antibiotic activity against multidrug-resistant Gram-negative bacteria, *Nat. Commun.*, 2020, 11(1), 3184, DOI: [10.1038/s41467-020-16950-x](#).
- 12 A. R. Kirtane, M. Verma, P. Karandikar, J. Furin, R. Langer and G. Traverso, Nanotechnology approaches for global infectious diseases, *Nat. Nanotechnol.*, 2021, 16, 369–384, DOI: [10.1038/s41565-021-00866-8](#).
- 13 A. Levin, T. A. Hakala, L. Schnaider, G. J. Bernardes, E. Gazit and T. P. Knowles, Biomimetic peptide self-assembly for functional materials, *Nat. Rev. Chem.*, 2020, 4(11), 615–634, DOI: [10.1038/s41570-020-0215-y](#).
- 14 L. Schnaider, S. Brahmachari, N. W. Schmidt, B. Mensa, S. Shaham-Niv, D. Bychenko, L. Adler-Abramovich, L. J. Shimon, S. Kolusheva, W. F. DeGrado and E. Gazit, Self-assembling dipeptide antibacterial nanostructures with membrane disrupting activity, *Nat. Commun.*, 2017, 8, 1365, DOI: [10.1038/s41467-017-01447-x](#).
- 15 L. Liu, K. Xu, H. Wang, P. K. Tan, W. Fan, S. S. Venkatraman, L. Li and Y. Y. Yang, Self-assembled cationic peptide nanoparticles as an efficient antimicrobial agent, *Nat. Nanotechnol.*, 2009, 4(7), 457–463, DOI: [10.1038/nnano.2009.153](#).
- 16 Z. Lyu, L. Ding, A. Tintaru and L. Peng, Self-Assembling Supramolecular Dendrimers for Biomedical Applications: Lessons Learned from Poly(amidoamine) Dendrimers, *Acc. Chem. Res.*, 2020, 53(12), 2936–2949, DOI: [10.1021/acs.accounts.0c00589](#).
- 17 Y. Liu, L. Shi, L. Su, H. C. van der Mei, P. C. Jutte, Y. Ren and H. J. Busscher, Nanotechnology-based antimicrobials and delivery systems for biofilm-infection control, *Chem. Soc. Rev.*, 2019, 48(2), 428–446, DOI: [10.1039/c7cs00807d](#).



- 18 M. A. Mintzer, E. L. Dane, G. A. O'Toole and M. W. Grinstaff, Exploiting dendrimer multivalency to combat emerging and re-emerging infectious diseases, *Mol. Pharmaceutics*, 2012, **9**(3), 342–354.
- 19 Z. Lyu, L. Ding, D. Dhumal, A. Y. Huang, C. L. Kao and L. Peng, Poly (amidoamine)(PAMAM) dendrimers: Synthesis and biological applications, *Dendrimer Chem.*, 2020, 85–113, DOI: [10.1039/9781788012904-00085](https://doi.org/10.1039/9781788012904-00085).
- 20 R. Duncan and L. Izzo, Dendrimer biocompatibility and toxicity, *Adv. Drug Delivery Rev.*, 2005, **57**(15), 2215–2237, DOI: [10.1016/j.addr.2005.09.019](https://doi.org/10.1016/j.addr.2005.09.019).
- 21 A. Castonguay, E. Ladd, T. G. van de Ven and A. Kakkar, Dendrimers as bactericides, *New J. Chem.*, 2012, **36**, 199–204, DOI: [10.1039/C1NJ20481E](https://doi.org/10.1039/C1NJ20481E).
- 22 K. Jain, P. Kesharwani, U. Gupta and N. K. Jain, Dendrimer toxicity: Let's meet the challenge, *Int. J. Pharm.*, 2010, **394**(1–2), 122–142, DOI: [10.1016/j.ijpharm.2010.04.027](https://doi.org/10.1016/j.ijpharm.2010.04.027).
- 23 M. Sowińska, A. Laskowska, A. Guśpiel, J. Solecka, M. Bochynska-Czyż, A. W. Lipkowski, K. Trzeciak and Z. Urbanczyk-Lipkowska, Bioinspired Amphiphilic Peptide Dendrimers as Specific and Effective Compounds against Drug Resistant Clinical Isolates of *E. coli*, *Bioconjugate Chem.*, 2018, **29**(11), 3571–3585, DOI: [10.1021/acs.bioconjchem.8b00544](https://doi.org/10.1021/acs.bioconjchem.8b00544).
- 24 T. N. Siriwardena, M. Stach, R. He, B. H. Gan, S. Javor, M. Heitz, L. Ma, X. Cai, P. Chen, D. Wei, H. Li, J. Ma, T. Köhler, C. van Delden, T. Darbre and J. L. Reymond, Lipidated Peptide Dendrimers Killing Multidrug-Resistant Bacteria, *J. Am. Chem. Soc.*, 2018, **140**(1), 423–432, DOI: [10.1021/jacs.7b11037](https://doi.org/10.1021/jacs.7b11037).
- 25 Z. Lai, Q. Jian, G. Li, C. Shao, Y. Zhu, X. Yuan, H. Chen and A. Shan, Self-Assembling Peptide Dendron Nanoparticles with High Stability and a Multimodal Antimicrobial Mechanism of Action, *ACS Nano*, 2021, **15**(10), 15824–15840, DOI: [10.1021/acsnano.1c03301](https://doi.org/10.1021/acsnano.1c03301).
- 26 M. Gide, A. Nimmagadda, M. Su, M. Wang, P. Teng, C. Li, R. Gao, H. Xu, Q. Li and J. Cai, Nano-Sized Lipidated Dendrimers as Potent and Broad-Spectrum Antibacterial Agents, *Macromol. Rapid Commun.*, 2018, **39**(24), e1800622, DOI: [10.1002/marc.201800622](https://doi.org/10.1002/marc.201800622).
- 27 W. Guo, Y. Wang, P. Wan, H. Wang, L. Chen, S. Zhang, C. Xiao and X. Chen, Cationic amphiphilic dendrons with effective antibacterial performance, *J. Mater. Chem. B*, 2022, **10**, 456–467, DOI: [10.1039/D1TB02037D](https://doi.org/10.1039/D1TB02037D).
- 28 J. Fernandez, G. Acosta, D. Pulido, M. Malý, J. L. Copa-Patiño, J. Soliveri, M. Royo, R. Gómez, F. Albericio, P. Ortega and F. J. de la Mata, Carbosilane Dendron-Peptide Nanoconjugates as Antimicrobial Agents, *Mol. Pharmaceutics*, 2019, **16**(6), 2661–2674, DOI: [10.1021/acs.molpharmaceut.9b00222](https://doi.org/10.1021/acs.molpharmaceut.9b00222).
- 29 R. Kannan, P. Prabakaran, R. Basu, C. Pindi, S. Senapati, V. Muthuvijayan and E. Prasad, Mechanistic study on the antibacterial activity of self-assembled poly(aryl ether)-based amphiphilic dendrimers, *ACS Appl. Bio Mater.*, 2019, **2**, 3212–3224, DOI: [10.1021/acsbm.9b00140](https://doi.org/10.1021/acsbm.9b00140).
- 30 S. R. Meyers, F. S. Juhn, A. P. Griset, N. R. Luman and M. W. Grinstaff, Anionic amphiphilic dendrimers as antibacterial agents, *J. Am. Chem. Soc.*, 2008, **130**, 14444–14445, DOI: [10.1021/ja806912a](https://doi.org/10.1021/ja806912a).
- 31 A. Chen, A. Karanastasis, K. R. Casey, M. Necelis, B. R. Carone, G. A. Caputo and E. F. Palermo, Cationic Molecular Umbrellas as Antibacterial Agents with Remarkable Cell-Type Selectivity, *ACS Appl. Mater. Interfaces*, 2020, **12**(19), 21270–21282, DOI: [10.1021/acsami.9b19076](https://doi.org/10.1021/acsami.9b19076).
- 32 A. Chen, E. Chen and E. F. Palermo, Guanidium-functionalized cationic molecular umbrellas as antibacterial agents, *Polym. Chem.*, 2021, **12**, 2374–2378, DOI: [10.1039/D1PY00071C](https://doi.org/10.1039/D1PY00071C).
- 33 T. Yu, X. Liu, A. L. Bolcato-Bellemin, Y. Wang, C. Liu, P. Erbacher, F. Qu, P. Rocchi, J. P. Behr and L. Peng, An amphiphilic dendrimer for effective delivery of small interfering RNA and gene silencing in vitro and in vivo, *Angew. Chem., Int. Ed.*, 2012, **51**(34), 8478–8484, DOI: [10.1002/anie.201203920](https://doi.org/10.1002/anie.201203920).
- 34 D. Dhumal, W. Lan, L. Ding, Y. Jiang, Z. Lyu, E. Laurini, D. Marson, A. Tintaru, N. Dusetti, S. Giorgio, J. Iovanna, S. Pricl and L. Peng, An ionizable supramolecular dendrimer nanosystem for effective siRNA delivery with a favorable safety profile, *Nano Res.*, 2021, **14**, 2247–2254, DOI: [10.1007/s12274-020-3216-8](https://doi.org/10.1007/s12274-020-3216-8).
- 35 X. Liu, C. Liu, J. Zhou, C. Chen, F. Qu, J. J. Rossi, P. Rocchi and L. Peng, Promoting siRNA delivery via enhanced cellular uptake using an arginine-decorated amphiphilic dendrimer, *Nanoscale*, 2015, **7**(9), 3867–3875, DOI: [10.1039/c4nr04759a](https://doi.org/10.1039/c4nr04759a).
- 36 C. Chen, P. Posocco, X. Liu, Q. Cheng, E. Laurini, J. Zhou, C. Liu, Y. Wang, J. Tang, V. D. Col, T. Yu, S. Giorgio, M. Fermeleglia, F. Qu, Z. Liang, J. J. Rossi, M. Liu, P. Rocchi, S. Pricl and L. Peng, Mastering Dendrimer Self-Assembly for Efficient siRNA Delivery: From Conceptual Design to In Vivo Efficient Gene Silencing, *Small*, 2016, **12**(27), 3667–3676, DOI: [10.1002/smll.201503866](https://doi.org/10.1002/smll.201503866).
- 37 X. Liu, J. Wu, M. Yammine, J. Zhou, P. Posocco, S. Viel, C. Liu, F. Ziarelli, M. Fermeleglia, S. Pricl, G. Victorero, C. Nguyen, P. Erbacher, J. P. Behr and L. Peng, Structurally flexible triethanolamine core PAMAM dendrimers are effective nanovectors for DNA transfection in vitro and in vivo to the mouse thymus, *Bioconjugate Chem.*, 2011, **22**(12), 2461–2473, DOI: [10.1021/bc200275g](https://doi.org/10.1021/bc200275g).
- 38 S. Bhattacharjee, DLS and zeta potential—what they are and what they are not?, *J. Controlled Release*, 2016, **235**, 337–351, DOI: [10.1016/j.jconrel.2016.06.017](https://doi.org/10.1016/j.jconrel.2016.06.017).
- 39 J. Aguiar, P. Carpena, J. A. Molina-Bolivar and C. C. Ruiz, On the determination of the critical micelle concentration by the pyrene 1:3 ratio method, *J. Colloid Interface Sci.*, 2003, **258**(1), 116–122, DOI: [10.1016/S0021-9797\(02\)00082-6](https://doi.org/10.1016/S0021-9797(02)00082-6).
- 40 S. P. Hawser and L. J. Douglas, Biofilm formation by *Candida* species on the surface of catheter materials in vitro, *Infect. Immun.*, 1994, **62**(3), 915–921, DOI: [10.1128/iai.62.3.915-921.1994](https://doi.org/10.1128/iai.62.3.915-921.1994).
- 41 E. A. Trafny, R. Lewandowski, I. Zawistowska-Marciniak and M. Stępińska, Use of MTT assay for determination of



- the biofilm formation capacity of microorganisms in metal-working fluids, *World J. Microbiol. Biotechnol.*, 2013, **29**(9), 1635–1643, DOI: [10.1007/s11274-013-1326-0](https://doi.org/10.1007/s11274-013-1326-0).
- 42 A. M. Holmes, J. R. Heylings, K. W. Wan and G. P. Moss, Antimicrobial efficacy and mechanism of action of poly (amidoamine)(PAMAM) dendrimers against opportunistic pathogens, *Int. J. Antimicrob. Agents*, 2019, **53**(4), 500–507.
- 43 M. A. Mintzer, E. L. Dane, G. A. O'Toole and M. W. Grinstaff, Exploiting dendrimer multivalency to combat emerging and re-emerging infectious diseases, *Mol. Pharm.*, 2012, **9**(3), 342–354, DOI: [10.1021/mp2005033](https://doi.org/10.1021/mp2005033).
- 44 K. Murzyn, T. Róg and M. Pasenkiewicz-Gierula, Phosphatidylethanolamine-phosphatidylglycerol bilayer as a model of the inner bacterial membrane, *Biophys. J.*, 2005, **88**(2), 1091–1103, DOI: [10.1529/biophysj.104.048835](https://doi.org/10.1529/biophysj.104.048835).

