

New catalytic model systems of tyrosinase: fine tuning of the reactivity with pyrazole-based N-donor ligands†

Jessica Nadine Hamann and Felix Tuczek*

Cite this: *Chem. Commun.*, 2014, 50, 2298Received 14th October 2013,
Accepted 8th January 2014

DOI: 10.1039/c3cc47888b


www.rsc.org/chemcomm

Two new Cu(I) complexes have been synthesized and investigated as model systems of the enzyme tyrosinase. The corresponding ligands are based on a combination of an imine function with two different pyrazole groups. The reactivity of the prepared systems with respect to the conversion of monophenols to the corresponding *ortho*-quinones is investigated. The resulting data are compared to results obtained for other catalytic model systems of tyrosinase.

The ubiquitous type 3 copper enzyme tyrosinase (Ty) catalyses the formation of *ortho*-quinones, starting from monophenols. The conversion of tyrosine to dopaquinone occurs in a two-step process, starting with an aromatic hydroxylation which is followed by two-electron oxidation.^{1,2} The formed dopaquinone polymerises spontaneously to the important pigment melanin.^{3,4} The active site of tyrosinase contains a binuclear copper centre, wherein each copper ion is coordinated by three histidine residues.^{5,6} In 2006 Matoba and coworkers published the first crystal structure of a tyrosinase derived from the bacterium *Streptomyces castaneoglobisporus*.⁵ In the meantime more tyrosinases have been structurally characterized.⁷

Hemocyanins (Hc) and catechol oxidases (CO) are also important type 3 copper proteins, but exhibit reactivities different to tyrosinase. Hemocyanins mediate the oxygen transport in arthropods and molluscs whereas catechol oxidases are responsible for the oxidation of catechols to the corresponding *ortho*-quinones.^{6,8,9} All of these copper type 3 enzymes have very similar active sites and bind dioxygen as peroxide in a typical $\mu^2-\eta^2:\eta^2$ (side-on bridging) geometry. In the *oxy* form the oxidation state of copper ions changes from +I to +II.¹


Tyrosinase exhibits monophenolase and diphenolase activity. The monophenolase cycle involves hydroxylation of monophenolic substrates to catechols, which are released as *ortho*-quinones (Scheme 1).^{1,6,10} In recent years, many low-molecular copper complexes have been synthesized as model systems of tyrosinase.^{1,11}



Scheme 1 Conversion of monophenolic substrates to *ortho*-quinones via catechols (monophenolase and diphenolase activity, respectively).

These compounds can be divided into systems which hydroxylate the ligand framework^{12–16} and those which are able to convert external monophenols to the corresponding *ortho*-quinones.^{11,17} The first model system exhibiting the latter reactivity in a catalytic fashion was published by Réglier and coworkers.¹⁸ This binuclear copper(i) complex generates 3,5-di-*tert*-butyl-*o*-quinone (DTBQ) from 2,4-di-*tert*-butyl phenol (DTBP-H) with a turnover number (TON) of 16. The formation of *ortho*-quinone was detected by using UV/Vis spectroscopy, because *ortho*-quinones show an intense absorption band in the range of 400–420 nm.¹⁹ In 1991 a further system exhibiting stoichiometric as well as weakly catalytic monooxygenation of external phenols was reported by Casella and coworkers.¹⁷ This year Herres-Pawlis *et al.* presented a new copper(i) catalyst mediating the hydroxylation of monophenols *via* a well-characterized peroxo intermediate.²⁰

In 2010 our group reported the first catalytic model system based on a mononuclear copper(i) complex.¹¹ This [Cu(I)L_{py}1(CH₃CN)₂]PF₆ system (*cf.* Scheme 2) was found to generate DTBQ from DTBP-H with a TON of 18 after 8 hours. More recently we presented a second model system containing a benzimidazole moiety.²¹ With the new




Scheme 2 (a) Ligand L_{py}1 of the first mononuclear catalytic model system, (b) ligand L_{bzm}1 and ligands L_{hpz}1 and L_{hpz}2 of this study (c and d).^{11,21}

Christian-Albrechts-Universität zu Kiel, Max-Eyth-Straße 2, 24118 Kiel, Germany.
E-mail: ftuczek@ac.uni-kiel.de; Fax: +49-(0)431-1520; Tel: +49-(0)431-1410

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c3cc47888b





Scheme 3 Syntheses and metalation of the new ligands.

L_{bzm}1 system, the TON could be increased to 31, along with a significantly increased turnover frequency.²¹ To further investigate the influence of the heterocyclic group on the reactive properties of our system, we developed two new pyrazole based copper(i) complexes as model systems.

The ligands of both systems contain an imine function terminated by a *tert*-butyl residue; they only differ with respect to the substituents of the pyrazole moiety (**L_{hpz}1** and **L_{hpz}2**; Scheme 2).

Synthesis and reactivity of the new Cu(i) complexes. The two ligands **L_{hpz}1** and **L_{hpz}2** were prepared in several steps by using similar procedures (Scheme 3 and ESI†). For the aminoethylation of the pyrazoles a modification of a literature procedure was used.²² The corresponding copper(i) complexes were prepared from tetrakis(acetonitrile)copper(i) hexafluorophosphate under anaerobic conditions.

To investigate the catalytic activity of the pyrazole-based model systems *in situ* UV/Vis spectroscopy was applied. To this end a 500 μ M solution of the respective copper(i) complex in dichloromethane was prepared and 50 eq. of DTBP-H and 100 eq. of triethylamine were added. Subsequent oxygenation at ambient temperature was found to result in the formation of DTBQ, as indicated by an absorption band appearing at 407 nm (Fig. 1).

As evident from Fig. 1, the complex $[\text{Cu}(\text{i})\text{L}_{\text{hpz}1}(\text{CH}_3\text{CN})_2]\text{PF}_6$ catalyses the formation of 3,5-di-*tert*-butyl-*o*-quinone (DTBQ) from the corresponding phenol DTBP-H. For the determination of the *turnover number* (TON), an extinction coefficient of $\epsilon = 1830 \text{ M}^{-1} \text{ cm}^{-1}$ at 407 nm was applied.¹⁹ Oxygenation of DTBP-H




Fig. 1 UV/Vis spectra of a 500 μ M solution of the complex $[\text{Cu}(\text{i})\text{L}_{\text{hpz}1}(\text{CH}_3\text{CN})_2]\text{PF}_6$ in CH_2Cl_2 after addition of 50 eq. DTBP-H, 100 eq. NEt_3 and oxygenation between 15 min and 6 h; $l = 1 \text{ mm}$.




Fig. 2 Turnover frequency as a function of time (15 min < t < 6 hours) for the systems **L_{hpz}1** and **L_{hpz}2**; inset: their turnover number per dicopper unit as a function of time.

occurred very fast during the first 100 minutes of reaction and then slowed down, leading to a TON of 29 after 6 hours.

The oxygenation was also performed using the complex $[\text{Cu}(\text{i})\text{L}_{\text{hpz}2}(\text{CH}_3\text{CN})_2]\text{PF}_6$. In analogy to the **L_{hpz}1** system, the formation of DTBQ happened quickly during the first two hours, reaching a TON of 23 after 6 hours. In Fig. 2 the TON and TOF (*turnover frequency*) of both systems are compared. Importantly, the **L_{hpz}1** system exhibits both a higher TON (29) and a higher TOF (0.85 min^{-1} after 15 min) as compared to its **L_{hpz}2** counterpart (TON = 23, TOF = 0.62 min^{-1}). Both systems form most of the quinone during the first 120 min and become inactive after ~ 6 hours.

In order to further prove the formation of DTBQ during the reaction of DTBP with molecular oxygen NMR spectra of the reaction mixture were recorded (Fig. 3). To this end the solutions were quenched with 6 M hydrochloric acid after 30 minutes of oxygenation and extracted with dichloromethane to eliminate the copper ions.

In agreement with similar model systems the resulting NMR spectra revealed signals for DTBQ, DTBP-H and the C-C coupling product 3,3',5,5'-tetra-*tert*-butyl-2,2'-biphenol ("biphenol", Fig. 3).^{11,21} The signals of DTBP-H, DTBQ and the biphenol are found at a ratio of 51 : 18 : 31. This result agrees with the TON of 18 measured after 30 minutes of oxygenation (Fig. 2). For the **L_{hpz}2** system the corresponding ratio was obtained as 70 : 15 : 15 (*cf.* ESI†).

Related small-molecule model systems of tyrosinase investigated before (*cf.* Scheme 2) have already indicated an important





Fig. 3 ^1H -NMR spectrum of the oxygenation solution after quenching with HCl (organic phase) for the **L_{hpz}1** system. Inset: resonances in the aliphatic region.



In comparison with the model systems **L_{bzm}1** (TON = 31) and **L_{py}1** (TON = 22) published earlier, the new systems have an

System	L_{py}^1	L_{hpy}^2	L_{hpy}^1	L_{bzm}^1
TON	22	23	29	31
TOF@15 min	0.56	0.62	0.85	0.98

The authors would like to thank Deutsche Forschungsgemeinschaft (DFG), COST and CAU Kiel for financial support as well as Holger Naggert for support with the measurements.

- 1 M. Rolff, J. Schottenheim, H. Decker and F. Tuczek, *Chem. Soc. Rev.*, 2011, **40**, 4077.
- 2 A. W. J. W. Tepper, E. Lonardi, L. Bubacco and G. W. Canters, in *Handbook of Metalloproteins*, ed. A. Messerschmidt, John Wiley, Chichester, 2010.
- 3 Á. Sánchez-Ferrer, J. N. Rodríguez-López, F. García-Cánovas and F. García-Carmona, *Biochim. Biophys. Acta, Protein Struct. Mol. Enzymol.*, 1995, **1247**, 1.
- 4 K. E. van Holde, K. I. Miller and H. Decker, *J. Biol. Chem.*, 2001, **276**, 15563.
- 5 Y. Matoba, T. Kumagai, A. Yamamoto, H. Yoshitsu and M. Sugiyama, *J. Biol. Chem.*, 2006, **281**, 8981.
- 6 H. Decker, T. Schweikardt and F. Tuczek, *Angew. Chem., Int. Ed.*, 2006, **45**, 4546.
- 7 (a) Y. C. Li, Y. Wang, H. B. Jiang and J. P. Deng, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 17002; (b) M. Sendovski, M. Kantev, V. S. Ben-Yosef, N. Adir and A. Fishman, *J. Mol. Biol.*, 2011, **405**, 227; (c) W. T. Ismaya, H. J. Rozeboom, A. Weijn, J. J. Mes, F. Fusetti, H. J. Wichers and B. W. Dijkstra, *Biochemistry*, 2011, **50**, 5477.
- 8 K. Selmeçzi, M. Réglier, G. Michel and G. Speier, *Coord. Chem. Rev.*, 2003, **245**, 191.
- 9 I. A. Koval, P. Gamez, C. Belle, K. Selmeçzi and J. Reedijk, *Chem. Soc. Rev.*, 2006, **35**, 814.
- 10 C. Eicken, C. Gerdemann and B. Krebs, in *Handbook of Metalloproteins*, ed. A. Messerschmidt, R. Huber, T. Poulos and K. Wieghardt, J. Wiley & Sons, Chichester, UK, 2001, vol. 2, p. 1319.
- 11 M. Rolff, J. Schottenheim, G. Peters and F. Tuczek, *Angew. Chem., Int. Ed.*, 2010, **49**, 6438.
- 12 (a) K. D. Karlin, S. Kaderli and A. D. Zuberbühler, *Acc. Chem. Res.*, 1997, **30**, 139; (b) M. Rolff, J. N. Hamann and F. Tuczek, *Angew. Chem., Int. Ed.*, 2011, **50**, 6924.
- 13 E. A. Lewis and W. B. Tolman, *Chem. Rev.*, 2004, **104**, 1047.
- 14 L. M. Mirica, X. Ottenwaelder and T. D. P. Stack, *Chem. Rev.*, 2004, **104**, 1013.
- 15 A. De, S. Mandal and R. Mukherjee, *J. Inorg. Biochem.*, 2008, **102**, 1170.
- 16 K. D. Karlin, J. C. Hayes, Y. Gultneh, R. W. Cruse, J. W. Mckown, J. P. Hutchinson and J. Zubieta, *J. Am. Chem. Soc.*, 1984, **106**, 2121.
- 17 L. Casella, M. Gullotti, M. Bartosek, G. Pallanza and E. Laurenti, *J. Chem. Soc., Chem. Commun.*, 1991, 1235.
- 18 M. Réglier, C. Jorand and B. Waegell, *J. Chem. Soc., Chem. Commun.*, 1990, 1752.
- 19 W. Flaig, Th. Ploetz and A. Küllmer, *Z. Naturforsch.*, 1955, **10b**, 668.
- 20 A. Hoffmann, C. Citek, S. Binder, A. Goos, M. Rübhausen, O. Troepfner, I. Ivanović-Burmazović, E. C. Wasinger, T. D. P. Stack and S. Herres-Pawlis, *Angew. Chem., Int. Ed.*, 2013, **52**, 5398.
- 21 J. Schottenheim, N. Fateeva, W. Thimm, J. Krahmer and F. Tuczek, *Z. Anorg. Allg. Chem.*, 2013, **639**, 1491.
- 22 K. Kenji, *Jp. Pat.* 2010-215610, 2010.