


 Cite this: *Chem. Commun.*, 2016, 52, 8115

 Received 19th April 2016,
 Accepted 25th May 2016

DOI: 10.1039/c6cc03289c

www.rsc.org/chemcomm

2-Methylresorcinarene: a very high packing coefficient in a mono-anion based dimeric capsule and the X-ray crystal structure of the tetra-anion†

 Rakesh Puttreddy,^a Ngong Kodiah Beyeh,^b Elina Kalenius,^a Robin H. A. Ras^b and Kari Rissanen^{*a}

Mono- and tetra-deprotonated 2-methylresorcinarene anions (1 and 2) as their *trans*-1,4-diammoniumcyclohexane (TDAC)²⁺ inclusion complexes are reported. The mono-anion forms a fully closed dimeric capsule [1·H₂O·MeOH]₂²⁻ with a cavity volume of 165 Å³ and (TDAC)²⁺ as the guest with an extremely high packing coefficient, PC = 84.2%, while the tetra-anion forms a close-packed structure with two structurally isomeric tetra-anions 2a and 2b with a 50 : 50 ratio in the crystal lattice.

Resorcinarenes are stable π -rich cavity containing host systems in the C_{4v} crown conformation.¹ Since the 1980's, resorcinarenes and their cavitand analogues have played a key role in molecular recognition processes in a multitude of host-guest systems.² The easy synthesis and selective functionalization either at the upper rim hydroxyl groups or the 2-position on the aromatic rings, and the lower rims have made them a very useful and accessible family of host compounds.² With the C_{4v} vase shaped cavity, resorcinarenes manifest selective guest complexation and in some cases encapsulation through a multitude of weak non-covalent interactions such as cation $\cdots \pi$, C-H $\cdots \pi$ and $\pi \cdots \pi$ interactions.³ In addition, the hydroxyl groups form strong hydrogen bonds with *exo*- or *endo*-complexed guests in the solid state.³ The construction of hydrogen bond based host-guest systems using resorcinarenes has been well-studied with alcohols,⁴ sugars,⁵ steroids,⁶ as well as with heterocyclic five- and six-membered ring compounds.⁷

Treating core resorcinarenes with even weak bases⁸ leads to the formation of deprotonated species in which the increased π -character of the resorcinarene cavity can be used for more efficient molecular recognition processes by enhancing the weak interactions.⁹ In resorcinarenes, the acidity of the first

proton is greater than that of the second and so on, and the cavity attains maximum stability upon tetra-deprotonation, retaining its crown C_{4v} conformation, and being stabilized by stronger circular intramolecular O-H \cdots O⁻ hydrogen bonds. The tetra-deprotonated resorcinarenes *viz.* tetraphenolate anions are known to exist in alkaline media, and are reported to have remarkably high binding constants towards alkyl ammonium salts when compared to the neutral non-deprotonated resorcinarenes.¹⁰ The strong electrostatic interactions between the π -delocalized negatively charged macrocyclic cavity and cationic guests have been reported to lead to high binding constants.⁹ Schneider and co-workers demonstrated the hydrolysis of acetylcholine to choline in the resorcinarene tetraphenolate cavity by ¹H NMR, supporting the hydrolysis mechanism with molecular models.¹¹ Other biological molecules have also been hosted by the resorcinarene tetraphenolate.¹²

Although the resorcinarene tetraphenolate is known in solution, the crystal structure of it has remained elusive. Along these lines, we have previously reported the structures of mono- and di-deprotonated core resorcinarenes using mono- and diamine bases in the solid state.¹³ In this contribution, we report the preparation, X-ray crystal structures and the electrospray ionization mass spectrometry (ESI-MS) characterization of mono- (1) and tetra-anions (2) as their *trans*-1,4-diammoniumcyclohexane, (TDAC)²⁺, complexes obtained by reacting 2-methylresorcinarene, 2-MeC2, and *trans*-1,4-diaminocyclohexane, TDAC, as shown in Fig. 1.

Single crystals of the complexes (TDAC)²⁺@[1·H₂O·MeOH]₂ and 2(2)·3(TDAC)²⁺·2(TDAC)⁺·15(MeOH)·0.5(H₂O) were prepared by mixing 1 : 1 and 1 : 4 molar ratios of 2-MeC2 and TDAC in methanol, respectively, followed by slow evaporation at room temperature. The reaction of 2-MeC2 and TDAC in a 1 : 1 molar ratio forms a tight neutral hydrogen bonded capsule, (TDAC)²⁺@[1·H₂O·MeOH]₂, where the cavity entraps (TDAC)²⁺, as the guest, as shown in Fig. 2. In this dimeric capsule, the capsule halves are held tightly together by hydrogen bonds *via* two methanol [(O-H)_{host} \cdots (O)_{MeOH} \cdots (H-O)_{host}] and two water molecules [(O-H)_{host} \cdots (O-H)_{water} \cdots (O-H)_{host}]. The *endo*-cavity (TDAC)²⁺ is encapsulated very tightly *via* additional N-H \cdots O

^a University of Jyväskylä, Department of Chemistry, Nanoscience Center, P.O. Box 35, FI-40014, Finland. E-mail: kari.t.rissanen@jyu.fi

^b Aalto University, Department of Applied Physics, Puumiehenkuja 2, 02150 Espoo, Finland

† Electronic supplementary information (ESI) available: Single crystal experimental details and mass spectrometry data are included. CCDC 1473236 and 1473237. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6cc03289c






Fig. 1 Synthesis of mono- (**1**) and tetra-anions (**2**) from 2-methyl-resorcinarene (**2-MeC2**) and *trans*-1,4-diaminocyclohexane (**TDAC**). Nomenclature: **TDAC** = *trans*-1,4-diaminocyclohexane; **(TDAC)⁺** = *trans*-1-ammonium-4-aminocyclohexane; **(TDAC)²⁺** = *trans*-1,4-diammonium-cyclohexane.

and N–H... π interactions to the hosting capsule (see ESI,† Fig. S3a). These concerted guest-to-host interactions resulted in an extremely high packing coefficient (PC) of 0.842, *viz.* 84.2%. The volume of **3**, 138.9 Å³, was obtained using Spartan¹⁴ at MM level, while the cavity volume, 165 Å³, was calculated using MSroll¹⁵ within X-Seed¹⁶ with a 1.2 Å probe, PC = 138.9 Å³/165 Å³ = 0.842.

Contrary to the mono-anion, the tetra-anion **2** is an open inclusion complex, with overall composition of 2(**2**)·3(**TDAC**)²⁺·2**TDAC**⁺·15(MeOH)·0.5(H₂O), the eight negative charges being counterbalanced with two *endo*-cavity (**TDAC**)²⁺, one *exo*-cavity (**TDAC**)²⁺ and two *exo*-cavity (**TDAC**)⁺ cations. The deprotonation of the hydroxyl groups in **2-MeC2** can, from very high quality single crystal X-ray diffraction data, be observed from the residual electron density map [in the case of **1**, all H-atoms are clearly visible and this unambiguously confirms that one of the hydroxyl groups does not have a H-atom in it, namely O6 (see ESI,† Fig. S1)]. Another way, which is more reliable with lower quality data (as in case of **2**), is the fact that a normal OH group both accepts and donates H-bonds, while a deprotonated,




Fig. 2 Dimeric capsule, **(TDAC)²⁺@[1-H₂O-MeOH]₂** shown in (a) as ball and stick with the calculated cavity in transparent grey colour and the guest **(TDAC)²⁺** as CPK to illustrate the very tight encapsulation and (b) as the CPK model.




Fig. 3 Chemdraw representation of tetra-anions **2a** and **2b**. The orientation of the resorcinarene core and the oxygen atoms are labelled as in the crystal structure shown in ESI,† Fig. S4.

viz. phenolate, group accepts three H-bonds. In the case of **2**, this is unambiguously verified for the hydroxyl O-atoms O13, O20 and O18 (see ESI,† Fig. S4). While the O-atoms O06 and O25 show a disorder of the hydrogen atoms with a disordered methanol molecule hydrogen bonded to these two O-atoms. Closer inspection reveals that indeed the crystal lattice contains two distinct structural isomers for the tetra-anion **2**, *viz.* **2a** and **2b**, as represented by the 2-D drawings in Fig. 3. The two isomers crystallize so that every second unit cell contains **2a** and **2b** (see ESI,† Fig. S4 for crystal structures), thus manifesting positional 50:50 disorder within the same unit cell. However both **2a** and **2b** have the same orientation in the unit cell so that the oxygen atoms O06 (50%) and O25 (50%) accept three hydrogen bonds¹³ as shown in Fig. 3.

The following observations were used to ascertain the structure of the mono-anion **1** and the two structural isomers **2a** and **2b** of the tetra-anion:

(1) The hydrogen atom positions were located unambiguously from the electron density map for the protonated -NH_3^+ groups in the **(TDAC)²⁺** and **(TDAC)⁺** cations for both **1** and **2**, as well as the upper rim O–H...O[−] hydrogen bonds for **1** (see ESI,† Fig. S1 and S2).

(2) Deprotonated, *viz.* the phenolate, oxygen atoms act as triple hydrogen bond acceptors in both **1** and **2**.

(3) The oxygen atoms, O06 and O25, in the two structural isomers **2a** and **2b** were confirmed to act as triple hydrogen bond acceptors at 50:50 ratios through careful analysis of the similarly disordered methanol molecules hydrogen bonded to them.

The effect of **TDAC** on the deprotonation of **2-MeC2** in the gas phase was studied through a series of ESI-MS analyses in MeCN/MeOH. Titration experiments showed that the deprotonated charge states of **2-MeC2** increased until four equivalents of **TDAC** were added (Fig. 4a). The mono- and di-deprotonated ions, $[\text{2-MeC2-H}]^-$ and $[\text{2-MeC2-2H}]^{2-}$, were clearly observed in the ESI-MS spectra (see ESI,† Fig. S5). Gas phase stability is known to be low for low molecular weight multiply charged ions in conditions where stabilizing solvent interactions are absent.¹⁷ Higher charge density, therefore, usually leads to coulombic explosion of the compound. Gas phase basicities for the multi-deprotonated ions $[\text{2-MeC2-3H}]^{3-}$ and $[\text{2-MeC2-4H}]^{4-}$,





Fig. 4 (a) Relative intensities (r.i.%) of deprotonated ions observed in (–)ESI-MS spectra as a function of added **TDAC** equivalents. Relative intensities (r.i.%) of the deprotonated ions observed in (–)ESI-MS spectra in aged samples; (b) **2-MeC2** in MeCN : MeOH (25 : 1 v/v) and (c) **2-MeC2** and **TDAC** 1 : 4 in MeCN : MeOH (25 : 1 v/v).

are assumed to be relatively high. Hence, the tri- and tetra-deprotonated ions can easily abstract a proton from methanol, which originates from the MeCN:MeOH (25:1 v/v) mixture used in the ESI-MS experiments due to too low solubility of **TDAC** in aprotic solvents. The stabilizing interactions between the deprotonated OH groups of the **2-MeC2** and **TDAC** molecules (which were present in the solid state structures) were not observed in gas phase measurements. However, interaction of the deprotonated ions with solvent molecules was clearly observed in time-dependent ESI-MS experiments, in which the samples of **2-MeC2** and **2-MeC2 + TDAC** (1 : 4) were aged (Fig. 4b and c). The interaction of **2-MeC2** with methanol or acetonitrile resulted in mono- and di-*O*-methylation products and methanol adducts (see ESI,† Scheme S1). The characterization of the *O*-methylation products was verified by collision induced dissociation (CID) experiments (see ESI,† Fig. S6).

In conclusion, we have utilized a simple organic diamine, *trans*-1,4-diaminocyclohexane, **TDAC**, to deprotonate the

2-methyl-resorcinarene and were able to isolate the mono- and tetra-anions. The mono-anion **1** forms, with the help of two water molecules and two methanol molecules, an unprecedented fully closed and charge-neutral hydrogen bonded dimeric capsule which has an extremely high packing coefficient PC of 84.2% for the *trans*-1,4-diammoniumcyclohexane as the guest. The X-ray crystal structure of **2** not only proves the existence of the tetra-anion in solid state, but reveals two structural isomers of it. The isomers, which occur in the same crystal lattice with a position of 50% occupancy each, show the same deprotonation pattern for the three hydroxyl groups and the difference between **2a** and **2b** occurring with the deprotonation of the fourth hydroxyl group at one “corner” of the resorcinarene skeleton in such a way that it does not affect the overall conformation of the tetra-anion, being C_{4v} for both isomers.

We gratefully acknowledge the Academy of Finland (KR: grant no. 263256, 292746, NKB: grant no. 258653, EK: grant no. 284562, 278743, RHAR: grant no. 272579), University of Jyväskylä and Aalto University for financial support.

References

- (a) P. Timmerman, W. Verboom and D. N. Reinhoudt, *Tetrahedron*, 1996, **52**, 2663–2704; (b) W. Sliwa and C. Kozłowski, *Calixarenes and Resorcinarenes*, Wiley, 2009; (c) J. L. Atwood and A. Szumna, *J. Supramol. Chem.*, 2002, **2**, 479–482.
- (a) D. Ajami, L. Liu and J. Rebek Jr., *Chem. Soc. Rev.*, 2015, **44**, 490–499; (b) K. Jie, Y. Zhou, Y. Yao and F. Huang, *Chem. Soc. Rev.*, 2015, **44**, 3568–3587; (c) T. S. Koblenz, J. Wassenaar and J. N. H. Reek, *Chem. Soc. Rev.*, 2008, **37**, 247–262; (d) A. Shivanyuk and J. Rebek, *J. Am. Chem. Soc.*, 2003, **125**, 3432–3433.
- (a) J. L. Atwood and A. Szumna, *Chem. Commun.*, 2003, 940–941; (b) J. L. Atwood, L. J. Barbour and A. Jerga, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 4837–4841; (c) L. R. MacGillivray and J. L. Atwood, *Chem. Commun.*, 1999, 181–182; (d) A. Shivanyuk, E. F. Paulus, K. Rissanen, E. Kolehmainen and V. Böhmer, *Chem. – Eur. J.*, 2001, **7**, 1944–1951; (e) N. K. Beyeh, M. Kogej, A. Åhman, K. Rissanen and C. A. Schalley, *Angew. Chem., Int. Ed.*, 2006, **45**, 5214–5218.
- (a) N. K. Beyeh, D. P. Weimann, L. Kaufmann, C. A. Schalley and K. Rissanen, *Chem. – Eur. J.*, 2012, **18**, 5552–5557; (b) L. Avram, Y. Cohen and J. Rebek Jr., *Chem. Commun.*, 2011, **47**, 5368–5375; (c) I. A. Koshets, Z. I. Kazantseva, A. E. Belyaev and V. I. Kalchenko, *Sens. Actuators, B*, 2009, **140**, 104–108; (d) O. Ugono and K. T. Holman, *Chem. Commun.*, 2006, 2144–2146; (e) O. D. Fox, J. F.-Y. Leung, J. M. Hunter, N. K. Dalley and R. G. Harrison, *Inorg. Chem.*, 2000, **39**, 783–790.
- (a) E. Kalenius, T. Kekäläinen, R. Neitola, K. Beyeh, K. Rissanen and P. Vainiotalo, *Chem. – Eur. J.*, 2008, **14**, 5220–5228; (b) M. He, R. J. Johnson, J. O. Escobedo, P. A. Beck, K. K. Kim, N. N. S. Luce, C. J. Davis, P. T. Lewis, F. R. Fronczek, B. J. Melancon, A. A. Mrse, W. D. Treleaven and R. M. Strongin, *J. Am. Chem. Soc.*, 2002, **124**, 5000–5009; (c) T. Evan-Salem, I. Baruch, L. Avram, Y. Cohen, L. C. Palmer and J. Rebek, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 12296–12300; (d) T. Rhlalou, M. Ferhat, M. A. Frouji, D. Langevin, M. Métayer and J.-F. Verchère, *J. Membr. Sci.*, 2000, **168**, 63–73.
- (a) A. Shivanyuk and J. Rebek, *Chem. Commun.*, 2001, 2374–2375; (b) J. D. Faull and V. K. Gupta, *Thin Solid Films*, 2003, **440**, 129–137; (c) J. D. Faull and V. K. Gupta, *Langmuir*, 2002, **18**, 6584–6592.
- (a) M. Nissinen, E. Wegelius, D. Falábu and K. Rissanen, *CrystEngComm*, 2000, **2**, 151; (b) M. Nissinen and K. Rissanen, *Supramol. Chem.*, 2003, **15**, 581–590.
- (a) A. Jasat and J. C. Sherman, *Chem. Rev.*, 1999, **99**, 931–968; (b) F. Farina, C. Talotta, C. Gaeta and P. Neri, *Org. Lett.*, 2011, **13**, 4842–4845, and references therein.
- H.-J. Schneider, *Angew. Chem., Int. Ed.*, 2009, **48**, 3924–3977.
- H.-J. Schneider, D. Güttes and U. Schneider, *Angew. Chem., Int. Ed.*, 1986, **25**, 647–649.



- 11 (a) H.-J. Schneider and U. Schneider, *J. Inclusion Phenom. Mol. Recognit. Chem.*, 1994, **19**, 67–83; (b) H. J. Schneider, D. Guettes and U. Schneider, *J. Am. Chem. Soc.*, 1988, **110**, 6449–6454; (c) H. J. Schneider and U. Schneider, *J. Org. Chem.*, 1987, **52**, 1613–1615; (d) F. Weinelt and H. J. Schneider, *J. Org. Chem.*, 1991, **56**, 5527–5535.
- 12 M. Inouye, K. Hashimoto and K. Isagawa, *J. Am. Chem. Soc.*, 1994, **116**, 5517–5518.
- 13 N. K. Beyeh, A. Valkonen and K. Rissanen, *CrystEngComm*, 2014, **16**, 3758–3764.
- 14 *Spartan'14*, Wavefunction, Inc., Irvine, USA.
- 15 M. L. Connolly, *J. Mol. Graphics*, 1993, **11**, 139–141.
- 16 (a) L. J. Barbour, *J. Supramol. Chem.*, 2001, **1**, 189–191; (b) J. L. Atwood and L. J. Barbour, *Cryst. Growth Des.*, 2003, **3**, 3–8.
- 17 R. B. Cole, *Electrospray Ionization Mass Spectrometry: Fundamentals, Instrumentation & Applications*, Wiley, New York, 1997, pp. 137–174.

