


Chemical Science

Volume 13
Number 19
21 May 2022
Pages 5419–5776

rsc.li/chemical-science



ISSN 2041-6539

EDGE ARTICLE

Uwe Grether, Marc Nazaré *et al.*
Detection of cannabinoid receptor type 2 in native cells
and zebrafish with a highly potent, cell-permeable
fluorescent probe

Cite this: *Chem. Sci.*, 2022, 13, 5539

All publication charges for this article have been paid for by the Royal Society of Chemistry

Detection of cannabinoid receptor type 2 in native cells and zebrafish with a highly potent, cell-permeable fluorescent probe†

Thais Gazzi,^a Benjamin Brennecke,^a Kenneth Atz,^b Claudia Korn,^b David Sykes,^{cd} Gabriel Forn-Cuni,^e Patrick Pfaff,^f Roman C. Sarott,^{ib} Matthias V. Westphal,^f Yelena Mostinski,^a Leonard Mach,^a Malgorzata Wasinska-Kalwa,^a Marie Weise,^a Bradley L. Hoare,^{cd} Tamara Miljuš,^{cd} Maira Mexi,^{cd} Nicolas Roth,^g Eline J. Koers,^{cd} Wolfgang Guba,^b André Alker,^b Arne C. Rufer,^b Eric A. Kuszniir,^b Sylwia Huber,^b Catarina Raposo,^b Elisabeth A. Zirwes,^b Anja Osterwald,^b Anto Pavlovic,^b Svenja Moes,^b Jennifer Beck,^b Matthias Nettekoven,^b Irene Benito-Cuesta,^h Teresa Grande,^h Faye Drawnel,^b Gabriella Widmer,^b Daniela Holzer,^b Tom van der Wel,ⁱ Harpreet Mandhair,^{ib} Michael Honer,^b Jürgen Fingerle,^b Jörg Scheffel,^{kl} Johannes Broichhagen,^{ib} Klaus Gawrisch,^m Julián Romero,^h Cecilia J. Hillard,ⁿ Zoltan V. Varga,^{mo} Mario van der Stelt,ⁱ Pal Pacher,^m Jürg Gertsch,^j Christoph Ullmer,^b Peter J. McCormick,^g Sergio Oddi,^{pq} Herman P. Spink,^{ib} Mauro Maccarrone,^{qr} Dmitry B. Veprintsev,^{cd} Erick M. Carreira,^f Uwe Grether,^{ab} and Marc Nazaré^{ib}*^a

Despite its essential role in the (patho)physiology of several diseases, CB₂R tissue expression profiles and signaling mechanisms are not yet fully understood. We report the development of a highly potent, fluorescent CB₂R agonist probe employing structure-based reverse design. It commences with a highly potent, preclinically validated ligand, which is conjugated to a silicon-rhodamine fluorophore, enabling cell permeability. The probe is the first to preserve interspecies affinity and selectivity for both mouse and human CB₂R. Extensive cross-validation (FACS, TR-FRET and confocal microscopy) set the stage for CB₂R detection in endogenously expressing living cells along with zebrafish larvae. Together, these findings will benefit clinical translatability of CB₂R based drugs.

Received 29th November 2021

Accepted 22nd January 2022

DOI: 10.1039/d1sc06659e

rsc.li/chemical-science

^aLeibniz-Forschungsinstitut für Molekulare Pharmakologie (FMP), Campus Berlin-Buch, 13125 Berlin, Germany. E-mail: nazare@fmp-berlin.de^bRoche Pharma Research & Early Development, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd., 4070 Basel, Switzerland. E-mail: uwe.grether@roche.com^cFaculty of Medicine & Health Sciences, University of Nottingham, Nottingham NG7 2UH, England, UK^dUnited Centre of Membrane Proteins and Receptors (COMPARE), University of Birmingham and University of Nottingham, Midlands, England, UK^eLeiden University, Einsteinweg 55, 2333 CC Leiden, the Netherlands^fLaboratorium für Organische Chemie Eidgenössische Technische Hochschule Zürich, Vladimir-Prelog-Weg 3, 8093 Zürich, Switzerland^gWilliam Harvey Research Institute, Barts and the London School of Medicine, Queen Mary University of London, London EC1M 6BQ, England, UK^hFaculty of Experimental Sciences, Universidad Francisco de Vitoria, Pozuelo de Alarcón, 28223, Madrid, SpainⁱDepartment of Molecular Physiology, Leiden Institute of Chemistry, Leiden University, 2333 CC, Leiden, the Netherlands^jInstitute of Biochemistry and Molecular Medicine, University of Bern, 3012 Bern, Switzerland^kDermatological Allergology, Allergie-Centrum-Charité, Department of Dermatology and Allergy, Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, Berlin, Germany^lAllergology, Fraunhofer Institute for Translational Medicine and Pharmacology ITMP, Berlin, Germany^mNational Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Rockville, MD 20852, USAⁿDepartment of Pharmacology and Toxicology, Neuroscience Research Center, Medical College of Wisconsin, Milwaukee, WI 53226, USA^oHCEMM-SU Cardiometabolic Immunology Research Group, Department of Pharmacology and Pharmacotherapy, Semmelweis University, 1085 Budapest, Hungary^pFaculty of Veterinary Medicine, University of Teramo, 64100 Teramo, European, Italy^qEuropean Center for Brain Research (CERC), Santa Lucia Foundation, 00179 Rome, Italy^rDepartment of Biotechnological and Applied Clinical Sciences, University of L'Aquila, 67100 L'Aquila, Italy† Electronic supplementary information (ESI) available: Supplementary figures and tables; general synthetic methods; compound synthesis and characterization; molecular docking; *in vitro* pharmacology; fluorescence spectroscopy; TR-FRET kinetic CB₂R binding assay; FACS analysis; time-lapse confocal imaging; *in vivo* zebrafish imaging; NMR spectra; and N-terminal SNAP-hCB₂R sequence. CCDC 1923120. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1sc06659e

Introduction

Cannabinoid type 1 and 2 receptors (CB₁R and CB₂R) are key transducers of extracellular stimuli in the endocannabinoid (eCB) system, a fundamental lipid signaling network in all vertebrates.^{1,2} Kidney, cardiovascular, gastrointestinal, lung, neurodegenerative and psychiatric disorders, along with pain and cancer are linked to eCB impairment.³⁻⁷ While CB₁R is mainly expressed in the central nervous system and to a lesser extent in peripheral tissue, CB₂R is found throughout the periphery and primarily expressed in immune cells.^{8,9} Tissue and cell-type specific receptor expression profiles remain largely uncharted because appropriate biological and chemical tools are lacking. This is aggravated by the absence of antibodies sufficiently specific for both human and rodent CB₂R; which even if available would be precluded for use in translational imaging and incapable of intracellular permeation.¹⁰⁻¹² The development of therapies involving novel CB₂R agonists requires understanding of molecular and cellular mechanisms of action,^{13,14} thus, chemical probes targeting CB₂R are needed. Herein we report the synthesis of a novel, selective, cell-permeable, high affinity CB₂R-agonist fluorescent probe (Fig. 1A) along with its pharmacological validation.

This is the first directly labeled small molecule probe to preserve binding affinity and agonistic efficacy at both human and mouse CB₂R. The probe is applied and cross-validated using time-resolved fluorescence resonance energy transfer (TR-FRET) by flow cytometry and cellular trafficking studies. We showcase its application with super resolution live cell imaging in native cells and zebrafish.





Fig. 1 CB₂R agonist **1** for the design of CB₂R-selective fluorescent probe and docking experiments into the co-crystal structure of active state CB₂R with agonist AM12033 (PDB 6KPC).¹⁵

Results and discussion

Probe design, synthesis, and pharmacological characterization

Fluorescent imaging probes have emerged as sensitive tools with high degree of spatiotemporal resolution.¹⁶ Accordingly, we sought to develop a probe that would provide a well-validated translational path from preclinical pharmacological animal data to clinic. In earlier work,^{17,18} we described fluorescent probes based on a phytocannabinoid that was lacking desirable pharmacokinetics for translational applications. Accordingly, here we employ a drug-derived, reverse-design approach from a pharmacologically well-validated *in vivo* active compound class that would satisfy a number of desirable boundary conditions: affinity, potency and selectivity, chemical stability, water solubility and membrane permeability.^{19,20} A ligand that meets these criteria is **1**,²¹⁻²³ displaying picomolar selective binding affinity on human and mouse CB₂R and full agonistic picomolar potency (Fig. 1B and ESI Table S1†). With this as a lead, a conjugation handle was required for fluorophore attachment.

Prospective SAR studies supported by molecular modelling using docking poses of **1** suggested an exit vector (Fig. 1B, green arrow and ESI Tables S2 and 3†) for thioether linked spacers reaching out into the extracellular space, ultimately leading to the design of (*S*)-**5** as a versatile lynchpin (Scheme 1). Its synthesis commenced with hydroboration of (*S*)-**2**, followed by thio-Mitsunobu reaction which allowed linker introduction. The absolute configuration of **2** was assigned by X-ray crystallography of a *N*-toluenesulfonyl-(*S*)-proline derivative (ESI-S4, see ESI, S23 and S24†). Following deprotection and coupling, (*S*)-**3** was isolated. Subsequent amide coupling of (*S*)-**3** with acid **4** and deprotection furnished (*S*)-**5**. This compound was used to



Scheme 1 Synthesis of probe candidates.






Fig. 2 (A) FACS analysis of the mean fluorescent intensity (MFI) of WT, hCB₂R, mCB₂R and hCB₁R overexpressing CHO cells at different concentrations of 8-SiR. (B) FACS analysis of the MFI of WT and hCB₂R-CHO cells pre-treated with JWH133 (10 μ M) and stained with different concentrations of 8-SiR; see ESI[†] for details. (C and D) TR-FRET characterization of 8-SiR binding association (C) and saturation analysis (D) using HEK-hCB₂R cell membranes. (E and F) Time-lapse confocal microscopy frames for hCB₂R (E) and hCB₁R (F). CHO cells co-stained with 8-SiR (red) and Hoechst 33342 (cyan, nucleus counter stain) at 1, 4, 6, 8 and 10 min; plasma and internal membranes are highlighted with white and yellow dashes, respectively. (G) Association curve of 0.4 μ M 8-SiR on plasma membrane and internal membranes of hCB₂R-CHO cells. See also ESI Videos S1 and S2.[†]

8-SiR enables CB₂R visualization in live cells

Finally, the ability of 8-SiR to visualize endogenous CB₂R expression in non-transfected cells was tested (Fig. 3). A challenge in CB₂R research is the very low expression level of the receptor in native cells, even when these are known to be responsive to CB₂R activation.^{34,35} To this end, primary cultures of human macrophages derived from healthy donors and of murine splenocytes from healthy C57BL/6J wild-type mice were isolated. Incubation of primary cell populations with 8-SiR resulted in increased mean fluorescence (Fig. 3A, B, ESI Videos S3, S4, and S6[†]).

Consistent with the results obtained previously with CHO-CB₂R cells, 8-SiR produced robust, time-dependent membrane labeling and effected receptor internalization. Probe specificity towards human and mouse CB₂R was evident by the abrupt reduction in observed fluorescence after co-incubating cells with 8-SiR with non-fluorescent JWH133, an agonist with high affinity for human and mouse CB₂R²⁵ (Fig. 3C, D and ESI Videos S5 and S7[†]). These competition studies demonstrate the capability of probe 8-SiR for real-time staining of CB₂R at native levels with high specificity. Additionally, the excellent cell permeability of 8-SiR allows accurate tracing of subcellular CB₂R receptor distribution and traffic dynamics as well as staining total cellular content of CB₂R in real-time, live-cell imaging.

Imaging CB₂R *in vivo* using 8-SiR

Having demonstrated the use of 8-SiR for detection of CB₂R in native cells, we then studied the probes in live animals.

Zebrafish embryos and larvae are an established model in biomedical research with high potential for clinical translation^{36–38} and show a remarkably high CB₂R binding site




Fig. 3 Super resolution fluorescence microscopy in native cells using probe 8-SiR. Time-lapse confocal microscopy frames with 8-SiR in murine splenocytes (A and C) and human macrophages (B and D). Cells were pre-stained with Hoechst 33342 (cyan) to counter stain the nuclei and incubated for 10 min with 0.4 μ M 8-SiR (A and B) or 0.6 μ M 8-SiR and 4 μ M JWH133 (C and D). See also ESI Videos S3 to S7.[†]






Fig. 4 Real-time *in vivo* visualization of CB₂R in zebrafish embryos. (A) **8-SiR** (red) injected in 3dpf Tg(fli1:EGFP) zebrafish larvae, which expresses EGFP fluorescent protein in the blood vasculature (cyan). **8-SiR** was found freely circulating in the blood vasculature after injection, with increased intensity in the CHT region (white triangles). (B) **8-SiR** injected in 3dpf Tg(mpeg1.1:mCherry-F) zebrafish larvae, which expresses mCherry-F fluorescent protein in macrophages (cyan), demonstrating its colocalization. (C) 7-AF488 (red) injected in 3dpf Tg(kdrl:mCherry) zebrafish larvae, which expresses mCherry fluorescent protein in the blood vasculature (cyan). See also ESI Videos S8 and S9.†

sequence identity to the human receptor^{39,40} (ESI Fig. S2†). Probes 7-AF488 and **8-SiR** were intravenously injected *via* the Duct of Cuvier at 3 days post fertilization (dpf), and their dynamics were analyzed using confocal microscopy (Fig. 4).

Both probes were freely circulating as evidenced by fluorescence within the blood vasculature, which slowly decayed during the first hours post injection due to diffusion throughout the body (Fig. 4A). Over time, probes were extensively taken up at the Caudal Hematopoietic Tissue (CHT), showing as a strong punctate pattern (Fig. 4A). The CHT is a highly perfused tissue wherein hematopoietic stem cells and immune precursors such as macrophages and neutrophils develop during zebrafish embryogenesis.⁴¹ Given CB₂R's implication in the immune system,^{42–44} we focused on CHT to study the possible uptake of the probes by the residing immune cells.^{45,46} However, the CHT is also the residing place for endothelial cells that remove macromolecular waste from the blood *via* scavenging receptors. Analysis revealed uptake of **8-SiR** from motile cells (immune-related) and non-motile (endothelial) cells. Indeed, **8-SiR** colocalized with local residing macrophages prior to its removal from blood circulation *via* endothelial cells (Fig. 4B, and ESI Videos S8 and S9†).

To avoid imaging of binding due to the unspecific scavenging function of endothelial cells, additional analyses were performed during the first 3 hours post injection. These demonstrated specific uptake of 7-AF488 and **8-SiR** from motile cells both inside CHT as well as crawling into the adjacent fin, putatively endocytosed after contact with CB₂R (Fig. 4C and ESI Video S9†). In contrast, injecting the fluorophores without the CB₂R recognizing element resulted in unspecific fluorescence signal in all cellular membranes in contact⁴⁷ (ESI Fig. S5 and Video S10†).

Conclusions

In conclusion, we have developed cell-permeable, fluorescent probe **8-SiR**, based on a preclinical CB₂R agonist **1**. It displays high affinity, specificity, and agonistic potency. Moreover, the probe shows unprecedented, highly consistent interspecies affinity and potency for both human and mouse CB₂R. Subsequent to validation in overexpressing systems, the probe was used for imaging in native cells and ultimately in live zebrafish. Specifically, in flow cytometry experiments we labelled CB₂R overexpressing cells and demonstrated target specificity by competition experiments. In fluorescence-based TR-FRET assays, kinetic and equilibrium binding profiles for **8-SiR** were determined. Fluorophore **8-SiR** was employed to monitor intracellular CB₂R distribution in real-time, live-cell imaging by super resolution confocal microscopy with overexpressing hCB₂R-CHO cells and endogenously expressing native cells from human macrophages and mouse splenocytes. In zebrafish larvae, **8-SiR** was found to be well tolerated after intravenous injection. It freely circulated inside the blood vasculature over prolonged times before colocalizing with CB₂R expressing macrophages. This demonstrates the implementation of the probe *in vivo* and attendant aspects such as biosafety, bio-distribution, and cellular uptake.

Probe **8-SiR** provides researchers a unique and novel platform to: (1) overcome the large interspecies differences; (2) access a high specificity and low nanomolar affinity probe for CB₂R with full agonist efficacy, and (3) perform an array of biologically and pharmacologically relevant experiments to detect CB₂R and study its function. We anticipate that this novel highly translational agonist CB₂R fluorescent probe will help to



elucidate CB₂R molecular and cellular mechanisms of action, as well as to unravel its expression levels in different disease states.

Data availability

Primary data for probe synthesis, characterization, photo-physical measurements, and fluorescence imaging, as well as atomic coordinates are provided in the ESI.†

Author contributions

M. N., U. G., E. M. C., C. U. and J. F. conceived the research. M. N. and U. G. acquired funding for this project. T. G., B. B., D. S., G. F.-C., W. G., A. C. R., N. G., M. H., J. F., K. G., J. R., C. J. H., P. J. MC., M. v. d. S., P. P., J. G., C. U., S. O., H. P. S., M. Ma., H. S., D. B. V., E. M. C., U. G. and M. N. designed the research approach. T. G., B. B., K. A., C. K., D. S., G. F.-C., Y. M., L. M., M. W., R. C. S., M. V. W., P. P., B. L. H., T. M., M. Me., N. R., E. J. K., W. G., A. A., E. A. K., S. H., C. R., E. A. Z., A. O., A. P., S. M., J. B., I. B.-C., F. D., G. W., D. H., T. v. d. W., H. M., Y. S., Z. V. V. G. F.-C., J. S., J. Br. and S. O. performed experiments, analyzed raw data, and worked on probe validation. T. G., C. K., D. S., G. F.-C., S. O., U. G., and M. N. analyzed all data generated, and wrote the manuscript. B. B., K. A., R. C. S., M. V. W., P. P., M. v. d. S., H. P. S., M. Ma., D. B. V., and E. M. C. provided useful comments and feedback for the manuscript. K. G. was not contactable during the submission process of the manuscript, but contributed to the work. All authors have given approval to the final version of the manuscript.

Conflicts of interest

The authors have no conflicts to declare.

Acknowledgements

M. N., T. G., B. B., Y. M., L. M. M. K.-W. and M. W. would like to thank Edgar Specker, Sandra Miksche and the NMR core facility of the FMP for their excellent support on compound characterization. We are grateful to Barth van Rossum for preparing graphical artwork. We greatly acknowledge the chiral separation of key building block 2 by Daniel Zimmerli and Erik Hunziker. Some aspects of developing TR-FRET assay were supported by the Swiss National Science Foundation grant 159748 to D. B. V., E. J. K. was funded by the IBSA Foundation for Scientific Research. M. Ma. and S. O. thank Lucia Scipioni and Antonio Totaro for cell culture and technical support, and Daunia Laurenti for her technical assistance in live imaging. They are also grateful to the Italian Ministry of Education, University and Research (MIUR) for partial financial support under the competitive grant PRIN 2017. The determination of solubility data by Catherine Karrer and log *D* values by Aynur Ekiciler is greatly acknowledged. We furthermore thank Björn Wagner, Virginie Micallef and Joelle Muller for the generation of PAMPA data. We thank Mathias Christmann, Rainer Haag and PD Daniel Häussinger for helpful discussions. This work was partially supported by the Sino-German Research Project

grant (GZ 1271) to M. N. K. G. Z. V. V. and P. P. are thankful for the support of the Intramural Research Program of NIAAA, NIH. E. M. C. is grateful to ETH Zürich for financial support. R. C. S. and P. P. are awardees of the Scholarship Fund of the Swiss Chemical Industry (SSCI).

References

- 1 S. Munro, K. L. Thomas and M. Abu-Shaar, *Nature*, 1993, **365**, 61–65.
- 2 W. A. Devane, F. A. Dysarz, M. R. Johnson, L. S. Melvin and A. C. Howlett, *Mol. Pharmacol.*, 1988, **34**, 605–613.
- 3 P. Pacher and R. Mechoulam, *Prog. Lipid Res.*, 2011, **50**, 193–211.
- 4 J. Guindon and A. G. Hohmann, *Br. J. Pharmacol.*, 2008, **153**, 319–334.
- 5 R. P. Picone and D. A. Kendall, *Mol. Endocrinol.*, 2015, **29**, 801–813.
- 6 A. M. Malfitano, S. Basu, K. Maresz, M. Bifulco and B. N. Dittel, *Semin. Immunol.*, 2014, **26**, 369–379.
- 7 R. G. Pertwee, *Philos. Trans. R. Soc.*, 2012, **367**, 3353–3363.
- 8 C. Turcotte, M. R. Blanchet, M. Laviolette and N. Flamand, *Cell. Mol. Life Sci.*, 2016, **73**, 4449–4470.
- 9 S. Galiegue, S. Mary, J. Marchand, D. Dussosoy, D. Carriere, P. Carayon, M. Bouaboula, D. Shire, G. Le Fur and P. Casellas, *Eur. J. Biochem.*, 1995, **232**, 54–61.
- 10 H. Y. Zhang, H. Shen, C. J. Jordan, Q. R. Liu, E. L. Gardner, A. Bonci and Z. X. Xi, *Acta Pharmacol. Sin.*, 2019, **40**, 398–409.
- 11 B. Cecyre, S. Thomas, M. Ptito, C. Casanova and J. F. Bouchard, *Naunyn Schmiedeberg's Arch. Pharmacol.*, 2014, **387**, 175–184.
- 12 Y. Marchalant, P. W. Brownjohn, A. Bonnet, T. Kleffmann and J. C. Ashton, *J. Histochem. Cytochem.*, 2014, **62**, 395–404.
- 13 D. S. Tyler, J. Vappiani, T. Caneque, E. Y. N. Lam, A. Ward, O. Gilan, Y. C. Chan, A. Hienzsch, A. Rutkowska, T. Werner, A. J. Wagner, D. Lugo, R. Gregory, C. Ramirez Molina, N. Garton, C. R. Wellaway, S. Jackson, L. MacPherson, M. Figueiredo, S. Stolzenburg, C. C. Bell, C. House, S. J. Dawson, E. D. Hawkins, G. Drewes, R. K. Prinjha, R. Rodriguez, P. Grandi and M. A. Dawson, *Science*, 2017, **356**, 1397–1401.
- 14 G. M. Simon, M. J. Niphakis and B. F. Cravatt, *Nat. Chem. Biol.*, 2013, **9**, 200–205.
- 15 T. Hua, X. Li, L. Wu, C. Iliopoulos-Tsoutsouvas, Y. Wang, M. Wu, L. Shen, C. A. Johnston, S. P. Nikas, F. Song, X. Song, S. Yuan, Q. Sun, Y. Wu, S. Jiang, T. W. Grim, O. Benchama, E. L. Stahl, N. Zvonok, S. Zhao, L. M. Bohn, A. Makriyannis and Z. J. Liu, *Cell*, 2020, **180**, 655–665.
- 16 L. A. Stoddart, L. E. Kilpatrick, S. J. Bridson and S. J. Hill, *Neuropharmacology*, 2015, **98**, 48–57.
- 17 R. C. Sarott, M. Westphal, P. Pfaff, C. Korn, D. A. Sykes, T. Gazzzi, B. Brennecke, K. Atz, M. Weise, Y. Mostinski, P. Hompluem, E. Koers, T. Miljus, N. J. Roth, H. Asmelash, M. C. Vong, J. Piovesan, W. Guba, A. Rufer, E. A. Kusznir, S. Huber, C. Raposo, E. A. Zirwes, A. Osterwald, A. Pavlovic, S. Moes, J. Beck, I. Benito-Cuesta, T. Grande, S. Ruiz de Martin, A. A. Yeliseev, F. Drawnel, G. Widmer,



- D. Holzer, T. van der Wel, H. Mandhair, C.-Y. Yuan, W. Drobyski, Y. Saroz, N. L. Grimsey, M. Honer, J. Fingerle, K. Gawrisch, J. Romero, C. Hillard, Z. Varga, M. van der Stelt, P. Pacher, J. Gertsch, P. McCormick, C. Ullmer, S. Oddi, M. Maccarrone, D. Veprintsev, M. Nazaré, U. Grether and E. M. Carreira, *J. Am. Chem. Soc.*, 2020, **142**, 16953–16964.
- 18 M. V. Westphal, R. C. Sarott, E. A. Zirwes, A. Osterwald, W. Guba, C. Ullmer, U. Grether and E. M. Carreira, *Chem.–Eur. J.*, 2020, **26**, 1380–1387.
- 19 J. Blagg and P. Workman, *Cancer Cell*, 2017, **32**, 9–25.
- 20 P. Workman and I. Collins, *Chem. Biol.*, 2010, **17**, 561–577.
- 21 C. Bissantz, U. Grether, P. Hebeisen, A. Kimbara, Q. Liu, M. Nettekoven, M. Prunotto, S. Roever, M. Rogers-Evans, T. Schulz-Gasch, C. Ullmer, Z. Wang and W. Yang, WO2012168350A1, 2012.
- 22 A. Haider, L. Gobbi, J. Kretz, C. Ullmer, A. Brink, M. Honer, T. J. Woltering, D. Muri, H. Iding, M. Burkler, M. Binder, C. Bartelmus, I. Knuesel, P. Pacher, A. M. Herde, F. Spinelli, H. Ahmed, K. Atz, C. Keller, M. Weber, R. Schibli, L. Mu, U. Grether and S. M. Ametamey, *J. Med. Chem.*, 2020, **63**, 10287–10306.
- 23 A. Haider, J. Kretz, L. Gobbi, H. Ahmed, K. Atz, M. Burkler, C. Bartelmus, J. Fingerle, W. Guba, C. Ullmer, M. Honer, I. Knuesel, M. Weber, A. Brink, A. M. Herde, C. Keller, R. Schibli, L. Mu, U. Grether and S. M. Ametamey, *J. Med. Chem.*, 2019, **62**, 11165–11181.
- 24 G. Lukinavičius, K. Umezawa, N. Olivier, A. Honigmann, G. Yang, T. Plass, V. Mueller, L. Reymond, I. R. Corrêa Jr, Z. G. Luo, C. Schultz, E. A. Lemke, P. Heppenstall, C. Eggeling, S. Manley and K. Johnsson, *Nat. Chem.*, 2013, **5**, 132–139.
- 25 M. Soethoudt, U. Grether, J. Fingerle, T. W. Grim, F. Fezza, L. de Petrocellis, C. Ullmer, B. Rothenhausler, C. Perret, N. van Gils, D. Finlay, C. MacDonald, A. Chicca, M. D. Gens, J. Stuart, H. de Vries, N. Mastrangelo, L. Xia, G. Alachouzos, M. P. Baggelaar, A. Martella, E. D. Mock, H. Deng, L. H. Heitman, M. Connor, V. Di Marzo, J. Gertsch, A. H. Lichtman, M. Maccarrone, P. Pacher, M. Glass and M. van der Stelt, *Nat. Commun.*, 2017, **8**, 13958–13972.
- 26 The binding data are generated from previously frozen membranes, while flow cytometry experiments are performed with viable cells. We hypothesize that membrane lysate preparation affects CB₂R and CB₁R structure differentially, with detrimental effects on the observed selectivity. In the cAMP assay, which is also performed on living cells, the behaviour is consistently similar to that in flow cytometry, confocal imaging. Therefore, in the subsequent much more relevant multiple whole cell evaluations targeting the agonist binding site in its native conformation we observed consistently a very high selectivity.
- 27 S. Singh, C. R. M. Oyagawa, C. Macdonald, N. L. Grimsey, M. Glass and A. J. Vernall, *ACS Med. Chem. Lett.*, 2019, **10**, 209–214.
- 28 F. Spinelli, R. Giampietro, A. Stefanachi, C. Riganti, J. Kopecka, F. S. Abatematteo, F. Leonetti, N. A. Colabufo, G. F. Mangiardi, O. Nicolotti, M. G. Perrone, J. Brea, M. I. Loza, V. Infantino, C. Abate and M. Contino, *Eur. J. Med. Chem.*, 2020, **188**, 112037.
- 29 S. Zhang, P. Shao and M. Bai, *Bioconjugate Chem.*, 2013, **24**, 1907–1916.
- 30 X. Ling, S. Zhang, P. Shao, W. Li, L. Yang, Y. Ding, C. Xu, N. Stella and M. Bai, *Biomaterials*, 2015, **57**, 169–178.
- 31 R. R. Petrov, M. E. Ferrini, Z. Jaffar, C. M. Thompson, K. Roberts and P. Diaz, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 5859–5862.
- 32 L. Wang, M. S. Frei, A. Salim and K. Johnsson, *J. Am. Chem. Soc.*, 2019, **141**, 2770–2781.
- 33 F. Reynolds and K. A. Kelly, *Mol. Imaging*, 2011, **10**, 407–419.
- 34 A. Cooper, S. Singh, S. Hook, J. D. A. Tyndall and A. J. Vernall, *Pharmacol. Rev.*, 2017, **69**, 316–353.
- 35 N. L. Grimsey, C. E. Goodfellow, M. Dragunow and M. Glass, *Biochim. Biophys. Acta*, 2011, **1813**, 1554–1560.
- 36 M. C. Gomes and S. Mostowy, *Trends Microbiol.*, 2020, **28**, 10–18.
- 37 E. E. Patton and D. M. Tobin, *Dis. Models Mech.*, 2019, **12**, dmm039370.
- 38 E. E. Patton, L. I. Zon and D. M. Langenau, *Nat. Rev. Drug Discovery*, 2021, **20**, 611–628.
- 39 R. G. Krug II and K. J. Clark, *Gene*, 2015, **570**, 168–179.
- 40 F. Oltrabella, A. Melgoza, B. Nguyen and S. Guo, *Dev., Growth Differ.*, 2017, **59**, 194–210.
- 41 E. Murayama, K. Kissa, A. Zapata, E. Mordelet, V. Briolat, H.-F. Lin, R. I. Handin and P. Herbomel, *Immunity*, 2006, **25**, 963–975.
- 42 V. Esain, W. Kwan, K. J. Carroll, M. Cortes, S. Y. Liu, G. M. Frechette, L. M. Sheward, S. Nissim, W. Goessling and T. E. North, *Stem Cells*, 2015, **33**, 2596–2612.
- 43 Y. J. Liu, H. B. Fan, Y. Jin, C. G. Ren, X. E. Jia, L. Wang, Y. Chen, M. Dong, K. Y. Zhu, Z. W. Dong, B. X. Ye, Z. Zhong, M. Deng, T. X. Liu and R. Ren, *J. Biol. Chem.*, 2013, **288**, 13551–13562.
- 44 I. Rodriguez-Martin, M. J. Herrero-Turrion, E. Marron Fdez de Velasco, R. Gonzalez-Sarmiento and R. E. Rodriguez, *Gene*, 2007, **389**, 36–44.
- 45 G. Arias-Alpizar, B. Koch, N. M. Hamelmann, M. A. Neustrup, J. M. J. Paulusse, W. Jiskoot, A. Kros and J. Bussmann, *Nanomedicine*, 2021, **34**, 102395.
- 46 F. Campbell, F. L. Bos, S. Sieber, G. Arias-Alpizar, B. E. Koch, J. Huwyler, A. Kros and J. Bussmann, *ACS Nano*, 2018, **12**, 2138–2150.
- 47 M. Guarin, R. Faelens, A. Giusti, N. De Croze, M. Leonard, D. Cabooter, P. Annaert, P. de Witte and A. Ny, *Sci. Rep.*, 2021, **11**, 12229.

