



Cite this: *Phys. Chem. Chem. Phys.*,
2024, 26, 24294

The strongest dative bond in main-group compounds. Theoretical study of OAeF^- ($\text{Ae} = \text{Be}-\text{Ba}$)†‡

Lei Qin,^{§a} Ruiqin Liu,^{§a} Filip Sagan,^{§b} Zhaoyin Zhang,^a Lili Zhao,^{id} ^{§a}
Mariusz Mitoraj^{id} ^{§b} and Gernot Frenking^{id} ^{§a,c}

Quantum chemical calculations of the anions OAeF^- ($\text{Ae} = \text{Be}-\text{Ba}$) have been carried out using *ab initio* methods at the CCSD(T)/def2-TZVPP level and density functional theory employing BP86 with various basis sets. The equilibrium structures have linear geometries for $\text{Ae} = \text{Be}$ and Mg but they are strongly bent for $\text{Ae} = \text{Sr}$ and Ba while the calcium species has a quasi-linear structure with a very low bending potential. The calculated bond dissociation energies suggest a record-high BDE of $D_e = 144.08 \text{ kcal mol}^{-1}$ for OBeF^- at the CCSD(T)/def2-TZVPP level, which is the strongest BDE for a dative bond that has been found so far. The BDE of the heavier homologues have a continuously decreasing order for Ae with $\text{Be} > \text{Mg}$ ($113.01 \text{ kcal mol}^{-1}$) $>$ Ca ($84.06 \text{ kcal mol}^{-1}$) $>$ Sr ($72.06 \text{ kcal mol}^{-1}$) $>$ Ba ($60.00 \text{ kcal mol}^{-1}$). The calculation of the charge distribution reveals a significant charge donation $\text{OAe} \leftarrow \text{F}^-$ with a declining sequence for the heavier atoms Ae . The oxygen atom in OAeF^- carries always a higher partial charge than the fluorine atom, which contradicts the standard electronegativities of the atoms. The surprising partial charges are explained with the bonding situation of the atoms in the actual electronic structure. The bonding analysis of the $\text{OAe}-\text{F}^-$ bonds using the EDA-NOCV method shows that the bonds have much more electrostatic character than the $\text{Ae}-\text{F}^-$ bonds in the diatomic anions. This finding is supported by the results of the LED partitioning approach. The dative interactions have three major and one minor component. The assignment of a quadruple bond for the heavier species with $\text{Ae} = \text{Ca}, \text{Sr}, \text{Ba}$ is not reasonable. The driving force for the bent geometries is the accumulation of electronic charge in the lone-pair region at the Ae atoms, which enhances the electrostatic attraction with the other atoms. An adequate description of the bonding situation is given by the formula $\text{O}^- - \text{Ae}^+ \leftarrow \text{F}^-$.

Received 7th May 2024,
Accepted 2nd September 2024

DOI: 10.1039/d4cp01909a

rsc.li/pccp

Introduction

Recently we reported theoretical work on the diatomic anions AeF^- ($\text{Ae} = \text{Be}-\text{Ba}$), which possess surprisingly strong dative bonds between the closed-shell species Ae and F^- .^{1,2} The calculated bond dissociation energies are between $87.5 \text{ kcal mol}^{-1}$ for BeF^- and $68.8 \text{ kcal mol}^{-1}$ for MgF^- and they show an unusual

increasing trend $\text{MgF}^- < \text{CaF}^- < \text{SrF}^- < \text{BaF}^-$. The strong bonds were explained by the inductive force of F^- , which heavily polarizes the $(n)s^2$ valence electrons of the Ae atoms, inducing in turn four dative covalent interactions^{3–5} of the heavier species involving the $(n-1)d$ AOs of $\text{Ca}-\text{Ba}$.^{1,2} The fluorine anion F^- being isoelectronic to Ne is a weak electron donor due to the high electronegativity of fluorine and the alkaline earth atoms are weak electron acceptors. We speculated that replacing the weak electron acceptors Ae with the stronger electron acceptors AeO could lead to even higher bond dissociation energies (BDEs). Earlier studies by one of us showed that diatomic BeO is likely the strongest neutral Lewis acid, which even binds the extremely weak Lewis base He with a BDE of $D_e \sim 3 \text{ kcal mol}^{-1}$.^{6–9}

The objective of this work is to search for compounds that possess even stronger dative bonds than diatomic AeF^- . A logical step is the replacement of the weak Lewis acceptor atom Ae by the strong Lewis acid AeO .

Here we present quantum chemical calculations using *ab initio* methods and density functional theory of the anions

^a State Key Laboratory of Materials-Oriented Chemical Engineering, School of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing 211816, China. E-mail: ias_llzhao@njtech.edu.cn

^b Department of Theoretical Chemistry, Faculty of Chemistry, Jagiellonian University, R. Gronostajowa 2, 30-387 Cracow, Poland. E-mail: mitoraj@chemia.uj.edu.pl

^c Fachbereich Chemie, Philipps-Universität Marburg, Hans-Meerwein-Strasse 4, D-35043 Marburg, Germany. E-mail: frenking@chemie.uni-marburg.de

† This paper is dedicated to Prof. Manfred Reetz on the occasion of his 80th birthday.

‡ Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4cp01909a>

§ L. Q., R. L. and F. S. contributed equally.



OAeF[−] (Ae = Be–Ba) and an analysis of the O Ae–F[−] bonds. None of the triatomic anions has yet been observed experimentally. The calculated results show that the compounds are very stable species with record-breaking strong dative bonds and unusual properties that can be observed in the gas phase. The theoretically predicted vibrational spectra are a helpful guide for identifying the molecules in experiments.

Methods

The bond lengths, vibrational frequencies and BDEs of O AeF[−] (Ae = Be–Ba) and in the electronic singlet ground state were calculated at the CCSD(T)¹⁰ and BP86^{11,12} level in conjunction with the basis sets def2-TZVPP.¹³ The CCSD(T) calculations were carried out to give quantitatively accurate numerical values and the DFT calculation were performed to provide a basis for the EDA-NOCV calculations and for investigating the performance of the BP86 functional, which was found to be very good. The calculations were carried out with the program Gaussian 16.¹⁴ The NBO calculations were performed using the program NBO 7.0.¹⁵ We also calculated atomic partial charges using the Hirshfeld¹⁶ and CM5¹⁷ methods with the program Gaussian 16, and calculated Voronoi¹⁸ charges and Mayer¹⁹ bond order with the program Multiwfn.²⁰

The bonding situation in the molecules was further analyzed by means of an energy decomposition analysis (EDA) which was introduced by Morokuma²¹ and by Ziegler and Rauk²² in conjunction with the natural orbitals for chemical valence (NOCV)^{23,24} method. The EDA-NOCV^{25,26} calculations were carried out with the ADF 2018.105 program package^{27,28} at the BP86-D3(BJ)²⁹ level with Slater-type basis function of DZP quality³⁰ using the BP86/def2-TZVPP optimized geometries. DZP is a double- ζ quality basis set augmented by a set of polarization functions. The use of larger basis sets does not lead to significantly different energy values, but can lead to unphysically large numerical contributions from atomic orbitals of higher quantum number such as $(n+1)s$ AOs of Ae atom, which represent numerical artifacts. In this analysis, the intrinsic interaction energy (ΔE_{int}) between two fragments can be divided into three energy components as follows:

$$\Delta E_{\text{int}} = \Delta E_{\text{elstat}} + \Delta E_{\text{Pauli}} + \Delta E_{\text{orb}} \quad (1)$$

The electrostatic ΔE_{elstat} term represents the quasiclassical electrostatic interaction between the unperturbed charge distributions of the prepared fragments, the Pauli repulsion ΔE_{Pauli} corresponds to the energy change associated with the transformation from the superposition of the unperturbed electron densities of the isolated fragments to the wavefunction, which properly obeys the Pauli principle through explicit antisymmetrization and renormalization of the production wavefunction. The orbital term ΔE_{orb} comprises the mixing of orbitals, charge transfer and polarization between the isolated fragments. The energy change involved in the latter step, which is the main difference between the Morokuma²¹ and Ziegler/Rauk²² approaches, is calculated with an extension of Slater's

transition state method³¹ for energy differences. It is often referred as ETS method. The orbital term ΔE_{orb} can be further decomposed into contributions from each irreducible representation of the point group of the interacting system as follows:

$$\Delta E_{\text{orb}} = \sum_r \Delta E_r \quad (2)$$

The combination of the EDA with NOCV enables the partition of the total orbital interactions into pairwise contributions of the orbital interactions which is very vital to get a complete picture of the bonding. The charge deformation $\Delta \rho_k(r)$, resulting from the mixing of the orbital pairs $\psi_k(r)$ and $\psi_{-k}(r)$ of the interacting fragments presents the amount and the shape of the charge flow due to the orbital interactions (eqn (3)), and the associated energy term ΔE_{orb} provides the size of stabilizing orbital energy originated from such interaction (eqn (4)) where $F_{-k,-k}^{\text{TS}}$ and $F_{k,k}^{\text{TS}}$ are the Fock matrix elements defined for the transition state electron density (the mid density between a molecule and fragments).^{25,26}

$$\Delta \rho_{\text{orb}}(r) = \sum_k \Delta \rho_k(r) = \sum_{k=1}^{N/2} \nu_k [-\psi_{-k}^2(r) + \psi_k^2(r)] \quad (3)$$

$$\Delta E_{\text{orb}} = \sum_k \Delta E_{\text{orb}}^k(r) = \sum_k \nu_k [-F_{-k,-k}^{\text{TS}} + F_{k,k}^{\text{TS}}] \quad (4)$$

The EDA-NOCV approach was previously criticized because its results are allegedly path-dependent.^{32–34} This was rebutted in a detailed discussion of the method.³⁵ More details about the EDA-NOCV method and its application are given in recent review articles.^{36–42}

Upon request of one referee we carried out additional calculations for the analysis of the chemical bonds. We employed the local energy decomposition^{43–45} (LED) method in conjunction with *ab initio* calculations at the DLPNO-CCSD(T)^{46–50} level. DLPNO-CCSD(T) recovers >95% of triples contribution and 99.8% of correlation energy.⁴⁹ The LED approach decomposes the interaction energy between the frozen fragments into the Hartree–Fock and correlation interaction energies:

$$\Delta E_{\text{int}} = \Delta E_{\text{int}}^{\text{HF}} + \Delta E_{\text{int}}^{\text{C}} + \Delta E_{\text{el-prep}} \quad (5)$$

Hartree–Fock part of interaction energy consists of the electrostatic and exchange terms.

$$\Delta E_{\text{int}}^{\text{HF}} = \Delta E_{\text{elstat}} + \Delta E_{\text{exch}} \quad (6)$$

Correlation interaction energy on the other hand is decomposed into:

$$\Delta E_{\text{int}}^{\text{C}} = \Delta E_{\text{CT}(A \rightarrow B)}^{\text{C-SP}} + \Delta E_{\text{CT}(B \rightarrow A)}^{\text{C-SP}} + \Delta E_{\text{disp}}^{\text{C}} + \Delta E^{\text{C-WP}} + \Delta E^{\text{C-T}} \quad (7)$$

where $\Delta E_{\text{CT}}^{\text{C-SP}}$ terms are the instantaneous strong-pair charge transfer terms, $\Delta E_{\text{disp}}^{\text{C}} = \Delta E_{\text{disp}}^{\text{C-SP}} + \Delta E_{\text{disp}}^{\text{C-WP}}$ is a dispersion term, consisting of strong-pair and weak-pair interaction, $\Delta E^{\text{C-WP}}$ is the rest of weak-pair interactions, while $\Delta E^{\text{C-T}}$ is the triples correction.



Table 1 Calculated vibrational frequencies ν and frequency shifts Δ with respect to AeO (cm^{-1}) and IR intensities I (km mol^{-1}) of OAeF^- and AeO (Ae = Be–Ba)

Molecule	CCSD(T)/def2-TZVPP [BP86/def2-TZVPP]			Ae–F stretch			O–Ae–F bent	
	Ae–O stretch ν	Δ	I^b	ν	Δ	I^b	ν	I^b
BeO	1434.0 [1454.5]	—	[4.2]	—	—	—	—	—
BeF [−]	—	—	—	1028.0 [1007.2]	—	[14.3]	—	—
OBe–F [−]	1519.3 [1481.2]	+85.3 [+26.7]	[204.9]	683.8 [665.0]	−344.2 [−354.7]	[4.6]	396.0 ^a [385.3] ^a	[34.0]
MgO	783.7 [801.8]	—	[13.8]	—	—	—	—	—
MgF [−]	—	—	—	577.7 [540.8]	—	[37.7]	—	—
OMg–F [−]	847.2 [827.4]	+33.9 [+25.6]	[73.3]	524.4 [502.8]	−53.3 [−45.5]	[10.9]	171.4 ^a [170.1] ^a	[48.6]
CaO	589.3 [751.8]	—	[68.5]	—	—	—	—	—
CaF [−]	—	—	—	485.6 [511.6]	—	[44.5]	—	—
OCa–F [−]	584.8 [601.6]	−4.5 [−150.2]	[93.9]	401.0 [406.2]	−84.6 [−111.1]	[91.8]	54.3 ^a [74.5]	[51.8]
SrO	407.8 [657.3]	—	[74.5]	—	—	—	—	—
SrF [−]	—	—	—	410.6 [428.5]	—	[48.6]	—	—
OSr–F [−]	402.1 [526.5]	−5.7 [−130.8]	[87.9]	306.0 [356.4]	−104.6 [−76.4]	[133.0]	43.4 [112.3]	[26.9]
BaO	618.9 [641.9]	—	[129.6]	—	—	—	—	—
BaF [−]	—	—	—	408.3 [409.1]	—	[56.9]	—	—
OBa–F [−]	485.6 [517.3]	−133.3 [−124.6]	[50.4]	319.7 [317.8]	−88.6 [−93.9]	[134.5]	88.8 [116.3]	[13.2]

^a Degenerate. ^b Intensities are not available at CCSD(T).

TZVPP. To our knowledge, this is the highest value for the BDE of a donor–acceptor bond between two closed-shell species for main group compounds and transition metal complexes that has been reported so far. The alternative electron-sharing rupture of the OBeF^- bond toward $\text{BeO}^- + \text{F}$ has an even higher BDE, because the calculated electron affinity of BeO (1.99 eV) is clearly lower than the electron affinity of fluorine (3.33 eV).⁵⁵ The BDEs for the reaction of the heavier homologues $\text{OAeF}^- \rightarrow \text{AeO} + \text{F}^-$ (Ae = Mg–Ba) are significantly smaller than for the beryllium anion. Unlike the diatomic anions AeF^- ,^{1,2} the triatomic anions OAeF^- exhibit a continuously decreasing trend $\text{Be} > \text{Mg} > \text{Ca} > \text{Sr} > \text{Ba}$, which concurs with the common trend of electron-sharing bonds of main-group compounds, typically becoming weaker for heavier-row atoms.

Table 2 gives also the BDEs for breaking the O–AeF[−] bonds. The bond rupture yielding the electronic ground state of the fragments O (³P) and AeF[−] (¹Σ⁺) at CCSD(T)/def2-TZVPP has slightly higher D_e values for the bonds O–BeF[−] (150.1 kcal mol^{−1}), O–SrF[−] (75.7 kcal mol^{−1}) and O–BaF[−] (70.3 kcal mol^{−1}), smaller D_e values for O–MgF[−] (97.5 kcal mol^{−1}) and a comparable D_e value for

O–CaF[−] (83.5 kcal mol^{−1}) than for the respective OAeF^- bonds. Both sets of O–AeF[−] and OAeF^- bonds have the same regular trend of the BDEs $\text{Be} \gg \text{Mg} > \text{Ca} > \text{Sr} > \text{Ba}$, which underlines the peculiar feature of the diatomic anions AeF^- .^{1,2} But the bond dissociation reaction $\text{OAeF}^- (\text{singlet}) \rightarrow \text{O} (^3\text{P}) + \text{AeF}^- (^1\Sigma^+)$ is spin-symmetry forbidden. The spin-symmetry allowed formation of O (¹D) + AeF[−] (¹Σ⁺) is thermodynamically 45.3 kcal mol^{−1} higher in energy. The BDE for the dissociation reaction $\text{OAeF}^- (\text{singlet}) \rightarrow \text{AeF}^- (^2\Sigma^+) + \text{O}^- (^2\text{P})$ is higher at CCSD(T)/def2-TZVPP than for the reaction yielding O (³P) and AeF[−] (¹Σ⁺), whereas BP86/def2-TZVPP gives slightly lower values for Ae = Ca–Ba. Thus, the strong O–Ae–F[−] bonds are the lowest energy symmetrically allowed reaction channels for fragmentation. All triatomic anions are thermodynamically very stable species.

Bonding analysis

We analyzed the electronic structure and bonding situation in the triatomic anions OAeF^- with a variety of methods. Table 3

Table 2 Calculated bond dissociation energies D_e of the $[\text{OAeF}]^-$ (Ae = Be–Ba) species

	$[\text{OAeF}]^- (\text{S}) \rightarrow \text{Oae} (\text{S}) + \text{F}^- (\text{S})$		$[\text{OAeF}]^- (\text{S}) \rightarrow \text{Oae}^- (\text{D}) + \text{F} (\text{D})$		$[\text{OAeF}]^- (\text{S}) \rightarrow [\text{AeF}]^- (\text{S}) + \text{O} (\text{T})$		$[\text{OAeF}]^- (\text{S}) \rightarrow \text{AeF} (\text{D}) + \text{O}^- (\text{D})$	
	CCSD(T)/def2-TZVPP	BP86/def2-TZVPP	CCSD(T)/def2-TZVPP	BP86/def2-TZVPP	CCSD(T)/def2-TZVPP	BP86/def2-TZVPP	CCSD(T)/def2-TZVPP	BP86/def2-TZVPP
$[\text{OBeF}]^-$	144.8	142.6	158.2	163.1	150.1 [201.3]	164.7 [229.8]	157.8	161.0
$[\text{OMgF}]^-$	113.1	114.9	142.9	143.2	97.5 [148.7]	113.3 [178.3]	111.4	116.5
$[\text{OCaF}]^-$	84.6	78.4	125.8	129.9	83.5 [134.7]	108.3 [173.4]	95.0	102.4
$[\text{OSrF}]^-$	72.6	72.1	118.6	125.1	75.7 [127.0]	101.5 [166.5]	94.4	95.7
$[\text{OBaF}]^-$	60.0	62.5	111.9	120.9	70.3 [121.5]	108.7 [173.8]	74.5	100.3

^a Using the experimental value for the ³P → ¹D excitation energy of oxygen atom 45.3 kcal mol^{−1}: Kramida, A., Ralchenko, Yu., Reader, J., and NIST ASD Team (2022). NIST Atomic Spectra Database (ver. 5.10), [Online]. Available: <https://physics.nist.gov/asd> [2023, August 21]. National Institute of Standards and Technology, Gaithersburg, MD. DOI: <https://doi.org/10.18434/T4W30F>.



Table 3 Calculated partial charges (q) and bond orders P of the OAe-F^- ($\text{Ae} = \text{Be-Ba}$) species at the respective equilibrium geometries

CCSD/def2-TZVPP [BP86/def2-TZVPP]					$P(\text{Ae-F}^-)$		
q							
Atoms	NBO ^a	Hirshfeld ^a	Voronoi ^b	CM5 ^a	Bond	Wiberg ^a	Mayer ^b
O	-1.66 [-1.67]	-0.75 [-0.69]	-0.81 [-0.75]	-0.96 [-0.90]	OBe-F ⁻	0.18 [0.22]	0.74 [0.87]
Be	1.54 [1.55]	0.14 [0.05]	0.19 [0.11]	0.48 [0.40]			
OBe	-0.12 [-0.12]	-0.61 [-0.64]	-0.62 [-0.64]	-0.48 [-0.51]	O-BeF ⁻	0.50 [0.60]	1.84 [2.08]
F	-0.88 [-0.88]	-0.39 [-0.36]	-0.39 [-0.36]	-0.52 [-0.49]			
O	-1.62 [-1.58]	-0.82 [-0.74]	-0.99 [-0.90]	-0.97 [-0.89]	OMg-F ⁻	0.10 [0.15]	0.53 [0.68]
Mg	1.55 [1.50]	0.34 [0.21]	0.49 [0.38]	0.58 [0.47]			
OMg	-0.07 [-0.08]	-0.48 [-0.53]	-0.50 [-0.52]	-0.39 [-0.43]	O-MgF ⁻	0.53 [0.74]	1.64 [1.93]
F	-0.93 [-0.92]	-0.51 [-0.47]	-0.50 [-0.48]	-0.61 [-0.57]			
O	-1.62 [-1.56]	-0.88 [-0.76]	-1.05 [-0.90]	-1.05 [-0.99]	OCa-F ⁻	0.08 [0.13]	0.36 [0.54]
Ca	1.61 [1.49]	0.50 [0.29]	0.59 [0.39]	0.76 [0.62]			
OCa	-0.01 [-0.07]	-0.38 [-0.47]	-0.46 [-0.51]	-0.29 [-0.37]	O-CaF	0.50 [0.82]	1.41 [1.82]
F	-0.99 [-0.93]	-0.62 [-0.53]	-0.55 [-0.48]	-0.71 [-0.63]			
O	-1.58 [-1.49]	-0.85 [-0.73]	-1.00 [-0.85]	-1.09 [-1.00]	OSr-F ⁻	0.07 [0.15]	0.31 [0.52]
Sr	1.53 [1.41]	0.46 [0.26]	0.50 [0.33]	0.81 [0.66]			
OSr	-0.05 [-0.08]	-0.39 [-0.47]	-0.50 [-0.52]	-0.28 [-0.34]	O-SrF ⁻	0.59 [0.91]	1.40 [1.77]
F	-0.95 [-0.92]	-0.60 [-0.53]	-0.50 [-0.48]	-0.72 [-0.66]			
O	-1.64 [-1.50]	-0.87 [-0.72]	-1.00 [-0.79]	-1.13 [-1.04]	OBa-F ⁻	0.06 [0.16]	0.24 [0.49]
Ba	1.58 [1.42]	0.50 [0.26]	0.50 [0.24]	0.85 [0.70]			
OBa	-0.06 [-0.08]	-0.37 [-0.46]	-0.50 [-0.55]	-0.28 [-0.34]	O-BaF ⁻	0.64 [0.90]	1.42 [1.74]
F	-0.94 [-0.92]	-0.64 [-0.54]	-0.51 [-0.45]	-0.71 [-0.66]			

^a Obtained at the CCSD/def2-TZVPP (density = current) level with Gaussian program. ^b Obtained at the CCSD/def2-TZVPP (density = current) level with Multiwfn program.

shows the atomic partial charges calculated with four different methods. It was previously found in our study of AeF^- that the charge distribution suggested by the popular NBO method is not reasonable, because the (n)p valence AOs of Ae are not treated as genuine valence orbitals, which leads to a negligible charge donation $\text{Ae} \leftarrow \text{F}^-$.^{1,2} The data in Table 3 show that the same problematic results occur for OAeF^- . The NBO method gives a miniscule charge donation $\text{OAe} \leftarrow \text{F}^-$ between 0.01–0.12e, whereas the other three methods suggest a much larger charge donation between 0.28–0.62e. Since a significant charge donation is supported by further analysis of the electronic structure, we think that the NBO charges are not a reliable indicator of the charge distribution in these molecules.

There is another interesting result concerning the charge distribution in the anions OAeF^- . All four methods suggest that the oxygen atom carries a higher negative partial charge than fluorine, which does not agree with the electronegativity of the atoms. The difference between the negative charges at the terminal atoms towards oxygen is very large. The result is not an artefact of the partitioning procedures for calculating partial charges. Fig. 2 shows the molecular electrostatic potential (MEP) of the molecules, which shows clearly that the oxygen atom carries a larger negative charge than fluorine. It has been shown before that the MEP results are the physically most reliable data for giving the charge distribution in a molecule.⁵⁶ The MEP result does not depend on the chosen value for the contour line. Fig. 2 is obtained with a value of 0.02 a.u. Fig. S2 of ESI,† shows the MEP with a much smaller value of 0.001 a.u. for the contour line. It also shows that oxygen carries a higher negative charge than fluorine.

The seemingly paradoxical result can be understood when one realizes that the common scales for atomic electronegativities are average values derived from compounds in standard electronic states. The electronegativity of an atom is the power




Fig. 2 Plot of the electrostatic potentials of OAeF^- at the BP86/def2-TZVPP level. Computed with value of 0.001 a.u. for the contour of the electronic density.



to attract electronic charge from another atom in a chemical bond, but it is affected by the actual bonding situation. A recent example concerns the carbon atom in the compound class of carbones CL_2 where the C atom has a ^1D ($2s^2 2p^2$) reference state. There are dative bonds $\text{L} \rightarrow \text{C} \leftarrow \text{L}$ where the ligands L donate electronic charge into the vacant 2p AOs of carbon. The bonding situation makes the carbon atom in carbones to have a much higher electronegativity than in standard organic molecules.^{41,42,57–61} There have been early studies where different values for the electronegativity of the valence orbitals in an atom were proposed, but they are hardly ever used and it seems that this approach is forgotten.^{62,63} The present results suggest that the charge distribution in OAeF^- is affected by the different nature of the two bonds, *i.e.* electron-sharing interactions between oxygen and Ae *versus* dative interactions between fluorine and Ae.

Table 3 gives also the bond order of the molecules using the frequently chosen Wiberg method⁶⁴ as well as the Mayer bond orders.^{19,65} The Wiberg bond orders (WBOs) are originally derived from semiempirical CNDO/2 calculations, which neglect the orbital overlap. It has been shown that this can lead to misleading values particularly for polar bonds.^{66–68} The data in Table 3 show that the WBO values are much smaller than the bond orders suggested by the Mayer partitioning method, which explicitly considers the interatomic overlap. The Mayer bond orders (MBOs) are significantly higher than the WBO data for the very polar OAe-F^- and particularly for the O-AeF^- bonds. The MBO values indicate for the latter bonds a

substantial multiple bond character whereas the WBO values are always < 1 . But we want to point out that the bond order values of polar bonds may not be directly taken as measure for the bond multiplicity. Polar bonds have lower bond orders < 1 for an electron-pair single bond due to the smaller overlap than non-polar bonds.⁶⁸ A polar triple bond can have an MBO value significantly smaller than 3 and yet it may possess three bonding electron pairs. This holds in particular for very polar bonds which were found in AeF^- (ref. 1 and 2) and which are present in OAeF^- . More sophisticated methods are needed to provide a meaningful answer to the question of bond multiplicity.

Much insight is gained by examination of the difference density maps where the charge distribution of free OAe and F^- is subtracted from the charge in OAe-F^- . This is shown in Fig. 3 for OBeF^- using four different values for the isosurface, which gives interesting information about the charge flow. The difference density map of the rather diffuse charge with the isosurface value of $0.001 \text{ e a.u.}^{-3}$ in Fig. 3 shows charge depletion from F^- and Be and charge enhancement in the F^- -BeO bonding region and at oxygen atom. More interesting information comes from Fig. 3b and c which displays the charge flow in the areas of higher charge density, which are directly associated with the formation of the F^- -BeO bond and the change in the O-Be bond. There is clearly an increase in the electronic charge between F^- and BeO as well as between O and BeF^- . The red and blue areas of decrease and increase of charge at Be nicely illustrate the $2s \rightarrow 2p_\sigma$ hybridization at Be. Note that the



Fig. 3 Charge density difference maps at the BP86/def2-TZVPP level. OAeF^- and the fragments $\text{OAe} + \text{F}^-$ (Ae = Be–Ba) with different values for the isodensity surfaces Iso given in e a.u.^{-3} . The direction of the charge flow is red $\rightarrow$ blue.



The breakdown of the total orbital term ΔE_{orb} into pairwise contributions gives deep insight into the formation of the covalent bonds. There are four occupied AOs of F^- which can serve as donor orbitals into the four vacant acceptor orbitals of AeO . For the lighter systems we found that there are indeed four pairwise contributions $\Delta E_{\text{orb1}} - \Delta E_{\text{orb4}}$ which provide $>90\%$ to ΔE_{orb} in the linear equilibrium structures of OAEF^- ($\text{Ae} = \text{Be}, \text{Mg}$). Fig. 4 shows the deformation densities $\Delta\rho$ and the associated most important interacting MOs of OBeF^- . The strongest orbital term ΔE_{orb1} comes from the σ donation $\text{OBe (LUMO)} \leftarrow \text{F}^- (2p_{\sigma}, 2s)$ with some further small mixing of OBe orbitals where the contribution of the fluorine $2s$ AO is small but important, because it allows a second albeit weak σ donation $\Delta E_{\text{orb4}} \text{ OBe (LUMO+2)} \leftarrow \text{F}^- (2p_{\sigma}, 2s)$. The stabilization due to the formation of the σ -bonding $\text{HOMO}-1$ of OBeF^- comes from the bonding overlap of the $2p_{\sigma}$ AO of F^- with the small inside green area at Be atom of OBe , which is hidden by the diffuse shaped green area. It becomes visible by inspection of the $\text{HOMO}-1$ of OBeF^- with different isodensity values, which are displayed in Fig. 5 along with the $\text{HOMO}-1$ of OMgF^- . The figure shows on top the 3D surfaces of the orbital with an isovalue of $0.05e \text{ a.u.}^{-3}$. Below are the contour line diagrams with different isovalues in the plane that contains the bond axis. The latter figure nicely shows the attractive interference of the fragment orbitals which lead to a stabilization of $\Delta E_{\sigma_1} = 24.4 \text{ kcal mol}^{-1}$ (OBeF^-) and $\Delta E_{\sigma_1} = 29.1 \text{ kcal mol}^{-1}$ (OMgF^-). A peculiar feature of the orbital mixing between the OAE and F^- fragments is the sign reversal in the $\text{Ae}-\text{F}$ region. There is a

Table 4 shows the numerical results for the anions OAE^- using AeO (S) and F^- (S) in the electronic singlet states as interacting fragments at the BP86/def2-TZVPP optimized

Fragments	OBe (S) + F ⁻ (S)	OMg (S) + F ⁻ (S)	OCa (S) + F ⁻ (S)	OSr (S) + F ⁻ (S)	OBa (S) + F ⁻ (S)	OBa (S) + F ⁻ (S)
Structure	Linear	Linear	Bent	Bent	Bent	Linear
ΔE_{int}	-147.5	-113.5	-79.5	-73.7	-69.8	-62.3
ΔE_{Pauli}	57.3	64.4	53.8	47.8	48.6	41.9
ΔE_{elstat}	-152.2 (74.3%)	-129.9 (73.0%)	-89.7 (67.3%)	-79.5 (65.4%)	-70.2 (59.3%)	-57.4 (55.1%)
ΔE_{orb}^a	-52.7 (25.7%)	-48.0 (27.0%)	-43.6 (32.7%)	-42.0 (34.6%)	-48.2 (40.7%)	-46.8 (44.9%)
$\Delta E_{\text{orb}1}^b$	-24.4 (46.3%)	-29.1 (60.6%)	-19.3 (44.3%)	-12.7 (30.2%)	-21.0 (43.6%)	-18.3 (39.1%)
$\Delta E_{\text{orb}2}^b$	-10.1 (19.2%)	-6.8 (14.2%)	-8.5 (19.5%)	-12.4 (29.5%)	-10.8 (22.4%)	-10.1 (21.6%)
$\Delta E_{\text{orb}3}^b$	-10.1 (19.2%)	-6.8 (14.2%)	-6.3 (14.4%)	-9.0 (21.4%)	-10.8 (22.4%)	-10.1 (21.6%)
$\Delta E_{\text{orb}4}^b$	-4.9 (9.3%)	-3.0 (6.3%)	-4.5 (10.3%)	-3.6 (8.6%)	-1.5 (3.1%)	-5.1 (10.9%)
ΔE_{rest}	-3.2 (6.0%)	-2.0 (4.7%)	-5.0 (11.5%)	-4.3 (10.3%)	-4.1 (8.5%)	-3.1 (6.7%)

This journal is © the Owner Societies 2024




Fig. 4 Plot of deformation densities $\Delta\rho$ of OBeF⁻. The pairwise orbital interactions and the associated interaction energies (ΔE_{orb}) between fragments OBe (S) and F⁻ (S) in OBeF⁻ at the equilibrium structure as well as the shape of the most important interacting MOs. The direction of the charge flow is red $\rightarrow$ blue. (Isovalue 0.03e a.u.⁻³).





Fig. 5 (Top) Plot of the 3D shape of the HOMO–1 orbitals with an isovalue of $0.03e \text{ a.u.}^{-3}$ of OBeF^- and OMgF^- and (bottom) contour line plot of the same orbital HOMO–1 in the plane containing the bond axis with different isovalues at the BP86/def2-TZVPP level. The isovalue below each plot represents the value of the outermost contour in the graph, and the values for the contour settings are in the order of 0.001, 0.002, 0.004, 0.008, 0.02, 0.04, 0.08, 2.0, $4.0e \text{ a.u.}^{-3}$.

diffuse bonding region (red lines) away from the Be–F axis and a blue bonding region close to the axis. The latter blue region is even interrupted by the red region, which does not affect the overall positive interference (bonding interaction) of the orbitals.

The orbital interactions ΔE_{orb2} and ΔE_{orb3} are due to the degenerate π donation $\text{OBe} (\text{LUMO}+1) \leftarrow \text{F}^- (2p_\pi)$. Although the stabilization of ΔE_{orb4} in OBeF^- is rather small, it contributes 9.3% to the covalent bonding and it builds up a different orbital (HOMO–3) than the σ interaction ΔE_{orb1} which leads to HOMO–1. The remaining orbital interactions ΔE_{rest} come from atomic polarization and O–Be interactions and they are not relevant for the OBe–F $^-$ bond. The HOMO of OBeF^- is a degenerate π orbital located mainly at oxygen atom which is only spuriously affected by the bonding interaction with F $^-$. The five energetically highest lying orbitals of OBeF^- along with those of the other OAeF^- anions are shown in Fig. S1 of ESI.†

It is amazing that four bonding orbitals between two main-group atoms of the first octal row elements can be formed. It was previously suggested that C_2 has a quadruple bond,^{71–73} but this claim was disputed by several workers.^{74–78} As mentioned in the introduction, a quadruple bond was suggested for AeF^- (Ae = Ca, Sr, Ba) because the heavier alkaline atoms use valence d-orbitals for covalent bonding. Although there are four bonding orbitals between OBe and F $^-$ we do not think that OBeF^- has a genuine quadruple bond. One reason is that the OBe–F $^-$ bond comes mainly from electrostatic attraction. According to the EDA-NOCV results, only 25.7% of the bond

comes from covalent (orbital) interaction. Another reason is the fact that the chemical bonds in OAeF^- are delocalized over three atoms and that a localized description is not appropriate for depicting the bonding situation. We refer to the chemical bonds in OCBbCO , which has a linear structure where all atoms are bonded by triple bonds.⁷⁹ The representation with a Lewis structure that has only triple bonds not only violates the octet rule, it also disregards the appearance of delocalized bonds. Bond delocalization over more than two atoms was not recognized by Gilbert Lewis and it limits the use of Lewis-type structures for describing chemical bonds.⁸⁰

The deformation densities $\Delta\rho$ and the interacting MOs of OMgF^- are very similar to those of OBeF^- and they are displayed in Fig. 6. The shape of the HOMO–1 hides like the HOMO–1 of OBeF^- the bonding character of the orbital. The bonding nature of the HOMO–1 becomes visible by the contour line diagrams shown in Fig. 5 where the isovalues of OMgF^- appear blue in the high-density areas. A close examination of the deformation density reveals the accumulation of electronic charge in the bonding region between Mg and F $^-$ with concomitant polarization at fluorine. The numerical results in Table 4 show that the energy contribution of the fourth bond ΔE_{orb4} provides only 6.3% of the total covalent bond, but the associated deformation density illustrates the accumulation of electronic charge in the OMgF^- bonding region, which is characteristic for a covalent interaction that comes from the interference of the wave functions.

The analysis of the bent equilibrium structures of OAeF^- (Ae = Ca, Sr, Ba) is very interesting, because the heavier alkaline earth atoms use their (n)d functions as valence orbitals besides the (n)s orbitals in contrast to the lighter homologues which have (n)p valence orbitals. All three complexes have four pairwise orbital interactions $\Delta E_{\text{orb1}} - \Delta E_{\text{orb4}}$ to the covalent OAeF^- bonding due to $\text{OAe} \leftarrow \text{F}^-$ charge donation like the linear structures of the lighter complexes, but the contribution of ΔE_{orb4} to the orbital interactions decreases from the calcium adduct (10.3%) to the barium complex where it is only 3.1% of ΔE_{orb} (Table 4). Fig. 7 shows that the orbital interactions ΔE_{orb1} , ΔE_{orb2} and ΔE_{orb4} in OCaF^- come from donation of F $^-$ into vacant orbitals of CaO which involve mainly valence d AOs of Ca as acceptor orbitals. The LUMO+2 of CaO is mainly a Ca sd_σ hybridized orbital whereas the degenerate LUMO+1/LUMO+1' are sd_π hybridized orbitals. Since the molecule has a bent geometry there is only one genuine occupied π orbital (HOMO–4) whereas the other occupied valence orbitals that are formed by the interaction between CaO and F $^-$ (HOMO, HOMO–1, HOMO–3) are by definition σ orbitals. The shape of the HOMO–3 indicates that it can be considered as in-plane π interaction, although this is not a structurally correct assignment. The mixing of the HOMO of CaO in ΔE_{orb3} and ΔE_{orb4} indicates the concomitant polarization along with the bond formation due to the interference of the fragment wave functions. The five highest lying MOs of OCaF^- are shown in Fig. S1 of ESI.†

The deformation densities $\Delta\rho$ and the interacting MOs of OSrF^- show interesting differences to those of OCaF^- and are






Fig. 6 Plot of deformation densities $\Delta\rho$ of OMgF⁻. The pairwise orbital interactions and the associated interaction energies (ΔE_{orb}) between fragments OMg (S) and F⁻ (S) in OMgF⁻ at the equilibrium structure, as well as the shape of the most important interacting MOs. The direction of the charge flow is red $\rightarrow$ blue. (Isovalue 0.03e a.u.⁻³).

shown in Fig. 8. The π -orbital interaction $\Delta E_{\text{orb}2}$ and the in-plane pseudo- π interaction $\Delta E_{\text{orb}3}$ of OCaF⁻ become $\Delta E_{\text{orb}3}$ and $\Delta E_{\text{orb}4}$ in OSrF⁻. The significantly stronger bending in OSrF⁻ (123.0°) than in OCaF⁻ (144.7°) enhances the second σ interaction in the former anion, which is $\Delta E_{\text{orb}4}$ in OCaF⁻ and becomes the second strongest orbital term $\Delta E_{\text{orb}2}$ in OSrF⁻, which is only slightly weaker ($-12.4 \text{ kcal mol}^{-1}$) than $\Delta E_{\text{orb}1}$ ($-12.7 \text{ kcal mol}^{-1}$). Examination of the shape of the valence orbitals clearly shows that the d-AOs of Sr are the dominant acceptor orbitals of the metal.

We analysed in OBaF⁻ both using the linear structure and with the bent equilibrium geometry in order to identify the

driving force for bending. Table 4 shows the numerical EDA-NOCV results for both structures. The bent equilibrium structure encounters stronger Pauli repulsion ΔE_{Pauli} than the linear form, but the total interaction energy ΔE_{int} is clearly more attractive in the energy minimum form. This is mainly due to the larger Coulomb attraction ΔE_{elstat} in the latter structure, which has a significantly shorter OBa-F⁻ bond than the former. But the orbital (covalent) interactions in the bent form are also slightly stronger than in the linear form. Stronger orbital interactions in the bent structure are found for $\Delta E_{\text{orb}1} - \Delta E_{\text{orb}3}$ but not for $\Delta E_{\text{orb}4}$, which has a negligible contribution in the equilibrium geometry. This makes the assignment of a






Fig. 7 Plot of deformation densities $\Delta\rho$ of OCaF^- . The pairwise orbital interactions and the associated interaction energies (ΔE_{orb}) between fragments OCa (S) and F^- (S) in OCaF^- at the equilibrium structure, as well as the shape of the most important interacting MOs. The direction of the charge flow is red $\rightarrow$ blue. (Isovalue 0.03e a.u.^{-3}).




Fig. 8 Plot of deformation densities $\Delta\rho$ of OSrF⁻. The pairwise orbital interactions and the associated interaction energies (ΔE_{orb}) between fragments OSr (S) and F⁻ (S) in OSrF⁻ at the equilibrium structure, as well as the shape of the most important interacting MOs. The direction of the charge flow is red $\rightarrow$ blue. (Isovalue 0.03e a.u.⁻³).






Fig. 9 Plot of deformation densities $\Delta\rho$ of OBaF^- . The pairwise orbital interactions and the associated interaction energies (ΔE_{orb}) between fragments Oba (S) and F^- (S) in OBaF^- of the bent equilibrium structure, as well as the shape of the most important interacting MOs. The direction of the charge flow is red $\rightarrow$ blue. (Isovalue $0.03e \text{ a.u.}^{-3}$).



quadruple bond between F^- and BaO inappropriate. The same conclusion holds for the bent structures of the Ca and Sr homologues.

Fig. 9 and 10 show the deformation densities $\Delta\rho$ and the interacting MOs for $\Delta E_{\text{orb1}} - \Delta E_{\text{orb4}}$ of the two structures. The orbitals of the acceptor fragment BaO suggest that the d-AOs are the most important acceptor orbitals of the metal. The strongest orbital interaction ΔE_{orb1} comes now from the donation of F^- into the $5d_\pi$ AO and (with smaller contribution) $6p_\pi$

AOs of Ba, which underline the strong contribution of the $(n-1)d$ AOs for covalent bond particularly for Ba. Close examination of the deformation densities in the linear and bent structures reveals the appearance of local charge accumulation at barium in the region of lone-pair, which comes from the charge donation into vacant d-AOs of the metal. This is clearly visible in the strong orbital interactions $\Delta E_{\text{orb1}} - \Delta E_{\text{orb3}}$. The results nicely agree with the finding of Kaupp *et al.* about the relevance of d-orbitals for the heavy alkaline earth atoms.⁵² The




Fig. 10 Plot of deformation densities $\Delta\rho$ of OBaF^- . The pairwise orbital interactions and the associated interaction energies (ΔE_{orb}) between fragments OBa (S) and F^- (S) in OBaF^- of the linear structure, as well as the shape of the most important interacting MOs. The direction of the charge flow is red $\rightarrow$ blue. (Isovalue $0.03e \text{ a.u.}^{-3}$).



accumulation negative charge toward the lone-pair area but away from the bonding region explains why the bent structure is mainly stabilized by electrostatic (Coulomb) interaction than by stronger covalent bonding. The detailed bonding analysis suggests that the OAe-F^- bonds have a higher electrostatic character than the Ae-F^- bonds in the diatomic anions.^{1,2} The covalent contributions to the OAe-F^- bonds come from dative interactions which have multiple-bond character, but the assignment of a quadruple bond like in the heavier AeF^- anion with $\text{Ae} = \text{Ca, Sr, Ba}$ is not reasonable. We suggest that the bonding situation is best described with a bond line O-AeF^- for the electron-sharing bond and an arrow $\text{OAe} \leftarrow \text{F}^-$ for the dative bond where the line above Ae in the bent structures signals the accumulation of local charge in the lone-pair region, although the overall charge of Ae is positive (Fig. 11). The situation may be compared with CO, where the carbon atoms carries a positive partial charge but the chemical reactivity and the dipole moment are dominated by the local charge accumulation in the lone-pair region.

There is a common feature in the deformation densities associated with the orbital interactions ΔE_{orb1} and ΔE_{orb4} and the MOs of the AeO fragments. Fig. 4 and 6–10 show that an occupied orbital of AeO is involved in the dative interactions $\text{OAe} \leftarrow \text{F}^-$ which suggests that there is concomitant polarization in the diatomic moiety. The mode of action between OAc and F^- involves a charge induced interaction of the F^- ion on the OAc moiety, which in turn yields a favorable hybridization of the vacant orbitals of Ae overlapping with the occupied AOs of F^- and leading to a stabilizing interference of the wave functions. This was already found in the diatomic anions AeF^- , where the charge induction of F^- leads to a polarization of the $(n)s$ valence electrons of Ae and the formation of a vacant $(n)sp(\sigma)$ AO. This is schematically shown in Fig. 12. A related mechanism is operative in OAeF^- .

Are the OAe-F^- bonds really dative bonds? The anion dissociates into the closed-shell species OAc and F^- , but this does not necessarily indicate that the finally formed bond has dative character. For example, rupture of the carbon-carbon bond of many olefins like $\text{F}_2\text{C}=\text{CF}_2$ gives carbenes CF_2 in the $^1\text{A}_1$ singlet state but the double bond comes from the interaction between CF_2 in the $^3\text{B}_1$ triplet state.⁸¹ Likewise dissociate the NaBH_3^- Cluster into the closed-shell fragments Na^- and BH_3 but the Na-B bond is better described as electron-sharing interactions between Na and BH_3^- both in their doublet state.⁸² There is a curve crossing of the ΔE_{orb} values along the



Fig. 11 Schematic representation of the bonding situation in the linear and bent structures of OAeF^- . The bonding line $\text{O}^- - \text{Ae}^+ \text{F}^-$ indicates electron-sharing bonding and the arrow $\text{O}^- - \text{Ae}^+ \leftarrow \text{F}^-$ indicates dative bonding both having multiple bond character. The line above Ae signals the accumulation of electronic charge in the lone-pair region in spite of the overall positive charge of Ae atom.



Fig. 12 Schematic representation of the σ bond formation in AeF^- in three steps. (a) Initial charge distribution in Ae atom and F^- where the arrows indicate the total electronic charge; (b) polarization of the valence electrons in Ae and $(n)s^2 \rightarrow (n)s/p(\sigma)$ hybridization; (c) concomitant formation of the vacant orthogonal $(n)s/p(\sigma)$ orbital at Ae atom which overlaps with the occupied σ orbitals of F^- which leads to a polar covalent Ae-F^- bond.

fragmentation pathway when the two sets of fragments are used in the EDA-NOCV calculations, which indicates that the fragments that provide the best description for the chemical bond in the molecule have a different electronic state than the fully separated fragments after dissociation. It could be envisaged that the charge donation $\text{OAe} \leftarrow \text{F}^-$ at the equilibrium geometry leads to an electron-sharing interaction between OAe^- anion and neutral F in the doublet states. We analysed the anions OAe-F^- with the EDA-NOCV method using OAe^- and neutral F as interacting fragments. The numerical results are shown in Table S1 of ESI.† The calculated values for the orbital interactions ΔE_{orb} are for all species at the equilibrium geometry significantly larger than using closed-shell fragments OAc and F^- . Since the final dissociation products are also OAc and F^- it becomes obvious that there is no curve crossing of the ΔE_{orb} , which clearly indicates that the OAe-F^- bonds are really dative bonds. It has been shown in numerous previous studies using the EDA-NOCV method that the relative amount of ΔE_{orb} is a useful indicator for the nature of the nature of the chemical bonds and for the best description of the interacting fragments.

We have further investigated the nature of OAe-F^- bonds with a completely different approach which is based on the electronic structures that come from the *ab initio* calculations at the DLPNO-CCSD(T)/def2-TZVP level using geometries optimized at CCSD(T)/TZVP. The numerical result of the local energy decomposition (LED) method^{43–45} are given in Table 5. The LED scheme breaks down the interaction energy between the fragments into different energy terms than the EDA-NOCV approach, but the main conclusion is the same. The electrostatic attraction $\Delta E_{\text{elst}}^{\text{HF}}$ is the dominant term of the attraction. The correlation interaction energy $\Delta E_{\text{int}}^{\text{C}}$ mostly covers here the charge-transfer term, which can be interpreted as the instantaneous ion pair formation. The most important information comes from the direction of the charge transfer terms ΔE_{CT} which show that the dynamic charge transfer $\text{OAe} \leftarrow \text{F}^-$ completely dominates while the opposite stabilization is



OSrF[−] and OBaF[−] have a red shift for the Ae–O mode. The calculated Ae–O frequency shifts and the Ae–F stretching vibration are a guideline for the hitherto unknown anions OAeF[−].

- The calculation of the charge distribution reveals a significant charge donation $\text{OAe} \leftarrow \text{F}^-$ with decreasing order for the heavier atoms Ae using the Hirshfeld, Voronoi and CM5 methods. In contrast, the NBO method suggests only negligible charge donation, which comes from the fact that only the (n)s but not the (n)p AOs of atoms Ae are treated as valence orbitals.

- The oxygen atom in OAeF[−] carries always a higher partial charge than the fluorine atom. This is suggested by all charge partitioning methods and by the molecular electrostatic map. The charge distribution contradicts the standard electronegativities of the atoms which assign fluorine to be more electronegative than oxygen. The surprising partial charges are explained with the bonding situation of the atoms in the actual electronic structure, which may lead to uncommon partial charges.

- The bonding analysis of the OAe–F[−] bonds using the EDA–NOCV method shows that the bonds have a much higher electrostatic character than the Ae–F[−] bonds in the diatomic anions. The dative interactions have three major and one minor component. The assignment of a quadruple bond for the heavier species with Ae = Ca, Sr, Ba is not reasonable. The driving force for the bent geometries is the accumulation of electronic charge in the lone-pair region at the Ae atoms, which enhances the electrostatic attraction with the other atoms. A reasonable description of the bonding situation is given by the formula $\text{O}^- - \text{Ae}^+ \leftarrow \text{F}^-$.

Note added in proof

The statement about the strongest dative bond refers to atoms heavier than hydrogen and helium. Proton affinities are dative bonds which can be much stronger than calculated in this work.⁹²

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

L. Z. and G. F. acknowledge the financial support from Nanjing Tech University (no. 39837123, 39837132 and the International Cooperation fund), the National Natural Science Foundation of China (no. 22373050), Natural Science Foundation of the Jiangsu province (no. BK20211587), and the State Key Laboratory of Materials-Oriented Chemical Engineering (no. SKL-MCE-23A06), and the Postgraduate Research & Practice Innovation Program of Jiangsu Province (KYCX23_1404). L. Q. and L. Z.

appreciate the Supported by Cultivation Program for The Excellent Doctoral Dissertation of Nanjing Tech University (2024-10). We appreciate the high performance center of Nanjing Tech University for supporting the computational resources. We also thank the PL-Grid Infrastructure and the Academic Computational Centre Cyfronet of the University of Science and Technology in Krakow for providing computational resources. GF is grateful to Holger Bettinger for pointing out the strength of proton affinities.

References

- 1 R. Liu, L. Qin, Z. Zhang, L. Zhao, F. Sagan, M. Mitoraj and G. Frenking, Genuine quadruple bonds between two main-group atoms. Chemical bonding in AeF[−] (Ae = Be–Ba) and isoelectronic EF (E = B–Tl) and the particular role of d orbitals in covalent interactions of heavier alkaline-earth atoms, *Chem. Sci.*, 2023, **14**, 4872–4887.
- 2 L. Qin, Y.-Q. Liu, R. Liu, X. Yang, Z.-H. Cui, L. Zhao, S. Pan, S. Fau and G. Frenking, Analysis of the Unusual Chemical Bonds and Dipole Moments of AeF[−] (Ae = Be–Ba): A Lesson in Covalent Bonding, *Chem. – Eur. J.*, 2024, e202304136.
- 3 The term “covalent” was suggested by Irving Langmuir in 1921 (Ref. 4) as expression for the two-electron bond introduced in 1921 by Gilbert Lewis (Ref. 5). There are two types of two-electron covalent bonds. One type is the electron-sharing bond A–B where both atoms contribute one electron each to the bond. The other type of covalent electron-pair bond is the dative bond A → B where both electrons are provided by one atom A. The two types of electron pair bonds were already distinguished by Lewis and the donor and acceptor fragments of a dative bond A → B are often referred to as Lewis acid and Lewis base. The symbol of an arrow → for a dative bond was introduced by Sidgwick: N. V. Sidgwick, Structure of Divalent Carbon Compounds, *Chem. Rev.*, 1931, **9**, 77–88.
- 4 I. Langmuir, Types of Valence, *Science*, 1921, **54**, 59–67.
- 5 G. N. Lewis, The Atom and the Molecule, *J. Am. Chem. Soc.*, 1916, **38**, 762–785.
- 6 W. Koch, J. R. Collins and G. Frenking, Are there neutral helium compounds which are stable in their ground state? a theoretical investigation of HeBCH and HeBeO, *Chem. Phys. Lett.*, 1986, **132**, 330–333.
- 7 W. Koch, G. Frenking, J. Gauss, D. Cremer and J. R. Collins, Helium chemistry: theoretical predictions and experimental challenge, *J. Am. Chem. Soc.*, 1987, **109**, 5917–5934.
- 8 G. Frenking, W. Koch, J. Gauss and D. Cremer, Stabilities and nature of the attractive interactions in HeBeO, NeBeO, and ArBeO and a comparison with analogs NGLiF, NGBN, and NGLiH (NG = He, Ar). A theoretical investigation, *J. Am. Chem. Soc.*, 1988, **110**, 8007–8016.
- 9 G. Frenking, W. Koch and J. R. Collins, Fixation of nitrogen and carbon monoxide by beryllium oxide: theoretical investigation of the structures and stabilities of NNBeO, OCBeO, and COBeO, *J. Chem. Soc., Chem. Commun.*, 1988, 1147–1148.



- 10 G. D. Purvis, III and R. J. Bartlett, A full coupled-cluster singles and doubles model: the inclusion of disconnected triples, *J. Chem. Phys.*, 1982, **76**, 1910–1918.
- 11 A. D. Becke, Density-functional exchange-energy approximation with correct asymptotic behavior, *Phys. Rev. A: At., Mol., Opt. Phys.*, 1988, **38**, 3098–3100.
- 12 J. P. Perdew, Density-functional approximation for the correlation energy of the inhomogeneous electron gas, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1986, **33**, 8822–8824.
- 13 F. Weigend and R. Ahlrichs, Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: design and assessment of accuracy, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 14 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16 Rev. C.01*, 2016.
- 15 E. D. Glendening, C. R. Landis and F. Weinhold, NBO 7.0: new vistas in localized and delocalized chemical bonding theory, *J. Comput. Chem.*, 2019, **40**, 2234–2241.
- 16 F. L. Hirshfeld, Bonded-atom fragments for describing molecular charge densities, *Theor. Chim. Acta*, 1977, **44**, 129–138.
- 17 A. V. Marenich, S. V. Jerome, C. J. Cramer and D. G. Truhlar, Charge Model 5: An Extension of Hirshfeld Population Analysis for the Accurate Description of Molecular Interactions in Gaseous and Condensed Phases, *J. Chem. Theory Comput.*, 2012, **8**, 527–541.
- 18 C. Fonseca Guerra, J.-W. Handgraaf, E. J. Baerends and F. M. Bickelhaupt, Voronoi deformation density (VDD) charges: assessment of the Mulliken, Bader, Hirshfeld, Weinhold, and VDD methods for charge analysis, *J. Comput. Chem.*, 2004, **25**, 189–210.
- 19 I. Mayer, Charge, bond order and valence in the AB initio SCF theory, *Chem. Phys. Lett.*, 1983, **97**, 270–274.
- 20 T. Lu and F. Chen, Multiwfn: a multifunctional wavefunction analyzer, *J. Comput. Chem.*, 2012, **33**, 580–592.
- 21 K. Morokuma, Molecular Orbital Studies of Hydrogen Bonds. III. C=O...H-O Hydrogen Bond in H₂CO...H₂O and H₂CO...2H₂O, *J. Chem. Phys.*, 2003, **55**, 1236–1244.
- 22 T. Ziegler and A. Rauk, On the calculation of bonding energies by the Hartree Fock Slater method, *Theor. Chim. Acta*, 1977, **46**, 1–10.
- 23 M. Mitoraj and A. Michalak, Donor-Acceptor Properties of Ligands from the Natural Orbitals for Chemical Valence, *Organometallics*, 2007, **26**, 6576–6580.
- 24 M. Mitoraj and A. Michalak, Applications of natural orbitals for chemical valence in a description of bonding in conjugated molecules, *J. Mol. Model.*, 2008, **14**, 681–687.
- 25 A. Michalak, M. Mitoraj and T. Ziegler, Bond Orbitals from Chemical Valence Theory, *J. Phys. Chem. A*, 2008, **112**, 1933–1939.
- 26 M. P. Mitoraj, A. Michalak and T. Ziegler, A Combined Charge and Energy Decomposition Scheme for Bond Analysis, *J. Chem. Theory Comput.*, 2009, **5**, 962–975.
- 27 G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, Chemistry with ADF, *J. Comput. Chem.*, 2001, **22**, 931–967.
- 28 E. J. Baerends, T. Ziegler, A. J. Atkins, J. Autschbach, O. Baseggio, D. Bashford, A. Bérces, F. M. Bickelhaupt, C. Bo, P. M. Boerrigter, C. Cappelli, L. Cavallo, C. Daul, D. P. Chong, D. V. Chulhai, L. Deng, R. M. Dickson, J. M. Dieterich, F. Egidi, D. E. Ellis, M. van Faassen, L. Fan, T. H. Fischer, A. Förster, C. Fonseca Guerra, M. Franchini, A. Ghysels, A. Giammona, S. J. A. van Gisbergen, A. Goez, A. W. Götz, J. A. Groeneveld, O. V. Gritsenko, M. Grüning, S. Gusarov, F. E. Harris, P. van den Hoek, Z. Hu, C. R. Jacob, H. Jacobsen, L. Jensen, L. Joubert, J. W. Kaminski, G. van Kessel, C. König, F. Kootstra, A. Kovalenko, M. V. Krykunov, P. Lafiosca, E. van Lenthe, D. A. McCormack, M. Medves, A. Michalak, M. Mitoraj, S. M. Morton, J. Neugebauer, V. P. Nicu, L. Noodleman, V. P. Osinga, S. Patchkovskii, M. Pavanello, C. A. Peeples, P. H. T. Philipsen, D. Post, C. C. Pye, H. Ramanantoanina, P. Ramos, W. Ravenek, M. Reimann, J. I. Rodríguez, P. Ros, R. Rüger, P. R. T. Schipper, D. Schlüns, H. van Schoot, G. Schreckenbach, J. S. Seldenthuis, M. Seth, J. G. Snijders, M. Solà, M. Stener, M. Swart, D. Swerhone, V. Tognetti, G. te Velde, P. Vernooijs, L. Versluis, L. Visscher, O. Visser, F. Wang, T. A. Wesolowski, E. M. van Wezenbeek, G. Wiesenekker, S. K. Wolff, T. K. Woo and A. L. Yakovlev, *ADF, SCM, Theoretical Chemistry*, Vrije Universiteit, Amsterdam, The Netherlands, 2019, <https://www.scm.com>.
- 29 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu, *J. Chem. Phys.*, 2010, **132**, 154104.
- 30 E. Van Lenthe and E. J. Baerends, Optimized Slater-type basis sets for the elements 1–118, *J. Comput. Chem.*, 2003, **24**, 1142–1156.
- 31 J. C. Slater, in *Adv. Quantum Chem.*, ed. P.-O. Löwdin, Academic Press, 1972, vol. 6, pp. 1–92.
- 32 D. M. Andrada and C. Foroutan-Nejad, Energy components in energy decomposition analysis (EDA) are path functions; why does it matter?, *Phys. Chem. Chem. Phys.*, 2020, **22**, 22459–22464.
- 33 M. Solà, M. Duran and J. Poater, The energy components of the extended transition state energy decomposition analysis



- are path functions: the case of water tetramer, *Theor. Chem. Acc.*, 2021, **140**, 33.
- 34 J. Poater, D. M. Andrada, M. Solà and C. Foroutan-Nejad, Path-dependency of energy decomposition analysis & the elusive nature of bonding, *Phys. Chem. Chem. Phys.*, 2022, **24**, 2344–2348.
 - 35 F. M. Bickelhaupt, C. Fonseca Guerra, M. Mitoraj, F. Sagan, A. Michalak, S. Pan and G. Frenking, Clarifying notes on the bonding analysis adopted by the energy decomposition analysis, *Phys. Chem. Chem. Phys.*, 2022, **24**, 15726–15735.
 - 36 L. Zhao, M. von Hopffgarten, D. M. Andrada and G. Frenking, Energy decomposition analysis, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2018, **8**, e1345.
 - 37 G. Frenking and F. Matthias Bickelhaupt, *The Chemical Bond*, 2014, pp. 121–157.
 - 38 L. Zhao, M. Hermann, N. Holzmann and G. Frenking, Dative bonding in main group compounds, *Coord. Chem. Rev.*, 2017, **344**, 163–204.
 - 39 L. Zhao, M. Hermann, W. H. E. Schwarz and G. Frenking, The Lewis electron-pair bonding model: modern energy decomposition analysis, *Nat. Rev. Chem.*, 2019, **3**, 48–63.
 - 40 L. Zhao, S. Pan and G. Frenking, in *Comprehensive Computational Chemistry*, ed. M. Yáñez and R. J. Boyd, Elsevier, Oxford, 1st edn, 2024, pp. 322–361.
 - 41 G. Frenking, R. Tonner, S. Klein, N. Takagi, T. Shimizu, A. Krapp, K. K. Pandey and P. Parameswaran, New bonding modes of carbon and heavier group 14 atoms Si–Pb, *Chem. Soc. Rev.*, 2014, **43**, 5106–5139.
 - 42 G. Frenking, M. Hermann, D. M. Andrada and N. Holzmann, Donor–acceptor bonding in novel low-coordinated compounds of boron and group-14 atoms C–Sn, *Chem. Soc. Rev.*, 2016, **45**, 1129–1144.
 - 43 A. Altun, M. Saitow, F. Neese and G. Bistoni, Local Energy Decomposition of Open-Shell Molecular Systems in the Domain-Based Local Pair Natural Orbital Coupled Cluster Framework, *J. Chem. Theory Comput.*, 2019, **15**, 1616–1632.
 - 44 A. Altun, R. Izsák and G. Bistoni, Local energy decomposition of coupled-cluster interaction energies: interpretation, benchmarks, and comparison with symmetry-adapted perturbation theory, *Int. J. Quantum Chem.*, 2021, **121**, e26339.
 - 45 W. B. Schneider, G. Bistoni, M. Sparta, M. Saitow, C. Riplinger, A. A. Auer and F. Neese, Decomposition of Intermolecular Interaction Energies within the Local Pair Natural Orbital Coupled Cluster Framework, *J. Chem. Theory Comput.*, 2016, **12**, 4778–4792.
 - 46 S. Saebo and P. Pulay, Local Treatment of Electron Correlation, *Annu. Rev. Phys. Chem.*, 1993, **44**, 213–236.
 - 47 F. Neese, F. Wennmohs and A. Hansen, Efficient and accurate local approximations to coupled-electron pair approaches: an attempt to revive the pair natural orbital method, *J. Chem. Phys.*, 2009, **130**, 114108.
 - 48 F. Neese, A. Hansen and D. G. Liakos, Efficient and accurate approximations to the local coupled cluster singles doubles method using a truncated pair natural orbital basis, *J. Chem. Phys.*, 2009, **131**, 064103.
 - 49 Y. Guo, C. Riplinger, U. Becker, D. G. Liakos, Y. Minenkov, L. Cavallo and F. Neese, Communication: an improved linear scaling perturbative triples correction for the domain based local pair-natural orbital based singles and doubles coupled cluster method [DLPNO-CCSD(T)], *J. Chem. Phys.*, 2018, **148**, 011101.
 - 50 C. Riplinger, B. Sandhoefer, A. Hansen and F. Neese, Natural triple excitations in local coupled cluster calculations with pair natural orbitals, *J. Chem. Phys.*, 2013, **139**, 134101.
 - 51 G. H. K. P. Huber, *Molecular Spectra and Molecular Structure*, Media, New York, 1979.
 - 52 M. Kaupp, P. V. R. Schleyer, H. Stoll and H. Preuss, The question of bending of the alkaline earth dihalides MX₂ (M = beryllium, magnesium, calcium, strontium, barium; X = fluorine, chlorine, bromine, iodine). An ab initio pseudopotential study, *J. Am. Chem. Soc.*, 1991, (113), 6012–6020.
 - 53 M. Hargittai, The molecular geometry of gas-phase metal halides, *Coord. Chem. Rev.*, 1988, **91**, 35–88.
 - 54 The calculated Ae–F bond lengths [Å] in AeF[−] at CCSD(T)/def2-TZVPP [BP86/def2-TZVPP] reported in ref. 1 are 1.429 [1.428] for BeF[−], 1.840 [1.847] for MgF[−], 2.104 [1.992] for CaF[−], 2.237 [2.134] for SrF, 2.339 [2.234] for BaF[−].
 - 55 *Computational Chemistry Comparison and Benchmark Data Base Release 22. Standard Reference Database 101*, National Institute of Standards and Technology, 2022.
 - 56 J. S. Murray and P. Politzer, Molecular electrostatic potentials and noncovalent interactions, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2017, **7**, e1326.
 - 57 R. Tonner, F. Öxler, B. Neumüller, W. Petz and G. Frenking, Carbodiphosphoranes: The Chemistry of Divalent Carbon(0), *Angew. Chem., Int. Ed.*, 2006, **45**, 8038–8042.
 - 58 R. Tonner and G. Frenking, C(NHC)₂: Divalent Carbon(0) Compounds with N-Heterocyclic Carbene Ligands—Theoretical Evidence for a Class of Molecules with Promising Chemical Properties, *Angew. Chem., Int. Ed.*, 2007, **46**, 8695–8698.
 - 59 R. Tonner and G. Frenking, Divalent Carbon(0) Chemistry, Part 1: Parent Compounds, *Chem. – Eur. J.*, 2008, **14**, 3260–3272.
 - 60 R. Tonner and G. Frenking, Divalent Carbon(0) Chemistry, Part 2: Protonation and Complexes with Main Group and Transition Metal Lewis Acids, *Chem. – Eur. J.*, 2008, **14**, 3273–3289.
 - 61 S. Klein, R. Tonner and G. Frenking, Carbodicarbenes and Related Divalent Carbon(0) Compounds, *Chem. – Eur. J.*, 2010, **16**, 10160–10170.
 - 62 J. Hinze, The concept of electronegativity of atoms in molecules, *Theor. Comput. Chem.*, 1999, **6**, 189–212.
 - 63 D. Bergmann and J. Hinze, Electronegativity and Molecular Properties, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 150–163.
 - 64 K. B. Wiberg, Application of the pople-santry-segal CNDO method to the cyclopropylcarbinyl and cyclobutyl cation and to bicyclobutane, *Tetrahedron*, 1968, **24**, 1083–1096.
 - 65 I. Mayer, Bond order and valence: Relations to Mulliken's population analysis, *Int. J. Quantum Chem.*, 1984, **26**, 151–154.
 - 66 A. J. Bridgeman, G. Cavigliasso, L. R. Ireland and J. Rothery, The Mayer bond order as a tool in inorganic chemistry, *J. Chem. Soc., Dalton Trans.*, 2001, 2095–2108.



- 67 I. Mayer, Bond order and valence indices: a personal account, *J. Comput. Chem.*, 2007, **28**, 204–221.
- 68 L. Zhao, S. Pan and G. Frenking, The nature of the polar covalent bond, *J. Chem. Phys.*, 2022, **157**, 034105.
- 69 K. Fukui, *Theory of Orientation and Stereoselection*, Springer Verlag, Berlin, 1975.
- 70 R. B. Woodward and R. Hoffmann, *The Conservation of Orbital Symmetry*, Verlag Chemie, Weinheim, 1970.
- 71 S. Shaik, D. Danovich, W. Wu, P. Su, H. S. Rzepa and P. C. Hiberty, Quadruple bonding in C₂ and analogous eight-valence electron species, *Nat. Chem.*, 2012, **4**, 195–200.
- 72 D. Danovich, P. C. Hiberty, W. Wu, H. S. Rzepa and S. Shaik, The Nature of the Fourth Bond in the Ground State of C₂: The Quadruple Bond Conundrum, *Chem. – Eur. J.*, 2014, **20**, 6220–6232.
- 73 S. Shaik, D. Danovich, B. Braida and P. C. Hiberty, The Quadruple Bonding in C₂ Reproduces the Properties of the Molecule, *Chem. – Eur. J.*, 2016, **22**, 4116–4128.
- 74 W. Zou and D. Cremer, C₂ in a Box: Determining Its Intrinsic Bond Strength for the X¹Σ_g⁺ Ground State, *Chem. – Eur. J.*, 2016, **22**, 4087–4099.
- 75 M. Hermann and G. Frenking, The Chemical Bond in C₂, *Chem. – Eur. J.*, 2016, **22**, 4100–4108.
- 76 M. Piris, X. Lopez and J. M. Ugalde, The Bond Order of C₂ from a Strictly N-Representable Natural Orbital Energy Functional Perspective, *Chem. – Eur. J.*, 2016, **22**, 4109–4115.
- 77 D. L. Cooper, R. Ponc and M. Kohout, New insights from domain-averaged Fermi holes and bond order analysis into the bonding conundrum in C₂, *Mol. Phys.*, 2016, **114**, 1270–1284.
- 78 D. W. O. de Sousa and M. A. C. Nascimento, Is There a Quadruple Bond in C₂?, *J. Chem. Theory Comput.*, 2016, **12**, 2234–2241.
- 79 L. C. Ducati, N. Takagi and G. Frenking, Molecules with All Triple Bonds: OCBBCO, N₂BBN₂, and [OB BBBO]^{2−}, *J. Phys. Chem. A*, 2009, **113**, 11693–11698.
- 80 L. Zhao, W. H. E. Schwarz and G. Frenking, The Lewis electron-pair bonding model: the physical background, one century later, *Nat. Rev. Chem.*, 2019, **3**, 35–47.
- 81 D. M. Andrada, J. L. Casals-Sainz, Á. Martín Pendás and G. Frenking, Dative and Electron-Sharing Bonding in C₂F₄, *Chem. – Eur. J.*, 2018, **24**, 9083–9089.
- 82 S. Pan and G. Frenking, Comment on “Realization of Lewis Basic Sodium Anion in the NaBH₃[−] Cluster”, *Angew. Chem., Int. Ed.*, 2020, **59**, 8756–8759.
- 83 K. Ruedenberg, The Physical Nature of the Chemical Bond, *Rev. Mod. Phys.*, 1962, **34**, 326–376.
- 84 W. Kutzelnigg, The Physical Mechanism of the Chemical Bond, *Angew. Chem., Int. Ed. Engl.*, 1973, **12**, 546–562.
- 85 F. Rioux, The Covalent Bond Examined Using the Virial Theorem, *Chem. Educ.*, 2003, **8**, 101.
- 86 S. Shaik and P. C. Hiberty, *A Chemist's Guide to Valence Bond Theory*, 2007, pp. 40–80.
- 87 L. Pauling, *The Nature of the Chemical Bond*, Cornell University Press, 1960.
- 88 S. Pan and G. Frenking, A Critical Look at Linus Pauling's Influence on the Understanding of Chemical Bonding, *Molecules*, 2021, **26**, 4695.
- 89 G. Frenking, Heretical thoughts about the present understanding and description of the chemical bond*, *Mol. Phys.*, 2022, **121**, e2110168.
- 90 J. H. Kim, X. Li, L.-S. Wang, H. L. de Clercq, C. A. Fancher, O. C. Thomas and K. H. Bowen, Vibrationally Resolved Photoelectron Spectroscopy of MgO[−] and ZnO[−] and the Low-Lying Electronic States of MgO, MgO[−], and ZnO, *J. Phys. Chem. A*, 2001, **105**, 5709–5718.
- 91 J. C. Wheeler, Electron Affinities of the Alkaline Earth Metals and the Sign Convention for Electron Affinity, *J. Chem. Educ.*, 1997, **74**, 123.
- 92 R. Tonner, G. Heydenrych and G. Frenking, First and Second Proton Affinities of Carbon Bases, *Chem. Phys. Chem.*, 2008, **9**, 1474.

