

Cite this: *Chem. Sci.*, 2024, 15, 765

All publication charges for this article have been paid for by the Royal Society of Chemistry

Mn(III)-mediated carbon-centered radicals generate an enhanced immunotherapeutic effect†

Jiaxuan Li,^{‡a} Baifei Hu,^{‡b} Zelong Chen,^{‡a} Jiahui Li,^a Wenjuan Jin,^a Yi Wang,^a Yichen Wan,^a Yinghua Lv,^{id a} Yuxin Pei,^{id *a} Hongtao Liu^{id *b} and Zhichao Pei^{id *a}

A strategy for designing cancer therapeutic nanovaccines based on immunogenic cell death (ICD)-inducing therapeutic modalities is particularly attractive for optimal therapeutic efficacy. In this work, a highly effective cancer therapeutic nanovaccine (denoted as MPL@ICC) based on immunogenic photodynamic therapy (PDT) was rationally designed and fabricated. MPL@ICC was composed of a nanovehicle of MnO₂ modified with a host–guest complex using amino pillar[6]arene and lactose–pyridine, a prodrug of isoniazid (INH), and chlorine e6 (Ce6). The nanovaccine exhibited excellent biosafety, good targeting ability to hepatoma cells and enrichment at tumor sites. Most importantly, it could modulate the tumor microenvironment (TME) to facilitate the existence of Mn(III) and Mn(III)-mediated carbon-centered radical generation with INH released from the prodrug *in situ* to further strengthen ICD. This is the first report on Mn(III)-mediated generation of carbon-centered radicals for successful anti-tumor immunotherapy using ICD, which provides a novel strategy for designing highly efficient cancer therapeutic nanovaccines.

Received 14th July 2023

Accepted 5th December 2023

DOI: 10.1039/d3sc03635a

rsc.li/chemical-science

Introduction

In recent years, cancer nanovaccine-based immunotherapy, due to its potential for high therapeutic efficacy, specificity, and ability to yield a long-term immune-memory effect, has exhibited promise to be one of the most effective ways of eradicating tumors and preventing tumor relapse, and alter the cancer treatment paradigm.¹ Cancer nanovaccines are nanoparticle-based vaccines, which can train and stimulate the immune system of the body to detect and eliminate tumors.² They can either consist of a careful combination of tumor antigens co-delivered with immunostimulatory adjuvants to induce potent T cell-mediated immune responses;³ or they can induce immunogenic cell death (ICD) to boost tumor-associated antigen release, which provides *in situ* tumor vaccination to enhance the anti-tumor T cell response against primary tumor growth.⁴ Research to date has indicated that several therapeutic modalities (such as chemotherapy,⁵ radiotherapy,⁶ phototherapy,⁷ etc.) are capable of eliciting ICD, a special type of cell death that can activate antitumor immune responses.⁸ However, ICD-mediated immunopotential achieved thus far is not sufficient for an ideal therapeutic result due to poor immunogenicity and the immunosuppressive tumor microenvironment (TME).⁹

Considering the unique advantages of ICD-inducing modalities,¹⁰ there is an urgent need to develop strategies to significantly increase their efficacy and durability, such as combining them with immune adjuvants and/or other ICD inducers. Adjuvants are indispensable components of vaccines that stimulate optimal immune responses towards non-replicating, inactivated, and subunit antigens.¹¹ In the search for adjuvants that can effectively activate cellular immunity, manganese oxide nanomaterials (MONs) have gained attention thanks to their unique physical and chemical properties, strengthening immunogenicity, modulation of the intrinsic TME, and excellent biodegradability and biosafety.¹² To date, the role of MONs in nanovaccines has been to induce ICD *via* Fenton-like reactions and ferroptosis.¹³ The most common method is to utilize MnO₂ in chemodynamic therapy (CDT), where it is first reduced to Mn(II) in tumor cells by the high concentration of glutathione (GSH), and then produces reactive oxygen species (ROS) in the form of [•]OH *via* a Mn(II)-mediated Fenton-like reaction. However, Mn(II)-mediated Fenton-like reactions rely on HCO₃[−] and the prevalence of H₂O₂, which is reduced in tumor cells, thus limiting the immunotherapeutic effect of Mn(II).¹⁴ Research has been conducted to mitigate these limitations, such as delivering HCO₃[−] salt into tumor cells to accelerate the generation of [•]OH,¹⁵ or by loading glucose oxidase into MONs to increase intracellular H₂O₂,¹⁶ while still using Mn(II) as the catalyst. Notwithstanding the improvements made on the efficiency of Mn(II), it is known that different manganese valence states among oxides exhibit different levels of toxicity.¹⁷ Compared with Mn(II), Mn(III), which is produced from MnO₂ through a single electron transfer (ET) process, has shown to be a stronger oxidant that can activate various organics

^aCollege of Chemistry & Pharmacy, Northwest A&F University, Yangling, Shaanxi 712100, P. R. China. E-mail: peiyx@nwfau.edu.cn; peizc@nwfau.edu.cn

^bCollege of Basic Medical Sciences, Hubei University of Chinese Medicine, Huangjiahui West Road 16, Wuhan 430065, P. R. China. E-mail: hongtaoliu@hbtcm.edu.cn

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3sc03635a>

‡ These authors contributed equally to this work.

(phenol, fullerene, isoniazid, *etc.*) to generate carbon-centered radicals ($\cdot\text{R}$),¹⁸ which are known to amplify ICD.¹⁹ For instance, an enhanced tumor immunotherapy can be realized *via* the combination of AIPH ($\cdot\text{R}$ producing reagent) and photothermal therapy (PTT) by triggering $\cdot\text{R}$ bursts.²⁰ Although Mn(III) has been identified to be a crucial component in certain reactions in biochemistry and bioinorganic chemistry,²¹ Mn(III)-mediated $\cdot\text{R}$ has not yet been explored in cancer immunotherapy because of the instability of Mn(III) in an intrinsic TME due to its acidity.²²


Photodynamic therapy (PDT), as a non-invasive and rapid tumor-ablation therapy, is one of the best candidates for constructing cancer therapeutic nanovaccines.²³ Tumors treated by PDT release tumor antigens, which act directly *in situ* as a patient-specific tumor vaccine.²⁴ However, the efficacy and immunity elicited by PDT alone are severely limited by the immunosuppressive TME, including the high concentration of GSH and the hypoxic environment.²⁵ Furthermore, the increased severity of hypoxia after PDT treatment, as a result of the photodynamic reactions, not only results in increased immunosuppression and impaired efficacy, but also causes rapid tumor recurrence.²⁶ Since MONs have the potential to reverse the immunosuppressive nature of the TME by increasing the pH, exhausting H₂O₂/GSH, and generating O₂,^{17a} there is potential to combine PDT with MON-based platforms to improve ICD.

Taken together, we hypothesize that by rationally designing a MON-based nanoplatform utilizing MONs that would modulate

the TME to improve the PDT induced ICD efficacy, as well as maximizing Mn(III)-mediated $\cdot\text{R}$ to further strengthen ICD, a highly effective cancer therapeutic nanovaccine can be achieved. Hence, as a proof of concept, we propose a cancer therapeutic nanovaccine (denoted as MPL@ICC) composed of an amino pillar[6]arene $\supset$ lactose-pyridine modified hollow MnO₂ nanovehicle, INH-CA, a prodrug consisting of isoniazid (INH) protected by cinnamic aldehyde (CA), and chlorine e6 (Ce6), a photosensitizer (Scheme 1a). INH was chosen as it is known to generate $\cdot\text{R}$ when it reacts with Mn(III)^{18c,27} and is already an FDA-approved drug commonly used against tuberculosis.²⁸ The use of INH in its prodrug state INH-CA should prevent the reaction between INH and MnO₂ before MPL@ICC enters tumor cells. After MPL@ICC enters tumor cells, MnO₂ will consume GSH and convert H₂O₂ to O₂, which triggers the release of INH-CA as well as Ce6. The acidic TME subsequently facilitates the rapid release of INH from INH-CA. INH and Ce6 then, together enhance the effect of PDT to induce ICD (Scheme 1b). In addition, the consumption of GSH and H₂O₂ by MnO₂ will increase the pH, which alters the TME to be more conducive for the existence of Mn(III). Subsequently, Mn(III) facilitates the generation of $\cdot\text{R}$ from INH to further potentiate immunogenic death of tumor cells.

Results and discussion

Before constructing the proposed nanovaccine, it was first necessary to confirm that the reaction between MnO_2 and INH



Scheme 1 (a) Scheme of the synthesis of the MPL@ICC nanovaccine. (b) Scheme of the nanovaccine used to induce ICD to elicit an anti-tumor immune response.




Fig. 1 (a) (b) UV-vis absorption spectra of MB after degradation in different solutions. (c) Mn 3s and (d) Mn 2p spectra of MnO₂ and MnO₂ + INH. MB: methylene blue; INH: isoniazid.

can produce $\cdot R$ through the Mn(III) intermediate. Theoretically, the spatial difference in the d_{z^2} and $d_{x^2-y^2}$ orbitals of Mn(IV) allows for a single electron transfer process during the reduction of Mn(IV).^{22b,29} Although early studies indicated that MnO₂ can react with INH to produce $\cdot OH$ in tumor cells through non-Fenton-like reactions,^{14b} a recent study demonstrated that in a buffer solution at pH 7.4, $\cdot R$ can also be generated in the same reaction.²⁷ It is therefore reasonable to hypothesize the formation of a Mn(III) intermediate in the reaction between MnO₂ and INH, which mediates the generation of $\cdot R$. To confirm this, the free radical production in different solutions was determined using methylene blue (MB) as a probe (Fig. 1a). The degradation of MB caused only by INH is negligible in the pH range from 6.5 to 7.4 (Fig. S1†). While clear absorption decreases were observed in the solutions containing H₂O₂, interestingly, the absorption decreased slightly more when the pH rose from 6.5 to 7.4, which is less conducive to Mn(II)-mediated Fenton-like reactions, indicating another free radical production mechanism that requires a more basic condition is involved. Furthermore, the elimination of H₂O₂, which is crucial for Mn(II)-mediated Fenton-like reactions, resulted in the lowest absorption and thus the most free radical production, with the maximum production observed at pH 7.4.

Given that previous studies indicated that Mn(III) can react with INH to produce $\cdot R$ at pH 7.4, MnO₂ was replaced with Mn(II)Cl₂, Mn(III)(OAc)₃ and the precipitate that was collected after the reaction of MnO₂ with INH (denoted as MnO_x) to verify the reason for the MB oxidation observed under conditions not conducive to Mn(II)-mediated Fenton-like reactions (Fig. 1b). As expected, no significant decrease in MB absorbance was observed when Mn(II) was combined with INH at pH 7.4; when Mn(III) was combined with INH at pH 7.4, the MB absorbance completely disappeared, and an obvious decrease in absorption was observed when MnO_x was combined with INH at pH 7.4, further confirming the existence of a separate oxidation mechanism from the well-known Mn(II)-mediated Fenton-like reaction. Next, XPS was used to examine the precipitate that was

collected after the reaction of MnO₂ with INH (Fig. 1c and d) in various solutions. The binding energy difference $\Delta E = 5.1$ eV in the Mn 3s spectrum of the precipitate indicates that the valence state of Mn was almost identical to that of MnOOH previously reported.³⁰ In the spectra of Mn 2p, the peaks located at 643.28 and 654.98 eV are attributed to the Mn 2p_{3/2} and Mn 2p_{1/2} for the MnO₂, which is in good agreement with the previous report on MnO₂.^{31a} After reacting with INH, the two peaks shifted to 641.88 and 653.68 eV, respectively, which agrees well with that reported for MnOOH.^{31b} This verifies the formation of Mn(III) from MnO₂ in the solutions, possibly through the ET process.^{29,32} Having confirmed the presence of Mn(III) in the reactions, the $\cdot R$ generation of MnO₂ and INH under different conditions was studied by electron spin resonance (ESR) using 3,4-dihydro-2,3-dimethyl-2H-pyrrole 1-oxide (DMPO) (Fig. S2†). Although the generation of $\cdot OH$ from the Fenton-like reaction was observed at varying degrees in all but one solution (MnO₂, INH, pH 6.5) and control, no significant $\cdot R$ signal was observed. The difficulty in detecting $\cdot R$ through ESR is likely due to its instability, which leads to a low steady-state concentration that is difficult to be trapped by DMPO. However, the presence of $\cdot R$ self-quenching products was detected after the reaction between MnO₂ and INH using HRMS (Fig. S3 and S4†), which indicates the formation of $\cdot R$ in this reaction. These combined results strongly suggest that INH reduces Mn(IV) to Mn(III), and then Mn(III) catalyzes INH to generate carbon-centered free radicals, although the oxidized product of INH in the reduction step of Mn(IV) to Mn(III) by INH is not clear since it is very difficult to distinguish from that in the step of oxidation of INH by Mn(III) (Fig. S5†). This would provide an alternative and complementary pathway to Mn(II)-mediated Fenton-like reactions for MnO₂ to generate free radicals and more effectively kill tumor cells *via* ICD.

Having determined the potential of Mn(III) to generate $\cdot R$ for inducing ICD, MPL@ICC was fabricated. First, all the required compounds (amino pillar[6]arene (P), lactose-pyridine (L) and INH-CA) were synthesized according to the method as described in the ESI† and characterized by NMR and HRMS (Fig. S6–S22†). Thereafter, the host-guest complexation between amino pillar[6]arene and lactose-pyridine was investigated by ¹H NMR. The chemical shift of pyridine protons clearly shifted to a high field due to the shielding effect of amino pillar[6]arene (Fig. S23†), which indicated the formation of a pillar[6]arene ⊃ lactose-pyridine complex (PL). Using a Job's plot method, the stoichiometry of the PL complexation was determined to be 1 : 1 for amino pillar[6]arene ⊃ lactose-pyridine (Fig. S24†).

HMnO₂ was prepared according to a previously reported method.^{12a,33} Transmission electron microscopy (TEM) images (Fig. 2a) clearly revealed the hollow structure of HMnO₂ and HMnO₂ modified with PL (MPL). The particle size of MPL (~170 nm) was observed to be significantly larger than that of HMnO₂ (~127 nm) due to the modification of PL onto the surface of HMnO₂, while the cavity sizes remained largely the same. Further, the hydrated particle size distribution of the different nanoparticles was analysed by dynamic light scattering (DLS). The HMnO₂ obtained after Na₂CO₃ etching had the largest particle size (average size = 1088 nm, PDI = 0.798), indicating






Fig. 3 (a) Confocal fluorescent microscopic images of different cells treated with MPL@ICC for 3 h (1×10^5 cells per dish, Hoechst 33258: Ex: 346 nm/Em: 460 nm; Ce6: Ex: 660 nm/Em: 710 nm). Scale bar: 30 μm. (b) The hypoxia detection of HepG2 cells cultured under different conditions for 3 h (1×10^5 cells per dish, pimonidazole as a hypoxia probe, FITC-labeled anti-pimonidazole antibody shows fluorescence. Ex: 488 nm/Em: 530 nm). Scale bar: 30 μm. (c) The mean fluorescence intensity of (a) ($n = 3$). (d) The mean fluorescence intensity of (b). (e) GSH content in HepG2 cells (1×10^7) after being cultured with MPL for different times ($n = 3$). Data are expressed as means $\pm$ SD.

strongest green fluorescence. Subsequently, the standard MTT assay was used to detect cell viability under different treatments. No decrease in cell viability was observed after co-incubating MPL with HL7702 cells, while the viability of HepG2 cells under the same conditions slightly decreased due to Mn(II)-mediated Fenton-like reactions (Fig. S30[†]). Neither free Ce6 nor MPL@Ce6 causes obvious damage to HepG2 cells

in the dark. In contrast, a rapid decrease of HepG2 cell viability with increasing concentration of MPL@ICC was observed (Fig. 4c). After light irradiation, free Ce6 exhibited strong tumor cytotoxicity. Compared with free Ce6, MPL@Ce6 + light treatment showed higher cytotoxicity likely due to the enhanced cellular uptake of MPL@Ce6 and disruption of the redox balance caused by GSH depletion. As expected, the combined




Fig. 4 (a) The oxidative stress in H22 cells with or without 660 nm light (1×10^5 cells per dish, DCF: Ex: 495 nm/Em: 530 nm). Scale bar: 50 μ m. (b) The mean fluorescence intensity of (a) ($n = 3$). (c) The relative cell viabilities of HepG2 cells after incubation with free Ce6, MPL@Ce6, and MPL@ICC at various doses in the dark ($n = 5$, 5×10^3 cells per well). (d) The relative cell viabilities of HepG2 cells after incubation with free Ce6, MPL@Ce6, and MPL@ICC at various doses in light (660 nm, 5 mW cm^{-2} , 30 min, $n = 5$, 5×10^3 cells per well). Data are expressed as means $\pm$ SD.

therapeutic effect resulted in the death of nearly all tumor cells after exposure to light in the presence of low concentrations of MPL@ICC (Fig. 4d). Next, MPL@ICC was co-incubated with HL7702 cells for 24 h, and the results revealed that cytotoxicity was only observed in the presence of high concentrations of the nanoparticles (Fig. S31†).

Building upon the positive results above, the potential enhancement of MPL@ICC on the induction of ICD was investigated. Three effectors of ICD,³⁵ calreticulin (CRT) exposure, high mobility group box-1 (HMGB-1) release, and ATP release, were evaluated in H22 tumor cells following different treatments. In the treatment groups without light irradiation, there was no obvious CRT exposure after 3 h incubation due to insufficient free radical production (Fig. 5a). After 24 h, CRT exposure to cell membranes was observed in the MPL@IC and MPL@ICC treatment groups in contrast to the MPL treatment group. Given the lack of PDT, CRT exposure can only be attributed to ICD induced by Mn(III) mediated $\cdot$ R. In the treatment groups exposed to 3 h of light irradiation, CRT exposure

was the weakest on the membrane of cells treated with free Ce6, which can be attributed to PDT alone. Cells treated with MPL@Ce6 displayed stronger CRT exposure, likely due to the enhanced cellular uptake and disruption of cellular redox balance by MnO_2 through Mn(II)-mediated Fenton-like reactions. The even stronger CRT exposure exhibited by the MPL@ICC treatment can therefore only be attributed to the additional Mn(III)-mediated $\cdot$ R generation, which further disrupts the cellular redox balance. The extracellular contents of HMGB-1 and ATP were examined and showed a similar trend to CRT expression (Fig. 5b and c, respectively). Based on all the results thus far, it can be concluded that MPL@ICC can successfully target hepatoma cells *in vitro* and induce a more robust ICD response by enhancing the PDT efficacy and generating $\cdot$ R.

Tumor cell rechallenge is the gold standard for detecting the ICD effect³⁶ *in vivo*. Therefore, to demonstrate the efficacy of MPL@ICC, we first treated H22 cells with MPL@ICC + light to obtain ICD cells. We then used these ICD cells as an immune






Fig. 5 (a) Calreticulin (CRT) exposure on the surface of H22 cells. (Anti-CRT antibody and FITC-labelled secondary antibody are used to label CRT. Hoechst 33258: Ex: 346 nm/Em: 460 nm; FITC: Ex: 488 nm/Em: 530 nm. scale bar: 50 μ m.). (b) High Mobility Group Box-1 (HMGB-1) and (c) ATP release from H22 tumor cells after various treatments ($n = 3$). (I: PBS, II: MPL, III: MPL@IC, IV: MPL@ICC, V: free Ce6 + light, VI: MPL@Ce6 + light, VII: MPL@ICC + light). (d) The scheme of mice vaccinated. (e) Survival time of mice after being rechallenged by tumor cells ($n = 6$). (f) The photo of fixed tumor tissues (below: necrotic cells and above: ICD cells). Data are expressed as means $\pm$ SD. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$.

vaccine and injected them into the right hindlimb of mice, while untreated necrotic H22 cells, obtained by freeze-thaw cycles, were inoculated into the right hindlimb of another group of mice as a control group. After 7 days, normal H22 cells were re-inoculated into the left hindlimb of mice and tumor growth was monitored (Fig. 5d). Compared with the control group, the survival time and the number of surviving mice in the test group significantly improved (Fig. 5e and f). These results confirmed that ICD obtained *via* treatment with MPL@ICC + light could

induce an immune response in mice and thus MPL@ICC can function as an effective tumor vaccine.

Furthermore, an *in vivo* imaging system was used to study the drug distribution of the nanovaccine in H22 tumor-bearing mice at different times after tail vein injection. The fluorescence of Ce6 gradually increased at the tumor site and reached a peak around 12 h, which could still be observed even after 48 h (Fig. 6a and b). 12 h and 24 h after vein injection, the tumor and main organs were collected to observe the Ce6 fluorescence




Fig. 6 (a) Real-time fluorescence images of H22 tumor-bearing mice after i.v. of MPL@ICC (fluorescence is attributed to Ce6, imaging mode: Ex: 675 nm/Em: Cy 5.5). (b) The corresponding mean fluorescence intensity of (a) ($n = 3$). (c) Ex vivo fluorescence images of major organs and tumors at 12 h and 24 h after intravenous injection of MPL@ICC (fluorescence is attributed to Ce6, imaging mode: Ex: 675 nm/Em: Cy 5.5). (d) The corresponding mean fluorescence intensity of (c) ($n = 3$). H: heart, Li: liver, S: spleen, L: lung, K: kidney, T: tumor. (e) Representative immunofluorescence images of the hypoxia area in tumors 12 h post injection of PBS, free drug, and MPL@ICC (DAPI: Ex: 360 nm/Em: 460 nm. Pimonidazole as a hypoxia probe, FITC-labeled anti-pimonidazole antibody shows fluorescence. Ex: 488 nm/Em: 530 nm.). Scale bar: 100 μ m. (f) The percentage of the hypoxia positive area in tumors ($n = 3$). (g) pH of tumor tissue 24 h post injection of PBS, MPL, and MPL@ICC ($n = 3$). Data are expressed as means $\pm$ SD. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$.

signal. MPL@ICC had the highest mean fluorescence intensity in tumors 12 h after intravenous injection (Fig. 6c and d). All these results suggest that MPL@ICC exhibits increased tumor accumulation, which can be attributed to the EPR effect and targeting ability of MPL@ICC. Subsequently, a hypoxia probe was used to assess hypoxia relief at the tumor site 12 h after vein injection of the nanoparticles (Fig. 6e and f). The intravenously

injected MPL@ICC group showed fewer hypoxia areas than the other groups, suggesting that MnO_2 , which catalyses the decomposition of H_2O_2 to release O_2 , alleviated hypoxia *in vivo*. The pH of tumor tissues was determined 24 h after intravenous injection of different nanoparticles. It was found that MPL and MPL@ICC upregulated the pH of tumor tissues (Fig. 6g).






Fig. 7 (a) Tumor growth curves during the treatment and (b) average tumor weight after various treatments ($n = 5$). n.s.: not significant, $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$. (c) Body weight change curves of mice during various treatments ($n = 5$). (d) The photo of the tumor at the end of treatment (red circles represent no tumor). (e) H&E staining of tumor slices after various treatments. (f) TUNEL staining of tumor slices after various treatments (nucleus stained with DAPI; Ex: 360 nm/Em: 460 nm. FITC-dUTP: Ex: 488 nm/Em: 530 nm). Scale bar: 100 μm data are expressed as means $\pm$ SD.

The therapeutic effects of nanoparticles were then evaluated in H22 tumor-bearing mice (Fig. 7a, b and 7d). When the tumor reached approximately 100–200 mm^3 , H22 bearing mice were randomly divided into seven groups: PBS, MPL, free drug (Ce6 and INH-CA) + light, MPL@IC, MPL@ICC, MPL@Ce6 + light, MPL@ICC + light ($\text{MnO}_2 = 9 \text{ mg kg}^{-1}$, Ce6 = 7.8 mg kg^{-1} , and INH-CA = 4.1 mg kg^{-1}). 660 nm light irradiation (0.4 W cm^{-2}) was applied at the tumor site for 5 min after 12 h injection. The MPL group exhibited a tumor suppressing effect consistent with previous studies. The tumor inhibition was significantly enhanced in the MPL@IC and MPL@ICC groups, which proves that Mn(III) mediated free radical production has a stronger antitumor effect than Mn(II) mediated CDT. Compared with the PBS group, the tumors in the free drug + light treated and the MPL@Ce6 + light treated showed rapid ablation after light therapy. Unfortunately, tumor recurrence occurred in both groups of mice, and was more rapid for the free drug + light group (Fig. 7a). MPL@ICC + light treatment completely eradicated the tumors, which may be attributed to the combined

effects of PDT and free radicals (Fig. 7d and S32[†]). H&E and TUNEL staining assays were performed after various treatments. The result is given in Fig. 7e, which shows that destructive cell necrosis occurred in the group treated with MPL@Ce6 + light irradiation. Only a few apoptotic cells were observed in the light-treated free drug, MPL@IC, and the unlighted MPL@ICC groups. All these values were higher than the control and MPL groups, which did not show significant cellular damage. The results of TUNEL staining were consistent with the results of H&E staining (Fig. 7f). The biosafety of MPL@ICC was assessed by body weight changes and H&E staining of major organs. After various treatments, the body weight of the mice did not change significantly, and no obvious damage was observed in the major organs, indicating that MPL@ICC exhibits good biosafety (Fig. 7c and S33[†]).

Previous studies have shown that the hypoxic TME promotes the recruitment of regulatory T cells (Treg) to tumor sites and activates M2 macrophages to induce immunosuppression.^{9,27,37} Considering that MPL@ICC can alleviate hypoxia and induce




Fig. 8 (a) Representative flow cytometry charts of cytotoxic T lymphocytes (CTLs, CD3⁺CD8⁺) in tumors. (b) Representative flow cytometry charts of regulatory T (Treg, CD25⁺FoxP3⁺) in CD4⁺ cells in tumors. (c) and (d) Quantification of CTLs and Treg in tumors analyzed by flow cytometry, respectively. Each group represents three independent mice ($n = 3$). (e) Quantification of M1 and M2 in tumors analyzed by immunofluorescence. Each group represents three independent mice ($n = 3$). (f) and (g) INF- γ and TNF- α in tumor tissue detected with RT-PCR (GAPDH as an internal reference gene), respectively. (h) The content of IL-6 in the blood of mice. I: PBS, II: MPL, III: MPL@IC, IV: MPL@ICC, V: free drug + light, VI: MPL@Ce6 + light, VII: MPL@ICC + light. Data are expressed as means $\pm$ SD. n.s.: not significant, * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. INF- γ : interferon- γ ; TNF- α : tumor necrosis factor- α ; IL-6: interleukin-6; GAPDH: glyceraldehyde-3-phosphate dehydrogenase.

TAA release from ICD tumor cells, the role of the nanovaccine in overcoming TME immune suppression and inducing systemic immune responses in mice was investigated. Tumors were collected on day 5 after the different treatments and the proportions of T cells were analysed by flow cytometry (Fig. 8a–d). There was only a small increase in cytotoxic T lymphocytes (CTLs) at the tumor site in mice treated with MPL (group II, about 0.45%) compared with the PBS group (group I, about

0.23%) (Fig. 8c). Notably, the tumors of the mice treated with MPL@IC (group III, about 1.10%) and MPL@ICC (group IV, about 1.16%) exhibited greater infiltration of CTL cells than that with MPL. These results suggest that Mn(III) mediated free radical production is more effective than Mn(II) mediated CDT in activating the immune system. After free drug + light (group V) treatment, CTLs in tumors were significantly increased (about 1.77%), indicating the potent role of PDT in inducing



- Q. Wang, H. Kam, B. Xie, S. Lee, G. Chen and R. Wang, *ACS Nano*, 2023, **17**, 4034–4049.
- 2 (a) L. Qin, H. Zhang, Y. Zhou, C. S. Umeshappa and H. Gao, *Small*, 2021, **17**, e2006000; (b) J. Pan, Y. Wang, C. Zhang, X. Wang, H. Wang, J. Wang, Y. Yuan, X. Wang, X. Zhang, C. Yu, S. K. Sun and X. P. Yan, *Adv. Mater.*, 2018, **30**, 1704408.
- 3 (a) M. A. Islam, J. Rice, E. Reesor, H. Zope, W. Tao, M. Lim, J. X. Ding, Y. H. Chen, D. Aduluso, B. R. Zetter, O. C. Farokhzad and J. J. Shi, *Biomaterials*, 2021, **266**, 120431; (b) G. Z. Zhu, F. W. Zhang, Q. Q. Ni, G. Niu and X. Y. Chen, *ACS Nano*, 2017, **11**, 2387–2392.
- 4 (a) Y. H. Wang, Q. F. Zhao, B. Y. Zhao, Y. S. Zheng, Q. Y. Zhuang, N. S. Liao, P. Y. Wang, Z. X. Cai, D. Zhang, Y. Y. Zeng and X. L. Liu, *Adv. Sci.*, 2022, **9**, 2105631; (b) X. Yang, Y. Yang, J. Y. Bian, J. J. Wei, Z. Wang, Z. W. Zhou, Z. T. Li and M. J. Sun, *Nano Today*, 2021, **38**, 101109.
- 5 (a) H. Inoue and K. Tani, *Cell Death Differ.*, 2014, **21**, 39–49; (b) X. Zhao, K. Yang, R. Zhao, T. Ji, X. Wang, X. Yang, Y. Zhang, K. Cheng, S. Liu, J. Hao, H. Ren, K. W. Leong and G. Nie, *Biomaterials*, 2016, **102**, 187–197; (c) D. W. Zheng, J. L. Chen, J. Y. Zhu, L. Rong, B. Li, Q. Lei, J. X. Fan, M. Z. Zou, C. Li, S. X. Cheng, Z. Xu and X. Z. Zhang, *Nano Lett.*, 2016, **16**, 4341–4347.
- 6 (a) S. C. Formenti and S. Demaria, *J. Natl. Cancer Inst.*, 2013, **105**, 256–265; (b) Y. Min, K. C. Roche, S. Tian, M. J. Eblan, K. P. McKinnon, J. M. Caster, S. Chai, L. E. Herring, L. Zhang, T. Zhang, J. M. DeSimone, J. E. Tepper, B. G. Vincent, J. S. Serody and A. Z. Wang, *Nat. Nanotechnol.*, 2017, **12**, 877–882; (c) Z. Huang, Y. Wang, D. Yao, J. Wu, Y. Hu and A. Yuan, *Nat. Commun.*, 2021, **12**, 145.
- 7 (a) G. Lan, K. Ni, Z. Xu, S. S. Veroneau, Y. Song and W. Lin, *J. Am. Chem. Soc.*, 2018, **140**, 5670–5673; (b) L. Li, S. Yang, L. Song, Y. Zeng, T. He, N. Wang, C. Yu, T. Yin, L. Liu, X. Wei, Q. Wu, Y. Wei, L. Yang and C. Gong, *Theranostics*, 2018, **8**, 860–873; (c) M. Li, R. Guo, J. Wei, M. Deng, J. Li, Y. Tao, M. Li and Q. He, *Acta Biomater.*, 2021, **136**, 546–557; (d) C. Gao, Q. Wang, J. Li, C. H. T. Kwong, J. Wei, B. Xie, S. Lu, S. M. Y. Lee and R. Wang, *Sci. Adv.*, 2022, **8**, eabn1805.
- 8 (a) T. L. Aaes and P. Vandenabeele, *Cell Death Differ.*, 2021, **28**, 843–860; (b) R. D. Mule, A. Kumar, S. P. Sancheti, B. Senthilkumar, H. Kumar and N. T. Patil, *Chem. Sci.*, 2022, **13**, 10779–10785; (c) S. Sen, M. Won, M. S. Levine, Y. Noh, A. C. Sedgwick, J. S. Kim, J. L. Sessler and J. F. Arambula, *Chem. Soc. Rev.*, 2022, **51**, 1212–1233.
- 9 (a) S. D. Jo, G. H. Nam, G. Kwak, Y. Yang and I. C. Kwon, *Nano Today*, 2017, **17**, 23–37; (b) S. K. Daniel, K. M. Sullivan, K. P. Labadie and V. G. Pillarisetty, *Clin. Transl. Med.*, 2019, **8**, e10; (c) A. Facciabene, X. Peng, I. S. Hagemann, K. Balint, A. Barchetti, L. P. Wang, P. A. Gimotty, C. B. Gilks, P. Lal, L. Zhang and G. Coukos, *Nature*, 2011, **475**, 226–230.
- 10 (a) Y. Fan and J. J. Moon, *Vaccines*, 2015, **3**, 662–685; (b) S. Zhang, J. Wang, Z. Kong, X. Sun, Z. He, B. Sun, C. Luo and J. Sun, *Biomaterials*, 2022, **282**, 121433; (c) J. Zhou, L. Rao, G. Yu, T. R. Cook, X. Chen and F. Huang, *Chem. Soc. Rev.*, 2021, **50**, 2839–2891.
- 11 (a) M. Singh and D. O'Hagan, *Nat. Biotechnol.*, 1999, **17**, 1075–1081; (b) B. Pulendran, P. S. Arunachalam and D. T. O'Hagan, *Nat. Rev. Drug Discovery*, 2021, **20**, 454–475; (c) W. Lee and M. Suresh, *Front. Immunol.*, 2022, **13**, 940047.
- 12 (a) G. Yang, L. Xu, Y. Chao, J. Xu, X. Sun, Y. Wu, R. Peng and Z. Liu, *Nat. Commun.*, 2017, **8**, 902; (b) R. Liang, L. Liu, H. He, Z. Chen, Z. Han, Z. Luo, Z. Wu, M. Zheng, Y. Ma and L. Cai, *Biomaterials*, 2018, **177**, 149–160; (c) L. Hou, C. Tian, Y. Yan, L. Zhang, H. Zhang and Z. Zhang, *ACS Nano*, 2020, **14**, 3927–3940; (d) X. Chen, S. S. Gao, X. Wang, M. Y. Guo, Y. Y. Cui, Z. W. Chen, Y. Liu and Y. L. Wang, *Nano Today*, 2022, **46**, 101583; (e) H. Wang, Y. Liu, C. Xu, X. Wang, G. R. Chen, T. D. James, Y. Zang, J. Li, X. Ma and X. P. He, *Chem. Commun.*, 2018, **54**, 4037–4040.
- 13 (a) B. Ding, P. Zheng, F. Jiang, Y. Zhao, M. Wang, M. Chang, P. Ma and J. Lin, *Angew. Chem., Int. Ed.*, 2020, **59**, 16381–16384; (b) Y. Yang, Z. Gu, J. Tang, M. Zhang, Y. Yang, H. Song and C. Yu, *Adv. Sci.*, 2021, **8**, 2002667; (c) B. Ding, J. Yue, P. Zheng, P. Ma and J. Lin, *J. Mater. Chem. B*, 2021, **9**, 7117–7131; (d) X. J. Di, Z. C. Pei, Y. X. Pei and T. D. James, *Coord. Chem. Rev.*, 2023, **484**, 215098.
- 14 (a) E. Ember, S. Rothbart, R. Puchta and R. van Eldik, *New J. Chem.*, 2009, **33**, 34–49; (b) Y. Cheng, F. Yang, K. Zhang, Y. Zhang, Y. Cao, C. Liu, H. Lu, H. Dong and X. Zhang, *Adv. Funct. Mater.*, 2019, **29**, 1903850.
- 15 Y. Feng, Y. Liu, X. Ma, L. Xu, D. Ding, L. Chen, Z. Wang, R. Qin, W. Sun and H. Chen, *J. Nanobiotechnol.*, 2022, **20**, 193.
- 16 B. B. Ding, P. Zheng, J. Tan, H. Chen, Y. L. Bian, C. Z. Yang, P. A. Ma and J. Lin, *Chem. Mater.*, 2022, **34**, 1800–1808.
- 17 (a) B. Ding, P. Zheng, P. a. Ma and J. Lin, *Adv. Mater.*, 2020, **32**, 1905823; (b) J. Xu, W. Han, P. Yang, T. Jia, S. Dong, H. Bi, A. Gulzar, D. Yang, S. Gai, F. He, J. Lin and C. Li, *Adv. Funct. Mater.*, 2018, **28**, 1803804; (c) L. Tian, Q. Chen, X. Yi, J. Chen, C. Liang, Y. Chao, K. Yang and Z. Liu, *Small*, 2017, **13**, 1700640.
- 18 (a) B. B. Snider and R. B. Smith, *Tetrahedron*, 2002, **58**, 25–34; (b) G. W. Wang, T. H. Zhang, X. Cheng and F. Wang, *Org. Biomol. Chem.*, 2004, **2**, 1160–1163; (c) L. Qin, C. H. Huang, D. Xu, L. N. Xie, J. Shao, L. Mao, B. Kalyanaraman and B. Z. Zhu, *Free Radical Biol. Med.*, 2019, **143**, 232–239.
- 19 (a) W. Wu, Y. Yang, Z. Liang, X. Song, Y. Huang, L. Qiu, X. Qiu, S. Yu and W. Xue, *Nanoscale*, 2021, **13**, 11169–11187; (b) Y. Zhang, Q. Wang, Y. Ji, L. Fan, B. Ding, J. Lin and L. Wang, *Chem. Eng. J.*, 2022, **435**, 134869.
- 20 B. Ning, Y. Liu, B. Ouyang, X. Su, H. Guo, Z. Pang and S. Shen, *J. Colloid Interface Sci.*, 2022, **614**, 436–450.
- 21 (a) S. M. Webb, G. J. Dick, J. R. Bargar and B. M. Tebo, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 5558–5563; (b) A. S. Madison, B. M. Tebo, A. Mucci, B. Sundby and G. W. Luther 3rd, *Science*, 2013, **341**, 875–878.
- 22 (a) H. Lin, N. H. Szeinbaum, T. J. DiChristina and M. Taillefert, *Geochim. Cosmochim. Acta*, 2012, **99**, 179–192; (b) E. Hu, Y. Zhang, S. Wu, J. Wu, L. Liang and F. He, *Water Res.*, 2017, **111**, 234–243.



- 23 (a) M. E. Aikins, C. Xu and J. J. Moon, *Acc. Chem. Res.*, 2020, **53**, 2094–2105; (b) L. Zhang, K. Wang, Y. Huang, H. Zhang, L. Zhou, A. Li and Y. Sun, *Biomaterials*, 2022, **282**, 121411; (c) W. Qian, M. Zuo, G. Sun, Y. Chen, T. Han, X. Y. Hu, R. Wang and L. Wang, *Chem. Commun.*, 2020, **56**, 7301–7304.
- 24 (a) S. Liu, G. Feng, B. Z. Tang and B. Liu, *Chem. Sci.*, 2021, **12**, 6488–6506; (b) X. Li, Y. Zhang, X. Wu, J. Chen, M. Yang, F. Ma and L. Shi, *Small*, 2022, **18**, 2203100.
- 25 (a) W. Feng, S. Zhang, Y. Wan, Z. Chen, Y. Qu, J. Li, T. D. James, Z. Pei and Y. Pei, *ACS Appl. Mater. Interfaces*, 2022, **14**, 20749–20761; (b) H. Zhu, Q. Li, B. Shi, F. Ge, Y. Liu, Z. Mao, H. Zhu, S. Wang, G. Yu, F. Huang and P. J. Stang, *Angew. Chem., Int. Ed.*, 2020, **59**, 20208; (c) N. Song, Z. Zhang, P. Liu, D. Dai, C. Chen, Y. Li, L. Wang, T. Han, Y. W. Yang, D. Wang and B. Z. Tang, *Adv. Funct. Mater.*, 2021, **31**, 2009924.
- 26 (a) W. R. Wilson and M. P. Hay, *Nat. Rev. Cancer*, 2011, **11**, 393–410; (b) S. Chouaib, M. Z. Noman, K. Kosmatopoulos and M. A. Curran, *Oncogene*, 2017, **36**, 439–445; (c) C. Gao, C. H. T. Kwong, Q. Wang, H. Kam, J. Wei, Q. Chen, J. Zhang, S. M. Y. Lee, D. Gu and R. Wang, *Mater. Today*, 2022, **58**, 57–70.
- 27 L. Qin, C. H. Huang, L. Mao, B. Shao and B. Z. Zhu, *Free Radical Biol. Med.*, 2020, **154**, 1–8.
- 28 C. Vilchèze and W. R. Jacobs, *Antimicrob. Agents Chemother.*, 2012, **56**, 5142–5148.
- 29 G. W. Luther, *Geomicrobiol. J.*, 2005, **22**, 195–203.
- 30 (a) Y. Zhao, X. Zhang, C. Jia, J. Wu, H. Tang, J. Shang, C. Yuan, Y. Wang and P. Zhang, *Colloids Surf., B*, 2021, **201**, 111627; (b) E. S. Ilton, J. E. Post, P. J. Heaney, F. T. Ling and S. N. Kerisit, *Appl. Surf. Sci.*, 2016, **366**, 475–485.
- 31 (a) M. Li, J. Yu, X. Wang and Z. Yang, *Appl. Surf. Sci.*, 2020, **530**, 147230; (b) S. Yang, X. Song, P. Zhang and L. Gao, *J. Mater. Chem. A*, 2015, **3**, 6136–6145.
- 32 (a) J. E. Johnson, P. Savalia, R. Davis, B. D. Kocar, S. M. Webb, K. H. Nealson and W. W. Fischer, *Environ. Sci. Technol.*, 2016, **50**, 4248–4258; (b) P. Wuamprakhon, A. Krittayavathananon, N. Ma, N. Phattharasupakun, T. Maihom, J. Limtrakul and M. Sawangphruk, *J. Electroanal. Chem.*, 2018, **808**, 124–132.
- 33 F. Gao, Y. Tang, W. L. Liu, M. Z. Zou, C. Huang, C. J. Liu and X. Z. Zhang, *Adv. Mater.*, 2019, **31**, e1904639.
- 34 (a) M. Oishi, Y. Nagasaki, K. Itaka, N. Nishiyama and K. Kataoka, *J. Am. Chem. Soc.*, 2005, **127**, 1624–1625; (b) P. a. Ma, S. Liu, Y. Huang, X. Chen, L. Zhang and X. Jing, *Biomaterials*, 2010, **31**, 2646–2654.
- 35 (a) Z. L. Li, X. Q. Lai, S. Q. Fu, L. Ren, H. Cai, H. Zhang, Z. W. Gu, X. L. Ma and K. Luo, *Adv. Sci.*, 2022, **9**, 2201734; (b) Q. Chen, C. Y. Li and Q. B. Wang, *Small Methods*, 2023, 2201457; (c) X. Duan, C. Chan and W. Lin, *Angew. Chem., Int. Ed.*, 2019, **58**, 670–680.
- 36 (a) O. Kepp, L. Senovilla, I. Vitale, E. Vacchelli, S. Adjemian, P. Agostinis, L. Apetoh, F. Aranda, V. Barnaba, N. Bloy, L. Bracci, K. Breckpot, D. Brough, A. Buque, M. G. Castro, M. Cirone, M. I. Colombo, I. Cremer, S. Demaria, L. Dini, A. G. Eliopoulos, A. Faggioni, S. C. Formenti, J. Fucikova, L. Gabriele, U. S. Gaip, J. Galon, A. Garg, F. Ghiringhelli, N. A. Giese, Z. S. Guo, A. Hemminki, M. Herrmann, J. W. Hodge, S. Holdenrieder, J. Honeychurch, H. M. Hu, X. Huang, T. M. Illidge, K. Kono, M. Korbelik, D. V. Krysko, S. Loi, P. R. Lowenstein, E. Lugli, Y. Ma, F. Madeo, A. A. Manfredi, I. Martins, D. Mavilio, L. Menger, N. Merendino, M. Michaud, G. Mignot, K. L. Mossman, G. Multhoff, R. Oehler, F. Palombo, T. Panaretakis, J. Pol, E. Proietti, J. E. Ricci, C. Riganti, P. Rovere-Querini, A. Rubartelli, A. Sistigu, M. J. Smyth, J. Sonnemann, R. Spisek, J. Stagg, A. Q. Sukkurwala, E. Tartour, A. Thorburn, S. H. Thorne, P. Vandenabeele, F. Velotti, S. T. Workenhe, H. Yang, W. X. Zong, L. Zitvogel, G. Kroemer and L. Galluzzi, *Oncoimmunology*, 2014, **3**, e955691; (b) J. Rios-Doria, J. Harper, R. Rothstein, L. Wetzel, J. Chesebrough, A. Marrero, C. Chen, P. Strout, K. Mulgrew, K. McGlinchey, R. Fleming, B. Bezabeh, J. Meekin, D. Stewart, M. Kennedy, P. Martin, A. Buchanan, N. Dimasi, E. Michelotti and R. Hollingsworth, *Cancer Res.*, 2017, **77**, 2686–2698.
- 37 D. Laoui, E. Van Overmeire, G. Di Conza, C. Aldeni, J. Keirsse, Y. Morias, K. Movahedi, I. Houbracken, E. Schouppe, Y. Elkrim, O. Karroum, B. Jordan, P. Carmeliet, C. Gysemans, P. De Baetselier, M. Mazzone and J. A. Van Ginderachter, *Cancer Res.*, 2014, **74**, 24–30.

