Volume 42 Number 3 March 2025 Pages 423-640

Natural Product Reports

rsc.li/npr



ISSN 0265-0568



REVIEW ARTICLE

Nunziatina De Tommasi, Angela Bisio *et al.* Sesterterpenoids: sources, structural diversity, biological activity, and data management

Natural Product Reports

REVIEW

Check for updates

Cite this: Nat. Prod. Rep., 2025, 42, 443



View Article Online

View Journal | View Issue

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence. Open Access Article. Published on 20 2025. Downloaded on 2025/8/21 23:50:35.

Sesterterpenoids: sources, structural diversity, biological activity, and data management[†]

Valeria lobbi, ^(b) ‡^a Valentina Parisi, ^(b) ‡^{bg} Mauro Giacomini, ^(b) ^c Francesco De Riccardis, ^(b) ^d Paola Brun, ^(b) ^e Laura Núñez-Pons, ^(b) ^{fg} Giuliana Drava,^a Paolo Giordani,^a Maria Chiara Monti, ^(b) ^h Roberto Poggi, ^(b) ⁱ Ylenia Murgia,^c Nunziatina De Tommasi ^(b) *^{bg} and Angela Bisio ^(b) *^a

Reviewing the literature published up to October 2024.

Sesterterpenoids are one of the most chemically diverse and biologically promising subgroup of terpenoids, the largest family of secondary metabolites. The present review article summarizes more than seven decades of studies on isolation and characterization of more than 1600 structurally novel sesterterpenoids, supplemented by biological, pharmacological, ecological, and geographic distribution data. All the information have been implemented in eight tables available on the web and a relational database https:// sesterterpenoids.unige.net/. The interface has two sections, one open to the public for reading only and the other, protected by an authentication mechanism, for timely updating of published results.

Received 3rd August 2024 DOI: 10.1039/d4np00041b

rsc.li/npr

- Introduction
 Information infrastructure
- 2.1 Conceptual model
- 2.2 Logical model
- 2.3 Data presentation
- 2.4 External connections
- 3 Sesterterpenoids isolated from marine and terrestrial organisms
- 4 Distribution of sesterterpenoids
- 4.1 Sesterterpenoids in animals
- 4.1.1. Sesterterpenoids in Porifera

^aDepartment of Pharmacy, University of Genova, Viale Cembrano 4, 16148 Genova, Italy. E-mail: angela.bisio@unige.it

- ^bDepartment of Pharmacy, Via Giovanni Paolo II, 132, 84084 Fisciano, Salerno, Italy. E-mail: detommasi@unisa.it
- ^cDepartment of Informatics, Bioengineering, Robotics and System Science, University of Genova, Via all'Opera Pia 13, 16146 Genova, Italy
- ^dDepartment of Chemistry and Biology "A. Zambelli", Via Giovanni Paolo II, 132, 84084 Fisciano, Salerno, Italy
- ^eDepartment of Molecular Medicine, Section of Microbiology, University of Padova, Via A. Gabelli, 63, 35121 Padova, Italy
- Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli, Italy
- ⁸NBFC, National Biodiversity Future Center, 90133 Palermo, Italy
- ^hDepartment of Pharmacy, University of Napoli "Federico II", Via T. De Amicis 95, 80131 Napoli, Italy
- ¹Museo Civico di Storia Naturale Giacomo Doria, Via Brigata Liguria 9, 16121 Genova, Italy
- \dagger Electronic supplementary information (ESI) available. See DOI: $\mbox{https://doi.org/10.1039/d4np00041b}$
- ‡ These authors share the first authorship.

- 4.1.2. Sesterterpenoids in Mollusca
- 4.1.3. Sesterterpenoids in Cnidaria
- 4.1.4. Sesterterpenoids in insects
- 4.2 Sesterterpenoids in microrganisms
- 4.2.1. Sesterterpenoids in fungi
- 4.2.2. Sesterterpenoids in bacteria
- 4.2.3. Sesterterpenoids drive symbiotic relationships in fungi and bacteria
- 4.3 Sesterterpenoids in plants
- 5 Biological and pharmacological properties of sesterterpenoids
- 5.1 Sesterterpenoids with anti-cancer activity
- 5.2 Sesterterpenoids with anti-inflammatory activity
- 5.3 Sesterterpenoids with anti-microbial activity
- 5.4 Sesterterpenoids with anti-neurodegenerative activity
- 5.5 Sesterterpenoids with phytotoxicity
- 5.6 Sesterterpenoids with nematocidal and anti-feedant activity
- 5.7 Sesterterpenoids with immunosuppressive activity
- 5.8 Miscellaneous
 - Conclusions

6

7

- Author contributions
- 8 Conflicts of interest
- 9 Acknowledgements
- 10 References

1 Introduction

Despite originating from a single biosynthetic pentaprenyl linear precursor, sesterterpenoids¹⁻⁹ epitomize the astounding

strive of nature towards molecular diversity and complexity. The incredibly vast chemical space covered by sesterterpenoids embodies a myriad of forms, skeletal architectures, and substitution patterns. To date, over 1600 structures have been reported, with tens of unique hetero- and carbocyclic ring systems.

Since the isolation of the first members, in the late fifties and early sixties,⁸ it has been clear that sesterterpenoids were widespread in several phyletic groups, including marine sponges, nudibranchs, bacteria, lichens, fungi, higher plants and insects (Fig. 1).^{7,10–12} Although only a few macrocategories of sesterterpenoids are known for some taxa, most phyla can synthesize a considerable variety of compounds, from the simplest, such as the acyclic linear (AI), to the most complex, such as the hexacarbocyclic (HC). In general, several evolutionary mechanisms have been described that can lead to biosynthetic diversity and cause biosynthetic pathways to converge or diverge within or between different groups of organisms. These include gene duplication (and gene loss), horizontal and endosymbiotic gene transfer, and gene fusion.¹³ With regard to sesterterpenoids, the picture of possible coevolution of biosynthetic pathways is still unclear, and this is certainly a gap worth exploring.

Among the natural sources, the Porifera phylum (sponges) is the most prolific.14-17 The original producers of most natural products, including sesterterpenoids, in sponges, but also in other pluricellular holobionts, are often suspected to be their associated microbes, with larger metabolic capabilities.18-23 This widely approved hypothesis has been at the moment scarcely proved.14,16,17 Nonetheless, technological advances in omics and biological spatial approaches, single cell analyses, and cultivation procedures may soon provide suitable tools to demonstrate a symbiotic implication in the synthesis or biotransformation of secondary metabolites.24 In the intricate interactions between microbes and their hosts and within microbial communities, sesterterpenoids act as chemical defence and communication systems to enhance beneficial associations,25 and thus they can be considered allelochemicals mediating species interactions.14,16,17 Marine sponges, algae, and terrestrial plants produce sesterterpenoids to disrupt



Fig. 1 Circular plots showing the relationships among referred compound categories and groups of organisms at phylum level. The dimension of color bands is proportional to the number of compounds found in each category. Abbreviations are as follow, for phyla: ARTHR = Arthropoda; CNID = Cnidaria; MOLL = Mollusca; PORIF = Porifera; ACTINO = Actinobacteria; CYANO = Cyanobacteria; BACIL = Bacillariophyta; PROT = Proteobacteria; OCHRO = Ochrophyta; ASCO = Ascomycota; BASID = Basidiomycota; MUCOR = Mucoromycota; TRACH = Tracheophyta. For sesterterpenes categories: AI = acyclic, linear; MC = monocarbocyclic; MH = monoheterocyclic; DC = dicarbocyclic; DH = diheterocyclic; TrC = tricarbocyclic; TeC = tetracarbocyclic; TH = triheterocyclic; PC = pentacarbocyclic; HC = hexacarbocyclic.

microbial membranes or ignite reactive species inside bacterial and fungal predators, whereas, in some plant-microbe interactions, sesterterpenoids produced by the plant can attract beneficial microbes that promote plant health and growth.²⁶ Certain spongivore molluses are known to specifically feed on sesterterpenoid rich prey, in order to bioaccumulate or biotransform them for their own defence.²⁷ In other instances, molluses and enidarians seem to produce these bioactive molecules *de novo*.^{14,16,17,27} In microbial communities, microbes produce sesterterpenoids to inhibit the growth of competing species, thereby securing their niche or influence quorum sensing, a mechanism that bacteria use to coordinate their behaviour and eventually control biofilm formation, virulence, and metabolism.²⁸

In other groups of organisms, such as insects and plants, the function of sesterterpenoids has been less well investigated. Future studies should ascertain the involvement of these compounds in mediating the relationships between organism and their environment.

The driving force behind sesterterpenoids research, besides the purely structural, and synthetic studies, has been the wide range of biological and pharmacological properties they often exhibit (*i.e.*: anticancer, antimicrobial, anti-inflammatory, antifeedant, and antiviral activities). Sesterterpenoids also play a key role in the modulation of neurodegenerative processes, they have been studied for the treatment of type-II diabetes, hypercholesterolemia and obesity and as potential immunosuppressive agents.

With their rich oxidation patterns and three-dimensional complexity, these pentaprenyl terpenoids constitute a vast chemical library that can be easily morphed in new chemical entities (*via* ingenious semisynthetic or synthetic approaches). However, no synthetic study is reported in this review (unless for structural/stereochemical confirmation of the isolated secondary metabolites).

This review considers the extensive literature on sesterterpenoids to identify the state of the art on the subject and to highlight gaps in knowledge that need to be addressed in future studies. Specifically, the review is structured as follows: Section 2 presents a database that collects and organises the available information on the over 1600 sesterterpenoids. Sections 3 and 4 review the main compounds found in the different organism groups, while Section 5 outlines the biological and pharmacological properties already tested for these compounds.

2 Information infrastructure

2.1 Conceptual model

Since the 1970s, the Entity-Relationship (ER) diagram has played a fundamental role in the initial phases of data modelling projects. In this project, the conceptual model based on ER diagrams played a crucial role in defining the common design of the project's information support. The diagrams' graphic nature facilitated effective collaboration among the project's experts, enabling the proper exchange of individual collaborators' skills. The diagrams depict compounds, organisms, and bibliographic resources as the primary entities, with production, biological activity, and corresponding bibliography descriptions as the primary relationships connecting them.

2.2 Logical model

To ensure a reliable and efficient database, we translated the ER diagram into a logical scheme using the well-known relational model. This allowed us to transfer the agreed-upon knowledge organization of all project participants and experts. The logical scheme was implemented in a relational database management system (DBMS). The Microsoft SQL Server 2022 DBMS²⁹ was chosen and mounted on a server running the Windows Server 2022 operating system.³⁰ The server is hosted within the IT structure of Genoa University, which is backed up daily. In this project, specific measures were applied in addition to the standard rules of the relational model. Among these measures, we included an identification code for the compounds specific to this project. This is necessary because different bibliographic sources do not refer to a common nomenclature standard. Additionally, translating the graphic peculiarities (such as the use of italics, superscripts, and subscripts) typical of the nomenclature rules of both compounds and organisms into HTML is necessary. Unicode coding was used to name the compounds to ensure the letters of the Greek alphabet are essential for their correct naming.

However, when it comes to organisms, scientific names that follow well-established taxonomic rules are used in the literature. These names have been stored in the database as they appear in the bibliographic resources used for this project. It is important to note that the taxonomy of many organisms can rapidly change, rendering some nomenclatures obsolete and introducing new names that are recognized internationally. All names listed in international standard nomenclators are stored in the database. For each organism, all taxonomic levels are stored in the database using a normalized relational structure. This allows for quick provision of descriptive statistics of the database contents.

2.3 Data presentation

Although the DBMS allows for adequate and efficient management of data storage, excessive normalization results in a high number of tables connected by numerical indexes. This can make table management difficult for non-experts and irrelevant to ordinary people. To address this issue, a web interface was developed, divided into two sections. The initial section is publicly accessible and read-only and is structured for easy navigation among the data collected for this research. It can be explored based on the three major entities listed above: compounds, organisms, and bibliographic resources. The second section is restricted to authorized personnel and requires first-level authentication. It enables regular maintenance of the database content. The need to feed the database during its development suggested implementing the possibility of feeding data stored in batch mode through a set of Ms Excel files. The format of these files is based on the results of the conceptual analysis explained above.

2.4 External connections

The working group frequently updates the database, but the taxonomies of the organisms involved in the research can frequently change, which can quickly render the stored data obsolete. To address this issue, automatic query mechanisms have been planned for the main global databases in the sector, often providing access *via* web services. The system enables interested users to access updated data on a particular organism by requesting the same interface used in this project to query the relevant services. The updated data is then presented on a page that is appropriately formatted for the purpose. This allows non-expert users to access the most recent data on the subject. The sources of the updates are clearly indicated on the page.

3 Sesterterpenoids isolated from marine and terrestrial organisms

The current review categorises sesterterpenoids into ten subgroups (Table S1, ESI,† Sections 1.1–1.10), based on their structural features and increasing molecular complexity, ranging from linear to hexacarbocyclic backbones. All previous reviews on the subject in Natural Product Reports have followed, directly or indirectly, this subdivision criterion.^{1–3,6,31,32} However, due to the large number of linear sesterterpenoids reported, a further partition has been implemented based on the growing number of heterocyclic nuclei present in the terpenoid backbone (Table S1, ESI,† Sections 1.1-1.4). Accordingly, Section 1.1 of Table S1[†] presents the structures of the simplest linear acyclic (AI) sesterterpenoids decorated with various functional groups (*i.e.*: AI-11 from *Oryza sativa*,³³ Fig. 2); in Section 1.2 linear sesterterpenoids incorporating a single heterocyclic nucleus are collected and labelled as linear monoheterocyclic (MH) sesterterpenoids (i.e.: granuloside, MH-52 from Charcotia granulosa, Fig. 2);³⁴ Section 1.3 includes the structures of linear diheterocyclic (DH) sesterterpenoids (i.e.: hippolide A, DH-80 from Hippospongia lachne, Fig. 2);35 Section 1.4 reports the structures of linear triheterocyclic (TH) sesterterpenoids (i.e.: ircinialactam F, TH-24 from Ircinia oros, Fig. 2)36 and their possible dimeric counterparts. From Sections 1.5 to 1.10 a variety of structurally diverse and morphologically complex sesterterpenoids including carbocyclic moieties have been reported. In particular, Section 1.5 reports monocarbocyclic (MC) sesterterpenoids (i.e.: manoalide, MC-13, from Luffariella variabilis, Fig. 3);³⁷ Section 1.6 includes dicarbocyclic (DC) sesterterpenoids (i.e.: terpestacin, DC-104, from Arthrinium sp.);³⁸ Section 1.7 reports tricarbocyclic (TrC) sesterterpenoids (i.e.: ophiobolin A, TrC-2 from Ophiobolus miyabeanus);39 Section 1.8 comprehends tetracarbocyclic (TeC) sesterterpenoids (*i.e.*: bipolarolide A, TeC-1 from *Bipolaris* sp.);⁴⁰ Section 1.9, pentacarbocyclic (PC) sesterterpenoids [i.e.: phyllofenone D, PC-13 from Carteriospongia (syn. of Phyllospongia folias*cens*];⁴¹ and Section 1.10, hexacarbocyclic (HC) sesterterpenoids (*i.e.*: niduterpenoid A, HC-1 from Aspergillus nidulans).⁴² For Table S1 (ESI[†]) consultation, it is important to consider the following relevant information: (1) the sesterterpenoids included in each of the ten sections have been listed without any specific structural, biogenetic, phylogenetic, or

 $\begin{array}{c} + & + & + \\ + & + \\ &$

Fig. 2 Representative structures of formally linear sesterterpenoids incorporating no heterocycles (AI-11), or incorporating one (MH-52), two (DH-80), and three heterocyclic rings (TH-24), as reported in Sections 1.1–1.4, respectively, of Table S1.† Sulawesin C (TH-25) and molliorin-B (TeC-362) represent rare dimeric sesterterpenoids.



Fig. 3 Representative structures of sesterterpenoids incorporating one (MC-13), two (DC-104), three (TrC-2), four (TeC-1), five (PC-13), and six carbocyclic rings (HC-1), as reported in Sections 1.5–1.10, respectively, of Table S1.†

chronological order. The only criterion followed is that they belong to the structural class indicated by the denomination of the section; (2) where a revision or a new stereochemical assignment has been published, the correct structures and configurations are given; (3) carbocycles are counted as single independent units, even if they have a different biogenetic origin (*i.e.*: see phenyl rings in sesterterpenoids MC-85,⁴³ TrC-112,44 HC-3 (ref. 45)); (4) dimeric structures are always reported in the section of the corresponding monomeric counterparts (*i.e.*: sulawesin C, TH-25, from *Psammocinia* sp. in Section 1.4,⁴⁶ and molliorin-B, TeC-362, from Cacospongia mollior in Section 1.8, Fig. 2);47 (5) heterocyclic and carbocyclic rings are counted as single independent units even when they are present as bridged units (two rings share more than two atoms: i.e.: DH-99,48 DC-205,49 TrC-177,50 PC-9 (ref. 51)); (6) Table S1 (ESI)† includes the sesterterpenoids isolated from natural sources and by biosynthetic experiments and reported since the sixties of the last century. In some cases, meroterpenoid derivatives have been included when the pentaprenyl co-substrate is easily recognisable in the structure. Unfortunately, in most of the cases, meroterpenoids are difficult to be classified and lack of proper biogenetic studies. This hampers proper skeleton recognition and structural sorting.

4 Distribution of sesterterpenoids

4.1 Sesterterpenoids in animals

Most soft-bodied and sessile marine organisms, lacking physical or other mechanical means of protection or locomotion for escape, have evolved chemical defences for survival. Often these defensive compounds are noxious to potential predators, toxic, or have some type of bioactivity that directly or indirectly interferes with the behaviour or biology of co-occurring competing species.^{16,17} Among these allelochemicals, terpenoids are the most abundant and important compounds of marine origin. In this large family, the sesterterpenoids form a relatively narrow group of molecules, found mainly in sponges, and to lesser extent in molluscs and cnidarians, as well as in a reduced group of terrestrial producers, as soft scales insects (Fig. 4a).

4.1.1. Sesterterpenoids in Porifera. Sponges, belonging to the phylum Porifera, are the oldest multicellular animals on Earth. This large taxon has been able to colonise all latitudes of the world's oceans and freshwater systems, thriving in tropical, temperate, and polar regions, including extensive depth ranges, from the intertidal to the deep sea. In all these habitats, sponges play critical functional roles in three-dimensional habitat formation, nutrient recycling, space competition, water clearance, symbiotic interactions, microbial shelters and secondary metabolite production, among other.52 The ubiquitous occurrence of these organisms makes them prominent members of the benthic fauna, comprising 9650 recognised species (source: World Porifera Database,53 and over 20 000 taxon names), distributed in four classes - Calcarea, Hexactinellida, Homoscleromorpha, and Demospongiae.54 Sponges have simple bodied anatomy, consisting of a diploblastic cellular organization with an intraepithelial mesenchyme called mesophyl, composed by collagen, amoeboid cells, and skeletal elements. Their body plan lacks true tissues and consists of a system of branched canals and choanocyte chambers, that produce a water flow for feeding, respiration, excretion, and reproduction. The majority of sponges are heterotrophic filter feeders, with some exceptions that are partially (mixotrophic) or completely (phototropic, chemotrophic) dependent on photosynthetic symbionts trophic exchange, and a few carnivorous species that feed on small invertebrates.18,52

Sponges form intimate associations with a wide range of microorganisms, including bacteria, archaea, fungi, protists, and viruses, which are essential for their health and survival. The meta-organismal systems formed by sponge hosts and their microbiota should be considered as the minimal functional



Fig. 4 Circular plots showing the relationships among referred compound categories and taxonomic categories in 4 groups: (A) animalia, (B) fungi, (C) bacteria, (D) plants. The dimension of color bands is proportional to the number of compounds found in each category. Abbreviations are as follow: animalia. Arthropoda: COCC = Coccidae. Cnidaria: CLADO = Cladocoridae. Mollusca: CHAL = Chalinidae; CHROM = Chromodorididae; CURN = Curnonidae. Porifera: AGEL = Agelasidae; ANCO = Ancorinidae; APLY = Aplysinellidae; CHAL = Chalinidae; CRAM = Crambeidae; DARW = Darwinellidae; DICT = Dictyodendrillidae; DISY = Dysideidae; HALI = Halichondriidae; HYME = Hymedesmiidae; IRCI = Irciniidae; LATR = Latrunculiidae; MICR = Microcionidae; MYCA = Mycalidae; OSCA = Oscarellidae; PACH = Pachastrellidae; PETR = Petrosiidae; PODO = Podospongiidae; SPON = Spongiidae; SUBE = Suberitidae; THOR = Thorectidae. Fungi: Ascomycota: APIO = Apiosporaceae; APLO = Aplosporellaceae: ASPE = Aspergillaceae: BOTR = Botryosphaeriaceae: CHAE = Chaetosphaeriaceae: DIAP = Diaporthaceae: DIAP = Diapormellaceae; GYPS = Gypsoplacaceae; HYPO = Hypocreaceae; LEPR = Leprocaulaceae; LOPH = Lophiostomataceae; MASS = Massarinaceae; MOLL = Mollisiaceae; NECT = Nectriaceae; NEOC = Neocamarosporiaceae; OPHI = Ophiocordycipitaceae; PARM = Parmeliaceae; PELT = Peltigeraceae; PLEO = Pleosporaceae; Pleosporineae: SACC = Saccharomycetaceae; TRIC = Trichocomaceae; VALS = Valsaceae; XYLA = Xylariaceae. Basidiomycota: HERIC = Hericiaceae. OMPH = Omphalotaceae; PLEUR = Pleurotaceae; STERE = Stereaceae. Mucoromycota: CUNN = Cunninghamellaceae. Bacteria and chromista: Actinobacteria: PSEU = Pseudonocardiaceae; STRE = Streptomycetaceae. Cyanobacteria: NOST = Nostocaceae; SCYT = Scytonemataceae. Proteobacteria: ENTE = Enterobacteriaceae; PSEUD = Pseudomonadaceae; RHIZ = Rhizobiaceae. Bacillariophyta: RHIZO = Rhizosoleniaceae; NAVI = Naviculaceae. Plants: Tracheophyta: APOC = Apocynaceae; ASPL = Aspleniaceae; ASTE = Asteraceae; BRAS = Brassicaceae; CHLO = Chloranthaceae; CUCU = Cucurbitaceae; EUPH = Euphorbiaceae; FABA = Fabaceae; GENT = Gentianaceae; HYPE = Hypericaceae; LAMI = Lamiaceae; LAUR = Lauraceae; MALV = Malvaceae; NART = Nartheciaceae; POAC = Poaceae; PTER = Pteridaceae; ROSA = Rosaceae; RUTA = Rutaceae; SAPI = Sapindaceae; SIMA = Simaroubaceae; SOLA = Solanaceae; ZING = Zingiberaceae. For sesterterpenes categories: AI = acyclic, linear; MC = monocarbocyclic; MH = monoheterocyclic; DC = dicarbocyclic; DH = diheterocyclic; TrC = tricarbocyclic; TeC = tetracarbocyclic; TH = triheterocyclic; PC = pentacarbocyclic; HC = hexacarbocyclic.

biological units.^{19,21} Microbial symbionts provide their hosts with nutrients, process waste products, and appear to be involved in nutrient recycling processes.^{22,23} Moreover, sponge microbiomes may enhance growth and competitive ability within benthic communities, through the exchange of certain bioactive molecules and precursors. In certain sponges, the accumulation of microbial-derived natural products has been shown to provide various chemical defence strategies, such as

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence. Open Access Article. Published on 20 2025. Downloaded on 2025/8/21 23:50:35. 8

deterrence of predatory fish from feeding, anti-fouling to prevent overgrowth and suffocation, or growth inhibition against competing or pathogenic microbes.19 Much of the repertoire of secondary metabolites in eukaryotic organisms, particularly sponges, is thought to be derived from associated microorganisms.^{55,56} These compounds include mostly terpenes, sterols, cyclic peptides, unusual nucleosides, alkaloids, fatty acids, peroxides, and amino acid derivatives. Many of these products show promising therapeutic potential due to their anti-inflammatory, anticancer, antimicrobial, antiatherosclerotic and antiherpetic properties (as seen in previous reviews of the series).10,15,57,58 However, apart from their biotechnological applicability, the presence of these molecules in the sponge host should primarily respond to ecological means in the first place, which in most cases have yet to be revealed. Although there is much evidence suggesting the symbiotic production of many sponge secondary metabolites, few studies have empirically demonstrated the microbial synthesis of these compounds, including the sesterterpenoid family.^{19,55,56} Recently, the discovery of type I terpene synthases in the sponge holobiomes may call into question the absolute production of secondary metabolites by microbiome associates, implying the involvement of the animal host in several terpenoid biosynthetic pathways.59 In the following lines, we will illustrate some examples of sesterterpenoids occurrence in Porifera (Table S2, ESI[†]). Tropical and temperate shallow-water Porifera provide most of the known bioactive molecules. The reason is probably due to the intense allelochemical interactions in the highly biodiverse tropical ecosystems (as seen in previous review of the series).10,15,17 The order Dictyoceratida represents the most prolific taxon of known secondary metabolites, while the scalarane tetracarbocyclic sesterterpenoids form by far the broadest structural group of the terpene family⁶⁰ (and this review). Scalaranes are mainly found in sponges, but

also in nudibranchs, largely due to trophic transfer (see below). In both organisms, they exert a deterrent effect against generalist consumers, serving as an efficient antipredator or multipurpose defence mechanism.17 These products further exhibit a wide array of pharmacological activities (anti-cancer, antiinflammatory and antimicrobial being the most frequently described).61 Scalarin (TeC-156, Fig. 5) was the pioneering compound giving name to this large group of sesterterpeneoids. It was originally elucidated from the Mediterranean sponge Scalarispongia scalaris.62 Since the first discovery, TeC-156 has been recurrently found in Dictyoceratida sponges from diverse geographic locations (e.g., Scalarispongia sp. from Korea, Hyrtios erectus from South China, Ircinia sp. and Dysidea sp. from China and New Zealand, Spongia (Spongia) matamata from Palau, Spongia (Spongia) tubulifera from Mexico, Spongia (Spongia) virgultosa from Spain, Hyattella intestinalis from Mexico).63-67 Subsequently, many other scalarane-type compounds have been unveiled. The widespread species Hyrtios erectus seems to be the most productive in tetracarbocyclic sesterterpenoids, revealing scalaradial (TeC-75), heteronemin (TeC-326) and numerous derivatives, together with hyrtial (TeC-48) products, hyrtiosins (TeC-305-TeC-309) or salmahyrtisols (TeC-123, TeC-373, TeC-342, and TeC-504), in specimens from various regions, including China, Japan, Egypt, Saudi Arabia, the Maldives, New Guinea, New Caledonia, and Tonga.68-85 Congeneric Hyrtios species from Fiji, Thailand, Paracel Islands, and New Caledonia contained scalaranes and other molecules such as thorectolide (DH-76), erectusolide (DH-79), sesterstamide (TeC-345).86-89 At the same time, monocarbocyclic thorectidaeolides (MC-5-MC-8), acantholide A (MC-12) and luffariellolide (MC-70) were documented from H. communis from Palau.90 Similarly, Indopacific Phyllospongia foliascens collected in several areas (e.g., South China, Japan, Indonesia, New Guinea, Australia, and India) revealed an extensive array of scalarane-related products



Fig. 5 Representative structures of sesterterpenoids isolated from sponges.

in addition to other metabolites.91-94 These included rare scalarane-derived pentacarbocyclic sesterterpenoids with an additional cyclobutene, like carteriofenones A-D (TeC-37-TeC-40) and E-K (TeC-247, PC-11, PC-20-PC-23, and TeC-258),95 and other sesterterpenes like foliaspongin (TeC-33) and derivatives as dehydrofoliaspongin (TeC-412), phyllofoliaspongin (TeC-413),96-98 phyllactones F-G (TeC-217-TeC-218) and phyllofolactones A-D and M (TeC-246, TeC-206-TeC-208).99-102 Phyllofolactones and scalaranes were additionally found in congeneric sponges Phyllospongia sp., P. lamellosa, P. vermicularis.¹⁰³⁻¹⁰⁷ Phyllactones were found in *P. papyracea* from specimens coming from Egypt, Madagascar, Indonesia, New Guinea and China.¹⁰⁸⁻¹¹⁰ Indian Hyattella cribriformis and a Korean Hyattella sp. were found to produce scalaranes, together with H. intestinalis from Australia and Mexico, which reported in addition to the repertoire of TeC-156 and TeC-326 relative metabolites, also norscalarals hyatolides (TrC-171, TrC-172 and TeC-299-TeC-301), mooloolabenes (TeC-66-TeC-70 and TeC-248-TeC-257), furoscalarol (TeC-329), and hyatelactam (TeC-344).^{66,111-114} Sponges Hvattella sp. from Indonesia were found to possess hyattellactones (TeC-239 and TeC-240) and phyllofolactones (TeC-246, TeC-205-TeC-214, TeC-302, TeC-303, TeC-492).115 Lendenfeldia sponges represent another taxon with an extended presence of scalaranes, homoscalaranes and related molecules, including furodendins, homoscalarates and homoscalaralactone from L. chondrodes from Palau, Australian L. dendyi and L. frondosa from the Salomon Islands and New Guinea, as well as sesterterpenes like lendenfeldaranes, felixins (TeC-100) or furanolipids, in Lendenfeldia sp. from Madagascar.116-123 The genus Spongia has afforded another notable repertoire of scalaranes, including the original TeC-156, TeC-75 and derivatives, as well as numerous other tetracyclic sesterterpenoids and different sesterterpene type molecules like scalalactams, furospongins, hyrtiosal (TrC-130), igernellin (MC-71), hipposulfates, ircinins, cometins, petrosaspongiolides, and plenty of others. These compounds were obtained from diverse species and locations such as Spongia (Spongia) agaricina, S. (S.) nintens and S. (S.) officinalis from Mediterranean, S. (S.) hispida and S. (S.) matamata from Papua New Guinea, S. (S.) oceanica from Hawaii, and from a number of unidentified Spongia sp. from the USA, Australia, Borneo, Japan, and Korea. 63,64,124-140 Other genera of sponge typically containing scalarane-related products include Smenospongia specimens from Korea,141,142 or the only species in the Collospongia genus, C. auris from Australia, containing TeC-326 and terpene derivatives.143 Strepsichordaia lendenfeldi from Australia revealed scalarane derivatives,144 whereas Indonesian Strepsichordaia aliena yielded honulactones, phyllofenone C (TeC-287) and phyllofolactones.145 Older investigations frequently described scalarane compound series in sponges whose terminology has changed after taxonomic revisions. TeC-75 and related scalardysins and scalarherbacins were described from Lamellodysidea herbacea from Gulf of Suez,146 and TeC-326, scalarolide (TeC-173), scalarafuran (TeC-331), furospinosulin-1 (MH-18), idiadione (MH-28) among others in Leiosella idia from USA.147 TeC-75 and scalarane-type molliorins were isolated from Mediterranean Cacospongia mollior, 47,148-155 and Pacific Cacospongia sp. unveiled

cacolic acid (MH-53) and several cacolides, which are mainly linear sesterterpene compounds.¹⁵⁶ Scalarane-type including TeC-326 and manoalide-type product (MC-27) were recovered from Thai Brachiaster sp. sponges.157 Tetracyclic scalarane-types and pentacyclic sesterterpenoids are commonly found in Dysidea genus. Scalarane compounds have been reported in Dysidea sp. from China,67 and in D. gumminae from Thailand, in addition to similan A, hyrtiolide, TeC-173 and scalafuran.¹⁵⁸ Chinese D. granulosa produced various tetracyclic and pentacyclic sesterterpenoids.¹⁵⁹ Bilosespenes were elucidated from Eritrean D. cinerea¹⁶⁰ and dysidiolide (DC-22, Fig. 5) from D. etheria from USA,¹⁶¹ while halisulfates 1, 3, and 5 (TrC-115, DC-38, and DC-40), dysideapalaunic acid (DC-23) and coscinoquinol (TrC-112) were found in Dysidea sp. from Palau and Micronesia.162,163 Psammocinia sp. sponges have been reported to produce variable suites of sesterterpenoids. For example, some specimens from Korea yielded scalarane-type products,164,165 or psammocinins A₁, A₂, and B (DH-30, DH-31, and DH-69) and variabilin (DH-16).¹⁶⁶ Meanwhile, Australian relatives were found to contain bicarbocyclic sesterterpenoids such as ircinianin (DC-159), ircinianin sulfate (DC-179, Fig. 5), ircinianin sulfate lactam (DC-177) and derivatives, as well as isopalinurin (DH-4),167 and linear sesterterpenes ircinins 1-2 (TH-8 and TH-9) and sulawesins A-C (MC-9, MC-10, and TH-25) were also isolated from Psammocinia sp. in Indonesia.46 Hippospongide A (TeC-385), hippospongide B (TeC-172) and other scalaranes were isolated from Taiwanese Hippospongia sp.168 Other unidentified Hippospongia provided furanoterpene hippospongins (TH-26-TH-31) from Australian specimens,169 or TrC-115 and DC-40 from Micronesia collections.¹⁷⁰ Linear and bicyclic hippolides A-J (DH-80, DH-81, MH-11-MH-16, DC-216, DC-217) and monocarbocyclic manoalides (MC-16 and MC-17) were isolated from H. lachne collected in South China.^{35,48,171}

Disparate tetracarbocyclic sesterterpenoids toxistylides A-B were identified in Mediterranean Clathria (Clathria) toxistyla,¹⁷² and in C. (C.) gombawuiensis from Korea, including ansellone C (TrC-206, Fig. 5), gombaspiroketals A-C (DC-195-DC-197) and phorone B (TeC-389).173 Related phorones A (TeC-386) and C (TeC-481) were also identified in Korean Phorbas sp. Anvilones A (TeC-390) and B (TeC-391), phorbadione (TrC-199), ansellones C-K (TrC-207-TrC-211 and TrC-323-TrC-326), and monocyclic phorbaketals A-C and L-M (MC-119-MC-121 and MC-130-MC-132) were purified from congenerics from British Columbia.¹⁷⁴⁻¹⁸¹ The antarctic collections of *P. areolatus*, in its place, revealed suberitenones (TeC-23-TeC-25, TeC-30) and suberitane derivatives, including isosuberitenone B (TeC-394), 19-epi-suberitenone (TeC-25), and isoxaspirosuberitenone (TeC-395).182 Furthermore, monocyclic spirocyclic MC-119-MC-129 and phorbin A (MC-4) were also obtained from Korean Monanchora sp. sponges.¹⁸³ A number of norsesterterpene peroxides have been reported in genus Diacarnus. Tasnemoxides A-C (MC-40-MC-42), muqubilin (MC-36-MC-37) and derivatives were found in D. erythraeanus from the Red Sea.184,185 Aikupikoxide A (MH-30) and sigmosceptrellin B (DC-17) were also identified in the same species. Sigmosceptrellins A-C (DC-60, DC-17, and DC-63) and diacarnoxides A-D (MC-49-MC-52) were respectively identified in D. laevis and D. levii both from Papua New

Guinea.¹⁸⁶ Diacarperoxides and **MC-36** were recovered in Indonesian *D. megaspinorhabdosa*.¹⁸⁷ Additionally, muqubilin relative products were also found in *D. spinipoculum* from Solomon Islands.¹⁸⁸ Other cyclic norsesterterpenes, known as mycaperoxides A–B (**DC-15** and **DC-16**), were retrieved from Thai *Mycale* sp.^{189,190} Tricarbocyclic sesterterpenoids called coscinolactams A–G (**TrC-200**, **TrC-154**, **TrC-201–TrC-205**), including suvanine (**TrC-143**) derivatives, were isolated from *Coscinoderma* sp. from Micronesia,^{191–193} and from *C. mathewsi* coming from Solomon Islands, along with coscinalactone (**TeC-402**) and coscinafuran (**TeC-403**).^{194–196}

The genus Ircinia probably comprises one of the most diversified in sesterterpene series, reporting a vast range of molecules including cyclic and linear furanosesterterpenoids, scalaranes, 24-homoscalaranes, tetronic acid related compounds, cheilanthane sesterterpenoids or C22-trinorsesterterpenoids, among others. Felixin scalaranes (TeC-100, TeC-101, TeC-165-TeC-169) were described from I. felix in Taiwan, ircinins (TH-8-TH-11) were detected in I. oros,^{36,197} and a numerous group of irciformonins (DH-10, DH-13, TH-19-TH-21) and ircinialactams (DH-96-DH-98, TH-24, TH-32) were isolated from Ircinia sp. from Taiwan and Australia.198,199 Moreover, strobilinins, felixinins and variabilins are frequently found in several congeneric species and regions: e.g., I. oros and I. variabilis from Mediterranean, I. campana from Colombia, I. strobilina from Brazil, and Ircinia sp. from several Pacific areas. DC-159 and wistarin (DC-160) were isolated from the Australian species I. wistarii.198,200-205 Chinese collections of Dactylospongia elegans provided γ -oxygenated butenolides, the dactylospenes A-E (MH-57, DC-224-DC-227).206 Jaspis sponges from New Guinea (Jaspis cf. johnstoni), China (Jaspis sp.), and Japan (J. stellifera) yielded jaspic acid (TrC-110), jaspolide F jaspiferals (TrC-135), and C-F (TrC-131-TrC-134), respectively.²⁰⁷⁻²⁰⁹ Acantholides C-E (MH-10, MC-1 and MC-2) were recorded from Pacific Acanthodendrilla sp.,²¹⁰ while agelisamines A and B (DC-29 and DC-30) and aplysinoplides A-C (MC-30, MC-33, and MC-34) were obtained from Agelas mauritiana; and Aplysinopsis elegans and Aplysinopsis sp. respectively all from Japanese collections.²¹¹⁻²¹³ New Caledonian Petrosaspongia nigra were found to possess numerous petrosaspongiolides, A-L (TrC-212-TrC-221, Fig. 5 and TrC-222), M-P (TrC-178-TrC-180), Q and R (TrC-182 and TrC-183),²¹⁴ and Petrosaspongia sp. from Fiji revealed several petrosaspongiolactams.215 Aurorals 1-4 (TrC-255-TrC-258) and globostelletins C-G (TrC-264-TrC-267) were isolated from Rhabdastrella globostellata from New Caledonia and South China, respectively.²¹⁶ Rhopaloic acids A-C (MH-7, MH-1-MH-3) were found in Rhopaloeides sp. from Japan.217,218

Luffariella representatives produce specific bicyclic, monocyclic, and acyclic sesterterpenoids, including luffarins A–O (DC-43–DC-57), P (MC-65), Q (MH-8), R (DH-58), S (DH-106), T and U (DH-59) and DH-60) from Australian *L. geometrica*,²¹⁹ luffariolides A–G (MC-77–MC-83, MC-70, luffalides A–F (MC-57– MC-62) or luffolide (TrC-280) from *Luffariella* sp. from several Pacific collections (*e.g.*, Australia, Micronesia, Japan, Taiwan, Palau). Additionally, *Luffariella* sp. produces a variable suite of product series from *L. variabilis* according to the collection site.

For example, specimens from Australia produce luffariellin A (MC-53), 25-acetoxyluffariellin (MC-55), 25-acetoxyluffariellin B (MC-69), and manoalide-type products (MC-13-MC-15, MC-28), while Palau sponges produce luffariellin B (MC-56), luffalactone (DC-162), several manoalides (MC-13-MC-16, MC-19, MC-21, MC-28) in Malaysian samples, or oshimalides A and B (MC-146 and MC-147) from deep sea populations.37,220-229 Sarcotragus sp. sponges are similarly sources of norsesterterpenoids, linear sesterterpenes and other secondary metabolites. Collections from Korea have revealed a notable number of bioactive compounds, including a list of sarcotins A-C (DH-36-DH-38), D and E (TH-6 and TH-7), F (DH-74), G and H (TH-22-TH-22), I and J (MH-46 and MH-47), M (DH-40), O and P (MH-20 and MH-21), sarcotrines (DH-87-DH-91), sarcotragins A-C (MC-85, MH-42, and MH-54). 43,230-234 Unidentified congenerics from New Zealand or Australia have yielded ircinialactams (DH-96, MH-32-MH-34, MH-55-MH-56) ircinialactone A (DH-62), DH-16 or MH-18.235,236 Antarctic Suberites sp. demonstrated to possess a series of suberitenones (TeC-23, TeC-24, TeC-27, and TeC-28, Fig. 5)^{237,238} and S. caminatus also from Southern Polar waters contained suberitenones derivatives (TeC-26 and TeC-30), together with caminatal (TrC-250).²³⁹ Pacific Thorectandra sp. from Palau unveiled the presence of thorectandrols (DC-8, DC-9, DC-11-DC-13), palauolol (DC-2) and DH-58.240 Thorectolide monoacetate (DH-77) was reported in Australian T. excavatus.²⁴¹ Among the few linear sesterterpene products known in marine habitats, a group of the so-called balibalosides and derivatives (MH-4-MH-6, and MH-9) were described in the Mediterranean Oscarella balibaloi from France.242

4.1.2. Sesterterpenoids in Mollusca. For the purposes of this review, a specific group of marine gastropod molluscs belonging to the subclass Opisthobranchia (from the Greek opisten = posterior and branchion = gills) were also considered (Table S3, ESI[†]). The classification of this relatively small group of molluscs, estimated to range from 5000 to 6000 species, is still widely debated.243,244 Opisthobranchia are characterized for having detorsioned nervous system and modified respiratory organs. Determining diagnostic traits further comprise notable shell reduction remaining vestigial in some cases, or only maintained during larval stages, as well as internal shell forms overgrown by soft tissue. These modified shell forms serve majorly as skeletal mechanical support, rather than for physical protection of the slugs. The Nudibranchia, the largest opisthobranch order with approximately 3000 species, lacks a shell and relies on chemical means for defence, often accompanied by shimmering or confusing colourations. This taxon exhibits significant diversification in shape and ecological traits related to trophic habits, reproduction, and protective strategies.²⁴⁴⁻²⁴⁸

Nudibranchs have a diverse diet, consuming various food items, including Chlorophyta, Ochrophyta, Rhodophyta (green, brown, and red algae, respectively), Porifera, Cnidaria, Bryozoa, Chordata (tunicates), and other Mollusca. However, their feeding behaviour is highly specialized, and almost each species predates on one or few prey.²⁴⁶ Such pattern is correlated with the ecological competence and defensive tactics of these animals, due to their ability to "steal" functional structures (cleptoplasty) or chemical products (cleptochemistry) from their

Natural Product Reports

prey. For example, they can incorporate chloroplasts or zooxanthellae from algae to obtain energy and camouflage,^{249,250} or nematocysts from cnidarian prey to use as protective weapons.²⁵¹ Cleptochemistry is the process of sequestrating natural products derived from food items, and then the usage for self-defence. Some nudibranchs can biotransform the dietary metabolites into less toxic compounds to allow bioaccumulation, or to more effective noxious products to deter predators. While a few species can uptake simple precursors and de novo biosynthesize defensive molecules, most rely on sequestering compounds from their diet.27,252,253 For efficient energetic and protective purposes, the resulting bioactive compounds are often translocated to susceptible anatomical sites, glands, or exposed parts, or they may be exhausted within mucous secretions.^{27,245,254,255} Moreover, in synchrony with allelochemistry warning (aposematic) colorations frequently advise the presence of chemical defences or toxicity towards putative predators in exposed habitats. Consumers learn the connection between bright colorations and bad taste.27,245,256

It has been hypothesized that the reduction of the shell in the evolution of Opisthobranchia was facilitated by modifications in the foraging habits towards food items containing secondary metabolites, which would have been sequestered and used as chemical defences. This also suggests that the protection based on dietary allelochemicals would have preceded de novo synthesized defensive mechanisms.27,252 Dietary sesterterpenoids are common in dorids, particularly in family Chromodorididae. Most of these compounds are bioactive tetracyclic sesterterpenoid products, which are assimilated from sponge prey.27,257 The scalarane compounds of dietary acquisition, specifically TeC-75 (a potent anti-inflammatory metabolite) and other related bioactive molecules present in demosponge prey are of particular relevance.²⁵⁸ Once trophically incorporated, these metabolites are usually further bio transformed into derivatives by in chromodorid slugs as part of

detoxification processes. They are frequently allocated towards the mantle border and dermal formation-like structures for the purpose of antipredation protection. Glossodoris sedna from Costa Rica besides reporting sponge scalaranes,²⁵⁹ it was previously described to possess sednolide (TeC-365).²⁶⁰ Similarly, luffariellin C (MC-63) and D (MC-64) were detected along scalarane derivatives (TeC-164, TeC-281, and TeC-432) in C. funerea from Palau.²⁶¹ TeC-326 is a recurrent scalarane-type sesterterpene trophically transferred from sponges (e.g. Heteronema erecta, several Spongia sp.) to nudibranch slugs [e.g. Glossodoris (syn. of Doriprismatica) atromarginata, atromarginata].262 This compound with antimycobacterial properties263 was allocated in the viscera of the Australian slugs G. hikuerensis and G. vespa, in contrast with TeC-75, 12-deacetyl-12oxoscalaradial (TeC-46, Fig. 6), and 12-deacetoxy-12-oxodeoxoscalarin (TeC-141), which were detected in the mantle.264 Ten norscalaranes mooloolabenes D-O (TeC-69, TeC-70, TeC-248-TeC-257), along with scalaranes (TeC-141 and TeC-433) were isolated from Australian D. atromarginata from diverse locations, indicating varied sponge diet and/or diversified enzymatic detoxification mechanisms in these slugs.²⁶⁵ Other dietary tetracyclic sesterterpenoids found in sea slugs include ansellone A (TrC-192, Fig. 6) from Cadlina luteomarginata from Canada and its prey Phorbas sp.,¹⁷⁷ as well as MH-28 and luteone (TrC-232).147,266 Hamiltonin E (TrC-142) was isolated from South African Chromodoris hamiltoni.267 Inorolides A, B and C (TeC-368, TeC-369, and TrC-281) were found in Japanese Chromodoris inornata,268 along with TeC-156 or scalaradial derivatives (TeC-428-TeC-431). Variabilin derivatives (DH-22 Fig. 6, DH-49, DH-107, and DH-109) were obtained from the South African Hypselodoris capensis and its demosponge prey Fasciospongia sp.²⁶⁹ Finally, a unique example of a non-dietary linear sesterterpenoid is granuloside (MH-52), isolated from the Antarctic cladobranch Charcotia granulosa.34,270



Fig. 6 Representative structures of sesterterpenoids isolated from Mollusca, Cnidaria and insects.

4.1.3. Sesterterpenoids in Cnidaria. Within the phylum Cnidaria (Table S4, ESI†), the anthozoan order Scleractinia includes the "true corals" or "stony corals", which are represented by ~1500 extant species. Scleractinian corals exist as solitary (single polyps), or clonal as colonies comprised of many individual polyps. The polyps' tentacles are covered with nematocysts or cnidocysts that are used to capture prey. As the polyp grows the aragonite skeleton deposition begins. Many corals living in the photic zone establish trophic obligate symbiotic relationships with dinoflagellate algae belonging to the family Symbiodiniaceae, known as zooxanthellae.271 These zooxanthellae provide the coral host with the majority of the energetic boost from photosynthesis and nutrient exchanges, while receiving protection within the stinging coral's tissues.²⁷² Scleractinians have been reef-building organisms for the past 240 million years, making them crucial benthic habitat bioconstructors. They form the coral reef frameworks in the tropics, as well as important bioconstructions in temperate and deep-sea bottoms, where other organisms find shelter and food.273-275

In the Mediterranean, the most significant bioconstructing zooxanthellate scleractinid coral is the madreporaria *Cladocora caespitosa*,²⁷³ which was found to contain bioactive sesterterpenes called cladocoranes A and B (**DC-6**, Fig. 6, and **DC**-7). These products revealed biologcal properties in the treatment of various diseases, including cancer, together with potential antitubercular and antibacterial activities inhibiting the growth of Gram-positive strains.²⁷⁶

4.1.4. Sesterterpenoids in insects. The Coccidae is a family of Insecta Hemiptera Sternorrhyncha commonly known as wax scales, soft scales or tortoise scales (Table S5, ESI†). The females are wingless and flat, with an oval body that is often heavily sclerotized and covered with wax, while the males may be winged or wingless. Members of the family are known to feed on a variety of plants belonging to different plant orders. Some are polyphagous, but most are oligophagous or monophagous. Host plants are mainly perennial plants and often woody. Many Coccidae species are important and serious pest in agriculture, horticulture, and forestry. Some taxa have been introduced to new regions through the movement of plant material, resulting in their cosmopolitan distribution. The Coccidae family includes around 170 genera and 1100 species. The genus Ceroplastes, described by Gray in 1828, includes more than 130 species. From Ceroplastes, a little group of sesterterpenoids has been isolated so far. It has been hypothesized that the significance of the presence of sesterterpenoids in the external waxy cover of insects could be related to their activity as kairomones. Specifically, cerorubenic acids I and II (TrC-173 and TrC-174, Fig. 6) and cerorubenols I and II (TrC-175 and TrC-176), could be responsible for the ovipositional behaviour of the parasitic wasp Anicetus beneficus (Encyrtidae) towards C. rubens.277 Linear, monocyclic, bicyclic, and tricyclic sesterterpenoids have been isolated from the external waxy cover of female of the Mexican species C. albolineatus. The linear acyclic geranylfarnesol (AI-1)278 was isomeric with geranylnerolidol (AI-3) isolated by the fungus Cochliobolus heterostrophus.279 ω-Hydroxygerany

lfarnesol (AI-2)²⁸⁰ has been also isolated from the sponge Fasciospongia fovea.281 Albocerol (MC-103)282 was a monocarbocyclic sesterterpenoid also isolated from C. albolineatus. Its structure belongs to ceriferene sesterterpenoids, macrocyclic compounds containing a 14-membered ring (1,14-cyclogeranylfarnesase scaffold).²⁵ Similar compounds were isolated from C. ceriferus, a worldwide distributed species (MC-100-MC-102, MC-104-MC-111),^{283,284} and from C. pseudoceriferus (MC-112, MC-113, MC-138, MC-139).285 Albolineol (DC-103) was the only bicyclic compound isolated from C. albolineatus, and described together with the tricyclic ceroplastanes ceroplastol-I (TrC-98, Fig. 6) and ceroplastol-II (TrC-99).286 Ceroplastanes belong to the ophiobolane type sesterpenoids. The junction between rings A and B and between B and C in ceroplasteric acid (TrC-97) and TrC-98, are both trans, while ophiobolins TrC-2, TrC-8, and TrC-13 show a cis-trans disposition.²⁸⁶⁻²⁸⁸ Other analogous, all isolated from C. albolineatus, were albolic acid (TrC-100),²⁸⁹ ceroplastodiol (TrC-101), ceralbic acids I and II (TrC-103 and TrC-104)²⁹⁰ and ceralbol (TrC-105).291 The wax of C. madagascariensis afforded the tricarbocyclic gascardic acid (TrC-1),²⁹²⁻²⁹⁵ one of the first described sesterterpenoids. Other tricarbocyclic sesterterpenoids were isolated from C. floridensis, a pest insect which infests orchards as persimmon, and tangerine. These compounds (TrC-123-TrC-126) show an 11-membered ring, derive and biogenetically from a head-to-tail cyclization of 2-(Z)geranylfarnesylpyrophosphate.296

4.2 Sesterterpenoids in microrganisms

Fungi and bacteria represent significant sources of natural sesterterpenoids (Table S6, ESI,† and Fig. 4b). In the microbiological world, sesterterpenoids are metabolites endowed with different physiological effects. They drive interactions (*i.e.*, symbiosis, competition, or proliferation) with neighbouring commensal or invader organisms, and some are endowed with anti-inflammatory and anticancer activities.²⁹⁷ This section will examine the significance of sesterterpenoids in the ecological setting of microbes.

4.2.1. Sesterterpenoids in fungi. The kingdom of fungi includes highly diverse lineages of microbial eukaryotes classified basing on phenotype identifications, physiological profiling, and DNA barcoding. It is estimated that there are approximately 5.1 million of fungi species worldwide, but up to date, only 2% of them have been classified. Environmental sequencing analysis is filling the gap. However, there is an increasing rate of ecologically cryptic groups, which are species known for their DNA sequence but lack morphological and cultural characterization and do not have an accepted name. This generates dark taxa that cannot be formally described under the current nomenclature of fungal taxonomy rules, thus complicating the classification process.298,299 Nevertheless, fungi are all characterized by chitinous cell walls, membranebound organelles, and clearly defined nuclei. They are exclusively osmotrophic, taking up organic matter externally and digesting it at the external hyphae before absorbing it in the mycelia. For this reason, many fungi form parasitic or symbiotic relationships with bacteria, plants, or animals, even if others are free-living organisms.^{298,300} Fungi display an array of ecological functions due to their metabolic potential. They act as principal decomposers in ecological systems, detect environmental cues for the biological quorum, and produce biochemicals for defence, allelopathy, or maintenance of symbiosis.³⁰¹ Sesterterpenoids primarily participate in fungalmediated ecological functions. Dai and colleagues calculated that, at the intracellular level, 21% of all produced sesterterpenoids inhibit enzymes, thus controlling fungal metabolism. Most sesterterpenoids (Table S6, ESI†) act extracellularly, with 19% of them exhibiting antimicrobial activity against other fungi, bacteria, or viruses.³⁰¹ Indeed, ophiobolin derivative 3anhydro-6-hydroxyophiobolin A (TrC-6, Fig. 7) produced by the phytopathogenic fungus Bipolaris oryzae and ophiobolin T (TrC-56) produced by endolichenic fungus Ulocladium sp. reported significant bacteriostatic effects against Bacillus subtilis, Staphylococcus aureus, and methicillin-resistant S. aureus (MRSA). Ophiobolins, tricarbocyclic sesterterpenoids with a 5/8/5-fused carbocyclic skeleton, were first described in the last '50s' and early '60s' of the last century.³⁰² Ophiobolin A (TrC-2) was firstly isolated from Bipolaris species, as B. leersiae, 303 B. maydis, 303,304 B. oryzae,^{39,305-312} B. panici-miliacei,³⁰³ B. setariae,³¹³ B. sorghicola.^{304,314} It was subsequently isolated by a large number of organisms, including Cochliobolus heterostrophus, 315,316 C. miyabeanus,³¹⁷ Drechslera gigantea,^{318,319} D. zizaniae,^{320,321} Helminthosporium turcicum,³²² Helminthosporium spp.³²³

Bipolaris species are known to produce ophiobolins. 6-*epi*-Ophiobolin A (**TrC-3**), 3-anhydroophiobolin A (**TrC-4**),³²⁴ and 3anhydro-6-*epi*-ophiobolin A (**TrC-5**),³⁰⁴ as well ophiobolin F (**TrC-17**),^{279,325} 25-hydroxyophiobolin I (**TrC-34**)³⁰⁴ has been firstly isolated from *B. maydis*, **TrC-6**,³²⁶ ophiobolin B (**TrC-8**),³²⁷ ophiobolin I (**TrC-32**),^{312,328} 6-*epi*-ophiobolin I (**TrC-33**),³²⁹ ophiobolin J (**TrC-35**),³²⁹ and 8-deoxyophiobolin J (**TrC-37**)³²⁹ have been isolated from *B. oryzae*. Bipolarolides, ophiobolin derived sesterterpenoids have been isolated from *Bipolaris* sp. (TJ403-B1).⁴⁰ Bipolarolides A and B (**TeC-1** and **TeC-2**) are characterized by a multicyclic caged oxapentacyclo[9.3.0.0 (ref. 1 and 6).0 (ref. 5 and 9).1 (ref. 8 and 12)]pentadecane-bridged system. Bipolarolides C and D (**TeC-3** and **TeC-4**) show a 5/5/5/ 5-fused core skeleton, and bipolarolide C also contains a C-3-C-14 oxygen bridge to construct the caged architecture. Bipolarolides E-G (DC-1, TrC-107, and TrC-108, Fig. 7) are highly modified pentacyclic oxaspiro[4.4]nonane-containing sesterterpene-alkaloid hybrids.⁴⁰ From the same strain, growing on fermented rice medium, others ophiobolin-type metabolites, bipolaricins A-I (TrC-87-TrC-95),330 and bipolarins A-H (TrC-79-TrC-86), tetracyclic ophiobolin-type sesterterpenes characterized by an oxaspiro[4.4]nonane moiety, together with ophiobotriol (TrC-78) have been characterized.331 Ophiobolin-type sesterterpenoids maydispenoid A (TrC-306) with a decahydro-3-oxacycloocta[cd]pentalene moiety, and maydispenoid B (and TrC-307), have been isolated from B. maydis collected from Anoectochilus roxburghii (Wall.) Lindl leaves.332

Ophiobolins are also commonly found in the genus Aspergillus.^{326,331,333,334} A large number of compounds have been first obtained from the crude extracts of the liquid and solid cultures of the mangrove fungus A. ustus, namely ophiobolins G and H (TrC-18 and TrC-28),335 ophiobolin K (TrC-38),336 ophiobolin O (TrC-49), ophiobolin P (TrC-52) ophiobolins U-Z (TrC-57, TrC-60-TrC-64),^{337,338} 21-epi-ophiobolin O (TrC-51),³³⁷ 21-dehydroophiobolin U (TrC-59),³³⁷ 21-epi-ophiobolin Z (TrC-65),³³⁷ 21deoxyophiobolin K (TrC-42),³³⁷ 6-epi-ophiobolin K (TrC-39),³³⁶ (6a)-18,19,21,21-O-tetrahydro-18,19-dihydroxyophiobolin (TrC-24),³³⁸ (6α)-21-deoxyophiobolin G (TrC-26),³³⁸ (6α)-16,17dihydro-21-deoxyophiobolin G (TrC-27),338 (5α,6α)-ophiobolin H (TrC-29),³³⁹ (5α,6α)-5-O-methylophiobolin H (TrC-30),³³⁹ 5-Omethylophiobolin H (TrC-31),³³⁹ and (6a)-21,21-O-dihydroophiobolin G (TrC-25).339 The marine fungus A. flocculosus afforded 14,15-dehydro-6-epi-ophiobolin K (TrC-41), 14,15dehydro-ophiobolin K (TrC-40),340 14,15-dehydro-6-epi-ophiobolin G (TrC-23),340 14,15-dehydroophiobolin G (TrC-22),340 and 14,15-dehydro-(*Z*)-14-ophiobolin (TrC-21),340 G and ophiobolin N (TrC-47).340 6-epi-Ophiobolin N (TrC-48) has been obtained from A. insuetus.341

Other ophiobolins and congeners have been isolated firstly from other fungi. Ophiobolin C (zizanin A) (**TrC-13**) has been



Fig. 7 Representative structures of sesterterpenoids isolated from fungi

obtained from *Drechslera zizaniae* (as *Helminthosporium zizaniae*),³²⁰ ophiobolin D (cephalonic acid) (**TrC-15**) from *Cephalosporium caerulens*,³⁴²⁻³⁴⁴ ophiobolin E (**TrC-16**) and 8-*epi*-ophiobolin J (**TrC-36**) from *D. gigantea*,^{318,319} and 6-*epi*-ophiobolin G (**TrC-19**) from *Emericella variecolor* obtained from a marine sediment.^{345,346} *Cochliobolus heterostrophus* yielded ophiobolin L (**TrC-43**),³¹⁵ ophiobolin M (**TrC-45**),³⁴⁷ 6-*epi*-ophiobolin M (**TrC-46**),³⁴⁷ the degradation products 3-anhydro-6-*epi*-ophiobolin B and the dimer di-3-anhydro-6-*epi*-ophiobolin B (**TrC-10** and **TrC-11**).³¹⁶

Variculanol (TrC-122, Fig. 7), a sesterterpenoid having a 5/12/ 5 ring system has been isolated from A. variecolor.348 This compound is probably produced from geranylfarnesylpyrophosphate after a requisite folding and cyclizations followed by a 1,5-hydride shift to the carbocation.³⁴⁸ Spectanoids A-G (TrC-311-TrC-317), showing the same ring system, have been later isolated from A. spectabilis.349 This unusual 5/12/5 ring system is also present in nitiol (TrC-184), isolated from Gentianella nitida.³⁵⁰ Spectanoid H (TeC-461), with a 5/8/6/5 ring system, and a different biosynthetic pathway compared to other spectanoids, has been isolated from A. spectabilis.349 Terretonin (TrC-160), a meroterpenoid with a heavily oxidized 25-carbon skeleton, produced from polyketide and terpenoid precursors,³⁵¹ was firstly isolated from *A. terreus* in 1979.³⁵² Further congeners terretonins A-D (TrC-161-TrC-164),³⁵³ terretonin G (TrC-168),³⁵⁴ and terretonins E and F (TrC-165 and TrC-166)³⁵⁵ have been subsequently isolated from the same species, form another Aspergillus sp. strain OPMF00272, and from A. insuetus, respectively. Terretonins H and I (TrC-169 and TrC-170)356 and 1,2-dihydroterretonin F (TrC-167)338 have been isolated from A. ustus. Halorosellinic acid (TrC-96) and 17-dehydroxyhalorosellinic acid (TrC-333) have been isolated from the culture broth of the marine fungus Halorosellinia oceanica.357,358 Clavaphyllene (DC-230), a byciclic hydrocarbon sesterterpenoid structurally distinct from the ophiobolanes, has been isolated from A. clavatus.359 Another byciclic sesterterpenoid, terpestacin (DC-104), was firstly isolated from the marine fungal strain Arthrinium sp.38 Oxidative derivatives of terpestacin have been isolated later from the same fungus, 21-hydroxyterpestacin (DC-107), terpestacin B (DC-109).360 11-epi-Terpestacin (siccanol) (DC-105)³⁶¹ and 11-epi-terpestacin glycoside (DC-106)³⁶² have been isolated from D. siccans and B. sorokiniana, respectively. Bipolarenic acid (DC-210) has been obtained from a marine isolated of the fungus Lophiostoma bipolare (BCC25910).363 Variecolin (TeC-5), a tetracyclic sesterterpenoid with a 5/8/6/5 ring system, was firstly isolated from A. variecolor.364 This compound was later isolated from other fungi, Emericella purpurea,³⁶⁵ E. aurantiobrunnea^{366,367} and Phoma sp.³⁶⁸ TeC-5 congeners, variecolol (TeC-6, Fig. 7) and variecolactone (TeC-11), were isolated from the mycelium of *E. purpurea*,³⁶⁹ and emericolins A-D (TeC-7-TeC-10) have been isolated from E. aurantiobrunnea,366 as well as variecoacetals A and B (TeC-12 and TeC-13).³⁶⁷ The asperanes are hydroxylated 7/6/6/5 tetracyclic sesterterpenoids featuring with a hydroxylated skeleton isolated from Aspergillus fungi. The first asperane-type sesterterpenoid aspergilloxide (TeC-22) was discovered in 2002, from Aspergillus sp.370 Later, six other asperane, asperunguisin

A-F (TeC-396-TeC-399 and TeC-353), were found in A. unguis, which inhabits the lichen Xanthoria sp.371 Aspergstressin (DC-253) hybrid polyketide sesterterpenoid, was discovered from A. sp. WU 243.372 Niduterpenoids A and B (HC-1 and HC-2), characterized by a highly congested hexacyclic 5/5/5/3/5 carbon skeleton, have been isolated from A. nidulans.42 The involvement of a cyclopropane in the ring system makes this skeleton uncommon. A hypothetical biosynthetic pathway from GFPP has been proposed, including a series of cyclization and Wagner-Meerwein hydride and alkyl shift reactions, to the formation of an intermediate with a hexacyclic 5/5/5/3/5 ring system, which after oxidation reactions forms HC-1 and HC-2.42 Several species of the diverged fungal classes Dothideomycetes (Bipolaris, Alternaria, Aplosporella, Pyrenophora) and Sordariomycetes (Arthrinium) are now known to possess highly conserved gene clusters that are mandatory for the sesterterpenoids biosynthesis.373 Co-evolution with host plants, however, has forced pathogenic fungi to acquire novel gene functions and pathways mining terpene scaffolds with new biological potential to overcome competitors or acquire novel physiological functions.374

The fungal terpene biosynthesis has been studied from the last decades of twentieth century. Sesterterpene synthases in fungi are all chimeric proteins consisting of a prenyltransferase (PT) domain at the C-terminus and a terpene cyclase (TC) domain at the N-terminus.^{297,375} Genome mining and heterologous expression of fungal bifunctional sesterterpene synthases have led to the discovery of new sesterpenoids.376-380 The first sesterterpene synthase, Aspergillus clavatus ophiobolin synthase (AcOS), responsible for the biosynthesis of TrC-17 was identified from the genome of A. clavatus in 2013. Two catalytically independent domains (prenyltransferase/terpene cyclase), homologous to those of diterpene synthase, fusicoccadiene synthase, were identified.³⁸¹ A single transformant with the ACLA_76850 gene from A. clavatus produced TrC-17 and three minor sesterterpene hydrocarbons, namely DC-230, 3,20-anhydroophiobolin F (ophiobolane 2) (TrC-297), and ophiobola-1,7,18-triene (ophiobolane 1) (TrC-298).297,359 Heterologous expression of bifunctional sesterfisherol synthase gene (NfSS) and cytochrome P450 monooxygenase (NfP450) from Neosartorya fischeri in A. oryzae system afforded the nitidasane sesterterpenoid sesterfisherol (TeC-20), containing a characteristic 5/6/8/5 tetracyclic ring system. TeC-20 is next modified by cytochrome P450 monooxygenase (NfP450) to sesterfisheric acid (TeC-21).382 An unified biogenesis for sesterterpenes branching from bicyclic (5/15), tricyclic (5/12/5), and tetracyclic (5/6/8/5) cation intermediates, distinct from that of separate class of sesterterpenes including ophiobolins, has been proposed.382

The bifunctional sesterterpene synthase StI-SS in the genome of *E. variecolor* has been identified as responsible for the biosynthesis of the tricyclic sesterterpenoid stellata-2,6,19-triene (**TrC-299**)³⁷⁸ with a 11/6/5 fused ring system. Investigation of the StI-SS gene revealed a gene encoding a cytochrome P450 monooxygenase located next to the StI-SS gene, that catalyzes three successive oxidation reactions on the C-20 methyl

group to generate the carboxylic acid stellatic acid (**TrC-121**),³⁸³ previously identified in *A. stellatus*.³⁷⁸

The sesterterpene synthase EvQS, obtained from E. variecolor NBRC 32302, heterologously expressed in A. oryzae NSAR1, afforded quiannulatene (PC-1), further oxidized to quiannulatic acid (PC-2) by the cytochrome P450 Qnn-P450.375 Genome mining of bifunctional terpene synthase PbTS1 (BtcAPb) against two phytopathogens, Phoma betae and Colletotrichum orbiculare resulted in the production of betaestacins I-IV, Va-c, and VI (TrC-224-TrC-231).³⁸⁴ Functional expression of a terpene synthase (EvAS) from E. variecolor NBRC 32302 in A. oryzae led to the production of astellifadiene (TeC-16), showing a 6/8/6/5 ring system.377 Heterologous expression of four clade-A bifunctional terpene synthases (BFTSs), BmTS1, BmTS2, and BmTS3 from B. maydis ATCC48331 and PbTS1 from Phoma betae PS-13 giving di/sesterterpenes with unique polycyclic carbon skeletons such as sesterfisherol, enabled the isolation of the sesterterpene 5/ 12/5 tricyclic hydrocarbons Bm1 (TrC-308) and betaestacin I (Pb1, TrC-224), the 5/6/8/5-tetracyclic hydrocarbon Bm2 (TeC-505), and of the sesterterpene 5/15 bicyclic alcohol Bm3 (DC-235).385

Based on the initial carbocation formation strategy, the cyclization mechanisms of terpene synthase have been classified into two types, namely A and B. Type A cyclization (C1-IV-V) is initiated between the C1-C15/C14-C18 of geranylfarnesyl diphosphate (GFPP) to yield a 5/15 ring system. Type B cyclization (C1-III-IV) is initiated between the C1-C11/C10-C14 of GFPP/geranylgeranyl diphosphate (GGPP) to yield a 5/11 ring system.386 Cyclization-based classification reflects the phylogenetic relationships among bifunctional terpene synthases. The known enzymes catalysing types A and B cyclization have been classified into two clades, namely clade I, catalysing type A (C1-IV-V) cyclization, and containing terpene synthase genes from five lineages, and clade II, catalysing type B (C1-III-IV) cyclization and containing terpene synthase genes from seven lineages of fungi.386 NfSS is a clade A enzyme, while PaFS and AcOS belong to clade B.382 Anyway, more than 90% of the chimeric terpene synthase genes is still functionally unknown.386 Systematic search of sequenced fungal genomes among diverse taxa revealed that chimeric terpene synthase genes were restricted to Dikarya subkingdom.386 Phylogenetic analysis led to the discovery of a sub-clade involved in the biosynthesis of a 5-15 trans-fused bicyclic sesterterpene, namely preterpestacin I (DC-245). A bifunctional terpene synthase, preterpestacin I synthase (BmTS3), from B. maydis, that catalyses a chain elongation and a cyclization to afford preterpestacin I, was identified. Oxidative modifications from DC-245 to DC-104 are catalysed by enzymes encoded by genes adjacent to BmTS3 (renamed as tpcA), two cytochrome P450 genes (tpcB and tpcC) and a single flavin-dependent oxidase gene (tpcD). The total biosynthesis of DC-104 was then obtained by artificial reconstitution of the biosynthetic machinery in A. oryzae. Heterologous expression in A. oryzae was applied to characterize the function of the putative modification enzyme genes tpcBCD, and this led to the isolation of two biosynthetic intermediates, preterpestacin II and preterpestacin III, and the natural product DC-104.387

Recently, a genomic organization analysis revealed a unique glycosyltransferase gene cluster in the graminaceous pathogen *B. sorokiniana*. This has resulted in the identification of two new metabolites, sestersorokinicin A (**DC-237**, Fig. 7) and sestersorokiniside A (**DC-238**), featuring glucosyl moieties that can enhance the pathogenic effects of bacterial lipopolysaccharide.³⁷³

In 2022, Yan and colleagues elegantly demonstrated that the oxidase $OblC_{Au}$ of *A. ustus* catalyzes dehydrogenation at the C16 and C17 sites of **TrC-17** and **TrC-13** which are intermediates of **TrC-38**. Subsequently, **TrC-38** is transported and stored in a space between the cell wall and membrane. Feedback mechanisms regulate the production of **TrC-38** and its precursors, as their excessive accumulation is closely related to cell toxicity.³⁸⁸

Genome analysis of Penicillium brasilianum NBRC 6234 and Penicillium verruculosum TPU1311 revealed the presence of two bifunctional StTPS genes with prenvl transferase (PT) and terpene synthase (TPS) domains, P. brasilianum sesterbrasiliatriene synthase (PbSS) and P. verruculosum preasperterpenoid A synthase (PvPS), that were heterologously expressed in A. oryzae NSAR1 affording sesterbrasiliatriene (Trc-293) and preasperterpenoid A (PC-79).379 Phylogenetic analysis of the TC domain of protein JNUA3651 from Talaromyces wortmannii ATCC 26942 with those derived from known sesterterpene synthases revealed that its closest neighbor was PvPS, suggesting that JNUA3651 is likely to play the same role as PvPS to synthesize PC-79. Stepwise reconstitution of this gene cluster in A. oryzae NSAR1 revealed that the terpene synthase AstC encodes a sesterterpene cyclase to synthesize PC-79. The P450 enzyme AstB oxidizes PC-79 to give PC-31 along with the minor product asperterpenoid B (PC-64). Subsequently another P450 enzyme AstA oxidizes PC-31 to asperterpenoid C (PC-65).389 Probable pathways catalyzed by AstB were then proposed, the oxidation order of C-19 and C-21 was revealed by quantum chemistry calculations and HPLC-MS analysis, and the intermediates, the three new asperterpenoids D-F (PC-66-PC-68) were obtained.390 Finally, other ten new asperterpenoids, namely asperterpenoids G-P (PC-69-PC-78), featuring a 5/7/3/6/ 5 skeleton, were obtained from two A. oryzae transformants with heterologous expression of a terpene cyclase gene AstC with one or two P450 genes AstB and AstA, by using a molecular networking approach.³⁹¹ Heterologous expression of chimeric enzymes PTTS010 (ZbSS), isolated from Zymoseptoria brevis, and Colletotrichum orbiculare sesterorbiculene synthase (CoSS) in Saccharomyces cerevisiae afforded sesterorbiculene (DC-249) and the 5/8/6/5 tetracyclic sesterevisene (TeC-440, Fig. 7).³⁸⁶

In fungi, the *in vitro* production of ophiobolins changes with the conditions of the culture. For example, *B. maydis* produces **TrC-2**, 3-anhydroophiobolin A (**TrC-4**), ophiobolin B (**TrC-8**), and ophiobolin L (**TrC-43**) in liquid broth, whereas it generates ophiobolin M (**TrC-45**), 6-*epi*-ophiobolin M (**TrC-46**), **TrC-13**, 6*epi*-ophiobolin C (**TrC-14**), **TrC-38**, and 6-*epi*-ophiobolin K (**TrC-39**), when grown in agar media. Ophiobola-7,19-dien-25-oic acid (14,18(*R*)-epoxy-3,5-dihydroxy- γ -lactone) (**TrC-66**) is produced by *Cochliobolus miyabeanus* under modified fermentation conditions.³⁹² More importantly, adding specific substrates, such as methionine, to the culture media of *Bipolaris* spp.

increases the production of **TrC-2** precursor.^{315,347} Generally, sesterterpenoid production increases with the culture time of fermentation in liquid broth. Production of sclareol peaks in old-stage cultures of *Fusarium* spp., *Rhizopus* spp., and *Aspergillus* spp. cultured for more than six days in standard media, indicating that fungal conidia are mainly involved in sesterterpenoids biosynthesis.^{393,394}

Given the broad spectrum of biological activities of ophiobolins, against nematodes, fungi, and bacteria, studies have been done to predict the metabolic pathway of these compounds. Transformations of **TrC-3** with *Polyangium cellulosum* produced **TrC-32** and ophiobolin A lactone (**TrC-7**), while *Pseudomonas aeruginosa* produced ophiobolin B lactone (**TrC-12**). Resting-cell preparations of *Penicillium patulum* afforded **TrC-3**, and 6-*epi*-ophiobolin L (**TrC-44**).³¹⁵

The in vitro standardized production of sesterterpenoids vielded the possibility of better understanding their biological role in the ecology of fungi. Most of the research has been focused on the phytotoxic properties of ophiobolins produced by Bipolaris spp. and Alternaria spp. as they have been implicated in significant plant disease epidemics (the Bengal rice famine in India, 1943 and the spotted leaf blight).^{304,395} When applied to plants, ophiobolins cause detrimental effects, such as growth inhibition of roots and coleoptiles, reduced seed germination, and decreased photosynthesis. The effects of ophiobolins depend on proton extrusion alterations and membrane permeability changes. Indeed, TrC-2 alters the permeability of the plasma membrane to potassium and impairs transport processes resulting in the leakage of electrolytes and glucose from roots and the impaired synthesis of the primary cell wall in plants.³⁹⁶ The phytotoxic effects of ophiobolins require concentrations of approximately 100 µM, which are typically only reached during epidemics. However, during endophytic relationships, fungi produce ophiobolins at concentrations that allow them to obtain ions and sugars from the host without causing significant plant damage.397

4.2.2. Sesterterpenoids in bacteria. Genes involved in terpene, sesquiterpene, and sesterterpene synthase expression have also been identified in bacteria (Table S6, ESI[†]), which therefore represent promising sources for the discovery new

natural sesterterpenoids (Fig. 4c). Bacterial genome sequencing and bioinformatic analysis have recently revealed over 250 terpene synthase genes. However, most of these genes appear to be silent, which explains the relatively small fraction of sesterterpenoids isolated from bacteria compared to fungi, marine organisms, and plants. Environmental cues, symbiotic associations, or competition could activate signalling pathways or repress control circuits to increase sesterterpenoids production in prokaryotic cells.³⁹⁸ Indeed, volatile terpenes can be frequently recognized in odoriferous cultures of Actinomycetales, filamentous Cyanobacteria, Myxobacteria, and Streptomyces albidoflavus. In contrast, sesterterpenoids are uncommon in prokaryotes. Bacterial terpene synthases do not share significant amino acid sequences with those from plants and fungi, thus representing a challenge for the studies and sequence identification.398

In *Streptomyces* spp., cosmopolitan soil bacteria providing valuable secondary metabolites, especially antibiotics,³⁹⁹ a synthase with sesqui-, di-, and sesterterpene synthase activity, coupled with the cyclase StsC, catalyses the formation of a new dicarbocyclic sesterterpenoid, somaliensene A (**DC-239**, Fig. 8), and of one monocarbocyclic (–)-somaliensene B (**MC-149**, Fig. 8) from geranylfarnesyl pyrophosphate.⁴⁰⁰

Yang and colleagues reported that StsC is a membranebound sesterterpene cyclase belonging to the UbiA superfamily of proteins in bacteria. DC-239 and MC-149 were obtained by expressing the corresponding gene in an engineered Escherichia coli strain. Among the 990 homologues of the UbiA family proteins reported in nature, 28 homologues have been identified in the Streptomyces genus However, the sts operon in Streptomyces flanks specific repressors that make StsC and the related products unstable, at least in vitro.401 Sequence analysis revealed that the Bacillus clausii genome contains a promiscuous terpene synthase homologue (Bcl-TS) (a sesterterpene/ triterpene synthase), which potentially catalyses the conversion of linear C35 isoprenoid into monocyclic isoprenoid. However, the cyclization step does not result in the sesterterpenoids production, as revealed by GC-MS analysis of the culture media.402 Nevertheless, the authors were able to purify functional Bcl-TS protein that successfully converted GFPP into



Fig. 8 Representative structures of sesterterpenoids isolated from bacteria.

View Article Online Review

the linear sesterterpenoid β -geranylfarnesene, by introducing the Bcl-TS gene of B. clausii in stable transfected Escherichia coli.402 Recent reports suggest that culture-related factors, such as the media's oxygenation or pH, dramatically change cyclization and the profile production of sesterterpenoids in bacteria.403 Therefore, enzyme activities select the substrates and restrict the range of products, making the *in vitro* bacterial cultures unsuitable tools for the research of sesterterpenoids. In nature, bacteria grow in structured communities, forming microcolonies that bound to surfaces to form biofilms. Biofilm formation is a dynamic process initiated by the early colonizers, which determine the three-dimensional expansion of the bacterial communities.404 The colonizers elaborate the outer polymer matrix to capture nutrients and new microbes, leading to a complex, multi-species biofilm that represents a protected environment allowing cells to survive in hostile situations. In the biofilm, bacteria communicate through small diffusible molecules to regulate gene expression, control the production of secondary metabolites, organize the community's structure and relationships, and colonize new surfaces by dispersing bacteria.405 Streptomyces albus forms a compact biofilm in which cells are embedded in an extracellular protein matrix composed of a network of fimbriae.406 The biofilm allows S. albus to maintain its metabolic functions, including the catalysis of regio- and stereo-selective hydroxylation.407 The co-cultivation of S. albus with Bacillus amyloliquefaciens leads to an increase in the formation of fimbriae and biofilm stability. This suggests that secondary metabolites extend the half-life of bacteria in the biofilm.406,408

Improved yeast-based promoter engineering platform (mCRISTAR),⁴⁰⁹ has been used to activate the previously uncharacterized Class II cyclase-containing gene cluster, called atolypene (ato) gene cluster, cloned from the genome of the cultured actinomycete *Amycolatopsis tolypomycina*. Heterologous expression of ato gene cluster into *S. albus* led to the characterization of atolypenes A and B (**TrC-144** and **TrC-145**, Fig. 8).⁴¹⁰

Two clade III promiscuous terpene synthases (Fusarium graminearum mangicdiene synthase, FgMS and F. graminearum GJ1012 synthase, FgGS) from an endophytic fungus, F. graminearum, showed to produce variable terpenoids in vivo by converting precursor polyisoprenoid diphosphates of different lengths (C10, C15, C20, C25). Six Escherichia coli variants, obtained by combining two terpene synthases and three PTs, afforded 50 different terpenoids, including the 11-6 bicyclic variecoltetraene (DC-234), the 5/5/6/5 tetracyclic mangicdiene (TeC-482), and (2E)-α-cericerene (MC-114). Further exploitation of mutants F65L and F159G afforded the 5/8/6/6 tetracyclic sesterterpene Tec-509, Fig. 8.411 Recently, Gu et al. reported the discovery of the sesterterpene synthases Streptomyces violens sesterviolene synthase (SvSS) from S. violens, that converted GFPP into a sesterterpene hydrocarbon, sesterviolene A (TeC-464), and a few trace compounds. Enzyme engineering through site-directed mutagenesis gave access to a high-yielding enzyme variant that provided six additional minor products sesterviolenes B-G (TrC-331, TeC-466, TeC-467, DC-250-DC-252) and the main product TeC-464.412 The guanidine-containing scalarane

scytoscalarol (TeC-366) has been isolated from the cyanobacterium Scytonema sp. (Scytonemataceae).413 Other guanidinebearing compounds are cybastacines A and B (TeC-359 and TeC-360, Fig. 8), isolated from another cyanobacterium, Nostoc sp. (Nostocaceae).⁴¹⁴ Only compounds with simpler structures, such as the acyclic sesterterpenoids (AI, Table S1, ESI[†]), are known in benthic diatoms. C25 highly branched isoprenoid alkenes (haslenes) are ubiquitous in marine sediments.415 These polyunsaturated sesterterpene oils have been isolated from several species of Haslea (Naviculaceae), i.e. H. ostrearia (AI-14-AI-21, AI-23 and AI-24), H. crucigera, H. pseudostrearia (AI-18) and H. saltstonica (AI-16).416,417 Similar compounds (AI-12 and AI-13 rhizenes) have been isolated from the marine diatom Rhizosolenia setigera (Rhizosoleniaceae).418 A promising source of sesterterpenoid synthesis could be that of lichenised Ascomycota species. As these compounds have been shown to mediate trophic or defensive interactions in many fungi, it can be expected that a wide variety of sesterterpenoids will be present in lichen species that have to cope with biotic relationships between mycobionts, algal or cyanobacterial phototrophs and a major component of the microbiota. To date, however, knowledge in this regard is relatively scarce. From Gypsoplaca macrophylla (Gypsoplacaceae), gypmacrophin A (PC-34, Fig. 8), showing a skeleton similar to asperterpenoids isolated from Aspergillus and other fungal species, has been isolated.379,391 Retigeranic acid A (PC-7) has been purified from Lobaria isidiosa var. subisidiosa, L. retigera, and L. subretigera (Peltigeraceae), while the epimer retigeranic acid B (PC-56) has been described only from Lobaria isidiosa var. subisidiosa.419-421 Retigeran-11-ol (PC-32) and 4-hydroxyretigeran-11-ol (PC-33) isolated from Leprocaulon microscopicum have been (Leprocaulaceae).422

4.2.3. Sesterterpenoids drive symbiotic relationships in fungi and bacteria. Endophytic fungi and symbiotic bacteria have coevolved with higher plants or other microorganisms, providing an eclectic metabolic potential that enables adaptation to the specific host and complements missing synthetic pathways.423 Among fungi (Table S6, ESI†), Alternaria spp. are among the most widely distributed existing endophytes of plants. The glucosyl sesterterpene, 24-a-p-glucosyl-(-)-terpestacin (DC-108) has been isolated from Alternaria alternata, an endophytic fungus from the fresh root of Ceratostigma griffithii.424 The 5/8/6/5 tetracyclic sesteraltererol (TeC-490) and 10,11-epoxysesteraltererol (TeC-491), together with DC-245, have been identified in A. alternata living in symbiosis with Leucosceptrum canum.⁴²³ Emericellenes A-E (DC-205-DC-209), showing an emericellane-type bicarbocyclic ring system, were obtained from the endophytic Emericella sp. AST0036 isolated from Astragalus lentiginosus.49 16,17-Dihydro-(-)-terpestacin (DC-110) and terpestacin C (DC-111) have been isolated from Aplosporella javeedii, obtained from the stem tissue of Orychophragmus violaceus.425 Ophiobolin O (TrC-49) and 6-epi-ophiobolin O (TrC-50) have been isolated from endophytic Aspergillus sp. from the body of Zoanthus.426 Ophiobolins R-T (TrC-55-TrC-56) have been obtained from the endolichenic Ulocladium sp. isolated from Everniastrum sp.334 Several sesterterpenoids have been isolated from mangrove endophytic fungi. When

associated with mangroves, Fusarium spp. produce neomangicols (TeC-370-TeC-372) and mangicols A-G (TeC-374-TeC-380), which represent novel classes of sesterterpenoids with an unprecedented carbon skeleton, including spirotricyclic structure components.427 These sesterterpenoids have not been recovered from in vitro pure cultures of the fungi. Fusaproliferin (DC-112), a 15/5-membered ring system with three trans olefins in the 15-membered ring, was firstly isolated from Fusarium proliferatum.428-430 From the same fungus, obtained from the fresh tissue of the marine mangrove plant Bruguiera sexangular, the similar compounds fusaprolifins A and B (DC-113 and DC-114) have been isolated.431 DC-112 has been also characterized from a strain of Fusarium solani, isolated from the plant Aglaonema hookerianum.432 In 2013, a novel 5/7/3/6/5 pentacyclic sesterterpenoid, asperterpenoid A (PC-31) was isolated from a mangrove endophytic fungus Aspergillus sp.433 Asperterpenols A and B (TeC-97 and TeC-98), with a 5/8/6/6 tetracyclic carbon skeleton, were isolated from a mangrove endophytic fungus Aspergillus sp.434 Aspterpenacids A and B (PC-38 and PC-39), featuring a 5/3/7/6/5 ring system, have been isolated from the endophytic fungus A. terreus of the mangrove plant Kandelia obovata.435 In the proposed biosynthetic pathway an intermediate with a 15/5 ring system, deriving from GFPP by head-to-tail connection and cyclization, originates the 5/6/7/3/5 carbon skeleton. Finally, further oxidation, reduction, and acetylation can generate PC-38 and PC-39.435 The endophytic fungal strain Aspergillus sp. ZJ-68, collected from the leaves of the mangrove plant Kandelia candel, afforded the nitride ophiobolins asperophiobolins A-K (TrC-67-TrC-77), with an additional fivemembered lactam ring between C-5 and C-21. Asperophiobolins A-K (TrC-67-TrC-70) were isolated from the fermentation culture of the mangrove endophytic fungus Aspergillus sp. ZJ-68 as the first ophiobolin derivatives with a five-membered lactam unit between C-5 and C-21.333 The nitidasane sesterterpenoid sesteralterin (TeC-393) has been obtained from the culture extract of an Alternaria alternata strain (k21-1) isolated from the surface of the marine red alga Lomentaria hakodatensis. Behind the structural support, endophytic fungi exploit the cyclization catalyzes of critical bifunctional diterpene synthase to direct by-products to the cytochrome P450 oxidation, resulting in the production of conidiogenone, a diterpene inducer of conidiation. Upon production, conidiogenone rapidly accumulates at the surface of the hypha. Once it reaches the threshold concentration, it triggers conidiogenesis.436 Conidiogenesis induces the growth of hyphae to further penetrate the substrates of plants for anchoring the mycelia and mining nutrients.437 As endophytes reside in association with plants for at least a part of their life cycle, it is possible that fungi can detect plant-derived signals to adjust their metabolism and growth to environmental conditions. Plant homoserine and asparagine induce gene expression in fungi, whereas horizontal gene transfer (plant to endophyte genome or vice versa) explains the metabolic complementation during symbiosis.438,439 Multiple species of microbes colonize plant tissues or marine organisms, resulting in interactions between associated endophytes, including fungus-fungus, fungus-bacteria, and bacteria-bacteria relationships. These

intricate networks of microbial interactions significantly impact metabolites production.374 The networks involve metabolic elicitors, such as quorum-sensing signal molecules that trigger otherwise silent pathways.440,441 Co-culture of different microbes that mimic the competition of natural ecosystems rather than maintaining axenic cultures is increasingly practiced in microbial natural product research.442 This strategy can stimulate the transcription and translation of genes, and enhancing the production of constitutively present natural products, or triggering the expression of silent biosynthetic pathways with the production of new compounds.443-445 The fermentation broths of coculture of basidiomycetes Trametes robiniophila and Pleurotus ostreatus afforded the linear sesterterpenoids postredienes A-C (AI-5-AI-7).442 Anyway, dynamic ¹³C-labeling analysis showed that sesterterpenoids were synthesized by P. ostreatus instead of T. robiniophila.442 Physical interactions between A. nidulans and Streptomyces spp. induce epigenetic regulation of secondary metabolism as previously observed in lichens.441 In these organisms, specific interactions between microorganisms belonging to different domains produce physical interactions that induce otherwise silent biosynthesis genes in response to environmental factors or internal biological processes. These observations demonstrate that novel terpenoid structures can be isolated from different ecological environments. Marine fungi such as the Aspergillus genus typically coexist with different sponges. In the Mediterranean Sea, A. insuetus produces different secondary metabolites if associated with Petrosia spp. or Psammocinia spp. sponges.⁴⁴⁶ Marine sponges also host a bacterial population that has not yet been fully characterized. Metagenomic analysis of the sponge Ircinia ramosa revealed that its microbiota is composed of 32% sequence-associated Archaea and 41% sequence-associated Bacteria. The metabolic reconstruction showed extensive redundancy across taxa.447 Bacteria inside sponges have both commensal and parasitic relationships, and the density and diversity of resident bacteria control metabolic production. The density of chemosynthetic autotrophs or heterotrophs increases from the inner to the outer, while cyanobacteria preferentially reside at the borders.^{20,447,448} However, there is still limited literature on the interaction of bacteria and sponges and its impact on sesterterpenoids production.

4.3 Sesterterpenoids in plants

With over 350 000 known species, plants (Table S7, ESI†), are a megadiverse kingdom of organisms.⁴⁴⁹ They were among the first organisms to colonise terrestrial environments, and have acted as habitat builders, significantly modifying the chemical composition of the Earth's atmosphere and the chemical and physical properties of the soil. Plant species have developed a complex array of primary and secondary metabolic pathways in response to environmental pressures. These pathways facilitate to the synthesis of metabolites that are biologically active within the plant and potentially useful for regulating the physiology of other organisms. Plants have been a significant source of food and medicine since ancient times, owing to their properties. In a constant balance between an adaptive response to the pressures of the environment and a modifying effect on it, plant species have differentiated in their genome a very rich complexity of primary and secondary metabolic pathways capable of leading to the synthesis of different compounds. These compounds can be functionally active both for the plant healthiness and potentially for regulation of the physiology of other organisms. Regarding sesterterpenoids, vascular plants (Tracheophyta) are one of the richest phyla with 219 known compounds, following Porifera and Ascomycota (Fig. 1). The diversity of compound classes (6 out of the 10 considered in this review) is remarkable, and the most represented class being dicarbocycles (DCs) with 114 known compounds (Fig. 1). TrCs (Table S1, ESI,[†] and Fig. 4d) are the class of sesterterpenoids most shared between different plant families, even those that are phylogenetically distant from each other. In some families, this is the predominant or even exclusive class of sesterterpenoids, such as Pteridaceae.450,451 Cheilanthane sesterterpenoids have been isolated from Cheilanthes farinosa, cheilarinosin (TrC-102, Fig. 9)⁴⁵² and cheilanthatriol (TrC-138),⁴⁵³ Aleuritopteris khunii (TrC-236 and TrC-237),450 A. agetae (TrC-300-TrC-302).451 17-Oxo-18,19-bisnorcheilanth-13(24)-en-6a-ol, a 19,20bisnorcheilanthane (C23 sesterpenoid) (TrC-303) and 13,17dioxo-18,19,24-trisnorcheilanth-6a-ol, a 19,20,21-trisnorcheilanthane (C₂₂ sesterpenoid) (TrC-304), have been also isolated from A. agetae.451 Ancepsone A (TrC-320), isolated from A. anceps,454 and (17Z)-13,19-epoxycheilanth-17-en-6a-ol (Tec-506, Fig. 9), from A. mexicana,455 showed a 13,19-epoxycheilanthane skeleton. 18-epi-Scalar-16-ene-6a,19-diol (Tec-507) and 16a,19epi-dioxy-18-episcalar-17(25)-en-6a-ol (Tec-508) have been also isolated from A. mexicana. Other cheilantanes, cristasesterterpenoic acid and cristasesterterpinol glucoside (TrC-276 and TrC-277), have been obtained from Caesalpinia crista

(Fabaceae).456 Involudispirones A and B (TrC-278 and TrC-279, Fig. 9), containing a 1,2-dioxadispiro[5.2.5.2]hexadecane ring system have been isolated from Stahlianthus involucratus (Zingiberaceae).457 It has been speculated that this spiro system could be formed by an enzyme-catalyzed Diels-Alder addition between the derivative of cadalenequinone, a major constituent in S. involucratus, and myrcene.457,458 Similarly, it has been hypothesized that heliocide H2 (TrC-157), isolated together with heliocides B1 (TrC-155) and H1, H3 and H4 (TrC-156, TrC-158, and TrC-159) from Gossypium hirsutum (Malvaceae),459-462 could be biosynthesized by a Diels-Alder addition between the sesquiterpenoid gossypolone and myrcene.461 A Diels-Alder reaction has been used to synthesize TrC-155 and TrC-159 from Linderhemigossypolone and trans-β-ocimene.462 asesterterpenoids A and B (TrC-319 and TrC-319) with an unusual 7-cyclohexyldecahydroazulene carbon skeleton have been isolated from the roots of Lindera glauca (Lauraceae).463 2'-Isopicrasin A (TrC-288)⁴⁶⁴ and picrasin A (TrC-289),⁴⁶⁵ simaroubolides having the C25 simarolidane skeleton which consists of three carbocycles and two lactone rings, and is closely related to simarolide, have been isolated from Picrasma quassioides (Simaroubaceae).465 Vulgarosides 1-4 (TrC-245-TrC-248), sesterterpene esters whose base aglycon skeleton is quite similar to TrC-280 (except for the stereochemistry at C-13 and C-14), have been obtained from Cydonia vulgaris (Rosaceae). The distribution of TrCs in vascular plants suggests a more ancestral inheritance of the genes involved in the synthesis of these compounds. While a detailed discussion of the phylogeny of sesterterpenoids in plants is not a primary objective of this review, it is worth noting that there are several gaps in our knowledge regarding this topic. Based on current knowledge, Lamiaceae is the largest source of compounds among the 22

PC-40



TrC-279



Fig. 9 Representative structures of sesterterpenoids isolated from plants

TrC-278



ĥ ŌН

TeC-435

•OH

TrC-188

vascular plant families for which sesterterpenoids have been described, with DC and MC being the most important. Coleifolides A and B (MH-50 and MH-51), characterized by a β methyl- α , β -unsaturated- γ -lactone and structurally similar to manoalide derivatives, have been isolated from Scutellaria coleifolia.466 Leucosceptrane sesterterpenoids (leucosceptroids), colquhounoids and eurysoloids have been isolated from the Lamiaceae Leucosceptrum canum, Colquhounia coccinea var. mollis and Eurysolen gracilis. From the trichome exudates and the leaves of L. canum bicarbocyclic sesterterpenoids have been isolated, i.e. leucosceptroids A and B, E-Q (DC-168 and DC-169),467 leucosceptroids E-N (DC-185-DC-192, DC-138, and DC-229) showing an α,β -unsaturated γ -lactone moiety,⁴⁶⁸ leucosceptroids P and Q (DC-198-DC-199),469 leucosesterlactone (DC-164),⁴⁷⁰ 17α-hydroxyleucosceptrine (DC-166),⁴⁷¹ leucosceptrine (DC-211).⁴⁷² Leucosceptroid O (DC-193), possessing a spiro α , β unsaturated y-lactone moiety, has been isolated from the flowers of the same species.473 Additionally, tricarbocyclic compounds with antipodal cyclopentenones leucosceptroids C-D (TrC-190 and TrC-191),⁴⁷⁴ leucosesterterpenone (TrC-188, Fig. 9),⁴⁷⁰ and 14β-methylleucosesterterpenone (TrC-189),⁴⁷¹ have been found. Leucosceptroid degradation products, *i.e.* the C20 norleucosceptroids A-C (DC-172-DC-174), and the C21 norleucosceptroids D-H (DC-200-DC-204).469,475 have been isolated from L. canum of Nepalese and Chinese origin, respec-1α-Hydroxyleucosceptrine tively. (DC-212) and 8αhydroxyleucosceptrine (DC-213) were produced microbial transformation of DC-211 by Rhizopus stolonifer.394 Colquhounoids A-D (DC-182-DC-184 and DC-246), and 14-epi-colquhounoid D (DC-247) have been isolated from Colquhounia coccinea var. mollis.476,477 Eurysoloids A and B (TeC-441 and TeC-442), characterized by a pentacyclic 5/6/5/10/5 scaffold with an unusual macrocyclic ether system have been isolated by Eurysolen gracilis.478 The genus Salvia has received much attention within Lamiaceae. Sesterterpenoids have been found in both cultivated plants e.g. S. tingitana479 and wild species, such as in the case of S. dominica from subarid regions of Jordan.480,481 Prenyllabdane-type sesterterpenoids²⁵ have been isolated from several Salvia species, as S. syriaca (DC-59, Fig. 9),482 S. hypoleuca (DC-64, DC-67, DC-68, and DC-70),483,484 S. palaestina (DC-73-DC-75),485 S. dominica (DC-81-DC-91, DC-94-DC-96, DC-101 and DC-102),481 S. tingitana (DC-120, DC-127, DC-129-DC-132, DC-146, DC-147),479 S. yosgadensis (DC-134), and S. mirzayanii (DC-121).486 Prenyllabdane sesterterpenoids with a lactone, ester or acetal functionality between C6 and C23 have been found in various species, as S. hypoleuca (DC-65, DC-66, DC-69, and TrC-223),483,484 S. tingitana (DC-127, DC-143-DC-145),479 S. sahendica (DC-71 and DC-72),487 S. dominica (DC-76-DC-80, DC-92-DC93, DC-101 and DC-102),481 S. lachnocalyx (DC-97-DC-98).488 Other metabolites with a tetrahydropyran ring similar to manoyloxidetype diterpenoids between C-8 and C-13 were obtained from S. mirzayanii (DC-122-DC-126)489 as well as other hydroxymanoyloxide derivatives (DC-140-DC-142) from S. limbata,490 (14E)methylmanoyloxide-14,16,18-trien-19,16-oxide-23-carboxylate (DC-139) from S. tingitana,479 lachnocalyxolides C and D (DC-99-DC-100) from S. lachnocalyx,488 yosgadensolides A and B (DC-134 and DC-135) along with their epimers from S. yosgadensis,⁴⁹¹

salviaethiopisolide derivatives (DC-219-DC-220)492 and others hydroxymanoyloxides (DC-221-DC-223)493 from S. aethiopis. A norprenyllabdane-type sesterterpenoid, (13E)-4a,6a,8atrihydroxylabd-13(14),17(18)-dien-16,19-olide (DC-132), was isolated from S. tingitana.479 Compounds arising from degradation of C-19 and C-20, 19,20-dinorsesterterpenoids were yosgadensenol (DC-133) and 13-epi-yosgadensenol (DC-42) from S. yosgadensis,494 and 6-dehydroxyyosgadensenol (DC-137) and 6dehydroxy-13-epi-yosgadensenol (DC-136) from S. limbata.490 The sesterpene γ -lactone genepolide (DC-167, Fig. 9), isolated from Artemisia umbelliformis (Asteraceae) has been described as a formal Diels-Alder adduct of the exomethylene-y-lactone germacranolide costunolide and the diene myrcene.495 Another intermolecular Diels-Alder reaction between thujanone, a thujane-type monoterpene one of main constituents in the essential oil of A. argyi, and a guaianolide, has been hypothesized for the biosynthesis of isoartemisolide (PC-26), isolated from the same species.496 Raoulic acid (DC-218), isolated from Raoulia australis, probably biosynthesized by diprenylation of a sesquiterpenoid precursor of the germacrene type, common in other Asteraceae species.497 Dibritannilactone A (TeC-45), isolated from Inula britannica, may derive from a monoterpene and a sesquiterpenoid.^{25,61} The trinorsesterterpene glycoside 3-[6-(4,8-dimethyl-nona-1,3,7-trienyl)-4-hydroxy2,6-dimethyl-

cyclohex-1-enyl]-3-hydroxypropionic acid 1 glucoside (MC-99) has been obtained from the fern Woodwardia virginica (Aspleniaceae).498 The scalarane corallocarpscalarolide (TeC-410) has been isolated from the roots of Corallocarpus epigaeus (Cucurbitaceae).499 Biyoulactones D and E (DC-170 and DC-171), tricyclic meroterpenoids possessing an octahydroindene ring, a γ -butyrolactone ring, and a β -diketone moiety, have been isolated from Hypericum chinense (Hypericaceae).⁵⁰⁰ From the fruit of Phellodendron chinense var. glabriusculum (Rutaceae), phellogine (TeC-32), whose structure is similar to tirucalla-7-ene derivatives in the same species, has been obtained.⁵⁰¹ Goniocarpic acid (DC-214) has been described within the constituents of the leaves of Serjania goniocarpa (Sapindaceae). Nevertheless, the classification of DC-214 as a sesterterpenoid has been questioned as it cannot be derived from GFDP.25 Among Monocotiledonous species, Aletris farinosa (Nartheciaceae), afforded two sclarane (TeC-226 and TeC-227) and four cheilantane sesterterpenoids (TrC-274, TrC-275, TrC-290, and TrC-291) have been studied.^{502,503} The scalarane sesterterpenoid perisomalien A (TeC-34)⁵⁰⁴ and the monocyclic *n*-non-2'-en-1'-yl-13(15,19,19trimethyl-cyclohex-14,16-dienyl)-2,6,10-trimethyl-tetradec-6-ol-13-on-1-oate (hemidesmu sesterterpenoid ester) (MC-98), have been isolated from Periploca somaliensis and Hemidesmus indicus, respectively.⁵⁰⁵ The potential of sesterterpenoids in plants is still largely untapped, as evidenced by the case of Gentianaceae.²⁵ Nitidasin (TeC-352)⁵⁰⁶ was the first gentianellane-type²⁵ sesterterpenoid isolated from the Andean species Gentianella nitida. From the same species, nitiol was isolated (TrC-184).350 It has been speculated that the seco-gentianellane alborosin (TrC-109) from G. alborosea, could be derived from the cyclization of GFPP through TeC-352 as a key intermediate, and after the final oxidative cleavage of the C11/C12 bond.507 Later, other gentianelloids have been isolated from the Chinese G. turkestanorum,

Natural Product Reports

namely the C-6 epimeric sesterterpenoids gentianelloids A and B (TrC-286 and TrC-287), two compounds containing an unusual 10,11-seco-gentianellane skeleton,508 and 18-epi-nitidasin (PC-80), gentianelloids C-F (TeC-447, TeC-449, TeC-451, TeC-453), 18-epi-gentianelloids C-F (TeC-448, TeC-450, TeC-452, TeC-454), and 18-epi-alborosin (TeC-455).509 Another species belonging to Gentianaceae, Swertia bimaculata, afforded aspterpenacid C (PC-40, Fig. 9)⁵¹⁰ with the 5/3/7/6/5 pentacyclic skeleton similar to aspterpenacids A and B (PC-38 and PC-39) mangrove endophytic fungus Aspergillus terreus, isolated from Kandelia obovata (Rhizophoraceae).435 Their biosynthetic pathway could be derived from GFPP with a series of cyclizations, rearrangements, redox and acetylation reactions.435 Bolivianine (PC-53) and isobolivianine (PC-53), two sesterterpenoids with an unusual skeleton, have been isolated from the bark of Hedyosmum angustifolium.⁵¹¹ The authors hypothesized their biosynthesis starting from onoseriolide, a sesquiterpene lactone already isolated from other Chloranthaceae. Another species, H. brasiliense, afforded hedvosulide (TrC-20), probably

biogenetically related to the α -exo-methylene- γ -lactone precursor hedyosumin A and myrcene.⁵¹² In the cormophytes, acyclic sesterterpenoids (AI, Table S1, ESI†) have been obtained from Poaceae, Euphorbiaceae, and Solanaceae. 2,6,10,14-Tetramethyl-18-butanecarboxymethylene-henecos-12-en-17β-ol (AI-11) has been isolated from Oryza sativa,³³ and (2Z,6Z,10E,14E)-geranylfarnesol (AI-8) from Triticum aestivum.513 sesterterpenoids, 2E-3,7,11,15,19-Other linear pentamethyleicos-2-en-1-ol (AI-22) (2Z,6E,10E,14E)-geranylfarnesol (AI-9), have been isolated from the aerial parts of Croton hieronymi (Euphorbiaceae).⁵¹⁴ 3,7,11,15,19-Pentamethyl-2-cis-6-trans-eicosadien-1-ol (AI-4) has been obtained from the unsaponifiable matter of the leaves of Solanum tuberosum (Solanaceae).515

Among terrestrial organisms, vascular plants are therefore the most important source of sesterterpenoids.²⁵ It is interesting to examine the geographical distribution of the source species of sesterterpenoids on a global scale. Fig. 10 shows the most noteworthy areas for raw materials collection to be



Fig. 10 Worldwide distribution of plant species known to be sources of different macrocategories of sesterterpenoids. The intensity of the colours in the figure is proportional to the number of taxa. The abbreviations of the macrocategories of compounds are as follow: AI = acyclic; MC = monocarbocyclic; DC = dicarbocyclic; TrC = tricarbocyclic; TeC = tetracarbocyclic; PC = pentacarbocyclic. Monoheterocyclic (MH), diheterocyclic (DH), triheterocyclic (TH) and hexacarbocyclic (HC) compounds are not included in the analysis, as they have little or no representation for plants in the database.

exploited for a potential use in the pharmaceutical field. The spatial distribution patterns of the compounds also reveal regions where it may be beneficial to increase eco-metabolomic research on sesterterpenoids. Overall, China, India and, more generally, the Asian continent are the areas with the greatest wealth of sesterterpenoids rich vascular plants even if Australia (for MC, DC and TrC), the Andean regions of South America (for AI) and the Mediterranean areas (for DC) are also notable. It should be noted that this distribution partially overlaps with the world's floristic hotspots. However, some areas with high specific richness and high rates of endemism, such as Macaronesia, Amazonia, or the Alpine regions, appear surprisingly underrepresented and may still contain a considerable wealth of sesterterpenoid diversity.

Thanks to the investigation of new biosynthetic chemistry, undescribed plant sesterterpenoids have been defined. By genome mining and gene clusters analysis, it appears that many genes and gene clusters do not direct the production of known metabolites of the organism in which they are found and that novel gene clusters provide the biosynthesis of known natural products.516 Moreover, the induced expression of silent genes can lead to the isolation of new metabolites.382 Terpenoid diversity in plants is determined by short-chain prenyltransferases (SC-PTs) and terpene synthases (TPSs). Among the SC-PTs, geranyl-farnesyl pyrophosphate synthases (GFPPS) produce the sesterterpene base scaffold GFPP.^{28,381,517} Sesterterpene synthases (sester-TPSs), a clade of the TPSa subfamily, are a branch of class I TPSs, firstly identified in Brassicaceae,28 and widely distributed in the plant kingdom. Sester-TPSs catalyse diphosphate abstraction, cyclization of C1-C15 and C14-C18, producing a 5/15 bicyclic carbocation, which then undergoes structural modifications to yield diverse polycyclic sesterterpene scaffolds.516 Many sesterterpenoids contain a cyclopentane moiety, generated via type A (C1-IV-V) (between the C1 cation, the C14-C15 double bond (IV), and the C18-C19 double bond (V)) or type B (C1-III-IV) (between the C1 cation, the C10-C11 double bond (III), and the C14-C15 double bond (IV)) early-stage cyclization mechanisms of GFPP in the catalytic pocket of plant sesterTPSs.^{28,51,518} Type A cyclization is mainly represented in plants. Examples are (-)-caprudiene A (TeC-435, Fig. 9), produced by transient expression of individual STS with a GFPPS gene by "agro-infiltration", with Agrobacterium tumefaciens infiltrated into the undersides of leaves of Nicotiana benthamiana,⁵¹⁹ and (-)-variculatriene A (TrC-117).^{51,519} Type-B Sester-TPSs, considered as possible progenitors of Type-A Sester-TPSs, have been discovered in Brassicaceae. AtTPS06, identified in Arabidopsis thaliana, is the first gene found in plants which produces the E/Z isomer of flocerene (TrC-292) via type-B cyclization mechanism.^{28,517,520} In the genome of A. thaliana, AtsesterTPS1, responsible for the biosynthesis of the new tricyclic sesterterpene (+)-thalianatriene (TrC-120) with the unprecedented 11-6-5 fused ring system, and AtsesterTPS2, which produced (–)-retigeranin B (PC-4), with the characteristic 5/5/5/6/5 fused ring system, were identified.521 Transient expression of TPSs from A. thaliana, Capsella rubella, and Brassica oleracea in Nicotiana benthamiana afforded sesterterpenoids with various scaffolds, as (-)-caprutriene (TrC-

116), (-)-variculatriene A (TrC-117), (-)-aleurodiscalene A (TeC-17), (-)-ent-quiannulatene (PC-3), (+)-boleracene (PC-6), and (+)-astellatene (PC-9).⁵¹ Engineered Nicotiana benthamiana expressing LcTPS2, characterized from Leucosceptrum canum, produced two 18- and four 14-membered ring sesterterpenoids (S,2E,6E,10E,14E)-3,7,11,15-tetramethyl-18-(1including hydroxyisopropyl)cyclooctadecatetraene (MC-141), (S,2E,6E,10E,14E)-3,7,11,15-tetramethyl-18-(1-methylethenyl) cyclooctadecatetraene (MC-142), (S,2E,15Z)-a-cericerene (MC-143), (S,2E)-cericerene (MC-114), (S,2E)-α-cericerene and (14S,15R,2E)-15-hydroxy-α-cericerene (MC-145).⁵²² Engineered Escherichia coli expressing CcTPS1, obtained from Colquhounia coccinea var. mollis, produced (+)-a-geranylbisabolene (MC-150) and (+)-somaliensene B (MC-151), characterized by an alotane skeleton, not previously isolated from any plants.⁵²³ Trans-prenyltransferase (PT) and N-terminal terpene synthase (TPS) gene pairs from Arabidopsis thaliana have been reported to synthesize sesterterpenoids. AtsesterTPS2 produced the pentacyclic sesterterpenoid (-)-retigeranin B (PC-4) with the characteristic 5/5/ 5/6/5 fused ring system. AtsesterTPS1 produced the tricyclic (+)-thalianatriene (TrC-120).521

5 Biological and pharmacological properties of sesterterpenoids

Over the years, sesterterpenoids have been investigated for their action on a wide range of biological activities of pharmacological importance including suppression of cancer cell growth, anti-inflammatory properties, mainly counteracting PLA2 activity, and modulation of neurodegenerative processes (Fig. 11). Moreover, they were explored for their key functions such as antimicrobial effects against fungi, Gram + and - bacteria, viruses such as HIV or as anti-malarial and anti-tuberculosis agents. Sesterterpenoids have been also inspected as ichthyoand phyto-toxins, as nematocidals, and antifeedants from an ecological point of view. Few of them were tested for the involvement in the treatment of metabolic diseases such as type-II diabetes, hypercholesterolemia and obesity and as immunosuppressive agents. Here, we briefly discuss those sesterterpenoids whose mechanism of action has been thoroughly investigated, focusing on their target enzymes. In parallel, a more comprehensive list of sesterterpenoids biological activity can be found in the annexed Table S8, ESI,† and at https://sesterterpenoids.unige.net/. Selected sesterpenoids structures were numbered to help reading.

5.1 Sesterterpenoids with anti-cancer activity

The majority of sesterterpenoids were probed for their cytotoxicity, predominantly against cancer cells. Principally, ophiobolins were screened *versus* multiple cancer cell lines: among the approximately 80 identified ophiobolins, about 20 were reported for their cytotoxic capability. Mainly, **TrC-2** (Fig. 3) displayed growing inhibition against 25 human cancer cells: it principally counteracts apoptosis-resistant glioblastoma cells by inducing a non-apoptotic cell death *via* reaction with primary amines prompting pyrrolylation of lysine residues on



Fig. 11 Circular plots showing the relationships among referred compound categories and their biological activities. The dimension of color bands is proportional to the number of compounds found in each category. Abbreviations are as follow: BACT = anti-bacterial; CANC = anti-cancer; CYTO = cytotoxic; ENZ = enzyme inhibitor; FEED = anti-feedant; FUNG = anti-fungal; IMM = immunosuppressive; INFL = anti-inflammatory; MICR = anti-microbial; VIR = antiviral; OTHER = other activities; UNKNOWN = unknown effect. For sesterterpenes categories: AI = acyclic, linear; MC = mono-carbocyclic; MH = monoheterocyclic; DC = dicarbocyclic; DH = diheterocyclic; TC = tricarbocyclic; TC = tetracarbocyclic; TH = triheterocyclic; PC = pentacarbocyclic; HC = hexacarbocyclic.

its intracellular target protein.524 Intrigued by this property, Fujiwara studied the behaviour of TrC-2 in the L1210 cell line in 2000, showing a concentration-dependent cytotoxicity.⁵²⁵ This compound provoked paraptosis-like cell death in human glioblastoma cells by altering the cytoskeleton dynamic pathway.526 Furthermore, TrC-2 selectively inhibited the correct development of solid and haematological cancer cells with a very low IC₅₀; it also prompted apoptosis in MDA-MB-231 cells through the inhibition of PI3K/mTOR, Ras/Raf/ERK and CDK/RB pathways³¹³ and it has been discovered to significantly repress the mammosphere formation.527 This super active compound has also a role in fighting multi-drug resistant cancer cells, as it acts against HL60 cells that are resilient to combined chemotherapy, ovarian carcinoma cells that are resistant to cisplatin, small lung carcinoma cell line GLC4 that is resistant to adriamycin, and the others.526 Ophiobolins not only were extensively studied for their in vitro cell inhibition, but also reduced tumour development in in vivo animal models. For instance, TrC-2 was inoculated in a mouse model of glioblastoma⁵²⁸ and of melanoma.526 Later on, TeC-326, heteronemin acetate (TeC-349), hyrtiosin E (TeC-309), 12-deacetoxy-scalarin 19-acetate (TeC-176) and TrC-130 were tested against a large panel of cells such as HuCCA-1, HeLa, MDA-MB-231, MCF7, HT-29, and H69AR, KB, human colon adenocarcinoma (DLD-1 and HCT-116),

hormone-dependent breast cancer (T-47D) and human chronic myelogenous leukaemia (K562) cell lines and TeC-326, Fig. 12, was the most potent on all cell lines.88,529 On examining the mechanism of TeC-326 action, it was found to induce apoptosis in leukaemia Molt4 cells by acting on the oxidative stress pathway, on mitochondrial dysfunction and on talin expression. More in detail, it was able to upregulate both talin and phosphorylated talin expression and, additionally, to interfere with actin microfilament formation inducing morphological alteration.⁵³⁰ TeC-326 was found to inhibit the proliferation of the HCC cell lines HA22T and HA59T and to induce apoptosis via the caspase pathway. TeC-326 treatment also induced the formation of reactive oxygen species (ROS), which is associated with TeC-326-induced cell death, and to triggers ROS removal by mitochondrial SOD2 instead of cytosolic SOD1. The mitogen-activated protein kinase (MAPK) signalling pathway was linked to ROS-induced cell death, and TeC-326 reduced the expression of ERK, a MAPK that is associated with cell proliferation. Inhibitors of JNK and p38, which are MAPKs associated with apoptosis, restored TeC-326-induced cell death. In addition, TeC-326 treatment reduced the expression of GPX4, a protein that inhibits ferroptosis, which is a novel form of non-apoptotic programmed cell death. Treatment with a ferroptosis inhibitor also restored TeC-326-induced cell death.⁵³¹ TeC-326, 12-epi-heteronemin acetate (TeC-328), TeC-331 and TrC-130 were found to be toxic also against human epidermoid carcinoma KB cells532 and TeC-326 is also reported to inhibit the proteasome, promoting apoptotic cell death.533 A very deep analysis has been carried out by Lai et al. 2016 on two scalarane sesterterpenoids, 12β-(3'β-hydroxybutanoyloxy)-20,24-dimethyl-24-oxo-scalara-16-en-25-al (TeC-105) and 12β-(3'β-hydroxypentanoyloxy)-20,24-dimethyl-24-oxo-scalara-16-en-25-al (TeC-320) on various cancer cells. In leukaemia Molt 4 cells, TeC-105, Fig. 12, triggered mitochondrial membrane potential disruption, induced ROS production, calcium release and ER stress and inactivates topoisomerase IIa, determining apoptosis. Moreover, this molecule is able to target the molecular chaperone Hsp90 modulating its client proteins expression.534 In 2009, sesterterpene lactones isolated from Salvia dominica were discovered to interact with tubulin-tyrosine ligase (TTL), an enzyme involved in tubulin tyrosination in cancer cells, by chemical proteomics and surface plasmon resonance assays. An inhibition of this enzyme has also been assessed by 8a,15(S),23a-trihydroxy-23,6 α-epoxy-labd-13(14),17-dien-16(S),19-olide (DC-78), 8a,15(S)-dihydroxy-23a-O-ethyl-23,6α-epoxy-labd-13(14),17-dien-16(S),19-olide (DC-79), 8a-hydroxy-23a-O-ethyl-23,6a-epoxy-labd-13(14),17-dien-16(*R*),19-olide (**DC-80**), 6α,8α,15(*S*),23-tetrahydroxy-labd-13(14), 17-dien-16(S),19-olide (DC-81), 6a,8a,15(S)-trihydroxy-23-carbossi-labd-13(14),17-dien-16(S),19-olide (DC-83), 6α,8αdihydroxy-23-carbossi-labd-13(14),17-dien-16,19-olide (DC-84), 6a,8a,15(S)-trihydroxy-23-oxo-labd-13(14),17-dien-16(S),19-

olide (**DC-85**), $6\alpha,8\alpha$ -dihydroxy-23-oxo-labd-13(14),17-dien-16(*R*),19-olide (**DC-86**), $6\alpha,15(S),23$ -trihydroxy-labd-8(22),13(14),17-trien-16(*S*),19-olide (**DC-87**), $6\alpha,15(S)$ -dihydroxy-23-oxo-labd-8(22),13(14),17-trien-16(*S*),19-olide (**DC-88**), $6\alpha,8\alpha,23$ -trihydroxy-labd-13(14),15,17-trien-16,19-olide (**DC-89**),



6a,8a-dihydroxy-23-carbossi-labd-13(14),15,17-trien-16,19-olide (DC-90), 6a,8a-dihydroxy-23-oxo-13(14),15,17-trien-16,19-olide (DC-91), 8a-hydroxylabd-13(14),15,17-trien-6R,23-16,19-diolide (DC-92) and 8a-23-dihydroxy-23,6R-epoxy-labd-13(14),15,17trien-16,19-olide (DC-93).481 Also kohamaic acid A (DC-4, Fig. 12), besides other bioactivities, has been described to prevent the growth of human cancer cell (promyelocytic leukaemia cell line, HL-60) and its derivatives by a binding to four mammalian polymerase Beta residues (Leu11, Lys35, His51 and Thr79).535 Starting from 2010, Prof. Monti's group published some results on TrC-178's, new targets obtained by chemical proteomics. In brief, this molecule was able to interact with the proteasome inhibiting its function and thus becoming a candidate as an anticancer drug. It also altered the autophagy pathway.⁵³⁶⁻⁵³⁸ DH-80, Fig. 2, and DH-81, Fig. 12, presented an anti-proliferative effect against A549, HeLa, and HCT-116 cells with IC₅₀ values in the micromolar range through the inhibition of protein-tyrosine phosphatase 1B.35 Differently, MC-9, MC-10, and TH-25 targeted a deubiquitinating enzyme named USP7 blocking its function with IC_{50} values in the range of 2.7–4.6 μ M and are being considered as new anticancer lead compounds.46 Also the suvanines, sulfate-containing sesterterpenoids, 539 differently decorated on their scaffold, including TrC-143, Fig. 12, suvanine N,N-dimethyl-1,3-dimethylherbipoline salt (TeC-387), suvanine N,N-dimethylguanidium salt (TeC-388), and TrC-200-TrC-205, revealed moderate cytotoxicity against the K562 and A549 cell lines.¹⁹² More in detail, using a chemical proteomics approach, Cassiano et al. 540 revealed a direct interaction of suvanine and the heat shock protein 60, a key chaperonin involved in several tumoral and inflammatory diseases.

5.2 Sesterterpenoids with anti-inflammatory activity

Sesterterpenoids have an exceptional potential as antiinflammatory compounds. In the latter part of 80s, many compounds were isolated from the marine sponge Luffariella sp. with a consistent anti-inflammatory activity: the most wellknown is manoalide (MC-13, Fig. 3),⁵⁴¹⁻⁵⁴³ which significantly lowers chemically-induced inflammation in vivo and irreversibly blocks the in vitro hydrolysis of phosphotidylcholine by phospholipase A₂ (PLA₂).^{544,545} PLA₂ is a key step at the beginning of the inflammatory cascade releasing arachidonic acid from membranes. MC-13 is a non-specific inhibitor of phospholipases that modulates the activity of phospholipase C, human sPLA2 (IC $_{50}$ = 1.7 μM), cPLA2 (IC $_{50}$ = 10 μM) and snake venom sPLA₂ (IC₅₀ = 0.03 μ M).⁵⁴³ This latter activity has been confirmed by Dal Piaz et al.546 in deciphering the mechanism of MC-13 along with TrC-178, Fig. 12, binding to bee venom PLA₂: the γ -hydroxybutenolide ring with its masked aldehyde is able to covalently bind the N-terminus of PLA₂ giving rise to a Schiff base. The same inhibition mechanism has also been demonstrated for different petrosaspongiolides,547-549 21-hydroxypetrosaspongiolide K (TeC-65, Fig. 13), and 21-hydroxypetrosaspongiolide P (TrC-181).133 Prof. Gomez-Paloma group also extended this work to human synovial PLA2 and demonstrated that TrC-178 was able to covalently modify a certain lysine on the protein's surface, reducing the adhesion of this protein on the membrane and inhibiting its function.550 MC-13 was registered by Allergan pharmaceuticals and reached phase II clinical trials as a topical antipsoriatic. However, because of formulation hitches, it stopped in that phase. Several congeners of MC-13 have been extracted from different organisms. For example, three sesterterpenes fitting in the same family, *i.e.* (4E,6E)-dehydromanoalide (MC-68), deoxysecomanoalide (MC-29) and deoxymanoalide (MC-22), showed good inactivation degree on the snake venom PLA_2 at 0.2 to 0.5 μM even if lower than MC-13.551 Noteworthy, these metabolites are reduced forms of MC-13, and thus their PLA₂ inhibitory activity is much weaker.⁵⁵² MC-70 is a powerful antagonist of topical phorbol



myristate acetate promoted inflammation in the mouse ear. It inhibited in vitro hydrolysis of phosphatidyl choline by bee venom PLA₂, too (IC₅₀ = 0.23μ M).⁵⁵³ The determined inhibition achievable by MC-70 is only 80% as compared to MC-13 because its binding was partially reversible (around 30%). Indeed, a detailed kinetic analysis of the MC-70 reaction with PLA₂ confirmed also a non-competitive type of inhibition.220 Furthermore, the luffariellins (mainly MC-53, Fig. 13 and MC-56, Fig. 13) from the sponge Luffariella variabilis were also considered anti-inflammatory sesterterpenes by the research group of Kernan in 1987.223 They are powerful antagonists of topical phorbol myristate acetate-induced inflammation in the mouse ear and also bee venom PLA2 was significantly downregulated by MC-53 (IC₅₀ = 0.056μ M) and MC-56 (IC₅₀ = 0.060µM). Barbara Potts confirmed in 1992 that even DC-162 displayed approximately 50% inhibition of oedema in the mouse ear assay.554 Other PLA2-modulating sesterterpenes from marine origin included cacospongionolides B (DC-19, Fig. 13) and E (DC-20) (IC₅₀ = 300 nM against human and bee venom sPLA $_2$ ^{555,556} and DH-16 (IC $_{50}$ = 6.9 μ M against human sPLA $_2$ and cPLA2 557 De Marino et al. 133 in 2000 proposed that spongidines A-D (TrC-259-TrC-262) inhibited hsPLA₂ as well. Interestingly, studies on diastereoisomeric mixture of DC-6 and DC-7, both bearing a hydroxybutenolide moiety, demonstrated that this group itself is not sufficient for PLA₂ inhibitory activity but that non-covalent interactions between the counterparts also play key roles in terms of compound potency.558,559 TeC-75, a dialdehyde containing sesterterpene, is considered a good antiinflammatory compound,560,561 mainly inactivating PLA2s.562 In 2000, Prof. Cimino's group isolated many scalarane and homoscalarane compounds from the nudibranchs Glossodoris sedna and G. dalli and tested them against mammalian cytosolic PLA2. The di-aldehyde-bearing compounds exhibited a consistent inhibition of the enzyme although it occurred at high concentration.²⁵⁹ Next, Monti et al. assessed the role of dialdehyde moiety of TeC-75 and of 12-epi-scalaradial (TeC-78) on

secretory PLA₂.^{537,563,564} Also **DH-76** and its monoacetate analogue, isolated from the sponge *Hyrtios* sp., were tested on cobra venom PLA₂, showing good results in terms of inactivation.⁵⁶⁵ In 2009, Prof. Zampella isolated **TrC-200**, Fig. 13 and **TrC-154**, Fig. 13, two sesterterpenoids bearing nitrogen atoms together with suvanine (**TrC-143**), the latter exerting antiinflammatory activity by the inhibition of different secretory PLA₂ such as groups IIA, IA (Naja naja venom), IB (porcine pancreatic enzyme) and III (bee venom enzyme).¹⁹⁵

5.3 Sesterterpenoids with anti-microbial activity

Hyrtiosin B (TeC-306, Fig. 14) from Hyrtios erecta was found to modulate the protein isocitrate lyase from Candida albicans.566 This compound was a good inhibitor of this enzyme which plays a crucial role in fungal development. Furospongin-4 (MH-44, Fig. 14) and isofurospongin-4 (MH-25, Fig. 14) showed weak influence on *Staphylococcus aureus* at 100 μ g mL⁻¹ ¹³⁷ while **PC**-31 exhibited strong influence against Mycobacterium tuberculosis protein tyrosine phosphatase B.433 Several sesterterpenoids have been described to have bacteriostatic and bactericidal effects against tested pathogens. The antibiotic profile of MC-13 was investigated early, in 1980,⁵⁶⁷ alongside (6E)-neomanoalide (MC-16), and (6Z)-neomanoalide (MC-17) which were found to be effective selectively against Gram-positive bacteria.37 The proposed mechanism regards the modulation of cell-cell communication by MC-13 and by manoalide 25-monoacetate (MC-14).568 In the field of HIV inhibitors, phyllolactones were verified for their effect in downregulating HIV-1 envelopemediated fusion.¹⁰⁴ Additionally, TrC-130 inactivates HIV-1 integrase by binding to DNA as reported by surface plasmon resonance and docking analysis.569 Using a model of HIV-1 infection, TrC-192 and TrC-211, alotaketals C and D (MC-117 and MC-118) and TeC-390 were found to promote HIV proviral gene expression as well as being at least as potent as MC-117 and MC-118.179 PC-40 inhibits HIV-1 replication.510 Also PC-31



Fig. 14 Representative structures of sesterterpenoids with anti-micro, anti-neurodegenerative and anti-feedant activities

and **PC-64** had potent inhibition against *Mycobacterium tuberculosis* protein tyrosine phosphatase B with IC_{50} values of 3–6 μ M.³⁸⁹

5.4 Sesterterpenoids with anti-neurodegenerative activity

Few sesterterpenoids have been studied for their ability to counteract neurodegeneration. In 2014, Prof. Monti's group identified TeC-326, Fig. 12, a marine sesterterpenoids isolated from Hyrtios sponge,⁵⁷⁰ as an interactor of trans-activation response DNA-binding protein of 43 kDa (TDP-43) by using chemical proteomics. This metabolite can bind the protein and can alter its tendency to create insoluble aggregates in brain cells. After this, a small library of natural scalarane derivatives such as petrosaspongiolactams A-C (TeC-126-TeC-128) has been investigated for their ability to modulate the activity of TDP-43 giving the identification of protein aggregates inhibitors.87 In parallel, TrC-6 from Bipolaris oryzae has been discovered to strongly induce the autophagic degradation pathways of insoluble aggregates involved in neurodegenerative diseases.⁵⁷¹ In this case, the targeted protein was alpha-synuclein in PC12 cells. Also, erinacine S (TeC-392), isolated from the ethanol extract of the mycelia of Hericium erinaceus, downregulates the formation of AB insoluble precipitates in the brains of transgenic mice after an oral administration as reported by Chen et al. 572 Following this, in 2019, Hu et al. deeper investigated the mechanism of TeC-392, examining its capability in reducing amyloid plaque formation and improving neurogenesis in rats. Furthermore, this study showed that TeC-392 can penetrate the blood-brain barrier.573 DC-164 and DC-211 showed their antineurodegenerative profile by acting against prolylendopeptidase470,472 as well as MC-36 was tested against Alzheimer showing a good profile as PPAR α/γ -RXR α agonist and RAR α positive allosteric modulator.⁵⁷⁴ TeC-97 and TeC-98 ⁴³⁴ played as acetylcholinesterase inhibitors in rats whereas the compound YW3699 inhibited the glycosylphosphatidylinositol (GPI) synthesis.575 Finally, palinurin (DH-3) acted by an allosteric regulation on GSK-3.576

5.5 Sesterterpenoids with phytotoxicity

In 1984, **TrC-2**, Fig. 3, and its congeners were deeply analysed as photosynthesis inhibitors. Furthermore, their biological profile

was acutely evaluated using spinach leaf slices and chlorella. TrC-2 resulted in well counteracting photosynthesis causing brown spots in rice plants.577 Later on, other ophiobolins were investigated by a mitochondrial electron transport assay designed to identify malate oxidation inhibitors: TrC-2 and TrC-5 were found to inhibit oxidation in maize mitochondria.324 Moreover, using the so-called leaf spot assay, their influence has been measured against several plants such as sorghum, maize, bentgrass, sicklepod and morning glory based on the necrosis superficial amplitude: TrC-2 and TrC-3 were more phytotoxic than their anhydro derivatives.³¹⁴ Additionally, in 1991, Prof. Ballio deeply studied the mode of action of these sesterterpenoids. He postulated that the probable role of TrC-2 in the development of brown spot of rice is due to its involvement in a precise membrane transport process related to K⁺ permeability. Indeed, he demonstrated that in vitro TrC-2 irreversibly inhibits the calmodulins of bovine brain, maize and spinach by a covalent bond of the toxin to lysine 63, identifying this enzyme as ophiobolins target: the Ca²⁺-calmodulin complex is accountable for the regulation of many cellular functions.⁵⁷⁸ In 2006, ophiobolins were also investigated as phytotoxins by Prof. Evidente's group: in particular, TrC-2 instigated the appearance of large necrosis areas, even at low concentrations on grass weeds.318 Also, ophiobolins TrC-8, TrC-16 and TrC-35 were toxic to various weed species by the leaf-puncture bioassay.319

5.6 Sesterterpenoids with nematocidal and anti-feedant activity

In plants, also **TrC-38**, Fig. 14, and **TrC-39** were tested as nematocidal agents and they were toxic against the free-living worm *Caenorhabditis elegans*.³⁴⁸ In 1996, Tsipouras deepened the understanding of the mechanism of **TrC-45** and its congeners.³⁴⁷ Indeed, all of these compounds exerted their nematocidal property by altering the ivermectin binding to nematode membranes. The same sesterterpenoids along with **TrC-13** were also effective in the *C. elegans* motility assays and they were toxic in the low μ M range.³⁴⁷ Some defensive sesterterpenoids with unusual cyclopentenone rings were extracted and purified from the leaves of *L. canum*: they were leucosceptroids **DC-168**, Fig. 14, **TrC-190** and **TrC-191**.⁴⁶⁷ The potent antifeedant activity recommended them as self-protective molecules.⁴⁷⁴ The related DC-172–DC-174 showed the same bioactivity as well as DC-187 and DC-192 and, finally, in 2014 also DC-200–DC-202.⁴⁶⁸ Moreover, it was discovered that leucosceptrane sesterterpenoid degradation products could also be involved in the plant defence mechanisms against insects.⁴⁶⁹ In 2013, in a very innovative paper, Li group studied DC-182, DC-183 and DC-184 from leaves through a laser-microdissection coupled to mass spectrometry analysis and X-ray diffraction and planned a defensive role for these sesterterpenoids since they were closely related to the above-described leucosceptroid-class. All three compounds disincentivised the beet armyworm and the cotton boll worm.⁴⁷⁷ Finally, TrC-20 has been indicated as trypanosomicidal: mainly, its anti-protozoal activity was exerted against trypomastigote and amastigote forms of *Trypanosoma cruzi*.⁵¹²

5.7 Sesterterpenoids with immunosuppressive activity

In 2007, irregularasulfate (DC-26), hipposulfate A (TrC-113) and halisulfate-7 (DC-31) were discovered to be moderate inhibitors of the catalytic subunits of the mammalian Ser/Thr protein phosphatases calcineurin and could be used as immunosuppressive molecules.579 Also TrC-286 and TrC-287 were considered immunosuppressive by a different mechanism of action inhibiting the proliferation, activation, and cytokine IFN-y production on T cells.580 Also TeC-5, TeC-6, TeC-11, TeC-12, and TeC-13, studied by Fujimoto et al., were immunosuppressive, acting on the human chemokine receptor CCR5.367 It's intriguing that more and more sesterterpenoids have been revealed to have immunosuppressive properties after the Covid-19 epidemic. Initially, the activity of leucosceptrane-type sesterterpenoids was assessed by preventing the release of cytokines in LPS-induced macrophages RAW264.7 and 3-H-2,17adihydroxy-leucosceptroid N (DC-270) demonstrated immunosuppressive activity against TNF-a and IL-6, with IC50 values of 13.39 and 19.34 µM, respectively.581 Then, a deep analysis of leucosceptroid N (DC-192), 5,13-dehydro-leucosceptroid A (DC-306), and 5a,16a-epoxy-leucosceptroid R (DC-291) showed their block on T cells and macrophages activations thought the inhibition of AKT-mTOR, JNK, p38 MAPK, and ERK pathway. Furthermore, DC-291 and DC-306 caused T cell G0/G1 cell arrest, whereas DC-192 markedly reduced IL-6 and TNF-α levels in peripheral blood serum and lessened the multiorgan damage in mice with LPS-induced sepsis, at 25 mg per kg dose. Additionally, there was a dose-dependent suppression of T cell proliferation and IFN- γ release when activated by anti-CD3/ CD28.582 The TNF-α secretion was also effectively inhibited by leucosceptrodine (MC-153) and nor-leucosceptroid L (DC-264), with IC₅₀ values of 11.21 µM and 12.19 µM, respectively.⁵⁸³ Lastly, few ophiobolin-like substances were studied in this area, including maydispenoids and undobolins: maydispenoids A (TrC-306) and B (TrC-307) demonstrated proliferation inhibitory activity on differently stimulated murine splenocytes332 and undobolin F (TrC-362) established significant inactivation against ConA-induced T lymphocyte proliferation with an IC₅₀ value of 2.3 µM.584

5.8 Miscellaneous

In 2006, Nam and coworkers isolated several sesterterpenoids from the sponge Spongia sp. including TeC-109 and TeC-110 which have a structure similar to guggulsterone and have been tested for their ability to modulate the transactivation of the farnesoid-X-receptor (FXR). These scalaranes inhibited FXR and were not cytotoxic for cells opening the way to treat hypercholesterolemia and, more in general, metabolic diseases.135 An anti-obesity profile has been reported for DC-169, based on a different mechanism of action: this metabolite acts reducing fat amassing through destroying unsaturated fatty acid biosynthesis. More deeply, it downregulates the expression of two stearoyl-CoA desaturase (SCD) genes fat-6 and fat-7, and a fatty acid elongase gene elo-2 in wild-type C. elegans.580 Inhibition of protein tyrosine phosphatase 1B has also been explored by a few sesteterpenoids as, for instance, hyattellactones A and B (TeC-239 and TeC-240), in an interesting paper¹¹⁵ since this protein can be considered a therapeutic target for the treatment of type-II diabetes and obesity. The same inactivation has been reported for phyllofolactones characterized by an alpha-beta-unsaturated lactone.¹¹⁵ In 2011, phorbasones were discovered to induce osteoblast differentiation, specifically phorbasone A (TrC-195) revealed a stimulatory effect on the calcium deposition activity in C3H10T1/2 cells.174 MC-119, also, significantly excites osteoblast differentiation by activating the ERK pathway.585

6 Conclusions

Within the terpenoids family, which is the biggest group of secondary metabolites, sesterterpenoids represent an incredible chemically varied and pharmacologically relevant subgroup, widespread in more than a few phyletic groups, including marine sponges, nudibranchs, bacteria, lichens, fungi, higher plants and insects. Over seven decades of research on their isolation and characterization, around 1600 structurally unique sesterterpenoids are summarized in the current review paper, which is supported by information on biological, pharmacological, ecological, and geographic distribution. It is highly probable that in the next years novel understudied taxa groups will bring up new interesting members of these pentaprenyl terpenes. Moreover, it is expected that the synthetic origin of some of these compounds will be disentangled, unveiling interconnecting mechanisms of symbiotic or trophic exchanges, bioaccumulation, biotransformation, and de novo synthesis. By regulating microbial metabolism, sesterterpenoids have facilitated the co-speciation of bacteria with fungi and the interaction of fungi with marine and terrestrial plants, thus generating complex interactions essential for microbial survival. Recently, there has been a growing recognition of the role of specific sesterterpenoids in establishing the composition of leaf and root microbiota in which the dynamic complementation of enzymatic pathways controls plant health.⁵¹ From a biological point of view, microorganisms as well as humans use them as antimicrobial, antiviral, antifeedants, and to combat neurodegeneration, hypercholesterolemia, diabetes,

and obesity, in addition to their regal role as anti-cancer and anti-inflammatory (ophiobolin and manoalide derivatives, about everything). Recently, tests for the inhibition of T cell proliferation and the release of IFN- γ , TNF- α , and IL-6 have been conducted in an effort to counterbalance overactive immune system components. This could potentially be a future avenue for the repurposing of these poly-pharmacological metabolites. Furthermore, it should be emphasized that genome mining and heterologous expression approaches of sesterterpene synthases, increasingly in recent years, have led to the identification of new sesterpenoids, drawing a path that will be surely implemented in the future for the discovery of new molecular scaffolds. In the present review, we reported selected examples of naturally occurring marine and terrestrial sesterterpenoids within their structures, ecological, geographical and biological details. But without a question, the most important component of this enormous undertaking is the exhaustive list of known metabolites, which is included in the annexed Tables, and at the friendly consultable https:// sesterterpenoids.unige.net/. The authors chose to publish the results of their work using an architecture formed by the integration between a relational database and a web-based interface. This architecture allows for rapid updating of research contents which will be implemented through two channels. The first is the direct intervention of the authors who will be able, after their authentication on the site, to manually insert new evidence relating to the subject matter present in new publications. The second concerns the interaction with public databases, selected by the authors of this work, for their evident authority, which will provide automatic access to updated in-depth content on the description of the individual organisms involved in the research. Pipelines based on Natural Language Processing to obtain automatic insertion of new evidence detected from scientific articles will be applied.

7 Author contributions

Conceptualization: A. B., M. G., and N. De T.; data curation: V. I., A. B., N. De T., V. P., and F. De R.; formal analysis: A. B., N. De T. and M. G.; funding acquisition: A. B. and F. De R.; investigation: V. I., M. G., F. De R., P. B., L. N.-P., G. D., P. G., M. C. M., R. P., Y. M., V. P., A. B.; methodology: A. B., M. G. and N. De T.; project administration: A. B., and N. De T.; resources: A. B. and N. De T.; software: M. G.; supervision: A. B., M. G., and N. De T.; validation: A. B., M. G., F. De R., and N. De T.; writing – original draft: M. G., F. De R., P. B., L. N.-P., G. D., P. G., M. C. M., R. P., and A. B.; writing – review & editing: V. P., M. C. M, A. B., G. D., P. G., and N. De T.

8 Conflicts of interest

There are no conflicts to declare.

9 Acknowledgements

FDR: Financial support from the University of Salerno (FARB), from PRIN 2020: "Natural Products-Assisted Organic Synthesis"

(2020AEX4TA). "Graphical Abstract was Created in BioRender. Núñez-Pons, L. (2024) BioRender.com/j70k119". NDT and VP: Finacial support by National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of December 16, 2021, rectified by Decree n.3175 of December 18, 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU; Award Number: Project code CN_00000033, Concession Decree No. 1034 of June 17, 2022 adopted by the Italian Ministry of University and Research, CUP: D43C22001260001, Project title "National Biodiversity Future Center - NBFC".

10 References

- 1 J. R. Hanson, Nat. Prod. Rep., 1986, 3, 123-132.
- 2 J. R. Hanson, Nat. Prod. Rep., 1992, 9, 481-489.
- 3 J. R. Hanson, Nat. Prod. Rep., 1996, 13, 529-535.
- 4 T. Mitsuhashi and I. Abe, in *Progress in the Chemistry of Organic Natural Products*, ed. A. D. Kinghorn, H. Falk, S. Gibbons, J. Kobayashi, Y. Asakawa and J. K. Liu, Springer Nature, Cham, Switzerland, 2020, vol. 111, pp. 1–79.
- 5 S. K. Talaprata, in *Chemistry of Plant Natural Products*, ed. S. K. Talapatra and B. Talapatra, Springer-Verlag, Berlin Heidelberg, 2015, pp. 511–515, DOI: **10.1007/978-3-642-45410-3_9**.
- 6 L. Wang, B. Yang, X.-P. Lin, X.-F. Zhou and Y. Liu, *Nat. Prod. Rep.*, 2013, **30**, 455–473.
- P. Crews and S. Naylor, in Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products, ed. W. Herz, H. Grisebach, G. W. Kirby and C. Tamm, Springer, Vienna, 1985, vol. 48, pp. 204–269.
- 8 G. A. Cordell, *Phytochemistry*, 1974, **13**, 2343–2364.
- 9 J. R. Hanson, *Terpenoids and Steroids*, London, UK, 1974, vol. 4.
- 10 S. S. Ebada, W. Lin and P. Proksch, *Mar. Drugs*, 2010, **8**, 313–346.
- 11 R. Ebel, Mar. Drugs, 2010, 8, 2340-2368.
- 12 P. Máximo and A. Lourenço, *Curr. Org. Chem.*, 2018, 22, 2381–2393.
- 13 A. J. Michael, Biochem. J., 2017, 474, 2277-2299.
- 14 A. R. Carroll, B. R. Copp, R. A. Davis, R. A. Keyzers and M. R. Prinsep, *Nat. Prod. Rep.*, 2019, 36, 122–173.
- 15 A. R. Carroll, B. R. Copp, R. A. Davis, R. A. Keyzers and M. R. Prinsep, *Nat. Prod. Rep.*, 2023, 40, 275–325.
- 16 L. Nunez-Pons and C. Avila, *Nat. Prod. Rep.*, 2015, **32**, 1114–1130.
- 17 V. J. Paul, K. E. Arthur, R. Ritson-Williams, C. Ross and K. Sharp, *Biol. Bull.*, 2007, 213, 226–251.
- 18 M. Maldonado and P. R. Bergquist, in Atlas of Marine Invertebrate Larvae, A. Press, 2002, ch. 2, pp. 21–50.
- 19 M. W. Taylor, R. Radax, D. Steger and M. Wagner, *Microbiol. Mol. Biol. Rev.*, 2007, 71, 295–347.
- 20 N. S. Webster and L. L. Blackall, ISME J., 2009, 3, 1-3.
- 21 N. S. Webster and M. W. Taylor, *Environ. Microbiol.*, 2012, 14, 335–346.

- M. Hudspith, L. Rix, M. Achlatis, J. Bougoure,
 P. Guagliardo, P. L. Clode, N. S. Webster, G. Muyzer,
 M. Pernice and J. M. De Goeij, *Microbiome*, 2021, 9, 44.
- 23 L. Pita, L. Rix, B. M. Slaby, A. Franke and U. Hentschel, *Microbiome*, 2018, **6**, 46.
- 24 J. M. Deutsch, M. O. Green, P. Akavaram, A. C. Davis, S. S. Diskalkar, I. A. Du Plessis, H. A. Fallon, E. M. Grason, E. G. Kauf, Z. M. Kim, J. R. Miller, A. L. Neal, T. Riera, S.-E. Stroeva, J. Tran, V. Tran, A. V. Coronado, V. V. Coronado, B. T. Wall, C. m. Yang, I. Mohanty, N. H. Abrahamse, C. J. Freeman, C. G. Easson, C. L. Fiore, A. E. Onstine, N. Djeddar, S. Biliya, A. V. Bryksin, N. Garg and V. Agarwal, *Mar. Drugs*, 2023, 21, 53.
- 25 K. Guo, Y. Liu and S.-H. Li, *Nat. Prod. Rep.*, 2021, **38**, 2293–2314.
- 26 F. Bibi, M. Faheem, E. I. Azhar, M. Yasir, S. A. Alvi, M. A. Kamal, I. Ullah and M. I. Naseer, *Curr. Drug Metab.*, 2017, 18, 11–15.
- 27 C. Avila, L. Núñez-Pons and J. Moles, in *Chemical ecology:* the ecological impacts of marine natural products, ed. M. P. Puglisi and M. A. Becerro, CRC Press, Boca Raton, FL, USA, 1st edn, 2018, pp. 71–163.
- 28 Q. Chen, J. Li, Y. Ma, W. Yuan, P. Zhang and G. Wang, *Plant Commun.*, 2021, 2, 100184.
- 29 Microsoft Corporation, Microsoft SQL Server, https:// www.microsoft.com/en-us/sql-server/sql-server-2022.
- 30 Microsoft Corporation, Windows Server, 2022, https:// learn.microsoft.com/en-us/windows-server/get-started/ whats-new-in-windows-server-2022.
- 31 K. Li and K. R. Gustafson, Nat. Prod. Rep., 2021, 38, 1251– 1281.
- 32 Y. Liu, L. Wang, J. H. Jung and S. Zhang, *Nat. Prod. Rep.*, 2007, 24, 1401–1429.
- 33 Y. S. Yang, I. M. Chung, S. H. Kim and A. Ahmad, *Asian J. Chem.*, 2014, **26**, 7792–7794.
- 34 A. Cutignano, J. Moles, C. Avila and A. Fontana, J. Nat. Prod., 2015, 78, 1761–1764.
- 35 S. J. Piao, H. J. Zhang, H. Y. Lu, F. Yang, W. H. Jiao, Y. H. Yi, W. S. Chen and H. W. Lin, *J. Nat. Prod.*, 2011, 74, 1248–1254.
- 36 G. Chianese, J. Silber, P. Luciano, C. Merten, D. Erpenbeck,
 B. Topaloglu, M. Kaiser and D. Tasdemir, *J. Nat. Prod.*, 2017,
 80, 2566–2571.
- 37 E. D. De Silva and P. J. Scheuer, *Tetrahedron Lett.*, 1981, 22, 3147–3150.
- 38 S. Iimura, M. Oka, Y. Narita, M. Konishi, H. Kakisawa, Q. Gao and T. Oki, *Tetrahedron Lett.*, 1993, 34, 493-496.
- 39 S. Nozoe, M. Morisaki, K. Tsuda, Y. Iitaka, N. Takahashi, S. Tamura, K. Ishibashi and M. Shirasaka, J. Am. Chem. Soc., 1965, 87, 4968–4970.
- 40 M. Liu, W. Sun, L. Shen, Y. He, J. Liu, J. Wang, Z. Hu and Y. Zhang, *Angew. Chem., Int. Ed.*, 2019, 58, 12091–12095.
- 41 H. J. Zhang, H. F. Tang, Y. H. Yi and H. W. Lin, *Helv. Chim. Acta*, 2009, **92**, 762–767.
- 42 Q. Li, C. Chen, M. Wei, C. Dai, L. Cheng, J. Tao, X. N. Li, J. Wang, W. Sun, H. Zhu and Y. Zhang, *Org. Lett.*, 2019, 21, 2290–2293.

- 43 J. Shin, J. R. Rho, Y. Seo, H. S. Lee, K. W. Choa and C. J. Simb, *Tetrahedron Lett.*, 2001, 42, 3005–3007.
- 44 G. Alea, A. Carroll and B. Bowden, *Aust. J. Chem.*, 1994, 47, 191–194.
- 45 G. Cimino, P. De Luca, S. De Stefano and L. Minale, *Tetrahedron*, 1975, **31**, 271–275.
- 46 A. H. Afifi, I. Kagiyama, A. H. El-Desoky, H. Kato,
 R. E. P. Mangindaan, N. J. de Voogd, N. M. Ammar,
 M. S. Hifnawy and S. Tsukamoto, *J. Nat. Prod.*, 2017, 80, 2045–2050.
- 47 F. Cafieri, L. De Napoli, E. Fattorusso and C. Santacroce, *Experientia*, 1977, **33**, 994–995.
- 48 S. J. Piao, W. H. Jiao, F. Yang, Y. H. Yi, Y. T. Di, B. N. Han and H. W. Lin, *Mar. Drugs*, 2014, **12**, 4096–4109.
- 49 Y. M. Xu, P. Espinosa-Artiles, M. X. Liu, A. E. Arnold and A. A. Gunatilaka, *J. Nat. Prod.*, 2013, **76**, 2330–2336.
- 50 E. Fattorusso, V. Lanzotti, S. Magno, L. Mayol and M. Pansini, J. Org. Chem., 1992, 57, 6921–6924.
- 51 A. C. Huang, S. A. Kautsar, Y. J. Hong, M. H. Medema, A. D. Bond, D. J. Tantillo and A. Osbourn, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, E6005–E6014.
- 52 J. J. Bell, Estuar. Coast Shelf Sci., 2008, 79, 341-353.
- 53 N. J. de Voogd, B. Alvarez, N. Boury-Esnault, P. Cárdenas, M.-C. Díaz, M. Dohrmann, R. Downey, C. Goodwin, E. Hajdu, J. N. A. Hooper, M. Kelly, M. Klautau, S. C. Lim, R. Manconi, C. Morrow, U. Pinheiro, A. B. Pisera, P. Ríos, K. Rützler, C. Schönberg, T. Turner, J. Vacelet, R. W. M. van Soest and J. Xavier, The World Porifera Database, <u>https://www.marinespecies.org/porifera/</u>, (accessed February 20th 2024).
- 54 R. W. M. Van Soest, N. Boury-Esnault, J. Vacelet, M. Dohrmann, D. Erpenbeck, N. J. De Voogd, N. Santodomingo, B. Vanhoorne, M. Kelly and J. N. A. Hooper, *PLoS One*, 2012, 7, e35105.
- 55 U. Hentschel, J. Piel, S. M. Degnan and M. W. Taylor, *Nat. Rev. Microbiol.*, 2012, **10**, 641–654.
- 56 S. Zhang, W. Song, L.-F. Nothias, S. P. Couvillion, N. Webster and T. Thomas, *Microbiome*, 2022, **10**, 22.
- 57 D. Varijakzhan, J.-Y. Loh, W.-S. Yap, K. Yusoff, R. Seboussi, S.-H. E. Lim, K.-S. Lai and C.-M. Chong, *Mar. Drugs*, 2021, 19, 246.
- 58 T. Hughes, A. Robinson, S. Rogers and V. Tillett, MarinLit Dedicated to marine natural products research, https:// marinlit.rsc.org, (accessed February 20th 2024).
- 59 K. Wilson, T. De Rond, I. Burkhardt, T. S. Steele, R. J. B. Schäfer, S. Podell, E. E. Allen and B. S. Moore, *Proc. Natl. Acad. Sci. U. S. A.*, 2023, **120**, e2220934120.
- 60 E. R. Abdelaleem, M. N. Samy, S. Y. Desoukey, M. Liu,
 R. J. Quinn and U. R. Abdelmohsen, *RSC Adv.*, 2020, 10, 34959–34976.
- 61 X.-F. Zhang, J. Ren, X.-R. Cheng, H.-Z. Jin and W.-D. Zhang, *RSC Adv.*, 2015, **5**, 1979–1982.
- 62 E. Fattorusso, S. Magno, C. Santacroce and D. Sica, *Tetrahedron*, 1972, **28**, 5993–5997.
- 63 Q. Lu and D. J. Faulkner, J. Nat. Prod., 1997, 60, 195-198.
- 64 G. Cimino, S. De Stefano, L. Minale and E. Trivellone, J. Chem. Soc., Perkin Trans. 1, 1977, (13), 1587–1593.

- 65 R. C. Cambie, C. E. F. Rickard, P. S. Rutledge and X. S. Yang, Acta Crystallogr., 1999, C55, 112–114.
- 66 C. J. Hernández-Guerrero, E. Zubía, M. J. Ortega and J. L. Carballo, *Tetrahedron*, 2006, 62, 5392–5400.
- 67 X. Yang, Z. Shao and X. Zhang, Z. Naturforsch., 2010, 65b, 625–627.
- 68 P. Crews, P. Bescansa and G. J. Bakus, *Experientia*, 1985, **41**, 690–691.
- 69 P. Crews and P. Bescansa, J. Nat. Prod., 1986, 49, 1041–1052.
- 70 Y. Kashman and A. Rudi, *Tetrahedron*, 1977, **33**, 2997–2998.
- 71 R. Kazlauskas, P. T. Murphy, R. J. Quinn and R. J. Wells, *Tetrahedron Lett.*, 1976, **17**, 2631–2634.
- 72 G. R. Pettit, Z. A. Cichacz, R. Tan, M. S. Hoard, N. Melody and R. K. Pettit, *J. Nat. Prod.*, 1998, **61**, 13–16.
- 73 N. Tsuchiya, A. Sato, T. Hata, N. Sato, K. Sasagawa and T. Kobayashi, *J. Nat. Prod.*, 1998, **61**, 468–473.
- 74 H. Miyaoka, S. Nishijima, H. Mitome and Y. Yamada, *J. Nat. Prod.*, 2000, **63**, 1369–1372.
- 75 D. T. A. Youssef, R. K. Yamaki, M. Kelly and P. J. Scheuer, *J. Nat. Prod.*, 2002, **65**, 2–6.
- 76 D. T. A. Youssef, L. A. Shaala and S. Emara, J. Nat. Prod., 2005, 68, 1782–1784.
- 77 Y. Qiu, Z. Deng, Y. Pei, H. Fu, J. Li, P. Proksch and W. Lin, J. Nat. Prod., 2004, 67, 921–924.
- 78 Z. G. Yu, K. S. Bi and Y. W. Guo, *Helv. Chim. Acta*, 2005, 88, 1004–1009.
- 79 A. R. Diaz-Marrero, T. Matainaho, R. van Soest, M. Roberge and R. J. Andersen, *Nat. Prod. Rep.*, 2008, 22, 1304–1309.
- 80 L. Yin, H. Li, X. Chen and Y. Qiu, *Rec. Nat. Prod.*, 2014, 8, 417-421.
- 81 S. S. Elhady, A. M. El-Halawany, A. M. Alahdal, H. A. Hassanean and S. A. Ahmed, *Molecules*, 2016, 21(82), 81–89.
- 82 A. M. Alahdal, H. Z. Asfour, S. A. Ahmed, A. O. Noor, A. M. Al-Abd, M. A. Elfaky and S. S. Elhady, *Molecules*, 2018, 23, 1–11.
- 83 P. Francis and K. Chakraborty, *Med. Chem. Res.*, 2021, 30, 886-896.
- 84 M.-Z. Su, Q. Zhang, L.-G. Yao, B. Wu and Y.-W. Guo, J. Asian Nat. Prod. Res., 2022, 1–7, DOI: 10.1080/ 10286020.2022.2150614.
- 85 H. N. K. Tran, M. J. Kim and Y.-J. Lee, *Mar. Drugs*, 2022, **20**, 604.
- 86 S. Jaisamut, S. Thengyai, S. Yuenyongsawad, C. Karalai,
 A. Plubrukarn and K. Suwanborirux, *Pure Appl. Chem.*, 2009, 81, 1019–1026.
- 87 C. Festa, C. Cassiano, M. V. D'Auria, C. Debitus, M. C. Monti and S. De Marino, *Org. Biomol. Chem.*, 2014, **12**, 8646–8655.
- 88 F. Yang, J.-H. Gan, X.-Y. Liu and H.-W. Lin, Nat. Prod. Commun., 2014, 9, 1934578X1400900608.
- 89 W. Kaweetripob, C. Mahidol, P. Tuntiwachwuttikul, S. Ruchirawat and H. Prawat, *Mar. Drugs*, 2018, **16**(474), 471-415.
- 90 J. Li, L. Du, M. Kelly, Y. D. Zhou and D. G. Nagle, J. Nat. Prod., 2013, 76, 1492–1497.
- 91 J. C. Braekman and D. Daloze, *Pure Appl. Chem.*, 1986, 58, 357-364.

- 92 J. C. Braekman, D. Daloze, M. Kaisin and B. Moussiaux, *Tetrahedron*, 1985, **41**, 4603–4614.
- 93 R. J. Quinn and D. J. Tucker, Aust. J. Chem., 1989, 42, 751– 755.
- 94 D. E. Williams, I. Hollander, L. Feldberg, E. Frommer, R. Mallon, A. Tahir, R. van Soest and R. J. Andersen, J. Nat. Prod., 2009, 72, 1106–1109.
- 95 F. Cao, Z. H. Wu, C. L. Shao, S. Pang, X. Y. Liang, N. J. de Voogd and C. Y. Wang, *Org. Biomol. Chem.*, 2015, 13, 4016–4024.
- 96 H. Kikuchi, Y. Tsukitani, I. Shimizu, M. Kobayashi and I. Kitagawa, *Chem. Pharm. Bull.*, 1981, **29**, 1492–1494.
- 97 H. Kikuchi, Y. Tsukitani, I. Shimizu, M. Kobayashi and I. Kitagawa, *Chem. Pharm. Bull.*, 1983, **31**, 552–556.
- 98 I. Kitagawa, M. Kobayashi, N.-K. Lee, Y. Oyama and Y. Kyogoku, *Chem. Pharm. Bull.*, 1989, **37**, 2078–2082.
- 99 X. Fu, L. Zeng, J. Su, M. Païs and P. Potier, J. Nat. Prod., 1993, 56, 1985–1988.
- 100 X. Fu, L. Zeng, J. Su and F. J. Schmitz, *J. Nat. Prod.*, 1999, **62**, 644–646.
- 101 L. Zeng, X. Fu, J. Su, E. O. Pordesimo, S. C. Traeger and F. J. Schmitz, J. Nat. Prod., 1991, 54, 421–427.
- 102 H. J. Zhang, Y. H. Yi, F. Yang, W. S. Chen and H. W. Lin, *Molecules*, 2010, **15**, 834-841.
- 103 M. Jaspars, E. Jackson, E. Lobkovsky, J. Clardy, M. C. Diaz and P. Crews, *J. Nat. Prod.*, 1997, **60**, 556–561.
- 104 L. C. Chang, S. Otero-Quintero, G. M. Nicholas and C. A. Bewely, *Tetrahedron*, 2001, **57**, 5731–5738.
- 105 M. C. Roy, J. Tanaka, N. de Voogd and T. Higa, *J. Nat. Prod.*, 2002, **65**, 1838–1842.
- 106 M. H. A. Hassan, M. E. Rateb, M. Hetta, T. A. Abdelaziz, M. A. Sleim, M. Jaspars and R. Mohammed, *Tetrahedron*, 2015, **71**, 577–583.
- 107 S. Hayes, A. C. Taki, K. Y. Lum, J. J. Byrne, J. M. White, M. G. Ekins, R. B. Gasser and R. A. Davis, *J. Nat. Prod.*, 2022, **85**, 1723–1729.
- 108 L. P. Ponomarenko, A. I. Kalinovsky and V. A. Stonik, *J. Nat. Prod.*, 2004, **67**, 1507–1510.
- 109 H. J. Li, T. Amagata, K. Tenney and P. Crews, *J. Nat. Prod.*, 2007, **70**, 802–807.
- 110 H. Zhang, P. Crews, K. Tenney and F. A. Valeriote, *Med. Chem.*, 2017, **13**, 295–300.
- 111 P. Karuso, R. C. Cambie, B. F. Bowden and P. R. Bergquist, *J. Nat. Prod.*, 1989, **52**, 289–293.
- 112 M. J. Somerville, J. N. A. Hooper and M. J. Garson, *J. Nat. Prod.*, 2006, **69**, 1587–1590.
- 113 M. M. Kumar, N. Krishna, P. Muralidhar, B. S. Sastry and D. V. Rao, *Indian J. Chem.*, 2008, **47B**, 325–328.
- 114 J. E. Jeon, J. Bae, K. J. Lee, K. B. Oh and J. Shin, *J. Nat. Prod.*, 2011, **74**, 847–851.
- 115 D. B. Abdjul, H. Yamazaki, O. Takahashi, R. Kirikoshi, R. E. Mangindaan and M. Namikoshi, *Bioorg. Med. Chem. Lett.*, 2015, 25, 904–907.
- 116 R. Kazlauskas, P. T. Murphy and R. J. Wells, *Aust. J. Chem.*, 1982, **35**, 51–59.
- 117 K. A. Alvi and P. Crews, J. Nat. Prod., 1992, 55, 859–865.

- 118 Y. Sera, K. Adachi and Y. Shizuri, J. Nat. Prod., 1999, 62, 152–154.
- 119 C. C. Stessman, R. Ebel, A. J. Corvino and P. Crews, *J. Nat. Prod.*, 2002, **65**, 1183–1186.
- 120 L. Chill, A. Rudi, M. Aknin, S. Loya, A. Hizi and Y. Kashman, *Tetrahedron*, 2004, **60**, 10619–10626.
- 121 J. Dai, Y. Liu, Y.-D. Zhou and D. G. Nagle, *J. Nat. Prod.*, 2007, **70**, 1824–1826.
- 122 Y. Liu, R. Liu, S. C. Mao, J. B. Morgan, M. B. Jekabsons,
 Y. D. Zhou and D. G. Nagle, *J. Nat. Prod.*, 2008, 71, 1854–1860.
- 123 B. R. Peng, K. H. Lai, Y. Y. Chen, J. H. Su, Y. M. Huang, Y. H. Chen, M. C. Lu, S. S. Yu, C. Y. Duh and P. J. Sung, *Mar. Drugs*, 2020, **18**, 76–86.
- 124 G. Cimino, S. De Stefano and L. Minale, *Tetrahedron*, 1972, 28, 5983–5991.
- 125 G. Cimino, S. De Stefano, L. Minale and E. Fattorusso, *Tetrahedron*, 1972, **28**, 267–273.
- 126 G. Cimino, S. De Stefano and A. Di Luccia, *Experientia*, 1979, **35**, 1277–1278.
- 127 B. Terem and P. J. Scheuer, *Tetrahedron*, 1986, **42**, 4409–4412.
- 128 A. De Giulio, S. De Rosa, G. Di Vincenzo and N. Zavodnik, *J. Nat. Prod.*, 1989, **52**, 1258–1262.
- 129 R. Davis and R. J. Capon, Aust. J. Chem., 1993, 46, 1295-1299.
- 130 H. He, P. Kulanthaivel and B. J. Baker, *Tetrahedron Lett.*, 1994, **35**, 7189–7192.
- 131 P. A. Searle and T. F. Molinski, *Tetrahedron*, 1994, **50**(33), 9893–9990.
- 132 A. Rueda, E. Zubía, M. J. Ortega, J. L. Carballo and J. Salvá, *J. Nat. Prod.*, 1998, **61**, 258–261.
- 133 S. De Marino, M. Iorizzi, F. Zollo, C. Debitus, J.-L. Menou, L. F. Ospina, M. J. Alcaraz and M. Payá, *J. Nat. Prod.*, 2000, 63, 322–326.
- 134 S. Tsukamoto, S. Miura, R. W. M. van Soest and T. Ohta, *J. Nat. Prod.*, 2003, **66**, 438–440.
- 135 S. J. Nam, H. Ko, M. K. Ju, H. Hwang, J. Chin, J. Ham, B. Lee,
 J. Lee, D. H. Won, H. Choi, J. Ko, K. Shin, T. Oh, S. Kim,
 J. R. Rho and H. Kang, *J. Nat. Prod.*, 2007, 70, 1691–1695.
- 136 S. J. Nam, H. Ko, M. Shin, J. Ham, J. Chin, Y. Kim, H. Kim, K. Shin, H. Choi and H. Kang, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 5398–5402.
- 137 E. Manzo, M. L. Ciavatta, G. Villani, M. Varcamonti, S. M. Sayem, R. van Soest and M. Gavagnin, *J. Nat. Prod.*, 2011, 74, 1241–1247.
- 138 C.-S. Phan, T. Kamada, T. Hamada and C. S. Vairappan, *Rec. Nat. Prod.*, 2018, **12**, 643–647.
- 139 I. Yang, J. Lee, J. Lee, D. Hahn, J. Chin, D. H. Won, J. Ko,
 H. Choi, A. Hong, S. J. Nam and H. Kang, *Molecules*, 2018,
 23(3187), 3181–3110.
- 140 C. Hertzer, S. Kehraus, N. Böhringer, F. Kaligis, R. Bara, D. Erpenbeck, G. Wörheide, T. F. Schäberle, H. Wägele and G. M. König, *Beilstein J. Org. Chem.*, 2020, 16, 1596– 1605.
- 141 J. R. Rho, H. S. Lee, H. J. Shin, J. W. Ahn, J. Y. Kim, C. J. Sim and J. Shin, *J. Nat. Prod.*, 2004, **67**, 1748–1751.

- 142 J. Song, W. Jeong, N. Wang, H.-S. Lee, C. J. Sim, K.-B. Oh and J. Shin, *J. Nat. Prod.*, 2008, **71**, 1866–1871.
- 143 P. R. Bergquist, R. C. Cambie and M. R. Kernan, *Biochem. Syst. Ecol.*, 1990, **18**, 349–357.
- 144 B. F. Bowden, J. C. Coll, H. Li, R. C. Cambie, M. R. Kernan and P. R. Bergquist, *J. Nat. Prod.*, 1992, 55, 1234–1240.
- 145 J. I. Jiménez, W. Y. Yoshida, P. J. Scheuer, E. Lobkovsky, J. Clardy and M. Kelly, *J. Org. Chem.*, 2000, **65**, 6837–6840.
- 146 Y. Kashman and M. Zviely, *Tetrahedron Lett.*, 1979, 20, 3879–3882.
- 147 R. P. Walker, J. E. Thompson and D. J. Faulkner, *J. Org. Chem.*, 1980, 45, 4976–4979.
- 148 G. Cimino, S. De Stefano and L. Minale, *Experientia*, 1974, 30, 846–847.
- 149 G. Cimino, A. Fontana, F. Giménez, A. Marin, E. Mollo, E. Trivellone and E. Zubía, *Experientia*, 1993, **49**, 582–586.
- 150 F. Cafieri, L. De Napoli, E. Fattorusso, C. Santacroce and D. Sica, *Tetrahedron Lett.*, 1977, **18**, 477–480.
- 151 F. Cafieri, L. De Napoli, E. Fattorusso, C. Santacroce and D. Sica, *Gazz. Chim. Ital.*, 1977, **107**, 71–74.
- 152 F. Cafieri, L. De Napoli, A. Iengo and C. Santacroce, *Experientia*, 1978, 34, 300–301.
- 153 F. Cafieri, L. De Napoli, A. Iengo and C. Santacroce, *Experientia*, 1979, **35**, 157–158.
- 154 M. Puliti, C. A. Mattia and L. Mazzarella, *Acta Crystallogr.*, 1995, **C51**, 1703–1707.
- 155 S. De Rosa, S. De Stefano and N. Zavodnik, *J. Org. Chem.*, 1988, **53**, 5020–5023.
- 156 S. Khushi, L. Nahar, A. A. Salim and R. J. Capon, *Mar. Drugs*, 2018, **16**, 456.
- 157 S. Wonganuchitmeta, S. Yuenyongsawad, N. Keawpradub and A. Plubrukarn, *J. Nat. Prod.*, 2004, **67**, 1767–1770.
- 158 C. Mahidol, H. Prawat, S. Sangpetsiripan and S. Ruchirawat, *J. Nat. Prod.*, 2009, **72**, 1870–1874.
- 159 Q. Wang, Y. Sun, L. Yang, X. Luo, N. J. de Voogd, X. Tang, P. Li and G. Li, *J. Nat. Prod.*, 2020, **83**, 516–523.
- 160 A. Rudi, T. Yosief, M. Schleyer and Y. Kashman, *Org. Lett.*, 1999, **1**, 471–472.
- 161 S. P. Gunasekera, P. J. McCarthy, M. Kelly-Borges, E. Lobkovsky and J. Clardy, *J. Am. Chem. Soc.*, 1996, **118**, 8759–8760.
- 162 H. Hagiwara and H. Uda, J. Chem. Soc., Chem. Commun., 1988, 815–817, DOI: 10.1039/C39880000815.
- 163 D. Lee, J. Shin, K. M. Yoon, T. I. Kim, S. H. Lee, H. S. Lee and K. B. Oh, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 5377–5380.
- 164 L. Murray, H. Hamit, J. Hooper, L. Hobbs and R. Capon, Aust. J. Chem., 1995, 48, 1899–1902.
- 165 D. Hahn, D. H. Won, B. Mun, H. Kim, C. Han, W. Wang, T. Chun, S. Park, D. Yoon, H. Choi, S. J. Nam, M. Ekins, J. Chin and H. Kang, *Bioorg. Med. Chem. Lett.*, 2013, 23, 2336–2339.
- 166 K. Choi, J. Hong, C.-O. Lee, D.-k. Kim, C. J. Sim, K. S. Im and J. H. Jung, *J. Nat. Prod.*, 2004, 67, 1186–1189.
- 167 W. Balansa, R. Islam, F. Fontaine, A. M. Piggott, H. Zhang, X. Xiao, T. I. Webb, D. F. Gilbert, J. W. Lynch and R. J. Capon, *Org. Biomol. Chem.*, 2013, 11, 4695–4701.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

Open Access Article. Published on 20 2025. Downloaded on 2025/8/21 23:50:35.

- 168 Y. C. Chang, S. W. Tseng, L. L. Liu, Y. Chou, Y. S. Ho, M. C. Lu and J. H. Su, *Mar. Drugs*, 2012, **10**, 987–997.
- 169 S. Rochfort, D. Atkin, L. Hobbs and R. Capon, *J. Nat. Prod.*, 1996, **59**, 1024–1028.
- 170 H. S. Lee, T. H. Lee, S. H. Yang, H. J. Shin, J. Shin and K. B. Oh, *Bioorg. Med. Chem. Lett.*, 2007, 17, 2483–2486.
- 171 W.-H. Jiao, L.-L. Hong, J.-B. Sun, S.-J. Piao, G.-D. Chen,
 H. Deng, S.-P. Wang, F. Yang and H.-W. Lin, *Eur. J. Org Chem.*, 2017, 2017, 3421–3426.
- 172 G. Cimino, S. De Stefano, L. Minale and R. Riccio, *Tetrahedron Lett.*, 1979, **38**, 3619–3622.
- 173 J. K. Woo, C. K. Kim, S. H. Kim, H. Kim, D. C. Oh, K. B. Oh and J. Shin, *Org. Lett.*, 2014, **16**, 2826–2829.
- 174 J.-R. Rho, B. S. Hwang, S. Joung, M. R. Byun, J.-H. Hong and H.-Y. Lee, *Org. Lett.*, 2011, **13**, 884–887.
- 175 J. R. Rho, B. S. Hwang, C. J. Sim, S. Joung, H. Y. Lee and H. J. Kim, *Org. Lett.*, 2009, **11**, 5590–5593.
- 176 J. Daoust, M. Chen, M. Wang, D. E. Williams, M. A. Garcia Chavez, Y. A. Wang, C. E. Merchant, A. Fontana, T. J. Kieffer and R. J. Andersen, *J. Org. Chem.*, 2013, **78**, 8267–8273.
- 177 J. Daoust, A. Fontana, C. E. Merchant, N. J. De Voogd,
 B. O. Patrick, T. J. Kieffer and R. J. Andersen, *Org. Lett.*, 2010, 12, 3208–3211.
- 178 W. Wang, Y. Lee, T. G. Lee, B. Mun, A. G. Giri, J. Lee, H. Kim, D. Hahn, I. Yang, J. Chin, H. Choi, S.-J. Nam and H. Kang, *Org. Lett.*, 2012, 14, 4486–4489.
- 179 M. Wang, I. Tietjen, M. Chen, D. E. Williams, J. Daoust, M. A. Brockman and R. J. Andersen, *J. Org. Chem.*, 2016, 81, 11324–11334.
- 180 M. Wang, A. Sciorillo, S. Read, D. N. Divsalar, K. Gyampoh, G. Zu, Z. Yuan, K. Mounzer, D. E. Williams, L. J. Montaner, N. de Voogd, I. Tietjen and R. J. Andersen, *J. Nat. Prod.*, 2022, 85, 1274–1281.
- 181 Y. Lee, W. Wang, H. Kim, A. G. Giri, D. H. Won, D. Hahn, K. R. Baek, J. Lee, I. Yang, H. Choi, S. J. Nam and H. Kang, *Bioorg. Med. Chem. Lett.*, 2014, 24, 4095–4098.
- 182 H. Solanki, C. Angulo-Preckler, K. Calabro, N. Kaur, P. Lasserre, B. Cautain, M. de la Cruz, F. Reyes, C. Avila and O. P. Thomas, *Tetrahedron Lett.*, 2018, **59**, 3353–3356.
- 183 W. Wang, B. Mun, Y. Lee, M. Venkat Reddy, Y. Park, J. Lee, H. Kim, D. Hahn, J. Chin, M. Ekins, S. J. Nam and H. Kang, *J. Nat. Prod.*, 2013, **76**, 170–177.
- 184 D. T. A. Youssef, W. Y. Yoshida, M. Kelly and P. J. Scheuer, J. Nat. Prod., 2001, 64, 1332–1335.
- 185 D. T. A. Youssef, J. Nat. Prod., 2004, 67, 112–114.
- 186 J. Dai, Y. Liu, Y.-D. Zhou and D. G. Nagle, *J. Nat. Prod.*, 2007, 70, 130–133.
- 187 S. R. M. Ibrahim, R. Ebel, V. Wray, W. E. G. Müller, R. A. Edrada-Ebel and P. Proksch, *J. Nat. Prod.*, 2008, **71**, 1358–1364.
- 188 S. Sperry, F. A. Valeriote, T. H. Corbett and P. Crews, *J. Nat. Prod.*, 1998, **61**, 241–247.
- 189 P. Phuwapraisirisan, S. Matsunaga, N. Fusetani, N. Chaitanawisuti, S. Kritsanapuntu and P. Menasveta, J. Nat. Prod., 2003, 66, 289–291.

- 190 J. Tanaka, T. Higa, K. Suwanborirux, U. Kokpol,
 G. Bernardinelli and C. W. Jefford, *J. Org. Chem.*, 1993, 58, 2999–3002.
- 191 J. Bae, J. E. Jeon, Y. J. Lee, H. S. Lee, C. J. Sim, K. B. Oh and J. Shin, *J. Nat. Prod.*, 2011, **74**, 1805–1811.
- 192 C. K. Kim, I. H. Song, H. Y. Park, Y. J. Lee, H. S. Lee, C. J. Sim, D. C. Oh, K. B. Oh and J. Shin, *J. Nat. Prod.*, 2014, 77, 1396–1403.
- 193 J.-W. Lee, H.-S. Lee, J. Shin, J. S. Kang, J. Yun, H. J. Shin, J. S. Lee and Y.-J. Lee, Arch. Pharmacal Res., 2015, 38, 1005–1010.
- 194 J. Kimura, E. Ishizuka, Y. Nakao, W. Y. Yoshida, P. J. Scheuer and M. Kelly-Borges, *J. Nat. Prod.*, 1998, **61**, 248–250.
- 195 S. De Marino, C. Festa, M. V. D'Auria, M.-L. Bourguet-Kondracki, S. Petek, C. Debitus, R. M. Andrés, M. C. Terencio, M. Payá and A. Zampella, *Tetrahedron*, 2009, 65, 2905–2909.
- 196 F. S. Di Leva, C. Festa, C. D'Amore, S. De Marino, B. Renga, M. V. D'Auria, E. Novellino, V. Limongelli, A. Zampella and S. Fiorucci, *J. Med. Chem.*, 2013, 56, 4701–4717.
- 197 G. Cimino, S. De Stefano and L. Minale, *Tetrahedron*, 1972, 28, 331–341.
- 198 W. Balansa, R. Islam, F. Fontaine, A. M. Piggott, H. Zhang, T. I. Webb, D. F. Gilbert, J. W. Lynch and R. J. Capon, *Bioorg. Med. Chem.*, 2010, 18, 2912–2919.
- 199 Y.-C. Shen, K.-L. Lo, Y.-C. Lin, A. T. Khalil, Y.-H. Kuo and P.-S. Shih, *Tetrahedron Lett.*, 2006, **47**, 4007–4010.
- 200 D. J. Faulkner, Tetrahedron Lett., 1973, 39, 3821–3822.
- 201 I. Rothberg and P. Shubiak, *Tetrahedron Lett.*, 1975, 16, 769–772.
- 202 A. Martinez, C. Duque, N. Sato and Y. Fujimoto, *Chem. Pharm. Bull.*, 1997, **45**, 181–184.
- 203 S. De Rosa, A. Milone, A. De Giulio, A. Crispino and C. Iodice, *Nat. Prod. Lett.*, 1996, 8, 245–251.
- 204 U. Höller, G. M. König and A. D. Wright, J. Nat. Prod., 1997, 60, 832–835.
- 205 A. Fontana, I. Fakhr, E. Mollo and G. Cimino, *Tetrahedron:* Asymmetry, 1999, **10**, 3869–3872.
- 206 H. B. Yu, B. B. Gu, A. Iwasaki, W. L. Jiang, A. Ecker, S. P. Wang, F. Yang and H. W. Lin, *Mar. Drugs*, 2020, 18(10), 491.
- 207 L. M. Murray, A. Johnson, M. C. Diaz and P. Crews, J. Org. Chem., 1997, 62, 5638–5641.
- 208 S. Tang, Y. Pei, H. Fu, Z. Deng, J. Li, P. Proksch and W. Lin, *Chem. Pharm. Bull.*, 2006, **54**, 4–8.
- 209 J. i. Kobayashi, K. Yuasa, T. Kobayashi, T. Sasaki and M. Tsuda, *Tetrahedron*, 1996, **52**, 5745–5750.
- 210 E. Elkhayat, R. Edrada, R. Ebel, V. Wray, R. van Soest, S. Wiryowidagdo, M. H. Mohamed, W. E. G. Müller and P. Proksch, *J. Nat. Prod.*, 2004, **67**, 1809–1817.
- 211 M. Ohba, K. Iizuka, H. Ishibashi and T. Fujii, *Tetrahedron*, 1997, **53**, 16977–16986.
- 212 R. Ueoka, Y. Nakao, S. Fujii, R. W. M. van Soest and S. Matsunaga, *J. Nat. Prod.*, 2008, **71**, 1089–1091.
- 213 P. Crews, C. Jiménez and M. O'Neil-Johnson, *Tetrahedron*, 1991, 47, 3585–3600.

- 214 L. Gomez Paloma, A. Randazzo, L. Minale, C. Debitus and C. Roussakis, *Tetrahedron*, 1997, **53**, 10451–10458.
- 215 A. Randazzo, C. Debitus, L. Minale, P. García Pastor, M. J. Alcaraz, M. Payá and L. Gomez-Paloma, *J. Nat. Prod.*, 1998, 61, 571–575.
- 216 J. Li, B. Xu, J. Cui, Z. Deng, N. J. de Voogd, P. Proksch and W. Lin, *Bioorg. Med. Chem.*, 2010, 18, 4639–4647.
- 217 S. Ohta, M. Uno, M. Yoshimura, Y. Hiraga and S. lkegami, *Tetrahedron Lett.*, 1996, 37, 2265–2266.
- 218 M. Yanai, S. Ohta, E. Ohta and S. Ikegami, *Tetrahedron*, 1998, 54, 15607–15612.
- 219 M. S. Butler and R. J. Capon, *Aust. J. Chem.*, 1992, **45**, 1705–1743.
- 220 K. F. Albizati, T. Holman, D. J. Faulkner, K. B. Glaser and R. S. Jacobs, *Experientia*, 1987, **43**, 949–950.
- 221 G. M. Köenig, A. D. Wright and O. Sticher, *J. Nat. Prod.*, 1992, 55, 174–178.
- 222 P. Ettinger-Epstein, C. A. Motti, R. de Nys, A. D. Wright,C. N. Battershill and D. M. Tapiolas, *J. Nat. Prod.*, 2007,70, 648–651.
- 223 M. R. Kernan, D. J. Faulkner and R. S. Jacobs, *J. Org. Chem.*, 1987, **52**, 3081–3083.
- 224 M. R. Kernan, D. J. Faulkner, L. Parkanyi, J. Clardy, M. S. de Carvalho and R. S. Jacobs, *Experientia*, 1989, **45**, 388–390.
- 225 J. i. Kobayashi, C.-m. Zeng, M. Ishibashi and T. Sasaki, *J. Nat. Prod.*, 1993, **56**, 436–439.
- 226 M. Tsuda, T. Endo, Y. Mikami, J. Fromont and J. Kobayashi, *J. Nat. Prod.*, 2002, **65**, 1507–1508.
- 227 T.-R. Su, M.-C. Lu, Y.-J. Wu and J.-H. Su, *Bull. Chem. Soc. Jpn.*, 2015, **88**, 176–182.
- 228 P. Ahmadi, M. Higashi, N. J. Voogd and J. Tanaka, *Mar. Drugs*, 2017, **15**, 1–8.
- 229 D. Kanki, K. Imai, Y. Ise, S. Okada and S. Matsunaga, *J. Nat. Prod.*, 2021, **84**, 1676–1680.
- 230 Y. Liu, B. H. Bae, N. Alam, J. Hong, C. J. Sim, C.-O. Lee,K. S. Im and J. H. Jung, *J. Nat. Prod.*, 2001, 64, 1301–1304.
- 231 Y. Liu, J. Hong, C. O. Lee, K. S. Im, N. D. Kim, J. S. Choi and J. H. Jung, *J. Nat. Prod.*, 2002, **65**, 1307–1314.
- 232 Y. Liu, T. A. Mansoor, J. Hong, C. O. Lee, C. J. Sim, K. S. Im, N. D. Kim and J. H. Jung, *J. Nat. Prod.*, 2003, **66**, 1451–1456.
- 233 J.-K. Woo, J. E. Jeon, B. Kim, C. Sim, D.-C. Oh, K.-B. Oh and J. Shin, *Nat. Prod. Sci.*, 2015, **21**, 237.
- 234 W. He, X. Lin, T. Xu, J. H. Jung, H. Yin, B. Yang and Y. Liu, *Chem. Nat. Compd.*, 2012, **48**, 208–210.
- 235 C. J. Barrow, J. W. Blunt, M. H. G. Munro and N. B. Perry, *J. Nat. Prod.*, 1988, **51**, 275–281.
- 236 P. Prasad, A. Zhang, A. A. Salim and R. J. Capon, *Fitoterapia*, 2018, **126**, 83–89.
- 237 J. Shin, Y. Seo, J.-R. Rho, E. Baek, B.-M. Kwon, T.-S. Jeong and S.-H. Bok, *J. Org. Chem.*, 1995, **60**, 7582–7588.
- 238 H. S. Lee, J. W. Ahn, Y. H. Lee, J. R. Rho and J. Shin, *J. Nat. Prod.*, 2004, **67**, 672–674.
- 239 A. R. Díaz-Marrero, I. Brito, M. Cueto, A. San-Martín and J. Darias, *Tetrahedron Lett.*, 2004, **45**, 4707–4710.
- 240 R. D. Charan, T. C. McKee and M. R. Boyd, *J. Nat. Prod.*, 2001, **64**, 661–663.

- 241 R. C. Cambie, P. A. Craw, P. R. Bergquist and P. Karuso, *J. Nat. Prod.*, 1988, **51**, 331–334.
- 242 C. Audoin, D. Bonhomme, J. Ivanisevic, M. de la Cruz,B. Cautain, M. C. Monteiro, F. Reyes, L. Rios, T. Perez and O. P. Thomas, *Mar. Drugs*, 2013, 11, 1477–1489.
- 243 W. F. Ponder and D. R. Lindberg, *Zool. J. Linn. Soc.*, 1997, **119**, 83–265.
- 244 H. Wägele, A. Klussmann-Kolb, E. Verbeek and M. Schrödl, *Org. Divers. Evol.*, 2014, **14**, 133–149.
- 245 H. Wägele and A. Klussmann-Kolb, *Front. Zool.*, 2005, **2**, 1–18.
- 246 K. Göbbeler and A. Klussmann-Kolb, *Thalassas*, 2011, 27, 121–153.
- 247 M. Medina, S. Lal, Y. Vallès, T. L. Takaoka, B. A. Dayrat, J. L. Boore and T. Gosliner, *Mar. Genomics*, 2011, 4, 51–59.
- 248 C. Avila, L. Núñez-Pons and J. Moles, in *Chemical Ecology*, ed. M. P. Puglisi and M. A. Becerro, CRC Press, 1st edn, 2018, pp. 71–163.
- 249 H. Wagele, Org. Divers. Evol., 2004, 4, 175-188.
- 250 B. Burghardt, J. Evertsen, G. Johnsen and H. Wägele, Symbiosis, 2005, 38, 227–250.
- 251 P. G. Greenwood and R. N. Mariscal, *Tissue Cell*, 1984, **16**, 719–730.
- 252 G. Cimino and M. T. Ghiselin, *Chemoecology*, 1999, **9**, 187–207.
- 253 C. Avila, Oceanogr. Mar. Biol. Annu. Rev., 1995, 33, 487-559.
- 254 C. Avila and V. Paul, *Mar. Ecol. Prog. Ser.*, 1997, **150**, 171–180.
- 255 A. Fontana, F. Giménez, A. Marin, E. Mollo and G. Cimino, *Experientia*, 1994, **50**, 510–516.
- 256 T. M. Gosliner, Boll. Malacol., 2001, 37, 163-170.
- 257 C. Avila, Mar. Drugs, 2020, 18, 162.
- 258 R. S. Jacobs, M. A. Bober, I. Pinto, A. B. Williams, P. B. Jacobson and M. S. De Carvalho, in *Pharmaceutical* and *Bioactive Natural Products*, ed. D. H. Attaway and O. R. Zaborsky, Springer US, Boston, MA, 1993, pp. 77–99.
- 259 A. Fontana, E. Mollo, J. Ortea, M. Gavagnin and G. Cimino, *J. Nat. Prod.*, 2000, **63**, 527–530.
- 260 J. E. Hochlowski, D. J. Faulkner, L. S. Bass and J. Clardy, *J. Org. Chem.*, 1983, **48**, 1738–1740.
- 261 M. R. Kernan, E. B. Barrabee and D. John Faulkner, *Comp. Biochem. Physiol., B*, 1988, **89**, 275–278.
- 262 A. Fontana, P. Cavaliere, N. Ungur, L. D'Souza,
 P. S. Parameswaram and G. Cimino, *J. Nat. Prod.*, 1999,
 62, 1367–1370.
- 263 K. A. El Sayed, P. Bartyzel, X. Shen, T. L. Perry, J. K. Zjawiony and M. T. Hamann, *Tetrahedron*, 2000, **56**, 949–953.
- 264 A. E. Winters, A. M. White, A. S. Dewi, I. W. Mudianta, N. G. Wilson, L. C. Forster, M. J. Garson and K. L. Cheney, *J. Chem. Ecol.*, 2018, 44, 384–396.
- 265 K. W. L. Yong, I. W. Mudianta, K. L. Cheney, E. Mollo, J. T. Blanchfield and M. J. Garson, *J. Nat. Prod.*, 2015, 78, 421–430.
- 266 J. Hellou, R. J. Andersen, S. Rafii, E. Arnold and J. Clardy, *Tetrahedron Lett.*, 1981, 22, 4173–4176.
- 267 J. Pika and D. John Faulkner, *Tetrahedron*, 1995, **51**, 8189–8198.

View Article Online Natural Product Reports

- 268 T. Miyamoto, K. Sakamoto, H. Amano, Y. Arakawa, Y. Nagarekawa, T. Komori, R. Higuchi and T. Sasaki, *Tetrahedron*, 1999, 55, 9133–9142.
- 269 K. McPhail, M. T. Davies-Coleman and P. Coetzee, J. Nat. Prod., 1998, 61, 961–964.
- 270 J. Moles, H. Wägele, A. Cutignano, A. Fontana and C. Avila, *Mar. Biol.*, 2016, **163**, 54.
- 271 J. E. N. Veron, *Coral in Space and Time The Biogeography and Evolution of the Scleractinia*, Cornell University Press, Ithaca, NY, 1995.
- 272 S. K. Davy, D. Allemand and V. M. Weis, *Microbiol. Mol. Biol. Rev.*, 2012, **76**, 229–261.
- 273 E. Ballesteros, Oceanogr. Mar. Biol. Annu. Rev., 2006, 44, 123–195.
- 274 R. G. Waller, in *Cold-Water Corals and Ecosystems*, ed. A. Freiwald and J. M. Roberts, Springer Berlin Heidelberg, Berlin, Heidelberg, 2005, pp. 691–700.
- 275 Z. Dubinsky and N. Stambler, *Coral reefs: an ecosystem in transition*, Springer Netherlands, Dordrecht, 2011.
- 276 A. Fontana, M. L. Ciavatta and G. Cimino, J. Org. Chem., 1998, 63, 2845–2849.
- 277 T. Noda, C. Kitamura, S. Takahashi, K. Takagi, T. Kashio and M. Tanaka, *Appl. Entomol. Zool.*, 1982, **17**, 350–357.
- 278 T. Ríos and S. Pérez C, *J. Chem. Soc. D*, 1969, 214–215, DOI: 10.1039/C29690000214.
- 279 S. Nozoe, M. Morisaki, K. Fukushlma and S. Okuda, *Tetrahedron Lett.*, 1968, **42**, 4457–4458.
- 280 L. Quijano, J. S. Calderón and T. Ríos, *Chem. Lett.*, 1979, 1387–1388.
- 281 R. Kazlauskas, P. T. Murphy, R. J. Quinn and R. J. Wells, *Tetrahedron Lett.*, 1976, **30**, 2635–2636.
- 282 R. Veloz, L. Quijano, J. S. Calderon and T. Rios, *J. Chem. Soc., Chem. Commun.*, 1975, 191–192.
- 283 F. Miyamoto, H. Naoki, T. Takemoto and Y. Naya, *Tetrahedron*, 1979, **35**, 1913–1917.
- 284 J. K. Pawlak, M. S. Tempesta, T. Iwashita, K. Nakanishi and Y. Naya, *Chem. Lett.*, 1983, 1069–1072.
- 285 M. Toki, T. Ooi and T. Kusumi, *J. Nat. Prod.*, 1999, **62**, 1504–1509.
- 286 T. Rios and F. Colunga, Chem. Ind., 1965, 26, 1184-1185.
- 287 Y. Iitaka, I. Watanabe, I. T. Harrison and S. Harrison, *J. Am. Chem. Soc.*, 1968, **90**, 1092–1093.
- 288 T. Rios and L. Quijano, *Tetrahedron Lett.*, 1969, **17**, 1317–1318.
- 289 T. Rios and F. Gomez, *Tetrahedron Lett.*, 1969, 34, 2929–2930.
- 290 J. S. Calderon, L. Quijano and T. Rios, *Chem. Ind.*, 1978, **15**, 584–585.
- 291 J. S. Calderon, L. Quijano and T. Rios, *Experientia*, 1978, 34, 421–422.
- 292 R. K. Boeckman, D. M. Blum, E. V. Arnold and J. Clardy, *Tetrahedron Lett.*, 1979, **20**, 4609–4612.
- 293 R. K. Boeckman, D. M. Blum and S. D. Arthur, J. Am. Chem. Soc., 1979, 101, 5060–5062.
- 294 G. Brochere-Ferreol and J. Polonsky, Bull. Soc. Chim. Fr., 1960, 963–967.

- 295 R. Scartazzini, Doctoral thesis, ETH, 1966, DOI: 10.3929/ ethz-a-000088868.
- 296 Y. Naya, K. Yoshihara, T. Iwashita, H. Komura, K. Nakanishi and Y. Hata, *J. Am. Chem. Soc.*, 1981, **103**, 7009–7011.
- 297 P. Zhang, J. Qi, Y. Duan, J. M. Gao and C. Liu, *J. Fungi*, 2022, 8, 1080.
- 298 R. Lucking, M. C. Aime, B. Robbertse, A. N. Miller, T. Aoki, H. A. Ariyawansa, G. Cardinali, P. W. Crous, I. S. Druzhinina, D. M. Geiser, D. L. Hawksworth, K. D. Hyde, L. Irinyi, R. Jeewon, P. R. Johnston, P. M. Kirk, E. Malosso, T. W. May, W. Meyer, H. R. Nilsson, M. Opik, V. Robert, M. Stadler, M. Thines, D. Vu, A. M. Yurkov, N. Zhang and C. L. Schoch, *Nat. Microbiol.*, 2021, 6, 540–548.
- 299 T. A. Richards, G. Leonard and J. G. Wideman, *Microbiol.* Spectr., 2017, 5, DOI: 10.1128/microbiolspec.FUNK-0044-2017.
- 300 D. L. Hawksworth and R. Lucking, *Microbiol. Spectr.*, 2017,
 5, DOI: 10.1128/microbiolspec.funk-0052-2016.
- 301 Q. Dai, F. L. Zhang and T. Feng, J. Fungi, 2021, 7, 1026.
- 302 W. Tian, Z. Deng and K. Hong, *Mar. Drugs*, 2017, **15**(229), 221.
- 303 K. Ishibashi, J. Agric. J. Agric. Chem. Soc. Jpn., 1962, **36**, 226–228.
- 304 F. Sugawara, G. Strobel, R. N. Strange, J. N. Siedow,
 G. D. Van Duyne and J. Clardy, *Proc. Natl. Acad. Sci. U. S. A.*, 1987, 84, 3081–3085.
- 305 M. Orsenigo, Phytopathol. Z., 1957, 29, 189-196.
- 306 M. Orsenigo and D. Pavan, Ann. Facul. Agric., 1958, 6, 19–54.
- 307 M. Nakamura, K. Ishibashi and J. Agric, J. Agric. Chem. Soc. Jpn., 1958, 32, 739–744.
- 308 L. Canonica, A. Fiecchi, M. G. Kienle and A. Scala, *Tetrahedron Lett.*, 1966, 7, 1211–1218.
- 309 L. Radics, M. Kajtar-Peredy, S. Nozoe and H. Kobayashi, *Tetrahedron Lett.*, 1975, **49**, 4415–4418.
- 310 J. Z. Xiao, M. Tsuda, N. Doke and S. Nishimura, *Phytopathology*, 1991, **81**, 58–64.
- 311 P. Phuwapraisirisan, K. Sawang, P. Siripong and S. Tippyang, *Tetrahedron Lett.*, 2007, **48**, 5193–5195.
- 312 C.-H. Yun, F. Sugawara and G. A. Strobel, *Plant Sci.*, 1988, 54, 237–243.
- 313 D. R. Bhatia, P. Dhar, V. Mutalik, S. K. Deshmukh, S. A. Verekar, D. C. Desai, R. Kshirsagar, P. Thiagarajan and V. Agarwal, *Nat. Prod. Rep.*, 2016, **30**, 1455–1458.
- 314 L. M. Pena-Rodriguez and W. S. Chilton, J. Nat. Prod., 1989, 52, 1170–1172.
- 315 E. Li, A. M. Clark, D. P. Rotella and C. D. Hufford, J. Nat. Prod., 1995, 58, 74–81.
- 316 X. Shen, S. B. Krasnoff, S.-W. Lu, C. D. Dunbar, J. O'Neal,
 B. G. Turgeon, O. C. Yoder, D. M. Gibson and
 M. T. Hamann, *J. Nat. Prod.*, 1999, 62, 895–897.
- 317 H. Ohkawa and T. Tamura, *Agric. Biol. Chem.*, 1966, **30**, 285–291.
- 318 A. Evidente, A. Andolfi, A. Cimmino, M. Vurro, M. Fracchiolla and R. Charudattan, J. Agric. Food Chem., 2006, 54, 1779–1783.

- 319 A. Evidente, A. Andolfi, A. Cimmino, M. Vurro, M. Fracchiolla, R. Charudattan and A. Motta, *Phytochemistry*, 2006, 67, 2281–2287.
- 320 K. Ishibashi, J. Antibiot., 1962, 15, 88–92.
- 321 S. Nozoe, K. Hirai and K. Tsuda, *Tetrahedron Lett.*, 1966, 7, 2211–2216.
- 322 K. Ishibashi, J. Agric. Chem. Soc. Jpn., 1961, 35, 257-262.
- 323 J.-M. Kim, S.-B. Hyeon, A. Isogai and A. Suzuki, *Agric. Biol. Chem.*, 2014, **48**, 803–805.
- 324 M. W. Canales and G. R. Gray, *Phytochemistry*, 1988, 27, 1653–1663.
- 325 S. Nozoe and M. Morisaki, *Chem. Commun.*, 1969, 1319-1320.
- 326 Q. X. Wang, J. L. Yang, Q. Y. Qi, L. Bao, X. L. Yang, M. M. Liu, P. Huang, L. X. Zhang, J. L. Chen, L. Cai and H. W. Liu, *Bioorg. Med. Chem. Lett.*, 2013, 23, 3547–3550.
- 327 L. Canonica, A. Friecchi, K. U. Galli and A. Scala, *Tetrahedron Lett.*, 1966, 7, 1329–1333.
- 328 J. W. Ahn, M. K. Lee, S. U. Choi, C. O. Lee and B. S. Kim, *J. Microbiol. Biotechnol.*, 1998, **8**, 406–408.
- 329 F. Sugawara, N. Takahashi, G. Strobel, C. H. Yun, G. George,
 Y. Fu and J. Clardy, *J. Org. Chem.*, 1988, 53, 2170–2172.
- 330 M. Liu, W. Sun, L. Shen, X. Hao, W. H. Al Anbari, S. Lin, H. Li, W. Gao, J. Wang, Z. Hu and Y. Zhang, *J. Nat. Prod.*, 2019, **82**, 2897–2906.
- 331 M.-T. Liu, Y. He, L. Shen, Z.-X. Hu and Y.-H. Zhang, *Chin. J. Nat. Med.*, 2019, **17**, 935–944.
- 332 X. Duan, X. Tan, L. Gu, J. Liu, X. Hao, L. Tao, H. Feng, Y. Cao, Z. Shi, Y. Duan, M. Deng, G. Chen, C. Qi and Y. Zhang, *Bioorg. Chem.*, 2020, **99**, 103816.
- 333 R. Cai, H. Jiang, Y. Mo, H. Guo, C. Li, Y. Long, Z. Zang and Z. She, *J. Nat. Prod.*, 2019, **82**, 2268–2278.
- 334 Q. X. Wang, L. Bao, X. L. Yang, D. L. Liu, H. Guo, H. Q. Dai,
 F. H. Song, L. X. Zhang, L. D. Guo, S. J. Li and H. W. Liu, *Fitoterapia*, 2013, **90**, 220–227.
- 335 H. G. Cutler, F. G. Crumley, R. H. Cox, J. P. Springer, R. F. Arrendale, R. J. Cole and P. D. Cole, *J. Agric. Food Chem.*, 1984, 32, 778–782.
- 336 S. B. Singh, J. L. Smith, G. S. Sabnis, A. W. Dombrowski, J. M. Schaeffer, M. A. Goetz and G. F. Bills, *Tetrahedron*, 1991, 47, 6931–6938.
- 337 T. Zhu, Z. Lu, J. Fan, L. Wang, G. Zhu, Y. Wang, X. Li, K. Hong, P. Piyachaturawat, A. Chairoungdua and W. Zhu, J. Nat. Prod., 2018, 81, 2–9.
- 338 X.-H. Liu, F.-P. Miao, M.-F. Qiao, R. H. Cichewicz and N.-Y. Ji, *RSC Adv.*, 2013, **3**, 588–595.
- 339 H.-B. Liu, R. Edrada-Ebel, R. Ebel, Y. Wang, B. Schulz, S. Draeger, W. E. G. Müller, V. Wray, W.-H. Lin and P. Proksch, *Helv. Chim. Acta*, 2011, 94, 623–631.
- 340 B. K. Choi, P. T. H. Trinh, H. S. Lee, B. W. Choi, J. S. Kang, N. T. D. Ngoc, T. T. T. Van and H. J. Shin, *Mar. Drugs*, 2019, 17(346), 341–349.
- 341 T. T. Bladt, C. Dürr, P. B. Knudsen, S. Kildgaard, J. C. Frisvad, C. H. Gotfredsen, M. Seiffert and T. O. Larsen, *Molecules*, 2013, 18, 14629–14650.
- 342 A. Itai, S. Nozoe, K. Tsuda, S. Okuda and Y. Iitaka, *Tetrahedron Lett.*, 1967, **42**, 4111–4112.

- 343 S. Nozoe, A. Itai, K. Tsuda and S. Okuda, *Tetrahedron Lett.*, 1967, **8**, 4113–4117.
- 344 A. Itai, S. Nozoe, S. Okuda and Y. Iitaka, Acta Crystallogr. Sect. B Struct. Sci., 1969, 25, 872–881.
- 345 H. Wei, T. Itoh, M. Kinoshita, Y. Nakai, M. Kurotaki and M. Kobayashi, *Tetrahedron*, 2004, **60**, 6015–6019.
- 346 M. Arai, H. Niikawa and M. Kobayashi, J. Nat. Med., 2013, 67, 271–275.
- 347 A. Tsipouras, A. A. Adefarati, J. S. Tkacz, E. G. Frazier,
 S. P. Rohrer, E. Birzin, A. Rosegay, D. L. Zink,
 M. A. Goetz, S. B. Singh and J. M. Schaeffer, *Bioorg. Med. Chem.*, 1996, 4, 531–536.
- 348 S. B. Singh, R. A. Reamer, D. Zink, D. Schmatz, A. Dombrowski and M. A. Goetz, *J. Org. Chem.*, 1991, 56, 5618–5622.
- 349 M. Wei, P. Zhou, L. Huang, J. Yin, Q. Li, C. Dai, J. Wang, L. Gu, Q. Tong, H. Zhu and Y. Zhang, *Phytochemistry*, 2021, **191**, 112910.
- 350 N. Kawahara, M. Nozawa, A. Kurata, T. Hakamatsuka, S. Sekita and M. Satake, *Chem. Pharm. Bull.*, 1999, **47**, 1344–1345.
- 351 C.-J. Guo, B. P. Knox, Y.-M. Chiang, H.-C. Lo, J. F. Sanchez, K.-H. Lee, B. R. Oakley, K. S. Bruno and C. C. C. Wang, *Org. Lett.*, 2012, **14**, 5684–5687.
- 352 J. P. Springer, J. W. Dorner, R. J. Cole and R. H. Cox, *J. Org. Chem.*, 1979, **44**, 4852–4854.
- 353 G. Y. Li, B. G. Li, T. Yang, J. H. Yin, H. Y. Qi, G. Y. Liu and G. L. Zhang, *J. Nat. Prod.*, 2005, **68**, 1243–1246.
- 354 T. Fukuda, Y. Kurihara, A. Kanamoto and H. Tomoda, J. Antibiot., 2014, **67**, 593–595.
- 355 M. P. López-Gresa, N. Cabedo, M. C. González-Mas, M. L. Ciavatta, C. Avila and J. Primo, J. Nat. Prod., 2009, 72, 1348–1351.
- 356 G. K. Oleinikova, V. A. Denisenko, D. V. Berdyshev, M. A. Pushilin, N. N. Kirichuk, N. I. Menzorova, A. S. Kuzmich, E. A. Yurchenko, O. I. Zhuravleva and S. S. Afiyatullov, *Phytochem. Lett.*, 2016, 17, 135–139.
- 357 M. Chinworrungsee, P. Kittakoop, M. Isaka, A. Rungrod, M. Tanticharoen and Y. Thebtaranonth, *Bioorg. Med. Chem. Lett.*, 2001, **11**, 1965–1969.
- 358 M. Chinworrungsee, P. Kittakoop, M. Isaka, R. Chanphen, M. Tanticharoen and Y. Thebtaranonth, J. Chem. Soc., Perkin Trans. 1, 2002, 2473–2476, DOI: 10.1039/B207887M.
- 359 R. Chiba, A. Minami, K. Gomi and H. Oikawa, *Org. Lett.*, 2013, **15**, 594–597.
- 360 B. Ye, W. Ding, P.-M. Wang and J. Xu, Chem. Nat. Compd., 2019, 55, 281–284.
- 361 C. H. Lim, Agric. Chem. Biotechnol., 1996, 39, 241-244.
- 362 C.-H. Lim and Y. Nihashi, *J. Appl. Bio Chem.*, 2018, **61**, 79–82.
- 363 C. Intaraudom, S. Nitthithanasilp, P. Rachtawee,
 T. Boonruangprapa, S. Prabpai, P. Kongsaeree and
 P. Pittayakhajonwut, *Phytochemistry*, 2015, **120**, 19–27.
- 364 O. D. Hensens, D. Zink, J. M. Williamson, V. J. Lotti, R. S. L. Chang and M. A. Goetz, *J. Org. Chem.*, 1991, 56, 3399–3403.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

Open Access Article. Published on 20 2025. Downloaded on 2025/8/21 23:50:35.

- 365 K.-i. Kawai, K. Nozawa and S. Nakajima, J. Chem. Soc., Perkin Trans. 1, 1994, 1673–1674, DOI: 10.1039/ P19940001673.
- 366 K. Yoganathan, C. Rossant, R. P. Glover, S. Cao, J. J. Vittal, S. Ng, Y. Huang, A. D. Buss and M. S. Butler, *J. Nat. Prod.*, 2004, 67, 1681–1684.
- 367 H. Fujimoto, E. Nakamura, E. Okuyama and M. Ishibashi, *Chem. Pharm. Bull.*, 2000, **48**, 1436–1441.
- 368 Y. Tezuka, A. Takahashi, M. Maruyama, T. Tamamura, S. Kutsuma, H. Naganawa and T. Takeuchi, Novel antibiotics, AB5362-A, B, and C, their manufacture, their use as fungicides, and Phoma species, *Japan Pat.*, JP 10045662, 1998.
- 369 H. Takahashi, T. Hosoe, K. Nozawa and K.-i. Kawai, *J. Nat. Prod.*, 1999, **62**, 1712–1713.
- 370 M. Cueto, P. R. Jensen and W. Fenical, *Org. Lett.*, 2002, 4, 1583–1585.
- 371 Y.-L. Li, Y. Gao, C.-Y. Liu, C.-J. Sun, Z.-T. Zhao and H.-X. Lou, *J. Nat. Prod.*, 2019, **82**, 1527–1534.
- 372 C. Ding, X. Wu, B. N. Auckloo, C.-T. A. Chen, Y. Ye, K. Wang and B. Wu, *Molecules*, 2016, **21**, 105.
- 373 L. Jiang, G. Zhu, J. Han, C. Hou, X. Zhang, Z. Wang, W. Yuan, K. Lv, Z. Cong, X. Wang, X. Chen, L. Karthik, H. Yang, X. Wang, G. Tan, G. Liu, L. Zhao, X. Xia, X. Liu, S. Gao, L. Ma, M. Liu, B. Ren, H. Dai, R. J. Quinn, T. Hsiang, J. Zhang, L. Zhang and X. Liu, *Appl. Microbiol. Biotechnol.*, 2021, **105**, 5407–5417.
- 374 S. Kusari, C. Hertweck and M. Spiteller, *Chem. Biol.*, 2012, 19, 792–798.
- 375 M. Okada, Y. Matsuda, T. Mitsuhashi, S. Hoshino, T. Mori,
 K. Nakagawa, Z. Quan, B. Qin, H. Zhang, F. Hayashi,
 H. Kawaide and I. Abe, *J. Am. Chem. Soc.*, 2016, 138, 10011–10018.
- 376 K. D. Bauman, K. S. Butler, B. S. Moore and J. R. Chekan, *Nat. Prod. Rep.*, 2021, **38**, 2100–2129.
- 377 Y. Matsuda, T. Mitsuhashi, S. Lee, M. Hoshino, T. Mori, M. Okada, H. Zhang, F. Hayashi, M. Fujita and I. Abe, *Angew. Chem., Int. Ed.*, 2016, 55, 5785–5788.
- 378 Y. Matsuda, T. Mitsuhashi, Z. Quan and I. Abe, *Org. Lett.*, 2015, **17**, 4644–4647.
- 379 T. Mitsuhashi, J. Rinkel, M. Okada, I. Abe and J. S. Dickschat, *Chem. Eur J.*, 2017, **23**, 10053–10057.
- 380 O. Mosunova, J. C. Navarro-Muñoz and J. Collemare, in *Encyclopedia of Mycology*, ed. Ó. Zaragoza and A. Casadevall, Elsevier, Oxford, 2021, vol. 2, pp. 458–476.
- 381 D. W. Christianson, Chem. Rev., 2017, 117, 11570-11648.
- 382 Y. Ye, A. Minami, A. Mandi, C. Liu, T. Taniguchi, T. Kuzuyama, K. Monde, K. Gomi and H. Oikawa, J. Am. Chem. Soc., 2015, 137, 11846–11853.
- 383 I. H. Qureshi, S. A. Husain, R. Noorani, N. Murtaza, Y. Iitaka, S. Iwasaki and S. Okuda, *Tetrahedron Lett.*, 1980, 21, 1961–1962.
- 384 L. Gao, K. Narita, T. Ozaki, N. Kumakura, P. Gan, A. Minami, C. Liu, X. Lei, K. Shirasu and H. Oikawa, *Tetrahedron Lett.*, 2018, 59, 1136–1139.
- 385 K. Narita, H. Sato, A. Minami, K. Kudo, L. Gao, C. Liu, T. Ozaki, M. Kodama, X. Lei, T. Taniguchi, K. Monde,

M. Yamazaki, M. Uchiyama and H. Oikawa, Org. Lett., 2017, 19, 6696-6699.

- 386 R. Chen, Q. Jia, X. Mu, B. Hu, X. Sun, Z. Deng, F. Chen, G. Bian and T. Liu, *Proc. Natl. Acad. Sci. U. S. A.*, 2021, 118, e2023247118.
- 387 K. Narita, A. Minami, T. Ozaki, C. Liu, M. Kodama and H. Oikawa, *J. Org. Chem.*, 2018, **83**, 7042–7048.
- 388 J. Yan, J. Pang, J. Liang, W. Yu, X. Liao, A. Aobulikasimu, X. Yi, Y. Yin, Z. Deng and K. Hong, *Int. J. Mol. Sci.*, 2022, 23(3), 1903.
- 389 J. H. Huang, J. M. Lv, Q. Z. Wang, J. Zou, Y. J. Lu, Q. L. Wang, D. N. Chen, X. S. Yao, H. Gao and D. Hu, *Org. Biomol. Chem.*, 2019, **17**, 248–251.
- 390 H.-Y. Huang, J.-H. Huang, Y.-H. Wang, D. Hu, Y.-J. Lu, Z.-G. She, G.-D. Chen, X.-S. Yao and H. Gao, *Front. Chem.*, 2021, 9, 785431.
- 391 W. Yang, T. Chen, Y. Chen, Q. Tan, Y. Ou, G. Li, B. Wang, D. Hu, H. Yao and Z. She, *J. Org. Chem.*, 2022, 87, 16807– 16819.
- 392 C. Rossi and L. Tuttobello, *Tetrahedron Lett.*, 1978, **19**, 307– 308.
- 393 C. Heddergott, A. M. Calvo and J. P. Latge, *Eukaryot. Cell*, 2014, **13**, 1014–1025.
- 394 M. I. Choudhary, R. Ranjit, R. Atta Ur, K. P. Devkota, S. G. Musharraf and T. M. Shrestha, *Phytochemistry*, 2006, 67, 439–443.
- 395 A. C. Stierle, J. H. Cardellina and G. A. Strobel, *Proc. Natl. Acad. Sci. U. S. A.*, 1988, **85**, 8008–8011.
- 396 L. Gianani, S. Cocucci, D. Pardi and G. Randazzo, *Planta*, 1979, **146**, 271–274.
- 397 T. K. Au, W. S. Chick and P. C. Leung, *Life Sci.*, 2000, 67, 733–742.
- 398 Y. Yamada, T. Kuzuyama, M. Komatsu, K. Shin-Ya, S. Omura, D. E. Cane and H. Ikeda, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 857–862.
- 399 O. Genilloud, Nat. Prod. Rep., 2017, 34, 1203-1232.
- 400 Y. Yang, Y. Zhang, S. Zhang, Q. Chen, K. Ma, L. Bao, Y. Tao,
 W. Yin, G. Wang and H. Liu, *J. Nat. Prod.*, 2018, 81, 1089–1092.
- 401 J. Rinkel, L. Lauterbach and J. S. Dickschat, *Angew Chem. Int. Ed. Engl.*, 2017, **56**, 16385–16389.
- 402 T. Sato, H. Yamaga, S. Kashima, Y. Murata, T. Shinada, C. Nakano and T. Hoshino, *ChemBioChem*, 2013, 14, 822– 825.
- 403 F. Lopez-Gallego, S. A. Agger, D. Abate-Pella, M. D. Distefano and C. Schmidt-Dannert, *ChemBioChem*, 2010, 11, 1093–1106.
- 404 A. J. Paula, G. Hwang and H. Koo, *Nat. Commun.*, 2020, **11**, 1354.
- 405 L. Hall-Stoodley, J. W. Costerton and P. Stoodley, *Nat. Rev. Microbiol.*, 2004, **2**, 95–108.
- 406 M. Winn, E. Casey, O. Habimana and C. D. Murphy, *FEMS Microbiol. Lett.*, 2014, **352**, 157–164.
- 407 S. Schulz, M. Girhard and V. B. Urlacher, *ChemCatChem*, 2012, 4, 1889–1895.
- 408 K. G. Domínguez-González, J. J. Robledo-Medrano,
 - J. J. Valdez-Alarcón, O. Hernández-Cristobal,

H. E. Martínez-Flores, J. F. Cerna-Cortés, M. G. Garnica-Romo and R. Cortés-Martínez, *Appl. Sci.*, 2022, **12**, 11425.

- 409 H.-S. Kang, Z. Charlop-Powers and S. F. Brady, *ACS Synth. Biol.*, 2016, 5, 1002–1010.
- 410 S. H. Kim, W. Lu, M. K. Ahmadi, D. Montiel, M. A. Ternei and S. F. Brady, *ACS Synth. Biol.*, 2019, **8**, 109–118.
- 411 G. Bian, Y. Han, A. Hou, Y. Yuan, X. Liu, Z. Deng and T. Liu, *Metab. Eng.*, 2017, **42**, 1–8.
- 412 B. Gu, B. Goldfuss and J. S. Dickschat, *Angew. Chem., Int. Ed.*, 2023, **62**, e202215688.
- 413 S. Mo, A. Krunic, S. D. Pegan, S. G. Franzblau and J. Orjala, *J. Nat. Prod.*, 2009, **72**, 2043–2045.
- 414 A. H. Cabanillas, V. Tena Perez, S. Maderuelo Corral, D. F. Rosero Valencia, A. Martel Quintana, M. Ortega Domenech and A. Rumbero Sanchez, *J. Nat. Prod.*, 2018, 81, 410–413.
- 415 S. J. Rowland and J. N. Robson, *Mar. Environ. Res.*, 1990, **30**, 191–216.
- 416 W. G. Allard, S. T. Belt, G. Massé, R. Naumann, J. M. Robert and S. Rowland, *Phytochemistry*, 2001, **56**, 795–800.
- 417 S. J. Rowland, S. T. Belt, E. J. Wraige, G. Massé, C. Roussakis and J. M. Robert, *Phytochemistry*, 2001, **56**, 597–602.
- 418 S. T. Belt, G. Massé, W. G. Allard, J.-M. Robert and S. J. Rowland, *Tetrahedron Lett.*, 2003, **44**, 9103–9106.
- 419 M. Kaneda, Y. Iitaka and S. Shibata, *Acta Crystallogr.*, 1974, **B30**, 358–364.
- 420 P. S. Rao, K. G. Sarma and T. R. Seshadi, *Curr. Sci.*, 1966, 6, 147–148.
- 421 H. Sugawara, A. Kasuya, Y. Iitaka and S. Shibata, *Chem. Pharm. Bull.*, 1991, **39**, 3051–3054.
- 422 M. Millot, M. Martin-de-Lassalle, M. Chollet-Krugler, Y. Champavier, L. Mambu, J.-A. Chulia and M.-A. Lacaille-Dubois, *Helv. Chim. Acta*, 2016, **99**, 169–173.
- 423 D. Li, M. Yang, R. Mu, S. Luo, Y. Chen, W. Li, A. Wang, K. Guo, Y. Liu and S. Li, *Chin. Chem. Lett.*, 2023, 34, 107469.
- 424 D.-L. Guo, M. Zhao, S.-J. Xiao, B. Xia, B. Wan, Y.-C. Gu, L.-S. Ding and Y. Zhou, *Phytochem. Lett.*, 2015, **14**, 260–264.
- 425 Y. Gao, F. Stuhldreier, L. Schmitt, S. Wesselborg, L. Wang,
 W. E. G. Müller, R. Kalscheuer, Z. Guo, K. Zou, Z. Liu and
 P. Proksch, *Fitoterapia*, 2020, 146, 104652.
- 426 D. Zhang, S. Fukuzawa, M. Satake, X. Li, T. Kuranaga, A. Niitsu, A. Niitsu, K. Yoshizawa and K. Tachibana, *Nat. Prod. Commun.*, 2012, 7, 1411–1414.
- 427 M. K. Renner, P. R. Jensen and W. Fenical, *J. Org. Chem.*, 2000, **65**, 4843–4852.
- 428 G. Randazzo, V. Fogliano, A. Ritieni, L. Mannina, E. Rossi,
 A. Scarallo and A. L. Segre, *Tetrahedron*, 1993, 49, 10883– 10896.
- 429 A. Ritieni, V. Fogliano, G. Randazzo, A. Scarallo,
 A. Logrieco, A. Moretti, L. Manndina and A. Bottalico, *Nat. Toxins*, 1995, 3, 17–20.
- 430 A. Santini, A. Ritieni, V. Fogliano, G. Randazzo, L. Mannina,
 A. Logrieco and E. Benedetti, *J. Nat. Prod.*, 1996, 59, 109–112.
- 431 D. Liu, X.-M. Li, C.-S. Li and B.-G. Wang, *Helv. Chim. Acta*, 2013, **96**, 437–444.

- 432 N. Hoque, C. M. Hasan, M. S. Rana, A. Varsha, M. H. Sohrab and K. M. Rahman, *Molecules*, 2018, 23(3288), 3281–3288.
- 433 X. Huang, H. Huang, H. Li, X. Sun, H. Huang, Y. Lu, Y. Lin, Y. Long and Z. She, *Org. Lett.*, 2013, **15**, 721–723.
- 434 Z. e. Xiao, H. Huang, C. Shao, X. Xia, L. Ma, X. Huang, Y. Lu, Y. Lin, Y. Long and Z. She, *Org. Lett.*, 2013, **15**, 2522–2525.
- 435 Z. Liu, Y. Chen, S. Chen, Y. Liu, Y. Lu, D. Chen, Y. Lin, X. Huang and Z. She, *Org. Lett.*, 2016, **18**, 1406–1409.
- 436 T. Shiina, K. Nakagawa, Y. Fujisaki, T. Ozaki, C. Liu, T. Toyomasu, M. Hashimoto, H. Koshino, A. Minami, H. Kawaide and H. Oikawa, *Biosci., Biotechnol., Biochem.*, 2019, 83, 192–201.
- 437 T. Roncal, S. Cordobes, O. Sterner and U. Ugalde, *Eukaryot. Cell*, 2002, **1**, 823–829.
- 438 S. Kusari and M. Spiteller, *Nat. Prod. Rep.*, 2011, **28**, 1203–1207.
- 439 C. A. Young, S. Felitti, K. Shields, G. Spangenberg, R. D. Johnson, G. T. Bryan, S. Saikia and B. Scott, *Fungal Genet. Biol.*, 2006, 43, 679–693.
- 440 L. Keller and M. G. Surette, *Nat. Rev. Microbiol.*, 2006, 4, 249–258.
- 441 V. Schroeckh, K. Scherlach, H. W. Nutzmann, E. Shelest, W. Schmidt-Heck, J. Schuemann, K. Martin, C. Hertweck and A. A. Brakhage, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, 106, 14558–14563.
- 442 T. T. Shen, X. H. Mo, L. P. Zhu, L. L. Tan, F. Y. Du, Q. W. Wang, Y. M. Zhou, X. J. Yuan, B. Qiao and S. Yanga, *Appl. Environ. Microbiol.*, 2019, **85**, e00293– e00219.
- 443 S. Bertrand, O. Schumpp, N. Bohni, M. Monod, K. Gindro and J.-L. Wolfender, *J. Nat. Prod.*, 2013, **76**, 1157–1165.
- 444 R. S. T. Kamdem, H. Wang, P. Wafo, W. Ebrahim,
 F. C. Özkaya, G. Makhloufi, C. Janiak,
 P. Sureechatchaiyan, M. U. Kassack, W. Lin, Z. Liu and
 P. Proksch, *Fitoterapia*, 2018, **124**, 132–136.
- 445 S. Bertrand, N. Bohni, S. Schnee, O. Schumpp, K. Gindro and J.-L. Wolfender, *Biotechnol. Adv.*, 2014, **32**, 1180–1204.
- 446 E. Cohen, L. Koch, K. M. Thu, Y. Rahamim, Y. Aluma, M. Ilan, O. Yarden and S. Carmeli, *Bioorg. Med. Chem.*, 2011, **19**, 6587–6593.
- 447 J. P. Engelberts, S. J. Robbins, J. M. de Goeij, M. Aranda, S. C. Bell and N. S. Webster, *ISME J.*, 2020, **14**, 1100–1110.
- 448 R. W. Thacker and C. J. Freeman, *Adv. Mar. Biol.*, 2012, **62**, 57–111.
- 449 WFO, World flora online, http://www.worldfloraonline.org, (accessed April 03 2024).
- 450 R. Kamaya and H. Ageta, *Chem. Pharm. Bull.*, 1990, **38**, 342–346.
- 451 R. Kamaya, K. Masuda, H. Ageta and H.-C. Chang, *Chem. Pharm. Bull.*, 1996, 44, 695–698.
- 452 R. T. Iyer, K. N. N. Ayengar and S. Rangaswami, *Indian J. Chem.*, 1972, **10**, 482–484.
- 453 H. Khan, A. Zaman, G. L. Chetty, A. S. Gupta and S. Dev, *Tetrahedron Lett.*, 1971, **12**, 4443–4446.
- 454 T. Jiang, X. Dai, T. Gao, L. Wang, F. Yang, Y. Zhang, N. Wang, G. Huang and J. Cao, *Tetrahedron Lett.*, 2022, 100, 153869.

- 455 R. Kamaya, K. Masuda, K. Suzuki, H. Ageta and H.-Y. Hsu, *Chem. Pharm. Bull.*, 1996, **44**, 690–694.
- 456 K. Zaman, D. Chetia and M. Ali, *Asian J. Chem.*, 2017, **29**, 485–488.
- 457 Q.-M. Li, J.-G. Luo, H.-J. Zhao, W.-Y. Yu, X.-B. Wang, M.-H. Yang, J. Luo, H.-B. Sun, Y.-J. Chen, Q.-L. Guo and L.-Y. Kong, *Asian J. Org. Chem.*, 2015, 4, 1366–1369.
- 458 Q.-M. Li, J.-G. Luo, Y.-M. Zhang, Z.-R. Li, X.-B. Wang, M.-H. Yang, J. Luo, H.-B. Sun, Y.-J. Chen and L.-Y. Kong, *Chem.-Eur. J.*, 2015, **21**, 13206–13209.
- 459 D. H. O'Brien and R. D. Stipanovic, *J. Org. Chem.*, 1978, 43, 1105–1111.
- 460 R. D. Stipanovic, A. A. Bell and M. J. Lukefahr, *Phytochemistry*, 1978, **17**, 151–152.
- 461 R. D. Stipanovic, A. A. Bell, D. H. O'Brien and M. J. Lukefahr, *Tetrahedron Lett.*, 1977, 18, 567–570.
- 462 R. D. Stipanovic, A. A. Bell, D. H. O'Brien and M. J. Lukefahr, J. Agric. Food Chem., 1978, 26, 115–118.
- 463 Z. Chen, X. Chen, Y. Tang, Y. Zhou, H. Deng, J. He, Y. Liu,Z. Zhao and H. Cui, *Org. Lett.*, 2022, 24, 3717–3720.
- 464 W. H. Jiao, H. Gao, F. Zhao, F. He, G. X. Zhou and X. S. Yao, *Chem. Biodivers.*, 2011, **8**, 1163–1169.
- 465 H. Hikino, T. Ohta and T. Takemoto, *Phytochemistry*, 1975, 14, 2473–2481.
- 466 S. Kurimoto, J. X. Pu, H. D. Sun, Y. Takaishi and Y. Kashiwada, *Chem. Biodivers.*, 2015, **12**, 1200–1207.
- 467 S.-H. Luo, Q. Luo, X.-M. Niu, M.-J. Xie, X. Zhao,
 B. Schneider, J. Gershenzon and S.-H. Li, *Angew. Chem.*, *Int. Ed.*, 2010, 49, 4471–4475.
- 468 S.-H. Luo, J. Hua, X.-M. Niu, Y. Liu, C.-H. Li, Y.-Y. Zhou, S.-X. Jing, X. Zhao and S.-H. Li, *Phytochemistry*, 2013, 86, 29–35.
- 469 S. H. Luo, C. L. Hugelshofer, J. Hua, S. X. Jing, C. H. Li, Y. Liu, X. N. Li, X. Zhao, T. Magauer and S. H. Li, *Org. Lett.*, 2014, **16**, 6416–6419.
- 470 M. I. Choudhary, R. Ranjit, A.-u. Rahman, S. Hussain,K. P. Devkota, T. M. Shrestha and M. Parvez, *Org. Lett.*, 2004, 6, 4139–4142.
- 471 M. I. Choudhary, R. R. Atta-ur-Rahman, K. P. Devkotaa and T. M. Shrestha, *Z. Naturforsch.*, 2007, **62b**, 587–592.
- 472 M. I. Choudhary, R. Ranjit, A.-u. Rahman, T. M. Shrestha,
 A. Yasin and M. Parvez, *J. Org. Chem.*, 2004, 69, 2906–2909.
- 473 S.-H. Luo, J. Hua, C.-H. Li, Y. Liu, X.-N. Li, X. Zhao and S.-H. Li, *Tetrahedron Lett.*, 2013, 54, 235–237.
- 474 S.-H. Luo, L.-H. Weng, M.-J. Xie, X.-N. Li, J. Hua, X. Zhao and S.-H. Li, *Org. Lett.*, 2011, **13**, 1864–1867.
- 475 S.-H. Luo, J. Hua, C.-H. Li, S.-X. Jing, Y. Liu, X.-N. Li, X. Zhao and S.-H. Li, *Org. Lett.*, 2012, **14**, 5768–5771.
- 476 S.-X. Jing, R. Fu, C.-H. Li, T.-T. Zhou, Y.-C. Liu, Y. Liu, S.-H. Luo, X.-N. Li, F. Zeng and S.-H. Li, *J. Org. Chem.*, 2021, **86**, 11169–11176.
- 477 C.-H. Li, S.-X. Jing, S.-H. Luo, W. Shi, J. Hua, Y. Liu, X.-N. Li,
 B. Schneider, J. Gershenzon and S.-H. Li, *Org. Lett.*, 2013, 15, 1694–1697.
- 478 L.-L. Teng, R.-F. Mu, Y.-C. Liu, C.-J. Xiao, D.-S. Li, J.-X. Gao, K. Guo, X.-N. Li, Y. Liu, F. Zeng and S.-H. Li, *Org. Lett.*, 2021, 23, 2232–2237.

- 479 A. Bisio, A. M. Schito, F. Pedrelli, O. Danton, J. K. Reinhardt, G. Poli, T. Tuccinardi, T. Burgi, F. De Riccardis, M. Giacomini, D. Calzia, I. Panfoli, G. C. Schito, M. Hamburger and N. De Tommasi, *J. Nat. Prod.*, 2020, 83, 1027–1042.
- 480 M. R. Hasan, H. I. Al-Jaber, M. A. Al-Qudah and M. H. Abu Zarga, *Phytochem. Lett.*, 2016, **16**, 12–17.
- 481 F. Dal Piaz, A. Vassallo, L. Lepore, A. Tosco, A. Bader and N. De Tommasi, *J. Med. Chem.*, 2009, **52**, 3814–3828.
- 482 A. Rustaiyan and A. Sadjadi, *Phytochemistry*, 1987, **26**, 3078–3079.
- 483 A. Rustaiyan, A. Niknejad, L. Nazarians, J. Jakupovic and F. Bohlmann, *Phytochemistry*, 1982, 21, 1812–1813.
- 484 A. Rustaiyan and S. Koussari, *Phytochemistry*, 1988, 27, 1767–1769.
- 485 G. Cioffi, A. Bader, A. Malafronte, F. Dal Piaz and N. De Tommasi, *Phytochemistry*, 2008, **69**, 1005–1012.
- 486 F. M. Moghaddam, R. Amiri, M. Alam, M. B. Hossain and D. van der Helm, *J. Nat. Prod.*, 1998, **61**, 279–281.
- 487 F. M. Moghaddam, M. M. Farimani, M. Seirafi, S. Taheri, H. R. Khavasi, J. Sendker, P. Proksch, V. Wray and R. Edrada, *J. Nat. Prod.*, 2010, 73, 1601–1605.
- 488 M. Moridi Farimani and Z. Mazarei, *Fitoterapia*, 2014, **98**, 234–240.
- 489 S. N. Ebrahimi, M. M. Farimani, F. Mirzania, M. A. Soltanipoor, M. De Mieri and M. Hamburger, *J. Nat. Prod.*, 2014, 77, 848–854.
- 490 A. Ulubelen, G. Topcu, U. Sönmez, C. Eriş and U. Özgen, *Phytochemistry*, 1996, **43**, 431–434.
- 491 G. Topcu, A. Ulubelen, T. C. M. Tam and T.-C. Chun, *Phytochemistry*, 1996, **42**, 1089–1092.
- 492 M. S. Gonzalez, J. M. San Segundo, M. C. Grande, M. Medarde and I. S. Bellido, *Tetrahedron*, 1989, 45, 3575–3582.
- 493 A. Malafronte, F. Dal Piaz, G. Cioffi, A. Braca, A. Leone and N. De Tommasi, *Nat. Prod. Commun.*, 2008, **3**, 877–880.
- 494 G. Topcu, A. Ulubelen, T. C.-M. Tam and C.-T. Che, *J. Nat. Prod.*, 1996, **59**, 113–116.
- 495 G. Appendino, O. Taglialatela-Scafati, A. Romano,
 F. Pollastro, C. Avonto and P. Rubiolo, *J. Nat. Prod.*, 2009,
 72, 340–344.
- 496 S. Wang, J. Li, J. Sun, K. W. Zeng, J. R. Cui, Y. Jiang and P. F. Tu, *Fitoterapia*, 2013, **85**, 169–175.
- 497 S. J. Bloor, Tetrahedron Lett., 1993, 34, 5617-5620.
- 498 L. O. Hanus, T. Rezanka and V. M. Dembitsky, *Phytochemistry*, 2003, **63**, 869–875.
- 499 M. Ali and J. Gupta, *J. Med. Aromat. Plant Sci.*, 1996, **18**, 791–794.
- 500 N. Tanaka, S. Abe and J. i. Kobayashi, *Tetrahedron Lett.*, 2012, **53**, 1507–1510.
- 501 C. Yan, Y. Wang and X. J. Hao, *Chin. Chem. Lett.*, 2008, **19**, 937–939.
- 502 V. L. Challinor, S. Chap, R. P. Lehmann, P. V. Bernhardt and J. J. De Voss, *J. Nat. Prod.*, 2013, **76**, 485–488.
- 503 V. L. Challinor, R. C. Johnston, P. V. Bernhardt, R. P. Lehmann, E. H. Krenske and J. J. De Voss, *Chem. Sci.*, 2015, 6, 5740–5745.

- 504 K. A. Jabal, H. M. Abdallah, G. A. Mohamed, I. A. Shehata, M. Y. Alfaifi, S. E. I. Elbehairi, A. A. Koshak and S. R. M. Ibrahim, *Nat. Prod. Res.*, 2019, 1–6, DOI: 10.1080/ 14786419.2019.1577842.
- 505 S. K. Roy, M. Sanjrani, M. Sharma and R. Ramachandram, Indian J. Chem., Sect. B:Org. Chem. Incl. Med. Chem., 2002, 41, 2390–2394.
- 506 N. Kawahara, M. Nozawa, D. Flores, P. Bonilla, S. Sekita, M. Satake and K. I. Kawai, *Chem. Pharm. Bull.*, 1997, 45, 1717–1719.
- 507 N. Kawahara, M. Nozawa, D. Flores, P. Bonilla, S. Sekita and M. Satake, *Phytochemistry*, 2000, **53**, 881–884.
- 508 K. Guo, X. Liu, T. T. Zhou, Y. C. Liu, Y. Liu, Q. M. Shi, X. N. Li and S. H. Li, *J. Org. Chem.*, 2020, **85**, 5511–5515.
- 509 K. Guo, Y.-C. Liu, Y. Liu, H. Zhang, W.-Y. Li, Q.-M. Shi, X.-N. Li, F. Zeng and S.-H. Li, *Phytochemistry*, 2021, 187, 112780.
- 510 M. Dong, L. Q. Quan, W. F. Dai, S. L. Yan, C. H. Chen, X. Q. Chen and R. T. Li, *Planta Med.*, 2017, 83, 1368–1373.
- 511 L. Acebey, M. Sauvain, S. Beck, C. Moulis, A. Gimenez and V. Jullian, *Org. Lett.*, 2007, **9**, 4693–4696.
- 512 C. Murakami, R. S. A. Cabral, K. S. Gomes, T. A. Costa-Silva, M. Amaral, M. Romanelli, A. G. Tempone, J. H. G. Lago, V. da S. Bolzani, P. R. H. Moreno and M. C. M. Young, *Phytochem. Lett.*, 2019, 33, 6–11.
- 513 T. Akihisa, K. Koike, Y. Kimura, N. Sashida, T. Matsumoto,M. Ukiya and T. Nikaido, *Lipids*, 1999, 34, 1151–1157.
- 514 C. A. N. Catalán, C. S. de Heluani, C. Kotowicz, T. E. Gedris and W. Herz, *Phytochemistry*, 2003, **64**, 625–629.
- 515 M. Toyoda, M. Asahina and H. Fukawa, *Tetrahedron Lett.*, 1969, 55, 4879–4882.
- 516 C. Corre and G. L. Challis, *Nat. Prod. Rep.*, 2009, **26**, 977–986.
- 517 C. Wang, Q. Chen, D. Fan, J. Li, G. Wang and P. Zhang, *Mol. Plant*, 2016, **9**, 195–204.
- 518 A. Minami, T. Ozaki, C. Liu and H. Oikawa, *Nat. Prod. Rep.*, 2018, **35**, 1330–1346.
- 519 A. C. Huang, Y. J. Hong, A. D. Bond, D. J. Tantillo and A. Osbourn, *Angew. Chem., Int. Ed.*, 2018, **57**, 1291–1295.
- 520 Q. Chen, J. Li, Z. Liu, T. Mitsuhashi, Y. Zhang, H. Liu, Y. Ma, J. He, T. Shinada, T. Sato, Y. Wang, H. Liu, I. Abe, P. Zhang and G. Wang, *Plant Commun.*, 2020, **1**, 100051.
- 521 J. Shao, Q.-W. Chen, H.-J. Lv, J. He, Z.-F. Liu, Y.-N. Lu, H.-L. Liu, G.-D. Wang and Y. Wang, *Org. Lett.*, 2017, **19**, 1816–1819.
- 522 Y.-G. Chen, D.-S. Li, Y. Ling, Y.-C. Liu, Z.-L. Zuo, L.-S. Gan, S.-H. Luo, J. Hua, D.-Y. Chen, F. Xu, M. Li, K. Guo, Y. Liu, J. Gershenzon and S.-H. Li, *Angew. Chem., Int. Ed.*, 2021, 60, 25468–25476.
- 523 D.-S. Li, J. Hua, S.-H. Luo, Y.-C. Liu, Y.-G. Chen, Y. Ling, K. Guo, Y. Liu and S.-H. Li, *Plant Commun.*, 2021, 2, 100233.
- 524 M. Masi, R. Dasari, A. Evidente, V. Mathieu and A. Kornienko, *Bioorg. Med. Chem. Lett.*, 2019, **29**, 859–869.
- 525 H. Fujiwara, K. Matsunaga, H. Kumagai, M. Ishizuka and Y. Ohizumi, *Pharm. Pharmacol. Commun.*, 2000, **6**, 427–431.
- 526 M. Bury, A. Girault, V. Megalizzi, S. Spiegl-Kreinecker, V. Mathieu, W. Berger, A. Evidente, A. Kornienko,

P. Gailly, C. Vandier and R. Kiss, *Cell Death Dis.*, 2013, 4, e561.

- 527 A. K. Najumudeen, A. Jaiswal, B. Lectez, C. Oetken-Lindholm, C. Guzman, E. Siljamaki, I. M. Posada, E. Lacey, T. Aittokallio and D. Abankwa, *Oncogene*, 2016, 35, 5248–5262.
- 528 R. Dasari, M. Masi, R. Lisy, M. Ferderin, L. R. English,
 A. Cimmino, V. Mathieu, A. J. Brenner, J. G. Kuhn,
 S. T. Whitten, A. Evidente, R. Kiss and A. Kornienko, *Bioorg. Med. Chem. Lett.*, 2015, 25, 4544–4548.
- 529 S. S. Elhady, A. M. Al-Abd, A. M. El-Halawany, A. M. Alahdal,
 H. A. Hassanean and S. A. Ahmed, *Mar. Drugs*, 2016, 14(130), 131–114.
- 530 Y. C. Chen, M. C. Lu, M. El-Shazly, K. H. Lai, T. Y. Wu, Y. M. Hsu, Y. L. Lee and Y. C. Liu, *Mar. Drugs*, 2018, 16(212), 211–222.
- 531 W. T. Chang, Y. D. Bow, P. J. Fu, C. Y. Li, C. Y. Wu, Y. H. Chang, Y. N. Teng, R. N. Li, M. C. Lu, Y. C. Liu and C. C. Chiu, Oxid. Med. Cell. Longev., 2021, 2021, 7689045.
- 532 Y. Doi, H. Shigemori, M. Ishibashi, F. Mizobe, A. Kawashima, S. Nakaike and J. Kobayashi, *Chem. Pharm. Bull.*, 1993, **41**, 2190–2191.
- 533 M. Schumacher, C. Cerella, S. Eifes, S. Chateauvieux,
 F. Morceau, M. Jaspars, M. Dicato and M. Diederich, *Biochem. Pharmacol.*, 2010, **79**, 610–622.
- 534 K. H. Lai, Y. C. Liu, J. H. Su, M. El-Shazly, C. F. Wu, Y. C. Du,
 Y. M. Hsu, J. C. Yang, M. K. Weng, C. H. Chou, G. Y. Chen,
 Y. C. Chen and M. C. Lu, *Sci. Rep.*, 2016, 6(36170), 36171–36114.
- 535 Y. Mizushina, D. Manita, T. Takeuchi, F. Sugawara,
 Y. Kumamoto-Yonezawa, Y. Matsui, M. Takemura,
 M. Sasaki, H. Yoshida and H. Takikawa, *Molecules*, 2008,
 14, 102–121.
- 536 L. Margarucci, M. C. Monti, A. Tosco, R. Riccio and A. Casapullo, Angew Chem. Int. Ed. Engl., 2010, 49, 3960– 3963.
- 537 L. Margarucci, A. Tosco, R. De Simone, R. Riccio, M. C. Monti and A. Casapullo, *ChemBioChem*, 2012, 13, 982–986.
- 538 M. C. Monti, L. Margarucci, R. Riccio, L. Bonfili, M. Mozzicafreddo, A. M. Eleuteri and A. Casapullo, *Biochim. Biophys. Acta*, 2014, **1844**, 713–721.
- 539 L. V. Manes, P. Crews, M. R. Kernan, D. J. Faulkner, F. R. Fronczek and R. D. Gandour, *J. Org. Chem.*, 1988, 53, 570–575.
- 540 C. Cassiano, M. C. Monti, C. Festa, A. Zampella, R. Riccio and A. Casapullo, *ChemBioChem*, 2012, **13**, 1953–1958.
- 541 J. C. de Freitas, L. A. Blankemeier and R. S. Jacobs, *Experientia*, 1984, **40**, 864–865.
- 542 C. F. Bennett, S. Mong, M. A. Clarke, L. I. Kruse and S. T. Crooke, *Biochem. Pharmacol.*, 1987, **36**, 733–740.
- 543 K. B. Glaser and R. S. Jacobs, *Biochem. Pharmacol.*, 1986, 35, 449–453.
- 544 K. B. Glaser and R. S. Jacobs, *Biochem. Pharmacol.*, 1987, **36**, 2079–2086.
- 545 K. B. Glaser, T. S. Vedvick and R. S. Jacobs, *Biochem. Pharmacol.*, 1988, **37**, 3639–3646.

- 546 F. Dal Piaz, A. Casapullo, A. Randazzo, R. Riccio, P. Pucci,
 G. Marino and L. Gomez-Paloma, *ChemBioChem*, 2002, 3, 664–671.
- 547 M. C. Monti, A. Casapullo, R. Riccio and L. Gomez-Paloma, *Bioorg. Med. Chem. Lett.*, 2004, **12**, 1467–1474.
- 548 M. C. Monti, A. Casapullo, R. Riccio and L. Gomez-Paloma, *FEBS Lett.*, 2004, **578**, 269–274.
- 549 M. C. Monti, R. Riccio and A. Casapullo, *Bioorg. Chem.*, 2009, **37**, 6–10.
- 550 M. C. Monti, A. Casapullo, C. N. Cavasotto, A. Tosco, F. Dal Piaz, A. Ziemys, L. Margarucci and R. Riccio, *Chemistry*, 2009, **15**, 1155–1163.
- 551 M. H. Uddin, M. Otsuka, T. Muroi, A. Ono, N. Hanif, S. Matsuda, T. Higa and J. Tanaka, *Chem. Pharm. Bull.*, 2009, 57, 885–887.
- 552 F. Folmer, M. Jaspars, M. Schumacher, M. Dicato and M. Diederich, *Biochem. Pharmacol.*, 2010, **80**, 1793–1800.
- 553 K. B. Glaser, M. L. Sung, D. A. Hartman, Y. W. Lock, J. Bauer, T. Walter and R. P. Carlson, *Skin Pharmacol.*, 1995, 8, 300–308.
- 554 B. C. M. Potts, R. J. Capon and D. J. Faulkner, *J. Org. Chem.*, 1992, 57, 2965–2967.
- 555 P. Garcia Pastor, S. De Rosa, A. De Giulio, M. Paya and M. J. Alcaraz, *Br. J. Pharmacol.*, 1999, **126**, 301–311.
- 556 R. P. Murelli, A. K. Cheung and M. L. Snapper, *J. Org. Chem.*, 2007, **72**, 1545–1552.
- 557 V. Escrig, A. Ubeda, M. L. Ferrandiz, J. Darias, J. M. Sanchez, M. J. Alcaraz and M. Paya, *J. Pharmacol. Exp. Ther.*, 1997, 282, 123–131.
- 558 H. Miyaoka, M. Yamanishi and H. Mitome, *Chem. Pharm. Bull.*, 2006, **54**, 268–270.
- 559 M. C. Monti, M. G. Chini, L. Margarucci, R. Riccio, G. Bifulco and A. Casapullo, *ChemBioChem*, 2011, 12, 2686–2691.
- 560 M. S. de Carvalho and R. S. Jacobs, *Biochem. Pharmacol.*, 1991, **42**, 1621–1626.
- 561 J. D. Winkler, C. M. Sung, W. C. Hubbard and F. H. Chilton, Biochem. Pharmacol., 1992, 44, 2055–2066.
- 562 M. S. Barnette, J. Rush, L. A. Marshall, J. J. Foley,
 D. B. Schmidt and H. M. Sarau, *Biochem. Pharmacol.*, 1994, 47, 1661–1667.
- 563 M. C. Monti, A. Casapullo, C. N. Cavasotto, A. Napolitano and R. Riccio, *ChemBioChem*, 2007, **8**, 1585–1591.
- 564 M. C. Monti, A. Casapullo, R. Riccio and L. Gomez-Paloma, Rapid Commun. Mass Spectrom., 2005, 19, 303–308.
- 565 M. L. Bourguet-Kondracki, C. Debitus and M. Guyot, *J. Chem. Res.*, 1996, (S), 192–193.
- 566 H. S. Lee, K. M. Yoon, Y. R. Han, K. J. Lee, S. C. Chung, T. I. Kim, S. H. Lee, J. Shin and K. B. Oh, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 1051–1053.

- 567 E. Dillp de Sllva and P. J. Scheuer, *Tetrahedron Lett.*, 1980, 21, 1611–1614.
- 568 M. E. Skindersoe, P. Ettinger-Epstein, T. B. Rasmussen, T. Bjarnsholt, R. de Nys and M. Givskov, *Mar. Biotechnol.*, 2008, **10**, 56–63.
- 569 L. Du, L. Shen, Z. Yu, J. Chen, Y. Guo, Y. Tang, X. Shen and H. Jiang, *ChemMedChem*, 2008, 3, 173–180.
- 570 C. Cassiano, R. Esposito, A. Tosco, A. Zampella, M. V. D'Auria, R. Riccio, A. Casapullo and M. C. Monti, *Chem. Commun.*, 2014, **50**, 406–408.
- 571 D. Xue, Q. Wang, Z. Chen, L. Cai, L. Bao, Q. Qi, L. Liu, X. Wang, H. Jin, J. Wang, H. Wu, H. Liu and Q. Chen, *Bioorg. Med. Chem. Lett.*, 2015, 25, 1464–1470.
- 572 C. C. Chen, T. T. Tzeng, C. C. Chen, C. L. Ni, L. Y. Lee, W. P. Chen, Y. J. Shiao and C. C. Shen, *J. Nat. Prod.*, 2016, 79, 438–441.
- 573 J. H. Hu, I. C. Li, T. W. Lin, W. P. Chen, L. Y. Lee, C. C. Chen and C. F. Kuo, *Molecules*, 2019, 24(1624), 1621–1613.
- 574 E. D'Aniello, F. A. Iannotti, L. G. Falkenberg, A. Martella,
 A. Gentile, F. De Maio, M. L. Ciavatta, M. Gavagnin,
 J. S. Waxman, V. Di Marzo, P. Amodeo and R. M. Vitale, *Mar. Drugs*, 2019, 17, 110.
- 575 Y. Wang, M. Dreyfuss, M. Ponelle, L. LukasOberer and H. Riezman, *Tetrahedron*, 1998, 54, 6415–6426.
- 576 A. Bidon-Chanal, A. Fuertes, D. Alonso, D. I. Perez,
 A. Martinez, F. J. Luque and M. Medina, *Eur. J. Med. Chem.*, 2013, 60, 479–489.
- 577 J.-M. Kim, S.-B. Hyeon, A. Isogai and A. Suzuki, Agric. Biol. Chem., 1984, 48, 803–805.
- 578 A. Ballio, Experientia, 1991, 47(8), 783-790.
- 579 G. Carr, M. Raszek, R. Van Soest, T. Matainaho, M. Shopik, C. F. B. Holmes and R. J. Andersen, *J. Nat. Prod.*, 2007, 70, 1812–1815.
- 580 Y. Ling, L.-L. Teng, J. Hua, D.-S. Li, S.-H. Luo, Y.-C. Liu, Y. Liu and S.-H. Li, *Chin. J. Nat. Med.*, 2019, **17**, 892–899.
- 581 K. Guo, L. Feng, H. Zhang, T.-T. Zhou, Y.-Y. Chen, C.-L. Tan, Y.-C. Liu, Y. Liu and S.-H. Li, *Fitoterapia*, 2024, **178**, 106158.
- 582 K. Guo, T.-T. Zhou, S.-H. Luo, Y.-C. Liu, Y. Liu and S.-H. Li, J. Med. Chem., 2024, 67, 513–528.
- 583 M. Li, L. Feng, H. Zhang, Y.-C. Liu, T.-T. Zhou, Y. Zheng, K. Guo, Y. Liu and S.-H. Li, *Org. Biomol. Chem.*, 2024, 22, 3019–3024.
- 584 Y. Zheng, Q. Li, M. Gu, H. Liao, Y. Liang, F. Liu, X.-N. Li, W. Sun, C. Chen, Y. Zhang and H. Zhu, *J. Nat. Prod.*, 2024, 87, 1965–1974.
- 585 M. R. Byun, A. R. Kim, J. H. Hwang, M. K. Sung, Y. K. Lee, B. S. Hwang, J. R. Rho, E. S. Hwang and J. H. Hong, *FEBS Lett.*, 2012, **586**, 1086–1092.