

including benzimidazole–naphthalene conjugate molecules,²⁹ silylated derivatives,³⁰ and 2-phenyl-2H-1,2,3-triazole derivatives.³¹ Besides, some Schiff base compounds can detect F⁻/CN⁻.^{32,33} However, sensors that can simultaneously recognize Al³⁺ and F⁻/CN⁻ are still relatively uncommon.^{34,35}

Therefore, we designed and synthesized a new Schiff base chemosensor bearing a julolidine unit; the sensor was easy to synthesize and could identify both Al³⁺ and F⁻/CN⁻. Detecting multiple targets with a single receptor is more efficient and less expensive than one-to-one analysis.³⁶ The julolidine group is an excellent fluorophore,³⁷ and the Schiff base compounds contain a benzhydrazone unit with donor sites such as N atom and O atom for Al³⁺.³⁸ Besides, benzhydrazone units contain an amide functional group, and julolidine possesses a phenolic hydroxyl group; the –NH proton and the phenolic hydroxyl group could be linked with F⁻/CN⁻ *via* a hydrogen bond to detect F⁻/CN⁻.^{39,40} The synthetic route of sensor HL was shown in Fig. 1. The HL molecule could be easily used as a fluorescent sensor to detect Al³⁺ and F⁻/CN⁻ and as a colorimetric sensor in acetonitrile to recognize F⁻/CN⁻.

2. Experimental

2.1. General methods

All the solvents were of analytical grade and were not purified before use. Other reagents were of spectroscopic grade. Metal ion solutions were obtained by dissolving the respective nitrates (0.1 M) in distilled water (2.0 mL) except for Hg²⁺ (its counter ion was chloride). All anion solutions were prepared from the dissolution of the corresponding potassium or sodium salts (0.1 M) in distilled water (2.0 mL). NMR spectra were recorded in a Bruker AV400 (400 MHz) spectrometer with DMSO-*d*₆ as solvent and tetramethylsilane (TMS) as internal standard. UV-vis spectra were measured on an Agilent 8453 UV-vis spectrophotometer. Fluorescence spectra were recorded with a Hitachi F-4600 fluorescence spectrophotometer. Fluorescence quantum yield was measured with an Absolute PL Quantum Yield Spectrometer QY C11347-11. Melting points were obtained on a WRS-1B melting point apparatus. High resolution mass spectra were recorded on an AB SCIEX Triple TOF 4600 instrument. Elemental analysis was performed with a PE CHN 2400 analyzer. Fluorescent cell imaging was obtained on an Olympus FV1000 confocal laser scanning microscope.

2.2. Synthesis of HL

8-Hydroxyjulolidine-9-carboxaldehyde (0.11 g, 0.5 mmol) and 2-thiophenecarboxylic hydrazide (0.07 g, 0.5 mmol) were added to ethanol (10 mL) and stirred for 10 min at room temperature. The mixture was heated to reflux for 24 h and then cooled to




Fig. 1 Synthetic route of HL.

room temperature; the light yellow precipitate was washed three times with cold ethanol and dried under vacuum to give product HL (0.08 g). Yield 47.06%. Mp 492–493 K. ¹H NMR (DMSO-*d*₆, 400 MHz), δ (ppm): 11.80 (s, 1H), 11.63 (s, 1H), 8.27 (s, 1H), 7.84–7.83 (d, 2H), 7.21–7.19 (t, 1H), 6.72 (s, 1H), 3.14–3.18 (m, 4H), 2.61–2.49 (m, 4H), 1.84 (s, 4H). ¹³C NMR (DMSO-*d*₆, 100 MHz): 157.54, 155.27, 151.43, 145.94, 138.63, 132.15, 129.27, 128.99, 128.81, 113.14, 106.96, 106.41, 49.98, 49.52, 27.20, 22.16, 21.34, 20.86. Anal. calcd for C₁₈H₁₉N₃O₂S (%): C, 63.32; H, 5.61; N, 12.31. Found: C, 62.24; H, 5.63; N, 12.34. HRMS: *m/z* = 364.1096 [M + Na]⁺ (calcd 364.1095). (Fig. S1–S3†).

2.3. Cytotoxicity assay

The HeLa cells were added to a 96-well plate. About 4000 cells per well were placed in a carbon dioxide cell incubator in an atmosphere of 5% CO₂ and 95% air at 37 °C. Two mg HL was dissolved in DMSO (40 μ L) and diluted with DMEM to obtain 5 different concentrations (100, 33.3, 11.1, 3.7, 1.2 μ g mL⁻¹). Different HL solutions were added to the 96-well plate in a low to high concentration sequence except for that in the control well. Cells were incubated for 24 h to test cytotoxicity. After 24 hours incubation, 20 μ L [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] (MTT) (5 mg mL⁻¹) was added to each well with continuous incubation for another 3.5 h. After MTT was removed, DMSO was added until the compounds were completely dissolved. The absorbance values were measured using a microplate reader. The absorbance of the solvent control cells was considered to be 100%; cell viability and the absorbance of the treated cells were used to calculate the cell viability at different concentrations of HL solution.

2.4. Cell culture and imaging

The human cervical HeLa cancer cells were incubated in DMEM (Dulbecco's modified Eagle's medium) supplemented with 10% FBS (fetal bovine serum) and 1% penicillin–streptomycin in the atmosphere of 5% CO₂ and 95% air at 37 °C. The cells were plated in a cell culture dish overnight. Before the experiments, the cells were washed with phosphate-buffered saline (PBS) buffer. Then,




Fig. 2 The changes in absorption spectra and color of HL induced by Al³⁺ in acetonitrile (2.0 $\times 10^{-5}$ mol L⁻¹).






Fig. 3 Fluorescence spectral changes of HL induced by Al^{3+} in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$).

the cells were incubated with **HL** ($20 \mu\text{M}$) dissolved in DMEM medium for 30 min at 37°C . Next, the cells were observed using an Olympus FV1000 confocal laser scanning microscope. Finally, fluorescence images were obtained after incubating the cells with Al^{3+} ($50 \mu\text{M}$) for another 30 min at 37°C .

3. Results and discussion

3.1. Absorption and fluorescence spectral responses of HL toward Al^{3+}

The detection characteristics of the sensor **HL** for Al^{3+} were preliminarily tested in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$) by a UV-vis spectrometer. Upon addition of Al^{3+} into the solution of **HL**, a new absorption band appeared, and it was centered at 420 nm and 440 nm. As the concentration of Al^{3+} increased, the absorbance at 420 nm and 440 nm gradually increased, whereas the absorption peak at 384 nm declined slowly. The color of the solution changed from colorless to kelly green (Fig. 2). The curves of absorbance that depended on the equivalents of Al^{3+} increased and then remained stable until the amounts of Al^{3+} reached 7.0 equivalents (Fig. S4[†]). At the same time, there







Fig. 4 The limit of detection (LOD); LOD is 20.5 nM.



Scheme 1 The proposed binding mode of sensor HL with Al^{3+} .

appeared a clear isosbestic point at 401 nm, indicating the formation of **HL**- Al^{3+} complex.

To further research the interaction between **HL** and Al^{3+} , fluorescence response experiments were conducted in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$). As shown in Fig. 3, **HL** exhibited weak fluorescence emission intensity ($\Phi = 0.008$) at 420 nm excitation due to C=N isomerization and photoinduced electron transfer (PET) processes.^{41,42} The fluorescence emission intensity of **HL** at 521 nm enhanced gradually along with the increase in Al^{3+} concentration until the amount of Al^{3+} reached 5.0 equivalents with a high quantum yield ($\Phi = 0.464$). Meanwhile, the



Blank Al^{3+} Cu^{2+} Co^{2+} Mn^{2+} Pb^{2+} Cd^{2+} Ca^{2+} K^{+} Fe^{3+} Mg^{2+} Sr^{2+} Ba^{2+} Cr^{3+} Zn^{2+} Ag^{+} Ni^{2+}




Fig. 5 (a) Fluorescence emission spectral changes of HL induced by various metal ions (5.0 equivalents) in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$) by excitation at 420 nm; (b) photos of fluorescence changes in acetonitrile; (c) the fluorescence intensity at 521 nm of HL upon addition of various metal ions (black bars: HL with metal ions; red bars: HL with other metal ions and Al^{3+}) in acetonitrile.






Fig. 6 (a) Absorption spectral change of HL induced by various metal ions (5.0 equiv.) in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$); (b) photos of color changes in acetonitrile; (c) absorption spectra and color of HL induced by F⁻ in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$).





Fig. 7 (a) The change in fluorescence spectra of HL induced by various anions (5.0 equiv.) in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$) (excited at 420 nm); (b) photos of color changes in acetonitrile.

fluorescence color of HL changed from very faint orange to strong cyan. Compared with the result of HL, the fluorescence intensity of complex HL-Al³⁺ was dramatically enhanced. The phenomenon of fluorescence enhancement was ascribed to the inhibition of C=N isomerization and PET processes.

To understand the binding characteristics of HL to Al³⁺, the stoichiometry of HL to Al³⁺ was confirmed based on the Job's plot of the fluorescence spectrum ($\lambda_{\text{em}} = 521 \text{ nm}$). When the molar fraction of $(\text{Al}^{3+})/[(\text{HL}-\text{Al}^{3+})]$ was 0.3, the fluorescence emission intensity value reached its maximum (Fig. S5[†]); this showed that a 2 : 1 HL-Al³⁺ complex was generated. The association constant (K_{a}) of HL-Al³⁺ was calculated to be $6.35 \times 10^4 \text{ M}^{-1}$ ($R = 0.996$) (Fig. S6[†]) from the Benesi-Hildebrand equation of the fluorescence titration data.⁴³ In addition, the limit of detection (LOD) was measured to be 20.5 nM according to the equation $\text{LOD} = 3\sigma/s$, where σ is the standard deviation of the blank signal, and s is the slope of the linear calibration plot (Fig. 4). Compared with the detection limit of Al³⁺ Schiff base sensors reported in the literatures,⁴⁴⁻⁴⁹ the value was relatively low, as shown in Table S1.[†] These results indicated that HL could be used as a highly sensitive fluorescent sensor to detect Al³⁺. Moreover, high-resolution mass spectrometry analysis of HL in the presence of Al³⁺ was performed; the HRMS peak at $m/z = 705.1917$ of $[\text{2HL} + \text{Al}^{3+} - 4\text{H}^+]^-$ (Fig. S7[†]) confirmed the 2 : 1 stoichiometry between HL and Al³⁺ in acetonitrile. Thus, the proposed sensing mechanism for the detection of Al³⁺ by HL is the chelation of Al³⁺ with the O atom of Ar-OH, the N atom from the -CH=N- group and the O atom of the -C=O moiety, leading to hindering of PET processes. In addition, the isomerization of the C=N bond was suppressed, which stabilized the chelate complexation of HL with Al³⁺ and increased the rigidity of the molecule, thereby causing a chelation-enhanced fluorescence (CHEF) effect.⁵⁰⁻⁵² (Scheme 1).

The fluorescence spectra of HL in the presence of diverse metal ions (5.0 equiv.) were obtained in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$), and the ions included Cu²⁺, Zn²⁺, Cr³⁺, Al³⁺, Co²⁺, Mn²⁺, Pb²⁺, Cd²⁺, Ca²⁺, K⁺, Fe³⁺, Mg²⁺, Sr²⁺, Ba²⁺, Ag⁺ and Ni²⁺ (Fig. 5). After the addition of equal amounts of metal ions, the fluorescence emission intensity of HL at 521 nm increased




Fig. 8 The change in fluorescence emission intensity of HL induced by F⁻ in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$).






Fig. 9 ^1H NMR spectral changes of HL induced by F^- in $\text{DMSO}-d_6$.


sharply only when Al^{3+} was added to the solution at 420 nm excitation (Fig. 5a), and the fluorescence color of the solution changed from faint orange to strong cyan; however, its fluorescence intensity and the color of the solution hardly changed in the presence of other metal ions (Fig. 5b). Subsequently, a competitive experiment was carried out in the presence of the above mentioned metal ions (5.0 equiv.) (Fig. 5c). All these fluorescence emission intensities were similar to that of Al^{3+} except for the result of Cu^{2+} , which could be due to the paramagnetic effect on account of spin-orbit coupling. These results showed that **HL** had excellent selectivity for Al^{3+} in acetonitrile.

3.2. Absorption and fluorescence spectral responses of HL to F^- and CN^-

The selective experiment of **HL** to anions was investigated first by UV-vis absorption spectra. Miscellaneous anions including F^- , CN^- , Cl^- , Br^- , I^- , HSO_4^- , NO_3^- , SCN^- , SO_4^{2-} , AcO^- , HSO_3^{2-} , NO_2^- , HCO_3^- , CO_3^{2-} and SO_3^{2-} were analyzed individually with the sensor **HL** in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$) at room temperature (Fig. 6). Upon addition of 5.0 equivalents of other anions to **HL**, no clear changes in UV-vis absorption spectra were observed; only the addition of the F^-/CN^- solution of **HL** resulted in the appearance of two new absorption peaks centered at 439 nm and 467 nm (Fig. 6a). At the same time, the color of the **HL** solution changed from colorless to visible light yellow (Fig. 6b). Other anions caused inappreciable changes of

the absorption spectra, and the color of the solution of **HL** did not change. The absorption spectral change of **HL** induced by F^- in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$) is shown in Fig. 6c. When the amount of F^- increased from 0 to 5.0 equivalents, the absorption peaks at 439 nm and 467 nm gradually increased and reached a maximum at 5.0 equivalents of F^- , whereas the absorption peak at 382 nm decreased when the color of the solution changed to light yellow. Moreover, an isosbestic point was observed at 400 nm, indicating that **HL-F}^-** complex was generated. The changes in absorption spectra could be due to intramolecular charge transfer (ICT) mechanism *via* the deprotonation effect. The increased negative charge density of the phenol oxygen atom accelerated ICT from O atom (Ar-OH) to aromatic rings.^{53,54}

Similarly, CN^- was added to the solution of **HL**; the absorption spectra presented analogous changes to that observed with the addition of F^- (Fig. S8a†). The corresponding sensing mechanism was the same as that for F^- .^{55,56} The results



Scheme 2 Proposed sensing mechanism of F^-/CN^- by **HL**.




Fig. 10 Cell viability of HeLa cells incubated with various concentrations of **HL** for 24 h.






Fig. 11 Fluorescence images of HeLa cells. (a) Microscopy image of HeLa cells treated with HL (20 μM); (b) microscopy image of HeLa cells treated with HL (20 μM) in the presence of Al^{3+} (50 μM); (c) merge of (a) and (b).

clearly demonstrated the selective characteristic of the **HL** sensor for the rapid detection of F^- and CN^- , which could be observed by the naked eye.

In addition to the fluorescence selectivity of **HL** ($2.0 \times 10^{-5} \text{ mol L}^{-1}$) mentioned above, various anions were also explored in acetonitrile, as depicted in Fig. 7. After addition of F^- (5.0 equiv.) to **HL**, the fluorescence emission intensity enhanced nearly 29-folds at 540 nm ($\lambda = 420 \text{ nm}$). On the contrary, there was almost no clear fluorescence signal when other anions were added (Fig. 7a). Meanwhile, the fluorescence color of the solution of **HL-F}^- changed from faint orange to strong green (Fig. 7b). The experiment results indicated that **HL** could effectively distinguish F^- by fluorescence methods. Furthermore, the fluorescent titration interaction between **HL** and F^- was studied in acetonitrile ($2.0 \times 10^{-5} \text{ mol L}^{-1}$), as shown in Fig. 8. On addition of various equivalents of F^- (0–5.0 equiv.), the fluorescence emission intensity at 540 nm increased significantly and reached the maximum ($\Phi = 0.021$) when the amount of F^- reached 5.0 equivalents. Upon continuous addition of F^- to the **HL** solution, the fluorescence spectra hardly changed. The fluorescence spectra of **HL-CN}^- were similar to those of **HL-F}^- (Fig. S8b[†]). Based on the above results, we inferred that **HL** can be used as an efficient F^-/CN^- selective fluorescent sensor.******

The association constants (K_a) for **HL-F}^- and **HL-CN}^- were obtained as $1.25 \times 10^4 \text{ M}^{-1}$ ($R = 0.995$) (Fig. S9[†]) and $1.48 \times 10^4 \text{ M}^{-1}$ ($R = 0.995$) (Fig. S11[†]), respectively, from the fluorescence titration data. The detection limits of **HL** for F^- and CN^- were calculated to be 88.4 nM (Fig. S10[†]) and 61.0 nM (Fig. S12[†]), respectively.****

^1H NMR titrations were performed in $\text{DMSO-}d_6$ to prove the reaction between **HL** and F^- . When excess F^- was added to a solution of **HL**, the signal for $-\text{OH}$ (11.80 ppm) completely disappeared. The proton H2 ($-\text{N-H}$) shifted from 11.65 ppm to 11.70 ppm, and the peak intensity decreased (Fig. 9). ^1H NMR titrations of **HL** with CN^- showed similarities to the results shown above (Fig. S13[†]). This clearly indicated that the strong hydrogen bond interaction caused deprotonation. The proposed sensing mechanism of F^-/CN^- by **HL** is shown in Scheme 2.

3.3. Cell cytotoxicity and cell imaging

To measure the cytotoxicity of **HL**, a cell cytotoxicity experiment by the MTT assay on HeLa cells was carried out. As shown in

Fig. 10, the cell viability gradually decreased as the concentration of **HL** increased. The MTT assay showed that the cell viability was basically over 80%. This result demonstrated that the cytotoxicity of **HL** was relatively low for fluorescence imaging in living cells.

Next, to examine the biological applicability of **HL** to detect Al^{3+} in the biological systems, the sensor **HL** was applied for fluorescence imaging experiments of intracellular Al^{3+} in living cells. The fluorescence images were taken by a confocal laser microscope. The HeLa cells were incubated with **HL** (20 μM) for 30 min at 37 $^\circ\text{C}$; almost no fluorescence was observed (Fig. 11a). Nevertheless, the addition of Al^{3+} (50 μM) to the above HeLa cells triggered a striking fluorescence enhancement and resulted in green fluorescence (Fig. 11b). The results demonstrated that the sensor **HL** could play an indispensable role in intracellular fluorescence imaging of Al^{3+} .

4. Conclusion

In summary, we have designed and developed a new multiple-ion-responsive Schiff base chemosensor **HL**, which can simultaneously detect Al^{3+} and F^-/CN^- . **HL** exhibited excellent colorimetric selectivity to F^-/CN^- and highly specific fluorescence responses to Al^{3+} and F^-/CN^- . Upon addition of Al^{3+} or F^-/CN^- , **HL** showed clear fluorescence increase with different emission spectra. The results demonstrated that the compound **HL** could be used as a fluorescent chemosensor to detect Al^{3+} and F^-/CN^- . Besides, it could also serve as a colorimetric chemosensor to recognize F^-/CN^- via a clear color change from colorless to visible light yellow. Furthermore, the relatively low cytotoxicity of **HL** could be successfully used for cell imaging to identify Al^{3+} , indicating its promising application prospects in living cells.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors are grateful for the financial support from the National Natural Science Foundation of China (21363009), the “5511” science and technology innovation talent project of Jiangxi, the key project of Natural Science Foundation of Jiangxi



Province (20171ACB20025), the Science Funds of Natural Science Foundation of Jiangxi Province (20171BAB203011), and the Project of the Science Funds of Jiangxi Education Office (GJJ170659).

References

- 1 M. Iniya, D. Jeyanthi, K. Krishnaveni, A. Mahesh and D. Chellappa, *Spectrochim. Acta, Part A*, 2014, **120**, 40–46.
- 2 Z. Li, J. L. Zhao, Y. T. Wu and L. Mu, *Org. Biomol. Chem.*, 2017, **15**, 8627–8633.
- 3 A. L. Guo, R. T. Zhu, Y. H. Ren, J. L. Dong and L. H. Feng, *Spectrochim. Acta, Part A*, 2016, **153**, 530–534.
- 4 X. L. Yue, Z. Q. Wang, C. R. Li and Z. Y. Yang, *Tetrahedron Lett.*, 2017, **58**, 4532–4537.
- 5 H. Y. Jeong, S. Y. Lee, J. Han, H. Lim and C. Kim, *Tetrahedron*, 2017, **73**, 2690–2697.
- 6 Y. P. Ren, J. Han and Y. Wang, *Luminescence*, 2018, **33**, 15–21.
- 7 L. K. Kumawat, N. Mergu, M. Asif and V. K. Gupta, *Sens. Actuators, B*, 2016, **231**, 847–859.
- 8 A. H. Panhwar, T. G. Kazi, H. I. Afridi, S. A. Arain, M. S. Arain, K. D. Brahman, N. Ullah, J. Ali and S. S. Arain, *Environ. Monit. Assess.*, 2015, **187**, 37.
- 9 G. Zhang, L. Wang, X. Cai, L. Zhang, J. Yu and A. Wang, *Dyes Pigm.*, 2013, **98**, 232–237.
- 10 W. J. Xu, S. J. Liu, X. Y. Zhao, S. Sun, S. Cheng, T. C. Ma, H. B. Sun and W. Huang, *Chem.–Eur. J.*, 2010, **16**, 7125–7133.
- 11 S. D. Liu, L. W. Zhang, P. P. Zhou, Y. Yang and W. S. Wu, *Sens. Actuators, B*, 2018, **255**, 401–407.
- 12 T. Anand, G. Sivaraman, M. Iniya, A. Siva and D. Chellappa, *Anal. Chim. Acta*, 2015, **876**, 1–8.
- 13 World Health Organization, World Health Organization, 2004.
- 14 Y. Sun, Y. Li, X. Ma and L. Duan, *Sens. Actuators, B*, 2016, **224**, 648–653.
- 15 Y. Sun, S. W. Fan, L. Duan and R. F. Li, *Sens. Actuators, B*, 2013, **185**, 638–643.
- 16 J. B. Chao, Z. Q. Li, Y. B. Zhang, F. J. Huo, C. X. Yin and H. B. Tong, *Sens. Actuators, B*, 2016, **228**, 192–199.
- 17 S. Azimi and Z. Es'haghi, *Bull. Environ. Contam. Toxicol.*, 2017, **98**, 830–836.
- 18 B. K. Bansod, T. Kumar, R. Thakur, S. Rana and I. Singh, *Biosens. Bioelectron.*, 2017, **94**, 443–455.
- 19 A. F. Liu, L. Yang, Z. Y. Zhang, Z. L. Zhang and D. M. Xu, *Dyes Pigm.*, 2013, **99**, 472–479.
- 20 C. R. Chen, G. W. Men, W. H. Bu, C. S. Liang, H. C. Sun and S. M. Jiang, *Sens. Actuators, B*, 2015, **220**, 463–471.
- 21 E. T. Feng, R. M. Lu, C. B. Fan, C. H. Zheng and S. Z. Pu, *Tetrahedron Lett.*, 2017, **58**, 1390–1394.
- 22 K. S. Shen, S. S. Mao, X. K. Shi, F. Wang, Y. L. Xu, S. O. Aderinto and H. L. Wu, *Luminescence*, 2018, **33**, 54–63.
- 23 V. K. Gupta, A. K. Singh and L. K. Kumawat, *Sens. Actuators, B*, 2014, **195**, 98–108.
- 24 S. Chemate and N. Sekar, *Sens. Actuators, B*, 2015, **220**, 1196–1204.
- 25 E. Ozcan, S. O. Tümay, H. A. Alidagı, B. Çosut and S. Yesilot, *Dyes Pigm.*, 2016, **132**, 230–236.
- 26 S. L. Gui, Y. Y. Huang, F. Hu, Y. L. Jin, G. X. Zhang, L. S. Yan, D. Q. Zhang and R. Zhao, *Anal. Chem.*, 2015, **87**, 1470–1474.
- 27 J. B. Liu, W. H. Wang, G. D. Li, R. X. Wang, C. H. Leung and D. L. Ma, *ACS Omega*, 2017, **2**, 9150–9155.
- 28 A. Kumar and P. S. Chae, *Anal. Chim. Acta*, 2017, **958**, 38–50.
- 29 Y. C. Wu, J. Y. You, K. Jiang, J. C. Xie, S. L. Li, D. R. Cao and Z. Y. Wang, *Dyes Pigm.*, 2017, **140**, 47–55.
- 30 A. D. S. Schramm, C. R. Nicoletti, R. I. Stock, R. S. Heying, A. J. Bortoluzzi and V. G. Machado, *Sens. Actuators, B*, 2017, **240**, 1036–1048.
- 31 Y. B. Chen, W. Shi, Y. H. Hui, X. H. Sun, L. X. Xu, L. Feng and Z. F. Xie, *Talanta*, 2015, **137**, 38–42.
- 32 S. Sarveswari, A. J. Beneto and A. Siva, *Sens. Actuators, B*, 2017, **245**, 428–434.
- 33 S. K. Padhan, M. B. Podh, P. K. Sahu and S. N. Sahu, *Sens. Actuators, B*, 2018, **255**, 1376–1390.
- 34 W. H. Ding, D. Wang, X. J. Zheng, W. J. Ding, J. Q. Zheng, W. H. Mu, W. Cao and L. P. Jin, *Sens. Actuators, B*, 2015, **209**, 359–367.
- 35 T. G. Jo, J. J. Lee, E. Nam, K. H. Bok, M. H. Lim and C. Kim, *New J. Chem.*, 2016, **40**, 8918–8927.
- 36 S. A. Lee, G. R. You, Y. W. Choi, H. Y. Jo, A. R. Kim, I. Noh, S. J. Kim, Y. Kim and C. Kim, *Dalton Trans.*, 2014, **43**, 6650–6659.
- 37 Y. W. Choi, J. J. Lee, E. Nam, M. H. Lim and C. Kim, *Tetrahedron*, 2016, **72**, 1998–2005.
- 38 J. Zhang, N. Li, F. Dai, Q. Luo and Y. X. Ji, *Sens. Transducers J.*, 2015, **186**, 125–128.
- 39 B. Yu, C. Y. Li, Y. X. Sun, H. R. Jia, J. Q. Guo and J. Li, *Spectrochim. Acta, Part A*, 2017, **184**, 249–254.
- 40 N. Gupta, D. Singhal, A. K. Singh, N. Singh and U. P. Singh, *Spectrochim. Acta, Part A*, 2017, **176**, 38–46.
- 41 Y. X. Ji, C. W. Yu, S. B. Wen and J. Zhang, *Turk. J. Chem.*, 2016, **40**, 625–630.
- 42 H. H. Wang, B. Wang, Z. H. Shi, X. L. Tang, W. Dou, Q. X. Han, Y. G. Zhang and W. S. Liu, *Biosens. Bioelectron.*, 2015, **65**, 91–96.
- 43 H. A. Benesi and J. Hildebrand, *J. Am. Chem. Soc.*, 1949, **71**, 2703–2707.
- 44 E. T. Feng, C. B. Fan, N. S. Wang, G. Liu and S. Z. Pu, *Dyes Pigm.*, 2018, **151**, 22–27.
- 45 S. B. Roy and K. K. Rajak, *J. Photochem. Photobiol., A*, 2017, **332**, 505–514.
- 46 Y. Wang, Z. Y. Ma, D. L. Zhang, J. L. Deng, X. Chen and C. Z. Xie, *Spectrochim. Acta, Part A*, 2018, **195**, 157–164.
- 47 S. M. Hossain, K. Singh, A. Lakma, R. N. Pradhan and A. K. Singh, *Sens. Actuators, B*, 2017, **239**, 1109–1117.
- 48 D. Singhal, N. Gupta and A. K. Singh, *New J. Chem.*, 2016, **40**, 7536–7541.
- 49 S. Y. Li, D. B. Zhang, J. Y. Wang, R. M. Lu, C. H. Zheng and S. Z. Pu, *Sens. Actuators, B*, 2017, **245**, 263–272.
- 50 J. C. Qin, T. R. Li, B. D. Wang, Z. Y. Yang and L. Fan, *Spectrochim. Acta, Part A*, 2014, **133**, 38–43.
- 51 K. Boonkitpatarakul, J. F. Wang, N. Niammont, B. Liu, L. McDonald, Y. Pang and M. Sukwattanasinitt, *ACS Sens.*, 2016, **1**, 144–150.
- 52 W. He and Z. Liu, *RSC Adv.*, 2016, **6**, 59073–59080.



