

Featuring work from Associate Prof. Aurélien Deniaud at Laboratory of Chemistry and Biology of Metals of the University Grenoble Alpes, the CEA and the CNRS, France.

Nuclear translocation of silver ions and hepatocyte nuclear receptor impairment upon exposure to silver nanoparticles

Silver nanoparticles are broadly used in consumer products and medical devices leading to human exposure and eventually accumulation in the liver. Within hepatocytes, silver nanoparticles efficiently dissolve into toxic Ag(I) species that are distributed in all organelles including the nucleus, where they trigger an endocrine disruptor-like mechanism.

As featured in:



See Aurélien Deniaud *et al.*,
Environ. Sci.: Nano, 2020, **7**, 1373.



Cite this: *Environ. Sci.: Nano*, 2020, 7, 1373

Nuclear translocation of silver ions and hepatocyte nuclear receptor impairment upon exposure to silver nanoparticles†

Vanessa Tardillo Suárez, ^a Elizaveta Karepina, ^{bc} Mireille Chevallet, ^b Benoit Gallet, ^d Cécile Cottet-Rousselle, ^e Peggy Charbonnier, ^b Christine Moriscot, ^{df} Isabelle Michaud-Soret, ^{ib} Wojciech Bał, ^g Alexandra Fuchs, ^h Rémi Tucoulou, ^a Pierre-Henri Jouneau, ^c Giulia Veronesi ^{id} ^{ab} and Aurélien Deniaud ^{ib} ^{*b}

The impact on human health of long-term exposure to low doses of silver nanoparticles (AgNPs) remains understudied. Cellular studies have shown the intracellular dissolution of AgNPs within endolysosomes followed by Ag(I) binding to biomolecular thiolate-containing molecules. However, the precise subcellular distribution of Ag(I) and the nature of the disrupted physiological pathways remained unknown. Novel imaging approaches enabled us to visualize the trafficking of AgNP-containing lysosomes towards a perinuclear location and a nuclear transfer of Ag(I) species with accumulation in the nucleoli. These Ag(I) species impaired nuclear receptor activity, disrupting critical mechanisms of liver physiology in low dose exposure scenarios, thus justifying further research into defining a framework for the safe use of AgNPs.

Received 27th November 2019,
Accepted 1st April 2020

DOI: 10.1039/c9en01348b

rsc.li/es-nano

Environmental significance

Silver nanoparticles (AgNPs) are massively used as biocides in consumer products and medical devices. The biocidal activity arises from the long-lasting release of Ag(I) species that is due to AgNP oxidative dissolution. This process can occur under environmental and biological conditions leading to the exposure of various organisms, including humans, to Ag(I) species. The acute toxicity of AgNPs and Ag(I) has been extensively studied but the physiologic disruptions induced by chronic exposure to AgNPs remain understudied. Under conditions mimicking realistic exposures, an innovative elemental imaging technique enabled Ag(I) nuclear translocation to be visualized. Under these AgNP exposure conditions, specific nuclear receptors get inhibited, revealing an endocrine disruptor-like effect. Due to their tremendous affinity for thiolate, nuclear Ag(I) species most probably replace Zn(II) in Zn finger domains of nuclear receptors.

1 Introduction

Silver nanoparticles (AgNPs) are widely used in everyday products for their biocidal activity and increasing production has led to environmental and human safety concerns over the past few years. Their biocidal properties stem from the durable release of Ag(I) ions from the NP's Ag(0) core due to surface oxidation,^{1,2} ions that are also toxic to mammalian cells.³

In humans, the main and most extensively described exposure route for AgNPs is dietary.^{4,5} However, the impact of nanosilver-containing medical devices that can directly release Ag into lymphatic and blood vessels has been sorely disregarded. The leaching of AgNPs is a well-known drawback of nano-enabled products, which has been studied and quantified, for instance in fabrics and consumer products for children.^{6–8} AgNP-based medical devices, although designed to release biocidal Ag(I) ions, can also leach AgNPs. In order

^a ESRF, The European Synchrotron, 71 Avenue des Martyrs, 38000 Grenoble, France

^b Univ. Grenoble Alpes, CNRS, CEA, IRIG, Laboratoire de Chimie et Biologie des Métaux, F-38000 Grenoble, France. E-mail: aurelien.deniaud@cea.fr; Tel: + 33 (0)4 38 78 96 51

^c Univ. Grenoble Alpes, CEA, IRIG, MEM, F-38000 Grenoble, France

^d Institut de Biologie Structurale, CEA, CNRS, Univ. Grenoble Alpes, 71 Avenue des Martyrs, F-38042 Grenoble, France

^e Univ. Grenoble Alpes, INSERM U1055, Laboratory of Fundamental and Applied Bioenergetics (LBFA), and Environmental and System Biology (BEeSy), 38000 Grenoble, France

^f Integrated Structural Biology Grenoble (ISBG), CNRS, CEA, Université Grenoble Alpes, EMBL, 71 Avenue des Martyrs, F-38042, Grenoble, France

^g Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Pawinskiego 5a, 02-106 Warsaw, Poland

^h DRF/BIG-INSERM U1160, Hôpital St. Louis, F-75010 Paris, France

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9en01348b



to prevent this undesired effect, synthetic strategies that allow adhesion of the NPs to the products are under investigation.^{9,10} Considering that the commercially available AgNP-based materials for clinical use are catheters and wound dressings,¹¹ leaching from such products results in the release of AgNPs into the bloodstream.

The exposure route to AgNPs influences the biodistribution and body clearance of silver, as shown by an *in vivo* study where mice were exposed to a single dose of AgNPs either orally or intravenously.¹² In both cases, Ag preferentially accumulates in the liver. However, in the case of parenteral exposure, the Ag concentration in the blood after 24 h is ten times higher, and its clearance is much slower than that after ingestion. This is most probably due to the direct introduction of Ag(0) from AgNPs into the bloodstream in the case of injection, whereas in the case of oral exposure AgNPs are readily transformed in the gastrointestinal tract, and Ag enters the bloodstream mainly in the form of various Ag(I) species (ref. 13 and for a recent review see ref. 4). Besides, several studies in burn patients revealed Ag blood levels in the low to high micromolar range proving the relevance of Ag exposure in the medical context.^{14–16} Therefore, it is of paramount importance to better understand the AgNP fate in the liver to predict the risk for patients during hospital care. In spite of its relevance, this exposure scenario has been as yet disregarded in nanotoxicology studies. In this context, we decided to address the impact of an exposure over several days to low and non-toxic concentrations of pristine AgNPs on critical hepatocyte functions.

Over the past few years, several studies have described the fate of AgNPs in various mammalian cell models.^{17–21} A consensus has emerged showing AgNP dissolution in endosomes and lysosomes and the subsequent distribution of Ag(I) species throughout the cell mainly in the form of Ag-thiolate complexes.^{18,20–22} This type of complexation comes as no surprise as Ag has a picomolar or higher affinity for thiolates depending on the coordination chemistry offered by the biomolecular ligands.²³ However, recent imaging studies did not provide the subcellular distribution of Ag(I) species at the organelle level.^{3,21} The identification of cell compartments that can store Ag and/or that can be critically affected by this non-physiological and deleterious metal ion remains a critical question. Of particular importance is the long-standing controversy about AgNP entry into the nucleus. Indeed, in the literature, AgNPs have been shown in the nucleus only in few publications,^{24,25} while the majority of the studies did not. Moreover, the chemical composition of these NPs visualized in the nucleus was not proven. We described in detail the limits of those studies in a recent review.⁴

In the current study, we explored the interior of hepatic cells exposed to AgNPs by applying several high-resolution imaging methods to thin cell sections. The synergistic use of confocal microscopy, electron microscopy, elemental imaging by synchrotron X-ray fluorescence (XRF), and energy

dispersive X-ray spectroscopy (EDX), complemented by chemical analysis through X-ray absorption spectroscopy (XAS), allowed us to infer the subcellular distribution of AgNPs and Ag-thiolate complexes. These experiments were carried out under long-term low dose exposure in order to be as realistic as possible. We thus highlighted that Ag(I) species but not AgNPs are translocated to the nucleus and we further analyzed their impact on nuclear functions. Hepatocyte nuclear receptors, such as the liver X receptor (LXR) and the farnesoid X receptor (FXR), also known as the bile acid receptor (BAR), were identified as molecular targets that were inhibited by these translocated metal ions even in a very low dose exposure scenario mimicking chronic exposure. AgNPs could thus trigger an endocrine disruptor-like effect.

2 Experimental

2.1 AgNP characterization

Highly homogenous 20 nm citrate-coated AgNPs were purchased from NanoComposix (20 nm Citrate BioPure™ Silver) and 90 nm PVP-coated AgNPs were purchased from Sigma-Aldrich. Both were previously characterized.²¹ Dynamic light scattering measurements were carried out in cell culture medium supplemented with 10% fetal calf serum at 25 °C in a UVette (Eppendorf) on a Nanostar instrument (Wyatt). Zeta potential was determined using a Malvern Nano ZS in 10 mM Hepes and 2 mM sodium citrate pH 7. In cell culture media, citrate- and PVP-coated AgNPs formed agglomerates of 160 nm and homogenous 90 nm particles, respectively (Table S1†).

2.2 Cell culture

HepG2 cells were grown in MEM medium supplemented with 10% FBS and 1% antibiotics–antimycotic and exposed to cit-AgNPs or PVP-AgNPs at the indicated concentrations of total Ag and for the indicated times. The medium and AgNPs were renewed every 24 hours. When required, chenodeoxycholic acid (CDCA, Sigma-Aldrich) at 80 μM or the equivalent volume of DMSO was added for the last 17 hours of exposure or 22R-hydroxycholesterol (22R-HC, Sigma-Aldrich) at 10 μM or the equivalent volume of absolute ethanol was added for the last 20 hours of exposure.

2.3 Spheroid preparation

Micropatterns were produced on glass slides by deep-UV photolithography by which a non-toxic cell repellent PEG-PLL coating was degraded locally by deep-UV light passing through a photomask (as described in ref. 26). Micropatterns were then coated with fibronectin (20 μg mL⁻¹ in PBS) to form adhesive disks of 50 μm in diameter, surrounded by a cytophobic surface.

HepG2/C3a cells were seeded on micropatterned surfaces at a density of 1.25×10^5 cells per cm² and were allowed to attach. After 2 hours, unattached and floating cells were removed by extensively washing the surface and gentle



plasma atomic emission spectroscopy (ICP-AES) using a Shimadzu ICPE-9800 after overnight mineralization at 50 °C in 65% HNO₃.

2.6 RNA extraction and quantification

At the indicated timepoints, cells were harvested and mRNA was isolated using the Nucleospin RNA kit (Macherey-Nagel) for 2D cultures or the NucleoSpin RNA Plus XS kit (Macherey-Nagel) for the spheroid samples. The RNA concentration was determined using a NanoDrop spectrophotometer (ND-1000). RNA was reverse transcribed with the Affinity script qPCR cDNA synthesis kit (Agilent), according to the manufacturer's instructions using random primers. Quantitative PCR was performed as described in ref. 21, qPCRs were run in triplicate, and quantification was performed using comparative regression (Cq determination mode) using Cfx (Bio-Rad Cfx manager) with GAPDH and HPRT amplification signals as housekeeping genes to correct for total RNA content, while the untreated sample was used as the "calibrator". At least three biological replicates were performed for the presented data. The primers used are described in Table S2, ESI.†

2.7 Nuclear receptor activity

The activity of selected nuclear receptors was determined by measuring the amount of one of their target transcripts upon addition of a specific ligand of the receptor. This activity was compared in hepatocytes exposed or not to AgNPs in order to measure the impact of these NPs on nuclear receptor activity. We followed the activity of two nuclear receptors: FXR and LXR. For the former, CDCA was used to induce the expression of *bsep* (coding for Bile Salt Export Pump, BSEP),²⁸ and for the latter, 22R-HC²⁹ was added and *abcg5* (coding for the ATP-binding cassette subfamily G member 5) expression was monitored. In this case, the control (ctl) sample (unexposed to AgNPs) was set to 100% activity of FXR or LXR and it corresponds to the ratio:

$$\text{Ratio (ctl)} = \frac{\text{mRNA level with ligand (CDCA or 22 R-HC)}}{\text{mRNA level without ligand}}$$

$$\text{Ratio (condition Z)} = \frac{\text{mRNA level with ligand (CDCA or 22 R-HC)}}{\text{mRNA level without ligand}}$$
$$\% \text{ activity } Z = \left[\frac{(\text{Ratio}(\text{condition } Z) - 1)}{(\text{Ratio}(\text{ctl}) - 1)} \right] \times 100$$

Subtracting 1 takes into account that the value of a control in mRNA normalized relative expression is 1.

The amount of Ag present in the cells and in nuclei and nucleoli fractions was quantified by inductively coupled

2.8 Sample preparation for EM and nano-XRF

At the desired timepoint, monolayers of HepG2 cells were fixed overnight at room temperature in a 1:1 ratio mixture of 4% paraformaldehyde, 0.4% glutaraldehyde in 0.2 M PHEM pH 7.2 and culture medium, washed in 0.1 M PHEM pH 7.2, and fixed for 30 minutes in 2% paraformaldehyde, 0.2% glutaraldehyde in 0.1 M PHEM pH 7.2, washed in 0.1 M PHEM pH 7.2 and post-fixed in 1% OsO₄ in 0.1 M PHEM buffer for 1 h at room temperature. Cells were then dehydrated in graded ethanol series, and flat-embedded using an Epoxy Embedding Medium kit (Sigma-Aldrich). Sections (200 or 400 nm) were cut on a Leica UC7 ultra-microtome using a DiATOME 35° diamond knife and collected on 50 nm-thick Si₃N₄ grids (Oxford Instruments). The same preparation protocol was used for spheroids, where some sections were collected on formvar carbon coated 100 mesh copper grids for scanning transmission electron microscopy coupled with energy-dispersive X-ray spectroscopy (STEM-EDX) and the block was then used for focused ion beam scanning electron microscopy (FIB-SEM).

2.9 Nano-XRF data acquisition and analysis

X-ray fluorescence (XRF) experiments were carried out at the state-of-the-art hard X-ray nanoprobe beamline ID16B at the ESRF.³⁰ The incoming photon energy was set at 29.6 keV and the beam was focused using Kirkpatrick–Baez mirrors down to 50 × 50 nm². The fluorescence emission from the sample was recorded using two 3-element silicon drift detector (SDD) arrays positioned at 13° from the sample. The photon flux on the sample was ~10¹¹ photons per s. High resolution maps were recorded at room temperature, raster scanning the sample in the X-ray focal plane with 100 × 100 nm² step size and 500 ms dwell time/pixel.

Hyperspectral images were analyzed using the PyMCA software package (<http://pymca.sourceforge.net/>).³¹ The detector response was calibrated over a thin film multilayer sample from AXO (RF8-200-S2453). XRF data were energy calibrated, normalized by the incoming photon flux, and batch-fitted in order to extract spatially resolved elemental concentrations, assuming a biological matrix of light elements and density 1 g cm⁻³ according to NIST standards (<https://physics.nist.gov/cgi-bin/Star/compos.pl?matno=261>). At least three different cells per exposure condition were analyzed, and their averages and standard deviations are reported in Fig. 2A.

2.10 X-ray absorption spectroscopy (XAS) data acquisition and analysis

Ag K-edge X-ray absorption spectra were acquired on the bending-magnet beamline BM23 of the ESRF.³² HepG2 cells were exposed to AgNPs for the indicated times, pelleted and resuspended in PBS containing 20% glycerol (20 µL). Drops of cell suspensions or reference solutions (50 µL) were deposited in the XAS sample holder equipped with Kapton

windows and immediately frozen in liquid N₂, then transferred into the He cryostat and measured at 10 K. The incoming photon energy was scanned around the Ag K edge (25.514 keV) with a fixed exit double crystal Si(311) monochromator (Kohzu, Japan) cooled with liquid N₂. Harmonic rejection was achieved with Pt coated mirrors. Slits were opened to provide a beam size at sample of ~3 × 1 mm² and minimize the photon density, in order to avoid radiation damage. The energy was calibrated over an Ag foil mounted downstream to the sample. XAS spectra were acquired in fluorescence mode, with a 13-element Ge detector from Canberra. Data processing and linear combination analysis were performed with the Athena software;³³ the chosen fitting range for linear combination analysis was [-20; 80] eV with respect to the edge energy.

2.11 STEM-EDX

For the chemical identification of AgNPs, thin sections were observed in scanning/transmission electron microscopy (STEM) mode and analyzed by EDX using either a Zeiss Merlin microscope operated at 30 kV and fitted with two Bruker SDD detectors or a FEI/Tecnaï Osiris microscope operated at 80 kV.

2.12 FIB-SEM acquisition and data analysis

Focused ion beam (FIB) tomography was performed with a Zeiss NVision 40 dual-beam microscope. In a first step, the surface of the sample was protected by the deposition in the microscope of a 2 µm thick Pt or C layer, on a surface of typically 20 × 30 µm². The resin-embedded HepG2/C3a spheroids were then cut in cross-sections, slice by slice, with a Ga⁺ ion beam (typically with a current of 300 or 700 pA at a beam acceleration of 30 kV). After a thin slice was removed with the ion beam, the newly exposed surface was imaged in the SEM at 1.5 kV using the in-column EsB backscatter detector. For each slice, a thickness of 8 nm was removed, and the SEM images were recorded with a pixel size of 4 × 4 nm². The image stack was then registered by cross-correlation using the StackReg plugin in the Fiji software.³⁴ Manual segmentation was performed in Ilastik³⁵ and projected in three-dimension using AVIZO (FEI, USA).

2.13 Reflection confocal microscopy

HepG2 cells were seeded on a two-well Thermo Scientific™ Nunc™ Lab-Tek™ chambered cover glass suitable for confocal microscopy. At 60% confluency, cells were incubated with 1 µM Hoechst 33342 for 30 minutes at room temperature then washed with complete medium. Confocal microscopy was carried out in a perfusion chamber (POC Chamber, LaCom®, Erbach, Germany) and incubation system (O₂-CO₂-⁰C, PeCom®, Erbach, Germany) that was mounted on a Leica TCS SP2 AOBS inverted laser scanning confocal microscope using a 63× oil immersion objective (HCX PL APO 63.0 × 1.40 OIL). Cells were exposed to cit-AgNPs or PVP-AgNPs at the indicated concentrations of total Ag just before



Environ. Sci.: Nano, 2020, 7, 1373–1387 | 1377

staining (Fig. 1A). In HepG2 cells exposed to low concentrations of AgNPs over three days, we observed Ag hot spots inside vesicles, most probably corresponding to pristine or partially dissolved NPs (Fig. 1B, red-yellow pixels). Using confocal microscopy, both citrate and PVP AgNPs can be detected in reflection mode (Fig. S4, ESI†, red) and they were often observed to be co-localized with Lamp1 stained vesicles, thus identifying such vesicles as lysosomes (Fig. S4, ESI†, green, co-localization red-green highlighted with white arrows). In addition, under the same experimental conditions, similar perinuclear lysosome vesicles containing dark particles were observed by STEM (Fig. S5 and S6, ESI†). EDX showed that they contained Ag as well as sulfur in the

case of both cit-AgNPs and PVP-AgNPs (Fig. S5 and S6, ESI†). A variable proportion of the detected elements was observed within the lysosome, with some areas of stronger co-localization (for instance in Fig. S5B, ESI†, lower panels comparing Ag and S EDX maps). This lets us hypothesize the formation in lysosomes of transient species in-between the pristine AgNPs and the Ag(I)-thiolate complexes revealed by XAS, which would represent the first step of surface oxidation necessary for Ag(I) ion release.^{38,39} However, considering that XAS proved that Ag(I)-thiolate species represent 70–75% of all Ag atoms in cells, we must exclude that all the thiolates are simply adsorbed at the surface of AgNPs, in particular if we consider that the fraction of surface Ag atoms in a 20 nm





Fig. 1 Semi-thin sections of HepG2 cells analyzed by nano-XRF. (A) Os and (B) Ag areal density maps for a 200 nm section of the same cell exposed for 72 h to 12.5 μM cit-AgNPs. (C) Ag areal density map (upper panel) and two-color map (Ag in red and Os in green, lower panel) of the same region for a 400 nm section of cells exposed for 48 h to 12.5 μM cit-AgNPs. (D) Ag areal density map (upper panel) and two-color map (Ag in red and Os in green, lower panel) of the same region for a 400 nm section of cells exposed for 72 h to 12.5 μM cit-AgNPs. Areal densities are expressed in $\mu\text{g cm}^{-2}$; pixel size is $100 \times 100 \text{ nm}^2$. Scale bars are 10 μm . c: cytoplasm; n: nuclei; N: nucleoli. The upper areal density limit in color bars is set to values that allow the visualization of the low areal density signal.



In order to confirm these data, we quantified Ag in nuclei and bound to nucleoli following subcellular fractionation after 24, 48 and 72 hours of exposure. In order to avoid contamination of the nuclei fraction by endosomes and lysosomes loaded with dense AgNPs, sucrose/percoll density gradient centrifugation was performed. A continuous uptake of Ag in nuclei was observed over 72 hours (Fig. 2B). Nucleoli clearly accumulated Ag since elemental quantification normalized on the protein amount showed about five times



Altogether, our study is the first to show that Ag(I) species enter nuclei and are preferentially accumulated in nucleoli. These novel results are supported by the identification of proteins from the nucleoli possessing Ag-affinity *in vitro*, the so-called AgNOR (Nucleolar Organizer Region)^{42,43} that most probably bind Ag *via* stretches of glutamate or aspartate residues.⁴⁴ However, carboxylates do not possess a high affinity for Ag(I), making Ag-rich nucleoli a reservoir of Ag(I) that can be eventually released in the nucleus and bind higher affinity protein sites.

The initial phase of ribosome biogenesis takes place in nucleoli,^{43,45} in particular with the biosynthesis of the 45S pre-ribosomal RNA (pre-rRNA). We assessed a possible impact of Ag binding to nucleolar proteins on this function. The amount of 45S pre-rRNA was determined by quantitative real-time PCR (qPCR) in HepG2 cells exposed or not to cit-

AgNPs at 6.25 and 12.5 μM for up to 4 days (Fig. 3A). 45S pre-rRNA production remained almost constant under the tested conditions. We only observed a slight overexpression after 24 hours of exposure to cit-AgNPs at 12.5 μM . It is possible that Ag binding to AgNOR slightly disturbs nucleolar functions but as ribosomes are essential, cells have no doubt developed ways to efficiently correct for alteration in their biogenesis pathway. In addition, Ag binding to AgNOR occurs in domains with unknown functions for these proteins, which could explain a limited impact on ribosome biogenesis.

The nucleus carries out many critical functions and any disruption can play a role in the development of non-transmissible pathologies such as cancers. Environmental pollutants, those classified as endocrine disruptors for instance, impair the activity of some nuclear receptors (for review see ref. 46). These are ligand-activated transcription factors with finely tuned activity. They contain zinc finger

DNA binding domains that are structured by zinc ions coordinated by four cysteines (Cys). This amino-acid possesses thiol functions that are very good ligands for Cu(I) or Ag(I) ions and it has been observed *in vitro* that Cu(I) can displace Zn(II) from zinc finger domains.^{47–49} Recently, it was shown that an excess of intracellular Cu(I) in Wilson disease impairs the function of nuclear receptors in the liver.^{28,50,51} To assess the activity of crucial liver nuclear receptors, we determined transcript levels of target genes under the control of these receptors upon addition of their specific ligands under standard growth conditions (100% activation) *versus* those in cells exposed to AgNPs (Fig. 3B–D). Farnesyl X receptor (FXR) is critical in maintaining bile salt homeostasis, and the addition of chenodeoxycholic acid (CDCA) into the culture medium triggers the expression of bile salt export protein (BSEP) mRNA.²⁸ The exposure of HepG2 cells to cit-AgNPs at 6.25 or 12.5 μM for 24 hours




Fig. 3 Impact of AgNP exposure on 45S pre-rRNA expression and FXR and LXR activity. (A) Variation of 45S pre-rRNA expression levels in HepG2 cells exposed to 6.25 or 12.5 μM cit-AgNPs for 24, 48, 72 or 96 hours. The amount of RNA was determined by qPCR with unexposed cells (ctl) set at 1. (B) FXR activity followed by CDCA-induced BSEP expression. HepG2 cells were exposed or not to AgNPs for 24 or 72 hours and to CDCA at 80 μM or DMSO during the last 17 hours. The RNA levels of BSEP were quantified under the different conditions using qPCR. 100% FXR activity was set using control conditions (unexposed to AgNPs) by calculating the ratio: amount of BSEP-RNA in ctl + CDCA/amount of BSEP-RNA in ctl without CDCA. This ratio (+CDCA/-CDCA) was measured for each condition and divided by the ctl ratio in order to determine the percentage of FXR activity in cells exposed to AgNPs (for details see the experimental section). (C) LXR activity followed by 22R-HC-induced ABCG5 expression. HepG2 cells were exposed or not to AgNPs for 24, 48, 72 or 96 hours and to 22R-HC at 10 μM or ethanol during the last 20 hours. The RNA levels of ABCG5 were quantified under the different conditions using qPCR. 100% LXR activity was set using control conditions (unexposed to AgNPs) by calculating the ratio: amount of ABCG5-RNA in ctl + 22R-HC/amount of ABCG5-RNA in ctl without 22R-HC. This ratio (+22R-HC/-22R-HC) was measured for each condition and divided by the ctl ratio in order to determine the percentage of LXR activity in cells exposed to AgNPs (for details see the experimental section). (D) FXR and LXR activities measured as in B and C in HepG2 cells exposed or not for 10 days to 1 μM cit-AgNPs or 2 μM PVP-AgNPs. For A to D, each experiment was performed at least three times independently. * stands for data statistically different from the corresponding control with $p < 0.05$ and ** stands for data statistically different from the corresponding control with $p < 0.01$.



The concentration of cit-AgNPs was then decreased down to 3.1 μ M Ag in order to observe effects on LXR activity that could not be related to metal stress in hepatocytes. Under these conditions, MTs were only minimally overexpressed (Fig. S2B, ESI[†]) and LXR β was very stable (Fig. S11B, ESI[†]). Interestingly, such a low concentration of AgNPs led to about 80 and 90% inhibition of LXR after 24 and 72 hours of exposure (Fig. 3C), respectively. Thus, we prove that the

Our data have clearly demonstrated the rapid uptake by nuclei of a significant amount of Ag(I) ions that accumulate in nucleoli and induce dose-dependent effects on nuclear functions (Fig. 1–3). A debate is still ongoing concerning the possible uptake of the nanoparticulate form of AgNPs in the nuclei, but the methods used to demonstrate this hypothesis cannot be considered as conclusive (for detailed explanations, see the recent review in ref. 4). In order to deepen our understanding of AgNP trafficking towards nuclei, we performed combined confocal reflected light microscopy and 3D-EM analysis using FIB-SEM. The particle nature of AgNPs makes it possible to visualize their dynamics by reflectance confocal microscopy (Fig. 4 and S13–S15, ESI†).⁵² Cit-AgNPs could only be detected in cells three hours after exposure, while PVP-AgNPs were observed as clear single spots immediately following their addition (Fig. 4A and S13, ESI† *versus* 4B and S14, ESI†). This is consistent with our previous observations of isolated PVP-AgNPs as opposed to agglomerated cit-AgNPs during endocytosis²¹ and with the kinetics of uptake of both types of AgNPs (Fig. S10, ESI†). We also recently published electron microscopy micrographs




Fig. 4 Dynamics of cit-AgNPs and PVP-AgNPs in HepG2 cells. (A) Confocal fluorescence and reflectance microscopy of HepG2 cells exposed to 12.5 μM cit-AgNPs for the indicated times. Nuclei were labeled with Hoechst (blue) and AgNPs were observed in reflection mode (green). At $t = 3$ h, a bright field image is also shown. (B) Confocal fluorescence and reflectance microscopy of HepG2 cells exposed to 25 μM PVP-AgNPs for the indicated times. Particles are represented in green and nuclei in blue. At $t = 0$ h, a bright field image is also shown. White arrows pinpoint particles close to the nucleus and red arrow PVP-AgNPs bound to the cell membrane. The scale bars correspond to 20 μm .

revealing PVP-AgNPs bound to the cell membrane at the interface between cells.⁴ This observation was made on thin cell sections and was specific to PVP-AgNPs. Similarly, confocal reflectance microscopy images showed particles evenly distributed on lines or curves between cells (Fig. 4B and S14, red arrows, ESI†) in addition to intracellular particles.

Therefore, for an unknown reason, we confirmed that a fraction of the PVP-AgNPs but not of the cit-AgNPs binds to membranes between cells as described in Marchioni *et al.*⁴ For cit-AgNPs, the reflection signal in the cell increased over time with a tendency to accumulate in the perinuclear region (Fig. 4A and S13, white arrows, ESI†). In the case of PVP-AgNPs, the dynamics were faster and NPs that came in the

vicinity of nuclei disappeared rapidly (Fig. 4B and S14, white arrows, ESI†). Therefore, within just a few hours, AgNPs reached the vicinity of the nucleus where they promptly dissolved. In order to go a step further, the PVP-AgNP distribution within hepatocytes grown as spheroids was also obtained at nanometer resolution using FIB-SEM. This method provided the unique capability to observe individual NPs without any preparation-dependent artifacts that might lead to erroneous localization of NPs within the cell. Indeed, the biological sample is embedded in a resin and imaged 8 nm layer by 8 nm layer thanks to Ga gun nanomachining (for review see ref. 53) thus avoiding diamond knife alterations of the AgNP location. Moreover, for this experiment, a derivative of the HepG2 cell line, named HepG2/C3a,⁵⁴ was used to






Fig. 5 EM analysis of the subcellular distribution of AgNP-containing vesicles. (A) Original micrographs extracted from two different FIB-SEM stacks of images (see also Video S1† for stack 1 with the corresponding segmentation). Endolysosomes containing AgNPs close to the nucleus (left) and in the middle of the mitochondrial network (right) are indicated with white arrows. (B) 3D reconstruction of a nucleus (green) with bound vesicles (yellow) containing AgNPs (black) seen by transparency (C). (D) SEM micrograph from the stack used to make the 3D reconstruction (B and C). This micrograph shows a direct contact between the AgNP-containing vesicle and the nucleus (red arrow). Scale bars represent 2 μm . (E) Large field of view STEM micrograph showing the presence of a lysosome containing NPs in contact with a nucleus in HepG2 cells exposed for 72 hours to 12.5 μM cit-AgNPs. (F) STEM micrograph of the orange area in (E) at higher magnification. The red arrow pinpoints a region of close contact between the lysosome and the nuclei. The different colored boxes correspond to areas scanned by EDX with the spectra presented in (G). (G) EDX spectra of different regions in the energy range 1–3.5 keV. The positions of Si, S, Cl and Ag peaks are highlighted.



It must be considered that XAS analysis (section 3.1) proved that AgNPs represent only $\sim 30\%$ of the total Ag species encountered in cells 24 h after exposure, whereas Ag-thiolate complexes represent $\sim 70\%$ of species. This extent of dissolution is consistent with the observed perinuclear localization of AgNP-containing endolysosomes. Indeed, the closer the lysosomes are to the nucleus, the lower their internal pH (ref. 55) and lower pH values favor AgNP dissolution. Nonetheless, the combination of all our data prompted us to reconsider our vision on AgNP dissolution. Indeed, different forms of species containing Ag and S exist within the cell. XRF microscopy showed intense Ag signals within endosomes and lysosomes and low Ag signals in the cytosol and other organelles such as nuclei. However, STEM-EDX experiments revealed that lysosomes contain both AgNPs close to the pristine form and highly transformed AgNPs (Fig. 5 and S5 and S6, ESI†). This is particularly clear in Fig. 5E–G where darker particles have a high signal of Ag (pink square in Fig. 5F), while grey particle-like areas have a low Ag signal with the presence of S in a similar intensity (blue and green squares in Fig. 5F). Interestingly, in Fig. 5F, we can almost see a connection between a lysosome and the nucleus (red arrow) but no particle nor Ag signal was observed in the nucleus even in areas that look darker in STEM. Similarly, Ag was never detected in the cytosol by EDX, while it was by XRF both in the cytosol and nuclei. Species containing Ag and S thus exist in two forms, particle-like in lysosomes and soluble elsewhere. Therefore, our results constitute the first evidence of a rapid trafficking of endosomes maturing into lysosomes that contain AgNPs under conditions favoring their rapid transformation into

4 Conclusions

Based on previously published *in vitro* data^{47–49} and on our recent study⁵⁹ we can ascribe the observed effects on nuclear receptor activity to a displacement of Zn(II) by Ag(I) in the zinc finger DNA binding domains, as already proposed with Cu(I) in a mouse model of Wilson disease.^{28,50,51} Zinc fingers require a tetra-coordination geometry to be able to bind their specific DNA-binding domains upon ligand binding. We showed that Ag(I) ions displace Zn(II) from a model zinc finger triggering the formation of an Ag₄S₄ cluster, which alters the finger structure.⁵⁹ This *in vitro* data provides the basis to explain the inhibition of nuclear receptor activity observed upon AgNP exposure. The impairment of this activity is highly detrimental to liver metabolism as it will affect lipid, cholesterol and bile salt homeostasis, among others. Moreover, released Ag(I) ions could also interact with other nuclear receptors, possibly leading to developmental diseases⁴⁶ such as those observed with endocrine disruptors. It could therefore be of high relevance to study the impact of the combined exposure to AgNPs and endocrine disruptors that target the same protein family. In addition, the binding of released Ag(I) cannot be limited to nuclear receptors. Zinc finger proteins account for 3% of the human genome⁶⁰ and many sites with two or three exposed thiols exist in proteins. It would therefore be of high

Environ. Sci.: Nano, 2020, 7, 1373–1387 | 1385

- 10 K. W. Yu, Y. Yang, Y. Zhang, J. Deng and C. Lin, Antimicrobial activity and cytocompatibility of silver nanoparticles coated catheters via a biomimetic surface functionalization strategy, *Int. J. Nanomed.*, 2015, **10**, 7241–7252.
- 11 S. Nakamura, M. Sato, Y. Sato, N. Ando, T. Takayama, M. Fujita and M. Ishihara, Synthesis and Application of Silver Nanoparticles (Ag NPs) for the Prevention of Infection in Healthcare Workers, *Int. J. Mol. Sci.*, 2019, **20**, 3620.
- 12 K. Park, E. J. Park, I. K. Chun, K. Choi, S. H. Lee, J. Yoon and B. C. Lee, Bioavailability and toxicokinetics of citrate-coated silver nanoparticles in rats, *Arch. Pharmacol. Res.*, 2011, **34**, 153–158.
- 13 P. Bove, M. A. Malvindi, S. S. Kote, R. Bertorelli, M. Summa and S. Sabella, Dissolution test for risk assessment of nanoparticles: a pilot study, *Nanoscale*, 2017, **9**, 6315–6326.
- 14 E. C. M. Mimura, J. P. M. Favoreto, M. E. Favero, K. L. Bonifacio, T. S. Peixe, A. A. Morita, D. S. Barbosa, M. J. S. Yabe and A. J. F. Carrilho, Silver serum levels in burned patients treated with silver sulfadiazine and its toxicity on inflammatory cells, *Burns*, 2019, DOI: 10.1016/j.burns.2019.11.012.
- 15 M. Trop, M. Novak, S. Rodl, B. Hellbom, W. Kroell and W. Goessler, Silver-coated dressing acticoat caused raised liver enzymes and argyria-like symptoms in burn patient, *J. Trauma: Inj., Infect., Crit. Care*, 2006, **60**, 648–652.
- 16 A. T. Wan, R. A. Conyers, C. J. Coombs and J. P. Masterton, Determination of silver in blood, urine, and tissues of volunteers and burn patients, *Clin. Chem.*, 1991, **37**, 1683–1687.
- 17 A. R. Gliga, S. Skoglund, I. O. Wallinder, B. Fadeel and H. L. Karlsson, Size-dependent cytotoxicity of silver nanoparticles in human lung cells: the role of cellular uptake, agglomeration and Ag release, *Part. Fibre Toxicol.*, 2014, **11**, 11.
- 18 X. Jiang, T. Miclăuș, L. Wang, R. Foldbjerg, D. S. Sutherland, H. Autrup, C. Chen and C. Beer, Fast intracellular dissolution and persistent cellular uptake of silver nanoparticles in CHO-K1 cells: implication for cytotoxicity, *Nanotoxicology*, 2014, **9**, 181–189.
- 19 R. P. Singh and P. Ramarao, Cellular uptake, intracellular trafficking and cytotoxicity of silver nanoparticles, *Toxicol. Lett.*, 2012, **213**, 249–259.
- 20 G. Veronesi, C. Aude-Garcia, I. Kieffer, T. Gallon, P. Delangle, N. Herlin-Boime, T. Rabilloud and M. Carriere, Exposure-dependent Ag⁺ release from silver nanoparticles and its complexation in AgS₂ sites in primary murine macrophages, *Nanoscale*, 2015, **7**, 7323–7330.
- 21 G. Veronesi, A. Deniaud, T. Gallon, P. H. Jouneau, J. Villanova, P. Delangle, M. Carriere, I. Kieffer, P. Charbonnier, E. Mintz and I. Michaud-Soret, Visualization, quantification and coordination of Ag⁺ ions released from silver nanoparticles in hepatocytes, *Nanoscale*, 2016, **8**, 17012–17021.
- 22 L. Wang, T. Zhang, P. Li, W. Huang, J. Tang, P. Wang, J. Liu, Q. Yuan, R. Bai, B. Li, K. Zhang, Y. Zhao and C. Chen, Use of Synchrotron Radiation-Analytical Techniques To Reveal Chemical Origin of Silver-Nanoparticle Cytotoxicity, *ACS Nano*, 2015, **9**, 6532–6547.
- 23 N. W. H. Adams and J. R. Kramer, Potentiometric Determination of Silver Thiolate Formation Constants Using a Ag₂S Electrode, *Aquat. Geochem.*, 1999, **5**, 1–11.
- 24 P. V. AshaRani, G. Low Kah Mun, M. P. Hande and S. Valiyaveetil, Cytotoxicity and genotoxicity of silver nanoparticles in human cells, *ACS Nano*, 2009, **3**, 279–290.
- 25 H. R. Kim, M. J. Kim, S. Y. Lee, S. M. Oh and K. H. Chung, Genotoxic effects of silver nanoparticles stimulated by oxidative stress in human normal bronchial epithelial (BEAS-2B) cells, *Mutat. Res., Genet. Toxicol. Environ. Mutagen.*, 2011, **726**, 129–135.
- 26 A. Azioune, N. Carpi, Q. Tseng, M. Théry and M. Piel, in *Methods in Cell Biology*, ed. L. Cassimeris and P. Tran, Academic Press, 2010, vol. 97, pp. 133–146.
- 27 A. Scherl, Y. Couté, C. Déon, A. Callé, K. Kindbeiter, J.-C. Sanchez, A. Greco, D. Hochstrasser and J.-J. Diaz, Functional Proteomic Analysis of Human Nucleolus, *Mol. Biol. Cell*, 2002, **13**, 4100–4109.
- 28 C. R. Wooton-Kee, A. K. Jain, M. Wagner, M. A. Grusak, M. J. Finegold, S. Lutsenko and D. D. Moore, Elevated copper impairs hepatic nuclear receptor function in Wilson's disease, *J. Clin. Invest.*, 2015, **125**, 3449–3460.
- 29 E. Kaneko, M. Matsuda, Y. Yamada, Y. Tachibana, I. Shimomura and M. Makishima, Induction of intestinal ATP-binding cassette transporters by a phytosterol-derived liver X receptor agonist, *J. Biol. Chem.*, 2003, **278**, 36091–36098.
- 30 G. Martinez-Criado, J. Villanova, R. Tucoulou, D. Salomon, J. P. Suuronen, S. Laboure, C. Guilloud, V. Valls, R. Barrett, E. Gagliardini, Y. Dabin, R. Baker, S. Bohic, C. Cohen and J. Morse, ID16B: a hard X-ray nanoprobe beamline at the ESRF for nano-analysis, *J. Synchrotron Radiat.*, 2016, **23**, 344–352.
- 31 V. A. Solé, E. Papillon, M. Cotte, P. Walter and J. Susini, A multiplatform code for the analysis of energy-dispersive X-ray fluorescence spectra, *Spectrochim. Acta, Part B*, 2007, **62**, 63–68.
- 32 O. Mathon, A. Beteva, J. Borrel, D. Bugnazet, S. Gatla, R. Hino, I. Kantor, T. Mairs, M. Munoz, S. Pasternak, F. Perrin and S. Pascarelli, The time-resolved and extreme conditions XAS (TEXAS) facility at the European Synchrotron Radiation Facility: the general-purpose EXAFS bending-magnet beamline BM23, *J. Synchrotron Radiat.*, 2015, **22**, 1548–1554.
- 33 B. Ravel and M. Newville, ATHENA, ARTEMIS, HEPHAESTUS: data analysis for X-ray absorption spectroscopy using IFEFFIT, *J. Synchrotron Radiat.*, 2005, **12**, 537–541.
- 34 J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak and A. Cardona, Fiji - an Open Source platform for biological image analysis, *Nat. Methods*, 2012, **9**, 676–692.
- 35 C. Sommer, C. Straehle, U. Köthe and F. A. Hamprecht, Ilastik: Interactive learning and segmentation toolkit, *2011 IEEE International Symposium on Biomedical Imaging: From Nano to Macro*, Chicago, IL, 2011, pp. 230–233.

- 36 M. Chevallet, B. Gallet, A. Fuchs, P. H. Jouneau, K. Um, E. Mintz and I. Michaud-Soret, Metal homeostasis disruption and mitochondrial dysfunction in hepatocytes exposed to sub-toxic doses of zinc oxide nanoparticles, *Nanoscale*, 2016, **8**, 18495–18506.
- 37 M. Cuillel, M. Chevallet, P. Charbonnier, C. Fauquant, I. Pignot-Paintrand, J. Arnaud, D. Cassio, I. Michaud-Soret and E. Mintz, Interference of CuO nanoparticles with metal homeostasis in hepatocytes under sub-toxic conditions, *Nanoscale*, 2014, **6**, 1707–1715.
- 38 A. P. Gondikas, A. Morris, B. C. Reinsch, S. M. Marinakos, G. V. Lowry and H. Hsu-Kim, Cysteine-induced modifications of zero-valent silver nanomaterials: implications for particle surface chemistry, aggregation, dissolution, and silver speciation, *Environ. Sci. Technol.*, 2012, **46**, 7037–7045.
- 39 M. Marchioni, T. Gallon, I. Worms, P.-H. Jouneau, C. Lebrun, G. Veronesi, D. Truffier-Boutry, E. Mintz, P. Delangle, A. Deniaud and I. Michaud-Soret, Insights into polythiol-assisted AgNP dissolution induced by bio-relevant molecules, *Environ. Sci.: Nano*, 2018, **5**, 1911–1920.
- 40 J. Lee, M. M. Pena, Y. Nose and D. J. Thiele, Biochemical characterization of the human copper transporter Ctr1, *J. Biol. Chem.*, 2002, **277**, 4380–4387.
- 41 K. I. Farley, Y. Surovtseva, J. Merkel and S. J. Baserga, Determinants of Mammalian Nucleolar Architecture, *Chromosoma*, 2015, **124**, 323–331.
- 42 C. Goodpasture and S. E. Bloom, Visualization of nucleolar organizer regions in mammalian chromosomes using silver staining, *Chromosoma*, 1975, **53**, 37–50.
- 43 P. Roussel and D. Hernandez-Verdun, Identification of Ag-NOR Proteins, Markers of Proliferation Related to Ribosomal Gene Activity, *Exp. Cell Res.*, 1994, **214**, 465–472.
- 44 P. Roussel, P. Belenguer, F. Amalric and D. Hernandez-Verdun, Nucleolin is an Ag-NOR protein; This property is determined by its amino-terminal domain independently of its phosphorylation state, *Exp. Cell Res.*, 1992, **203**, 259–269.
- 45 J. L. Woolford and S. J. Baserga, Ribosome Biogenesis in the Yeast *Saccharomyces cerevisiae*, *Genetics*, 2013, **195**, 643–681.
- 46 C. E. Foulds, L. S. Trevino, B. York and C. L. Walker, Endocrine-disrupting chemicals and fatty liver disease, *Nat. Rev. Endocrinol.*, 2017, **13**, 445–457.
- 47 B. J. Deegan, A. M. Bona, V. Bhat, D. C. Mikles, C. B. McDonald, K. L. Seldeen and A. Farooq, Structural and Thermodynamic Consequences of the Replacement of Zinc with Environmental Metals on ER α -DNA Interactions, *J. Mol. Recognit.*, 2011, **24**, 1007–1017.
- 48 T. W. Hutchens, H. Allen Mark, M. Li Chee and T.-T. Yip, Occupancy of a C2-C2 type 'Zinc-finger' protein domain by copper Direct observation by electrospray ionization mass spectrometry, *FEBS Lett.*, 1992, **309**, 170–174.
- 49 P. F. Predki and B. Sarkar, Effect of replacement of "zinc finger" zinc on estrogen receptor DNA interactions, *J. Biol. Chem.*, 1992, **267**, 5842–5846.
- 50 J. P. Hamilton, L. Koganti, A. Muchenditsi, V. S. Pendyala, D. Huso, J. Hankin, R. C. Murphy, D. Huster, U. Merle, C. Mangels, N. Yang, J. J. Potter, E. Mezey and S. Lutsenko, Activation of Ixr/rxr pathway ameliorates liver disease in atp7b(–/–) (Wilson disease) mice, *Hepatology*, 2016, **63**, 1828–1841.
- 51 P. Wilmarth, K. Short, O. Fiehn, S. Lutsenko, L. David and J. L. Burkhead, A systems approach implicates nuclear receptor targeting in the Atp7b(–/–) mouse model of Wilson's Disease, *Metallomics*, 2012, **4**, 660–668.
- 52 C. S. Kim, X. Li, Y. Jiang, B. Yan, G. Y. Tonga, M. Ray, D. J. Solfiell and V. M. Rotello, Cellular imaging of endosome entrapped small gold nanoparticles, *MethodsX*, 2015, **2**, 306–315.
- 53 K. Narayan and S. Subramaniam, Focused ion beams in biology, *Nat. Methods*, 2015, **12**, 1021–1031.
- 54 K. Wrzesinski, M. C. Magnone, L. V. Hansen, M. E. Kruse, T. Bergauer, M. Bobadilla, M. Gubler, J. Mizrahi, K. Zhang, C. M. Andreasen, K. E. Joensen, S. M. Andersen, J. B. Olesen, O. B. Schaffalitzky de Muckadell and S. J. Fey, HepG2/C3A 3D spheroids exhibit stable physiological functionality for at least 24 days after recovering from trypsinisation, *Toxicol. Res.*, 2013, **2**, 163–172.
- 55 D. E. Johnson, P. Ostrowski, V. Jaumouille and S. Grinstein, The position of lysosomes within the cell determines their luminal pH, *J. Cell Biol.*, 2016, **212**, 677–692.
- 56 B. E. Black, J. M. Holaska, F. Rastinejad and B. M. Paschal, DNA binding domains in diverse nuclear receptors function as nuclear export signals, *Curr. Biol.*, 2001, **11**, 1749–1758.
- 57 M. S. Mavinakere, J. M. Powers, K. S. Subramanian, V. R. Roggero and L. A. Allison, Multiple Novel Signals Mediate Thyroid Hormone Receptor Nuclear Import and Export, *J. Biol. Chem.*, 2012, **287**, 31280–31297.
- 58 A. Helleboid-Chapman, S. Helleboid, H. Jakel, C. Timmerman, C. Sergheraert, F. Pattou, J. Fruchart-Najib and J.-C. Fruchart, Glucose regulates LXR α subcellular localization and function in rat pancreatic β -cells, *Cell Res.*, 2006, **16**, 661.
- 59 K. Kluska, M. D. Peris-Diaz, D. Plonka, A. Moysa, M. Dadlez, A. Deniaud, W. Bal and A. Krezel, Formation of highly stable multinuclear Ag_nSn clusters in zinc fingers disrupts their structure and function, *Chem. Commun.*, 2020, **56**, 1329–1332.
- 60 A. Klug, The discovery of zinc fingers and their applications in gene regulation and genome manipulation, *Annu. Rev. Biochem.*, 2010, **79**, 213–231.
- 61 M. Marchioni, G. Veronesi, I. Worms, W. L. Ling, T. Gallon, D. Leonard, C. Gateau, M. Chevallet, P. H. Jouneau, L. Carlini, C. Battocchio, P. Delangle, I. Michaud-Soret and A. Deniaud, Safer-by-design biocide made of tri-thiol bridged silver nanoparticle assemblies, *Nanoscale Horiz.*, 2020, **5**, 507–513.
- 62 Y. G. Yuan, Q. L. Peng and S. Gurunathan, Silver nanoparticles enhance the apoptotic potential of gemcitabine in human ovarian cancer cells: combination therapy for effective cancer treatment, *Int. J. Nanomed.*, 2017, **12**, 6487–6502.
- 63 F. Peng, M. I. Setyawati, J. K. Tee, X. Ding, J. Wang, M. E. Nga, H. K. Ho and D. T. Leong, Nanoparticles promote in vivo breast cancer cell intravasation and extravasation by inducing endothelial leakiness, *Nat. Nanotechnol.*, 2019, **14**, 279–286.

