



# Reengineering of cancer cell surface charges can modulate cell migration†

 Mattia Ghirardello,‡ Radhe Shyam  ‡ and M. Carmen Galan  \*

 Cite this: *Chem. Commun.*, 2022, 58, 5522

 Received 21st January 2022,  
Accepted 6th April 2022

DOI: 10.1039/d2cc00402j

rsc.li/chemcomm

The ability to modulate the cell surface structure provides a powerful tool to understand fundamental processes and also to elicit desired cellular responses. Here we report the development of a new class of 'clickable labels' to reengineer the cell surface charges of live cells. The method relies on the use of metabolic oligosaccharide engineering (MOE) combined with chemo selective labeling of cell surface azido-containing sialic acids with dibenzocyclooctyne (DBCO) ionic-probes. Using this strategy, we demonstrate that reducing the negative charge induced by the overexpression of cell surface sialic acids in cancer cells leads to a reduction in cell migration without affecting drug susceptibility.

Cell adhesion and migration is essential in cell communication and regulation processes. As such, it plays fundamental roles in many biological processes such as embryogenesis, haematopoiesis, inflammation, immune responses and metastasis.<sup>1,2</sup> Thus, the biochemical and mechanical interactions between cells and their extracellular matrix (ECM) influence their behaviour and ultimately their role and function.<sup>2-4</sup> Cancer metastasis entails the relocation of malignant cells from a primary tumour site to distant organs, creating new tumour lesions. It is the key cause of failure in cancer therapies and is the primary cause of death in most cancer patients.<sup>5</sup> To move from the primary tumour site, cancer cells have to change their adhesion properties increasing motility and invasiveness capabilities.<sup>6</sup> The whole metastatic process consists of a series of biological events involving multiple biochemical and physical interactions between cancer cells with the ECM.<sup>7-9</sup> However, despite the many advances in cancer research, the mechanisms behind this process are still not fully understood due to the complexity of the system.<sup>6</sup>

Altered glycosylation is one of the hallmarks of cancer,<sup>10</sup> the abundant and aberrant overexpression of sialic acid-terminated glycosides on the cell surface has been correlated with immunosuppression, cell motility, cancer progression and metastasis.<sup>11-16</sup> Evidence suggests that sialic acid-binding receptors found in immune cells, such as Siglecs and Selectins, are exploited by hypersialylated cancer cells to induce immunosuppression and to modulate key immune cell types in the tumour that are responsible for maintaining the appropriate inflammatory environment.<sup>17-19</sup> For instance, it has been shown that hypersialylation may augment colon tumour progression by altering cell preference for certain extracellular matrix milieus, as well as by stimulating cell motility.<sup>15</sup>

Sialic acid is a negatively charged nine-carbon atom monosaccharide featuring a carboxylic acid group at the anomeric carbon. Thus, the aberrant overexpression of sialylated glycans in cancer cells leads to a more negatively charged cell surface when compared to normal cells.<sup>16</sup> Moreover, many of the major functions in cells and organs of the human body are controlled by ionic currents, electric fields, ion flow, and voltage gradients produced by ion channels and pumps, which are also responsible for cell proliferation, migration, and differentiation processes.<sup>20</sup> On this basis, we hypothesize that cellular migration could be modulated by modifying the cell surface charges and in particular those resulting from the overexpression of cell surface sialic acid residues.

Metabolic oligosaccharide engineering (MOE) is a strategy that allows the incorporation of modified sugar residues bearing an unnatural chemical reporter onto glycoproteins.<sup>21,22</sup> The labelled sugars are converted by the cell biosynthetic machinery into activated nucleotide sugars that are transported into the Golgi and then transferred to glycoconjugates destined for secretion, delivery to cellular compartments or presentation on the cell surface. On this context, a number of metabolic glycan reporters have been successfully used to hijack glycan biosynthesis, chemically modify cell surfaces, probe intracellular metabolic flux inside cells, and to identify specific glycoprotein subtypes from the proteome.<sup>21,23-28</sup> *N*-Azidoacetylmannosamine (ManNAz) is most

School of Chemistry, University of Bristol, Cantock's Close, BS8 1TS Bristol, UK.  
E-mail: m.c.galan@bristol.ac.uk; Web: <https://www.galanresearch.com>

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2cc00402j>

‡ These authors contributed equally to this work.




commonly used to visualize sialic acid-containing glycoproteins in living cells.<sup>29</sup> ManNAz utilizes an azide functional group as the chemical reporter, which upon being metabolized, can be selectively derivatized using the Staudinger ligation,<sup>30</sup> the Cu(I) catalyzed<sup>31</sup> or the strain-promoted<sup>32–35</sup> azide–alkyne [3+2] cycloaddition.


Previously in our lab, we described the use of imidazolium tagged-mannosamine derivatives as a chemical reporter that could be metabolically incorporated into cell-surface sialic acids.<sup>22</sup> However, low levels of cationic-labelled cell surface sialic acids were detected as determined by the electrokinetic potential (zeta potential) and fluorescent measurements, hampering its use for our proposed study of the effect on cell surface charge editing. In order to evaluate the effect of modifying the ionic charges of sialylated cell surfaces, a higher degree of cell surface expressed chemical reporters is required. Thus, we proposed that chemo-selective modification of cell surface azido-containing sialic acids with a suitably “clickable” probe would be more efficient. To that end, the strain-promoted azide–alkyne cycloaddition (SPAAC) reaction employing tetracytlylated *N*-Azido-acetylmannosamine ( $Ac_4ManNAz$ ) as the MOE-chemical reporter<sup>25</sup> and a suitably functionalized DBCO probe was chosen. To that end, cationic DBCO **1**, neutral DBCO **2** and anionic DBCO **3** were prepared starting from commercial DBCO-NH<sub>2</sub> **4** in one step and in 46–60% yields after HPLC isolation. EDC/NHS promoted amide coupling of **4** with carboxylic-Imidazolium (ITag) **5** furnished **1**, while ring opening of succinic anhydride or  $\beta$ -propiolactone with **4** gave access to **2** and **3**, respectively (Scheme 1).

To determine optimal dosage and cell cytotoxicity of the new DBCO-labels, cervical cancer (HeLa) cells and human umbilical vein (EA-hy926) cells were exposed for 1 hour to a range of concentrations of DBCO-probes **1–3** (from 12.5  $\mu$ M to 100  $\mu$ M). Metabolic competence was then assessed using Alamar Blue and compared to untreated controls. It was found that 25  $\mu$ M for all the probes and across the two cell lines was optimal to maintain close to 100% viability, while at higher concentrations viability dropped to 50–65% (see details and Fig. S7 in ESI†).

Next, we evaluated the ability of our DBCO-probes to label cell surface sialoglycans expressing the N<sub>3</sub> reporter by using a competitive fluorometric assay with commercial DBCO-Cy5



**Scheme 1** Reagent and conditions: (i) **5**, EDC·HCl, NHS, CH<sub>3</sub>CN, 21 h, rt 46%; (ii)  $\beta$ -propiolactone, toluene, 24 h, rt 60%; (iii) succinic anhydride, DCM, 16 h, rt then K<sub>2</sub>CO<sub>3</sub>, MeOH/H<sub>2</sub>O, 2 h, rt 53%.



**Fig. 1** Relative fluorescence % of HeLa, MDA and EA-hy926 cells which have been incubated with  $Ac_4ManNAz$  (25  $\mu$ M) and labelled with DBCO-Cy5 (Control M); or cells incubated with  $Ac_4ManNAz$  (Group A) or untreated cells (Group B) which were labelled with DBCO-probes **1–3** and DBCO-Cy5 and compare to control M set to 100%. Data shown after background fluorescence subtraction from native cells exposed to DBCO-Cy5 treatment (blank control). not significant (ns)  $p > 0.05$ , \*\*\*:  $p < 0.001$ .

dye. Breast cancer MDA-MB231 cells, HeLa cells and EA-hy926 cells were used as model systems. Cells previously treated with 25  $\mu$ M  $Ac_4ManNAz$  for 3 days to metabolically express cell surface *N*-azidoacetylneuraminic acid were then exposed to 25  $\mu$ M of DBCO-probes **1–3**, respectively, for 1 hour at 37 °C, followed by a 2nd labelling step with DBCO-Cy5 before measuring the fluorescence (Fig. 1, group A, and Tables S1–S3, ESI†). Results were then compared to control cells, *e.g.* cells that had been treated with  $Ac_4ManNAz$  and underwent the 2nd labelling step with fluorescent DBCO-Cy5 alone (control M) and untreated cells exposed to DBCO-Cy5 (control). As expected, the fluorescence of cells that had been labelled with DBCO-probes **1–3** prior to DBCO-Cy5 exposure was significantly reduced. Additionally, to assess non-specific labelling, cells grown in the absence of  $Ac_4ManNAz$  were also exposed to the same 2-step labelling procedures and showed low levels of background fluorescence (Fig. 1, group B). These results confirm the effective labelling with the novel DBCO-probes (>70% of all azido motifs were reacted when compared with DBCO-Cy5 labelling alone (control M)).

Sialic acid expressed on the cell surface is a major contributor to the net negative charge on the surface of mammalian cells. To confirm the impact of the clickable tags on cell surface charges, the electrokinetic potential (Zeta potential) of HeLa, MDA and EA-hy926 cells which had been treated with 25  $\mu$ M  $Ac_4ManNAz$  and further labelled with 25  $\mu$ M of either cationic DBCO-**1**, neutral DBCO-**2** or anionic DBCO-**3**, as described before, was measured and compared to that of controls *e.g.* untreated cells, cells treated with  $Ac_4ManNAz$  (control M) and untreated cells exposed to DBCO-**1** (Fig. 2 and Tables S4–S6, ESI†). Interestingly, all cells incubated with cationic **1** showed a significant shift towards less negative values when compared to controls or cells treated with either neutral or anionic DBCO probes **2** and **3**. These differences can be ascribed to the partial neutralization of the cell surface negative charges upon







