



Cite this: *Analyst*, 2023, **148**, 2594

Understanding radiation response and cell cycle variation in brain tumour cells using Raman spectroscopy†

Iona E. Hill,^a Marie Boyd,^b Kirsty Milligan,^c Cerys A. Jenkins,^d Annette Sorensen,^b Andrew Jirasek,^c Duncan Graham^a and Karen Faulds^{*a}

Radiation therapy is currently utilised in the treatment of approximately 50% of cancer patients. A move towards patient tailored radiation therapy would help to improve the treatment outcome for patients as the inter-patient and intra-patient heterogeneity of cancer leads to large differences in treatment responses. In radiation therapy, a typical treatment outcome is cell cycle arrest which leads to cell cycle synchronisation. As treatment is typically given over multiple fractions it is important to understand how variation in the cell cycle can affect treatment response. Raman spectroscopy has previously been assessed as a method for monitoring radiation response in cancer cells and has shown promise in detecting the subtle biochemical changes following radiation exposure. This study evaluated Raman spectroscopy as a potential tool for monitoring cellular response to radiation in synchronised *versus* unsynchronised UVW human glioma cells *in vitro*. Specifically, it was hypothesised that the UVW cells would demonstrate a greater radiation resistance if the cell cycle phase of the cells was synchronised to the G₁/S boundary prior to radiation exposure. Here we evaluated whether Raman spectroscopy, combined with cell cycle analysis and DNA damage and repair analysis (γ-H2AX assay), could discriminate the subtle cellular changes associated with radiation response. Raman spectroscopy combined with principal component analysis (PCA) was able to show the changes in radiation response over 24 hours following radiation exposure. Spectral changes were assigned to variations in protein, specifically changes in protein signals from amides as well as changes in lipid expression. A different response was observed between cells synchronised in the cell cycle and unsynchronised cells. After 24 hours following irradiation, the unsynchronised cells showed greater spectral changes compared to the synchronised cells demonstrating that the cell cycle plays an important role in the radiation resistance or sensitivity of the UVW cells, and that radiation resistance could be induced by controlling the cell cycle. One of the main aims of cancer treatment is to stop the proliferation of cells by controlling or halting progression through the cell cycle, thereby highlighting the importance of controlling the cell cycle when studying the effects of cancer treatments such as radiation therapy. Raman spectroscopy has been shown to be a useful tool for evaluating the changes in radiation response when the cell cycle phase is controlled and therefore highlighting its potential for assessing radiation response and resistance.

Received 20th January 2023,
Accepted 30th April 2023

DOI: 10.1039/d3an00121k

rsc.li/analyst

Every year in the UK, there are over 12 000 new brain, central nervous system and intracranial tumour cases diagnosed making brain tumours the 9th most common type of cancer.¹

These tumours account for over 5000 deaths annually in the UK and the ten-year survival rate for brain tumours is only around 15%, which is one of the lowest of all types of cancer.^{1,2} Difficulty treating cancers in the brain accounts for the poor patient survival.

For brain tumours, surgery is most commonly used to remove the bulk of the tumour, however surgery is not always possible due to the location of the tumour and delicate nature of the brain.³ For this reason, surgery is often used in combination with radiation therapy and chemotherapy, in particular temozolomide treatment.⁴ Radiation therapy is common across all types of cancer with over 50% of cancer patients receiving radiotherapy as part of their treatment regime.^{3,5} In

^aCentre for Molecular Nanometrology, Department of Pure and Applied Chemistry, Technology and Innovation Centre, University of Strathclyde, 99 George Street, Glasgow, G1 1RD, UK. E-mail: karen.faulds@strath.ac.uk

^bStrathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, G1 1XQ, UK

^cDepartment of Physics, The University of British Columbia, Kelowna, Canada

^dSwansea University Medical School, Swansea University, Singleton Park, Swansea, SA2 8PP, UK

†Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3an00121k>



an organ like the brain, damage to healthy tissue surrounding the tumour must be minimised, as damage to healthy brain tissue could have detrimental effects for the patients' cognitive function. Therefore, the dose used in radiation therapy must be kept as low as possible by utilising conformal radiation therapy and extended treatment times. The total clinical dose of radiation administered is normally between 45 and 60 Gy, however, a patient will receive regular lower doses (1.5–3 Gy) over a long period of time to make up this total dose.³ This method of treatment is known as radiation fractionation, where dividing the overall dose into smaller doses reduces the toxic effects to healthy cells.⁶

The treatment limitations of brain tumours have led to a drive towards patient tailored radiation therapy. Specifically, a range of innovations have been developed to adapt the treatment regime to a patient's specific needs, including methods of predicting radiation resistance,^{7–9} ways to monitor radiation response in real time¹⁰ or the use of radiosensitising agents.^{11–14} However, currently there are no clinically-implemented methods for assessing tumour radiation response in patients during the course of their treatment. Radiation response and the radiobiology of tumours and tissue is normally assessed *in vitro* in cancer cell lines and 3D tumour models assessing cell survival,¹⁵ apoptosis,¹⁶ hypoxia,^{17,18} deoxyribonucleic acid (DNA) damage and repair^{19,20} and cell cycle arrest.²¹ These methods are often time-consuming and destructive to the sample and therefore pose difficulties for translation to the clinic. Due to the heterogeneous nature of tumours between, and within patients, there is significant variability between how patients respond to therapy. This limits the efficacy of treatment, as lower doses of therapy are given to reduce the toxicity to the most sensitive within the population. Thus, understanding the differences in patient response would aid in improving patient tailored therapy and combination therapies.

Raman spectroscopy has previously been evaluated as a tool for monitoring radiation response at a cellular level.^{22–31} Raman is a vibrational spectroscopic technique which is non-invasive and non-destructive therefore allowing analysis of cell and tissue samples while maintaining integrity of the sample.^{32,33} This technique can be used to gain information about the molecular composition of a sample and to evaluate subtle biochemical changes at a sub-cellular level. The breadth of information that can be gained makes Raman spectroscopy a promising method to use for assessing metabolic changes in cancer cells at both the cellular and molecular level when exposed to a stressor, such as ionising radiation.

Raman spectroscopy has successfully been applied to the study of radiation response in both single cells^{22–31} and tumour models.^{34–37} Early studies by Matthews *et al.* investigated the radiation response of prostate, breast, and lung cancer cell lines when irradiated with 15–50 Gy.^{22,23} Over 72 hours following irradiation, spectral changes arising from changes in cellular concentrations of amino acids, conformational protein structures, nucleic acids and lipid groups were evaluated using principal component analysis. These

changes were proposed to be radiation response mechanisms associated with the synthesis and degradation of structured proteins and the expression of anti-apoptotic factors or other survival signals. Later studies confirmed these findings at clinically relevant doses.^{24,34} In addition, a study by Van Nest *et al.* used Raman spectroscopy to demonstrate cellular response was observed as early as 2 hours following 15 Gy irradiation and Raman detectable differences remained even 10 days after treatment.³⁶

Previous studies have also shown that Raman spectroscopy could be used to measure changes to cellular glycogen levels that occurred following exposure to X-ray radiation.^{24,30,31,34,36} These studies demonstrated a radiation dose dependent accumulation of glycogen in cancer cells following X-ray irradiation. However, this response was dependent on tumour origin, sample type and the inherent radiosensitivity of the cells. The authors reported that glycogen accumulation was only observed for radiation resistant tumour cell lines compared to radiation sensitive tumour cell lines. Glycogen metabolism can be related to a host of signalling pathways associated with tumour progression. Furthermore, studies have suggested a link between glycogen accumulation and hypoxia³⁸ as intracellular glycogen is thought to protect tumour cells against hypoxia and other forms of cellular stress.³⁹

Raman spectroscopy has not been extensively used to assess radiation response in brain tumour cells. An early study by Lakshmi *et al.* showed the spectral changes of mice brain tissue following irradiation were similar to changes observed when the mice were subject to other external stressors, such as restraint and inoculation.⁴⁰ Further work by Kumar *et al.* used Raman spectral signatures to predict the radiosensitivity of glioma stem-like cells.²⁶ Their study used radiosensitising agents to alter the sensitivity of the cells to external beam radiation (XBR) treatment. Raman spectroscopy was then used to assess the radiation response of the cells between 6 to 48 hours following 8 Gy XBR treatment. This allowed Raman spectroscopy to be used as a method for predicting radiation resistance in the glioma cells.

Raman spectroscopy has been further used as a method to predict radiation resistance in tumour cells by evaluating radiation response.^{26,28,37,41} These studies used Raman spectroscopy to determine that radiation resistant and radiation sensitive cells have varying responses to X-ray radiation exposure. This highlights the potential for Raman spectroscopy to further investigate the factors which effect cellular radiation response such as the phase of the cell cycle during radiation exposure.

Cancer cells contain mutations which can affect growth signalling, allowing the cells to evade apoptosis.^{42,43} The control over proliferation exhibited in normal cells is lost in cancer cells, causing continuous proliferation and therefore continuous progression through the cell cycle. The aim of most cancer treatments is to control and halt the progression of malignant cells through the cell cycle in order to kill the cells.⁴⁴ The position of a cell in the cell cycle is an important internal factor that effects the cellular radiosensitivity, as cellular radiosensitivity is dependent upon stage within the cell cycle (Fig. 1).⁴⁵






Fig. 1 Radiation resistance and sensitivity changes through the cell cycle.^{45–47}

In general, cells display radiation resistance towards the end of the G₁ phase and into the S phase, whereas cells in the G₂ phase and those going through mitosis, will display greater radiosensitivity. In addition, cells may possess greater radiation resistance at the very beginning of the G₁ phase, but this is dependent on the overall length of this phase for specific cell types.⁴⁵

The radiosensitivity of different phases of the cell cycle has been studied by irradiating synchronised populations of cells.^{46–48} However, comparison of the results is difficult due to the variation observed between cell types. Raman spectroscopy has been shown to be useful in the study of radiation response and radiation resistance in cancer cells. In addition, Raman spectroscopy has been used to discriminate between cells at different stages of the cell cycle.^{49–53} Therefore Raman spectroscopy shows promise as a method for the study of cell cycle induced radiation resistance. Here we use Raman spectroscopy to study radiation resistance in UVW human glioma cells to gain a better understanding of the effects of cell cycle synchronization on Raman spectral variation and thereby as a method for predicting radiation resistance.

Materials & methods

Cell culture

This study used the UVW (human glioma) cell line which is a cell line derived in the Beatson Institute for Cancer Research, UK. The cells were cultured in 75 cm³ tissue culture flasks in minimum essential medium (MEM) supplemented with 10% foetal bovine serum (FBS), 200 mmol L⁻¹ L-glutamine, 100 µg mL⁻¹ penicillin-streptomycin and 2 µg mL⁻¹ amphotericin B (Gibco, UK). The cells were incubated at 37 °C in a humidified 5% CO₂ atmosphere. Prior to experiments, UVW cells were seeded at a concentration of 1 × 10⁵ cells into 6-well plates containing CaF₂ coverslips for attachment and incubated at 37 °C in a humidified 5% CO₂ atmosphere for 24 hours until exponential growth phase was reached at 70–80% confluency.

Double thymidine block (DTB) cell synchronisation

A double thymidine block (DTB) treatment was used to synchronise the UVW cells at the G₁/S boundary of the cell cycle.⁵⁴ UVW cells were incubated with cell culture medium containing 2 mM thymidine for 18 hours. The treatment was then removed, the cells were washed with phosphate buffered saline (PBS) and incubated with thymidine-free cell medium for 9 hours. Cells were then treated again with 2 mM thymidine for 18 hours.

X-ray treatment

Cells were irradiated at a 6 Gy dose in the 6-well plate while maintained in MEM. X-ray exposure was performed with an X-RAD 225 X-ray cell irradiation cabinet (Precision X-ray, USA) with a 225 kVp X-ray beam and a dose rate of 2.2 Gy min⁻¹.

Raman spectroscopy

UVW cells were fixed at 1 hour, 4 hours and 24 hours following radiation treatment with 4% paraformaldehyde (PFA) prior to Raman analysis. Individual Raman cells maps were collected using a Renishaw InVia confocal Raman microscope with a 532 nm Ng:YAG laser excitation source (maximum power of 50 mW at laser), 1800 l mm⁻¹ grating, edge filter and a charge-coupled device (CCD) detector and a Leica 50× objective (N.A.: 0.75). Renishaw WiRE 4 series software was used to collect sample spectra cell maps. All spectral maps were collected with 1 second per spot acquisition time, 21 mW laser power at sample (100%), 1 µm X/Y step size and spectral centre of 1300 cm⁻¹. For each sample condition, maps were taken of at least 10 individual cells with at least 900 Raman measurements collected per cell. All experiments were carried out with three independent replicates and in total 43 cell maps were collected per sample group.

Cell cycle analysis

UVW cells were detached from the flask into a single cell suspension and fixed with 70% EtOH. Cells were then incubated with 50 µg mL⁻¹ ribonuclease A (RNase A) and 10 µg mL⁻¹ propidium iodide (PI) to degrade the ribonucleic acid (RNA) and stain cellular DNA, respectively (full details in ESI†). Samples were incubated in a light-free environment at 4 °C for at least one hour prior to analysis. All flow cytometry experiments were performed using an Attune NxT flow cytometer with Attune NxT Software (Version 3.1.2). For cell cycle analysis, cells were analysed with 100 µl acquisition volume and a 100 µl min⁻¹ flow rate with a minimum of 10 000 events analysed. Cell fluorescence was assessed using a blue 488 nm 50 mW laser and forward scatter (FCS), side scatter (SSC) and BL2 detectors (574/26 channel). Detector voltages were 140 V, 300 V and 320 V for FSC, SSC and BL2, respectively.

γ-H2AX assay

UVW cells were detached from the flask into a single cell suspension and fixed with 4% PFA. UVW cells were permeabilised using Triton-X-100, resuspended in blocking buffer (0.1%



Triton-X-100 and 0.5% bovine serum albumin (BSA) in PBS) and incubated with 100 $\mu\text{g mL}^{-1}$ FITC-conjugated anti-phospho-histone H2AX (SER139) antibody (Millipore, UK) for 20 minutes at 4 °C, full details in the ESI.† Cells were then analysed by flow cytometry with a 90 μL acquisition volume and a 200 $\mu\text{L min}^{-1}$ flow rate with a minimum of 10 000 events analysed. Cell fluorescence was assessed using a blue 488 nm 50 mV laser and forward scatter (FCS), side scatter (SSC) and BL1 detector (530/30 channel). Detector voltages were 90 V, 300 V and 265 V for FSC, SSC and BL1, respectively.

Data processing

The Raman spectra collected were first pre-processed using WiRE 4.4 software to remove cosmic rays from the spectra using the built-in cosmic ray removal function. MATLAB software (R2017a) was used to further process the Raman spectra, using in-house MATLAB scripts. This software was used for noise reduction (nonlinear iterative partial least squares (NIPALS) decomposition), *x*-axis standardisation, baseline correction (rolling-circle filter (RCF) with radius 150 units), quality control and area under the curve (AUC) normalisation. The spectra were also truncated to 900–1770 cm^{-1} and the spectra were *x*-axis normalised to the phenylalanine peak at 1004 cm^{-1} . The pre-processed spectra from each sample group were averaged and principal component analysis (PCA) was then performed using the in-built MATLAB PCA function. Random forest (RF) modelling was performed on PCA scores using the open source randomForest package in R software (Version 3.6.1). Variable importance was assessed using GINI index. GINI is a measure of how each variable contributes to the homogeneity of the nodes in the random forest. For example, each time a particular variable is used to split a node, the boundary value set on this variable will influence the resultant purity of the data split using this node. Variables which have a high value GINI are important factors in classifying the data correctly. ANOVA statistical analysis was carried out using GraphPad Prism 8 software (Version 8.4.2, GraphPad Software Inc., USA) with Bonferroni post-tests at a 95% confidence interval (CI) for the cell cycle data and with Wilcoxon rank sum test at 99% CI for PCA loadings. Full details of the data processing methods are described in the ESI.†

Results & discussion

The position of cells in the cell cycle is known to influence the radiation response of cancer cells.^{45–47} Despite this, synchronising the cell cycle prior to the study of radiation response has not been extensively researched when assessing Raman spectroscopy as a method for monitoring cellular radiation response. Predicting the differences in radiation response resulting from different factors, including cell cycle variation, is an important consideration to patient tailored radiation therapy. In most types of cancer therapy, for example chemotherapy and radiation therapy, a typical treatment outcome is cell cycle arrest, leading to cell cycle synchronisation. As treat-

ment is typically given over multiple fractions or doses it is important to understand how variation in cell cycle can affect treatment response. Raman spectroscopy has previously been shown to be useful as a method for monitoring cellular radiation response.^{22–31} Here we aimed to compare the radiation response of unsynchronised UVW cells and UVW cells synchronised to the G₁/S boundary prior to radiation treatment using Raman spectroscopy. In doing so we aimed to determine whether Raman spectroscopy could be used as a tool for monitoring subtle cellular changes associated with cell cycle variation and radiation response within the first 24 hours following radiation treatment.

Cell cycle synchronisation can be used to harmonise a large population of cells at a particular stage of the cell cycle. Different methods of cell cycle synchronisation have been previously described including serum starvation,^{55,56} contact inhibition^{52,57} and mitotic selection.⁵² Cell synchronisation can also be achieved by the addition of agents such as thymidine,^{52,54} nocodazole⁵² and aphidicolin.⁵⁷ For this study, four different synchronisation methods were investigated. These methods synchronised cells to the G₁/S boundary by double thymidine block (DTB) treatment⁵⁴ (Fig. S1B†), to the S phase by DTB treatment followed by fresh medium reincubation^{52,54} (Fig. S1C†), to the G₂/M boundary by thymidine and nocodazole treatment^{52,54} (Fig. S1D†) and to the early G₁ phase by thymidine and nocodazole treatment followed by fresh medium reincubation^{52,54} (Fig. S1E†). The full details of these synchronisation methods are shown in the ESI.† However, the most successful method was the DTB treatment. Thymidine reversibly incorporates into the DNA of cells preventing the cell from progressing from the G₁ phase to the S phase of the cell cycle since it prevents DNA replication from occurring.⁵⁴ The cells therefore become synchronised at the G₁/S boundary of the cell cycle, which has been demonstrated to render the cells more radioresistant.⁴⁵ Cell cycle analysis using fluorescence activated cell sorting (FACS) demonstrated that following treatment with the DTB, 71% of cells occupied the G₁ phase of the cell cycle suggesting successful synchronisation (ESI – Fig. S1†). This was in comparison to a control, unsynchronised, cell population where approximately 50% of the cells were present in the G₁ phase of the cell cycle.

Both unsynchronised and synchronised cells were then treated with 6 Gy XBR before samples were collected and fixed at 1 hour, 4 hours and 24 hours following radiation treatment for cell cycle analysis, Raman measurements and γ -H2AX analysis. First, cell cycle analysis was used to confirm the cell cycle distribution of the control and 6 Gy groups irradiated at each time point. The statistical significance of changes to cell cycle distribution, comparing the control to the irradiated cells, was obtained using two-way ANOVA analysis.

The cell cycle distribution of the unsynchronised cells is shown in Fig. 2A. The unirradiated control sample at each time point showed a distribution of cells which would be considered normal for an asynchronous population with 59% in G₀/G₁ and 30% in G₂/M phase. This distribution of the UVW cells in the different phases of the cell cycle did not change






Fig. 2 Cell cycle analysis of (A). unsynchronised UVW cells, and (B). synchronised UVW cells. FACS results comparing cell cycle distribution of control and 6 Gy irradiated cells at 1 hour, 4 hours and 24 hours following irradiation. The coloured bars represent percentage of cells in sG₁ phase (pink), G₀/G₁ phase (blue), S phase (grey) and G₂/M phase (yellow). Data presented as an average of three independent replicates (mean $\pm$ standard deviation). Two-way ANOVA compared the mean cell cycle phase population at each time point following irradiation to the untreated control cells. Statistical analysis was performed using two-way ANOVA with Bonferroni post-tests at 95% confidence interval ($p > 0.05$ = ns (not significant) and $p < 0.0001$ = ****).

over the 24 hours assessed. On the other hand, the cells which were irradiated show a distinct change in cell cycle distribution over the 24 hours following irradiation. At the 1-hour time point the 6 Gy sample had a cell cycle distribution similar to that of the unirradiated control cells. After 4 hours following irradiation the 6 Gy sample showed a small, but not statistically significant, decrease in G₀/G₁ population and an increase in G₂/M population compared to the control ($p > 0.05$). At the 24-hour time point there was a statistically significant increase in G₂/M population between the control and irradiated samples ($p < 0.0001$), with 81% of cells occupying the G₂/M phase of the cell cycle. This confirmed that the UVW cells exhibited cell cycle arrest in the G₂/M phase 24 hours following radiation exposure.

The cell cycle distribution of the synchronised cells was then assessed using FACS analysis. These cells were first synchronised at the G₁/S boundary by DTB treatment before exposure to 6 Gy XBR. The cell cycle progression of the synchronised cells is shown in Fig. 2B. At the 1-hour time point, approximately 80% of both the control and irradiated cells were in the G₀/G₁ phase of the cell cycle which confirmed that the synchronisation had been successful. At the 4-hour time point the cells showed that the distribution of cells had changed. Now the cells exhibited a lower population in the G₀/G₁ phase and a higher population in the G₂/M phase demonstrating that cells had begun to move through the cell cycle. Finally, at the 24-hour time point the percentage of the control cell population in the G₀/G₁ phase had increased to 58% indicating that cells had continued to progress through the cell cycle. At this time point the cells displayed a distribution similar to the unsynchronised control cells observed in Fig. 2A. This could suggest that following G₁/S boundary synchronisation by DTB treatment, the effects are short lived and cells become unsynchronised in the 24 hours following treatment. This is likely to be a result of the variation in the rate at


which cells progress through the cell cycle when they are released from thymidine treatment. This is one factor that limits the use of cell cycle synchronisation as a method for the study of the different phases of the cell cycle. After 24 hours there was a statistically significant increase in the population of 6 Gy irradiated cells in the G₂/M phase (72%) ($p < 0.0001$) which indicated that cells had arrested at the G₂/M boundary because of radiation exposure.

Both the unsynchronised and synchronised UVW cells exhibited cell cycle arrest in the G₂/M phase 24 hours following irradiation of the cells. The passage of cells to the next phase of the cell cycle is regulated at checkpoints throughout the cycle. This ensures that the DNA integrity is maintained preventing genetic flaws to be passed on to the next generation.⁵⁸ Two checkpoints detect DNA damage in a cell, one at the G₁/S transition and one at the G₂/M transition.⁵⁹ The checkpoint at the G₂/M boundary prevents a cell entering mitosis if it is compromised. The extent of the G₂/M arrest was significantly different between the unsynchronised and synchronised sample groups ($p < 0.05$). For the unsynchronised cells, 81% of the population occupied the G₂/M phase 24 hours following radiation exposure compared to only 72% occupying the G₂/M phase for the synchronised cells. This showed that in the synchronised cells fewer cells were experiencing arrest before mitosis and more cells were able to move through to the G₁ phase. This suggested that the cells were more viable when irradiated following synchronisation at the G₁/S boundary of the cell cycle which could suggest less DNA damage was induced. This confirmed that these cells display radioresistance in the G₁/S phase of the cell cycle.⁴⁵

In combination with FACS analysis, Raman spectroscopy was assessed as a method for monitoring cellular changes associated with radiation response immediately following radiation exposure. Raman analysis was used to collect individual cell maps of both unsynchronised and synchronised UVW cells collected and fixed 1 hour, 4 hours and 24 hours following treatment (control and 6 Gy). Average Raman spectra were produced by averaging all spectra from three independent replicates totaling 43 cell maps per treatment group; each map contained at least 900 individual cell spectra. The average Raman spectra of each sample group are shown in Fig. 3 and Raman peak assignments are shown in Table 1.

The spectra of unsynchronised cells show variation in Raman intensity between the control and irradiated samples at all time points (Fig. 3). The greatest variation was observed at 1-hour and 4 hours post irradiation. Differences in signal intensity were observed between the control and 6 Gy cells for Raman peaks at 1004 cm⁻¹ and 1032 cm⁻¹ assigned to phenylalanine, 1320 cm⁻¹ and 1340 cm⁻¹ assigned to amide III and 1555 cm⁻¹ and 1656 cm⁻¹ assigned to tryptophan and amide I (Table 1). The least variation was seen in the 24-hour sample (Fig. 3, right upper panel), where the control and 6 Gy samples spectra were the most similar although small differences in signal intensity between the control and 6 Gy samples were observed at 1555 cm⁻¹ and 1656 cm⁻¹, associated with tryptophan and amide I.





Wavenumber	Assignment
963 cm ⁻¹	Protein, tyrosine ⁶¹
1004 cm ⁻¹	Phenylalanine ²³
1032 cm ⁻¹	C-H in-plane bending mode of phenylalanine ^{22,52}
1056 cm ⁻¹	Lipid ⁶²
1065 cm ⁻¹	Protein C-N stretch, skeletal C-C stretch of lipids ⁶³
1235 cm ⁻¹	Amide III ^{64,65}
1270 cm ⁻¹	Phospholipids, amide II (proteins), C-N alpha helix proteins, C=C fatty acids ^{60,66}
1320 cm ⁻¹	Guanine, amide III ^{22,34}
1330 cm ⁻¹	DNA, phospholipids ⁶⁷
1340 cm ⁻¹	Amide III & CH ₂ wagging vibrations (glycine backbone, proline side chain), C-H deformation (protein) ^{22,52,68}
1427 cm ⁻¹	CH ₂ and CH ₃ bending of methyl bonds in nucleic acids
1445 cm ⁻¹	Collagen, phospholipids, CH ₂ bending modes (protein & lipids), CH ₂ deformation ^{22,65}
1555 cm ⁻¹	Amide II, tryptophan ⁶⁰
1596 cm ⁻¹	Amide I of protein (C=O stretching) ⁶⁹
1614 cm ⁻¹	Tyrosine, tryptophan ^{22,52}
1631 cm ⁻¹	Amide III ⁶⁴
1656 cm ⁻¹	Amide I ν (C=O), C=C lipids, tryptophan ^{22,52}

This variation was observed with differences in peak intensities between the samples at 1004 cm^{-1} and 1656 cm^{-1} indicating different expression of phenylalanine and amide I signals. Variation between the control and irradiated sample spectra were still present in the 24-hour sample (Fig. 3). The sample spectra showed differences in peak intensities between the control and 6 Gy samples at 1445 cm^{-1} and 1656 cm^{-1} . This demonstrated variation in collagen, phospholipids, tryptophan and amide I at this time point.^{37,70}

Overall, the average cell spectra comparing the control and 6 Gy irradiated cells suggested that there was a difference between the sample spectra at each time point for both unsynchronised and synchronised cells. However, it was unclear if the variation observed was solely caused by radiation response in the cells or if there were contributions from differences in cell cycle distribution. Principal component analysis (PCA) combined with random forest (RF) classification and variable selection, was used to evaluate the variation associated with radiation response. The sample spectra (Fig. 3) showed a large feature at 1555 cm^{-1} which was assigned to instrumental background. A Raman peak at 1555 cm^{-1} in cell spectra could be assigned to tryptophan/amide I (Table 1). A study by Van Nest *et al.* reported changes to the tryptophan signal (1555 cm^{-1}) in the Raman spectra of non-small cell lung cancer (NSCLC) tissue xenografts as a result of radiation response to 15 Gy XBR.³⁶ Furthermore, tryptophan metabolism has also been associated with radiation-induced immunosuppression involving immune checkpoint reactivation.⁷⁰ However, the appearance of the peak at 1555 cm^{-1} was very sharp (Fig. 3) and this feature was dominant in the Raman spectrum. These characteristics would be unusual in a cell spectrum, therefore this feature was assigned to instrumental background as off-

PCA was then performed on the 4-hour time point data alone, the first ten PC loadings are shown in the ESI (Fig. S6†). The RF model, which was used to classify the data as control *vs.* 6 Gy irradiated for each group (unsynchronised and synchronised), showed that for the 4-hour time point the OOB error was $40 \pm 3\%$ therefore showing better classification com-

Fig. 5 Principal component analysis (PCA) of average Raman spectra for just the 4-hour time point showing the box plots and PC loadings for PC5 and PC1. Box plot compares control cells and 6 Gy irradiated cells for unsynchronised UVW cells (blue) and synchronised UVW cells (pink). Centre point of box represents median value, notches represent the 25th and 75th percentile, whiskers represent the 5th and 95th percentile and stars represent outliers. PC5 and PC1 loading from individual time point PCA comparing control and 6 Gy samples for the 24-hour time point following 6 Gy XBR exposure for unsynchronised and synchronised UVW cells. Statistical analysis was performed using a two-way ANOVA with Wilcoxon rank sum test at 99% confidence interval ($p > 0.05 = \text{ns}$ (not significant), $p < 0.01 = **$ and $p < 0.001 = ***$).

difference between the control and 6 Gy irradiated samples in the unsynchronised sample groups ($p < 0.01$). The PC2 loading was almost identical to the PC3 loading from the PCA described for the data all together (Fig. 4) and were shown to have high positive correlation ($r = 0.95$) using Pearson's linear correlation coefficient. This showed features at 1004, 1065, 1270, 1320, 1427, 1596 and 1656 cm^{-1} (Table 1). Overall, the features were predominantly from amides associated with proteins and some contribution from lipids. These results showed that at the 4-hour time point, PCA was able to determine a significant difference in the radiation response of the synchronised but not in the unsynchronised cells. This could suggest that by removing the variation in sample spectra caused by cell cycle position, we are able to evaluate the radiation response as early as 4 hours following XBR treatment.

The significant difference between the control and irradiated samples in the synchronised cells could be explained by the positioning in the cell cycle at the point of radiation exposure. The response to radiation exposure varies depending on the phase of the cell cycle in which the cell occupies.^{46–48} As the cells are synchronised in the same phase, this removes variation in response relating to position in the cell cycle. However, in the unsynchronised cells the degree of radiation response will differ depending on the phase in which they were irradiated. Thus, at an early time point, like 4 hours after radiation, the variation in the unsynchronised cells may lead to less successful classification.

UVW cells have a doubling time of approximately 29 hours.⁷⁷ The longest phase of the cell cycle is the G_1 phase and may last up to 12 hours, the S phase will last between 5–8 hours and the G_2 phase will be shorter at 4–5 hours in length.⁷⁸ Finally, mitosis (M) lasts only around 1–2 hours. Cell cycle analysis showed that at the 4-hour time point, a large proportion of the synchronised cell population (>50%) were in the G_2/M phase of the cell cycle (Fig. 2B). This confirmed that at this time point the cells had moved through the S phase. The mechanisms of DNA damage and repair will ultimately lead to cell cycle arrest at the G_2/M boundary of the cell cycle.⁷⁹ We observed in both the unsynchronised and synchronised sample groups that by 24 hours post radiation a large population of cells had reached cell cycle arrest at G_2/M (Fig. 2). When cells were synchronised to the G_1/S boundary prior to irradiation, the majority will reach the G_2/M phase quicker and therefore experience an earlier cell cycle arrest. This could result in a more prominent difference in radiation response at the 4-hour time point compared to the unsynchronised cells. For the unsynchronised cells, at the 4-hour time point a large proportion of the cells (>50%) remain in the G_1 phase of the cell cycle (Fig. 2A). At this stage, these cells may not have reached the checkpoint prior to the S phase, therefore radiation response, such as DNA repair, may not have been instigated. The degree of radiation response in these cells is likely to differ to cells at a later stage of the cell cycle. This could explain why a larger radiation response is observed in the synchronised cells 4 hours post irradiation and by synchronising the cells to the G_1/S boundary we are inducing a quicker radiation response.

PCA was then performed on the 24-hour time point data alone, the first ten PC loadings are shown in the ESI (Fig. S6†). Using the RF model, the 24-hour time point once again showed the best OOB error following classification of control *versus* 6 Gy irradiated cells for both unsynchronised and synchronised groups with an error of $35 \pm 4\%$. This showed that the model was able to classify the sample correctly into treatment groups 65% of the time. The ten most important PC loadings found by the RF model for classification are shown in the ESI (Fig. S6†). The RF model found PC5 loading as the most important variable for classifying the samples into each group, followed by PC1. A Tukey box plot was used to show the difference between the sample groups for the PC5 loading, the box plot is shown in Fig. 6. The PC5 loading box plot showed a significant difference between the control and irradiated cells in the unsynchronised cells ($p < 0.0001$) but no significant difference between the control and irradiated cells in the synchronised sample group ($p > 0.05$). The PC5 loading had Raman features assigned to protein (963 cm^{-1} & 1429 cm^{-1}), amide III (1235 cm^{-1} & 1631 cm^{-1}), DNA and phospholipids (1330 cm^{-1}) (Table 1). A Tukey box plot was used to show the difference between the sample groups for the PC1 loading, the box plot is shown in Fig. 6. The PC1 box plot did not show a significant difference between the control and irradiated cells for either the unsynchronised or synchronised samples ($p > 0.05$), therefore this PC loading was not used for evaluation of radiation response (Fig. 6).





Fig. 6 Principal component analysis (PCA) of average Raman spectra for just the 24-hour time point showing the box plots and PC loadings for PC5 and PC1. Box plot compares control cells and 6 Gy irradiated cells for unsynchronised UVW cells (blue) and synchronised UVW cells (pink). Centre point of box represents median value, notches represent the 25th and 75th percentile, whiskers represent the 5th and 95th percentile and stars represent outliers. PC5 and PC1 loading from individual time point PCA comparing control and 6 Gy samples for the 24-hour time point following 6 Gy XBR exposure for unsynchronised and synchronised UVW cells. Statistical analysis was performed using a two-way ANOVA with Wilcoxon rank sum test at 99% confidence interval ($p > 0.05 = \text{ns}$ (not significant) and $p < 0.0001 = \text{****}$).



The γ -H2AX levels were assessed at the same time points following irradiation as the Raman analysis for control and 6 Gy irradiated cells in both unsynchronised and synchronised sample groups (Fig. 7). The unsynchronised cells had a $\sim$ 14-fold increase in γ -H2AX levels at 1-hour following irradiation ($p > 0.05$) and a significant 31-fold increase in γ -H2AX levels 4 hours following irradiation compared to the non-irradiated control cells ($p < 0.005$). At 24 hours following irradiation, there were lower γ -H2AX levels, $\sim$ 6-fold change in γ -H2AX levels, in the irradiated samples compared to the non-irradiated control ($p > 0.05$). This showed that DNA damage increased within the first 4 hours following irradiation and then by the 24-hour time point DNA damage had been resolved. This decrease in DNA damage could be a result of DNA repair mechanisms being activated in the first 24 hours following irradiation or that cells which had sustained unreparable damage were removed from the sample population due to apoptosis. The standard deviation observed from these



The γ -H2AX results were compared to the results from the Raman analysis (Fig. 3–6), but the Raman analysis did not show a similar trend across the three time points. The Raman analysis showed changes primarily in cellular proteins and

lipid whereas changes to DNA signal were not observed. Raman spectra from cells is dominated by protein and lipid signals, therefore, it may be expected that subtle changes to DNA, such as damage induced by radiation exposure, would not be observed. The γ -H2AX analysis for the unsynchronised cells showed largest DNA damage at 4 hours post irradiation (Fig. 7A). This then subsequently decreased by the 24-hour time point, suggesting DNA repair took place between 4 hours and 24 hours post radiation. The mechanism of DNA damage repair requires a large number of proteins responsible for double strand break detection, chromatin remodeling and DNA repair therefore it would be expected that the expression of proteins across the cell would change as the DNA is being repaired.⁸³ The radiation response observed by Raman spectroscopy with PCA was significant at 24 hours in PC5 of the unsynchronised cells (Fig. 6). The main features present in the PC5 loading for the 24-hour time point were protein, amide III, DNA and phospholipids. Thus, the changes observed at this time point could account for the proteins required to facilitate the DNA repair between the 4-hour and 24-hour time point. A key feature that was shown to contribute to radiation response in some previous studies was glycogen.^{24,30,31,34,36} However, in this work no glycogen features were observed. Most previous studies observed glycogen accumulation 24 hours or longer after irradiation, although following a dose of 15 Gy Van Nest *et al.* did report glycogen accumulation as early as 2-hours following irradiation, although the dose used was much higher than that used in this study.³⁶ Additionally, the most commonly assigned glycogen peak is around 473 cm^{-1} which was outside of the Raman range analysed in this study. The results presented could suggest that the first 24 hours after radiation exposure may be too soon to observe glycogen accumulation for UVW cells. Glycogen accumulation is also linked to radiation resistance in cells and therefore may not be present in cells which display radiosensitivity.²⁴ The cell survival fraction (SF) describes the fraction of cells which maintain reproductive integrity following radiation exposure at a particular dose (*e.g.* 2 Gy – $\text{SF}_2\text{ Gy}$). Although not extensively studied, previous work considered UVW cells to be moderately radioresistant ($\text{SF}_2\text{ Gy} = 0.56$),⁸⁴ therefore the lack of glycogen accumulation in these cells contradicts previous studies which link glycogen accumulation to a cell's radiation resistance.^{24,30,31,34,36} The previous study by Van Nest *et al.* (2018) also did not find glycogen accumulation following radiation treatment of human breast adenocarcinoma xenografts. Their study, in agreement with this study, found spectral changes related to proteins including, phenylalanine and amide bands following irradiation, however these were not linked to specific biological processes.³⁵ This study of radiation response in UVW cells presents similar findings with changes to protein features in the Raman spectra. Using γ -H2AX analysis in combination with Raman analysis was not conclusive in allowing similarities to be drawn between the spectral response and the level of DNA damage. However, the results suggested that the cellular protein changes observed at the 24-hour time point could be associated with the DNA

damage pathways and repair mechanisms. Additionally, this work demonstrated that radiation resistance could be induced by controlling the cell cycle prior to radiation exposure which is a key consideration of cancer therapy and that by using Raman spectroscopy the differences in radiation response could be assessed.

Conclusion

In this study Raman spectroscopy was used to assess radiation response in UVW human glioma cells cultured *in vitro*. The sensitivity of Raman spectroscopy for detection of subtle biochemical changes within the cells allowed evaluation of the early stages of radiation response as early as four hours following 6 Gy XBR treatment. The Raman spectral changes observed following radiation exposure were assigned to specific amino acids associated with proteins, including phenylalanine, and protein signals from amides. Using Raman spectroscopy in combination with γ -H2AX analysis suggested the protein changes observed in the synchronised cells were unlikely directly associated with the DNA damage and repair mechanisms of the cells. However, the protein changes observed, particularly between 4 hours and 24 hours post radiation treatment in unsynchronised cells could be associated with DNA repair pathways. Although Raman spectroscopy has been used previously to assess radiation response, few studies focus on the initial effects of this treatment compared to the longer-term radiation induced changes in cells. In the clinic, radiation treatment is often given daily in fractionated doses therefore monitoring the effects even after a short time, for example the first 24 hours following treatment, is equally important in assessing the long-term effects. In addition, cancer treatment often causes cell cycle arrest and therefore induces cell cycle synchronisation. Changes to the cell cycle during the course of a treatment regime will therefore influence the patients' response, highlighting cell cycle position as an important factor to consider and monitor when studying radiation response. Raman spectroscopy was used to demonstrate the difference in radiation response between cells occupying different phases of the cell cycle at the time of irradiation. This work was able to show different levels of radiation resistance between cells and confirm that cell cycle variation influences the response of UVW cells to radiation exposure. To the best of our knowledge, this is the first report using Raman spectroscopy to monitor radiation response in synchronised cells and the results suggested that cell synchronisation is an important factor to take into account in future studies. By evaluating factors that induce radiation resistance in cells, a better understanding can be gained about radiation treatment. In addition, Raman spectroscopy has been shown to be a useful tool in differentiating different degrees of treatment response, which is important when investigating patient tailored radiation therapy. This highlights a need to further investigate the radiation response of cells at different phases of the cell cycle and other factors which induce radiation resistance.



- 23 Q. Matthews, A. Jirasek, J. J. Lum and A. G. Brolo, Biochemical signatures of in vitro radiation response in human lung, breast and prostate tumour cells observed with Raman spectroscopy, *Phys. Med. Biol.*, 2011, **56**, 6839.
- 24 Q. Matthews, M. Isabelle, S. J. Harder, J. Smazynski, W. Beckham, A. G. Brolo, A. Jirasek and J. J. Lum, Radiation-Induced Glycogen Accumulation Detected by Single Cell Raman Spectroscopy Is Associated with Radioresistance that Can Be Reversed by Metformin, *PLoS One*, 2015, **10**, e0135356.
- 25 A. G. Brolo, A. Jirasek, J. J. Lum, M. Isabelle, Q. Matthews and S. J. Harder, A Raman Spectroscopic Study of Cell Response to Clinical Doses of Ionizing Radiation, *Appl. Spectrosc.*, 2015, **69**(2), 193–204.
- 26 S. Kumar, A. Visvanathan, A. Arivazhagan, V. Santhosh, K. Somasundaram and S. Umapathy, Assessment of Radiation Resistance and Therapeutic Targeting of Cancer Stem Cells: A Raman Spectroscopic Study of Glioblastoma, *Anal. Chem.*, 2018, **90**, 12067–12074.
- 27 S. Qiu, Y. Weng, Y. Li, Y. Chen, Y. Pan, J. Liu, W. Lin, X. Chen, M. Li, T. Lin, W. Liu, L. Zhang and D. Lin, Raman profile alterations of irradiated human nasopharyngeal cancer cells detected with laser tweezer Raman spectroscopy, *RSC Adv.*, 2020, **10**, 14368–14373.
- 28 K. Pansare, S. R. Singh, V. Chakravarthy, N. Gupta, A. Hole, P. Gera, R. Sarin and C. M. Krishna, Raman Spectroscopy: An Exploratory Study to Identify Post-Radiation Cell Survival, *Appl. Spectrosc.*, 2020, **74**(5), 553–562.
- 29 M. Roman, T. P. Wrobel, A. Panek, C. Paluszkiwicz and W. M. Kwiatek, Exploring subcellular responses of prostate cancer cells to clinical doses of X-rays by Raman microspectroscopy, *Spectrochim. Acta, Part A*, 2021, **255**, 119653.
- 30 X. Deng, R. Ali-Adeeb, J. L. Andrews, P. Shreeves, J. J. Lum, A. Brolo and A. Jirasek, Monitor Ionizing Radiation-Induced Cellular Responses with Raman Spectroscopy, Non-Negative Matrix Factorization, and Non-Negative Least Squares, *Appl. Spectrosc.*, 2020, **74**, 701–711.
- 31 K. Milligan, X. Deng, P. Shreeves, R. Ali-Adeeb, Q. Matthews, A. Brolo, J. J. Lum, J. L. Andrews and A. Jirasek, Raman spectroscopy and group and basis-restricted non negative matrix factorisation identifies radiation induced metabolic changes in human cancer cells, *Sci. Rep.*, 2021, **11**, 3853.
- 32 H. J. Butler, L. Ashton, B. Bird, G. Cinque, K. Curtis, J. Dorney, K. Esmonde-White, N. J. Fullwood, B. Gardner, P. L. Martin-Hirsch, M. J. Walsh, M. R. McAinsh, N. Stone and F. L. Martin, Using Raman spectroscopy to characterize biological materials, *Nat. Protoc.*, 2016, **11**, 664–687.
- 33 I. Pence and A. Mahadevan-Jansen, Clinical instrumentation and applications of Raman spectroscopy, *Chem. Soc. Rev.*, 2016, **45**, 1958–1979.
- 34 S. J. Harder, M. Isabelle, L. Devorkin, J. Smazynski, W. Beckham, A. G. Brolo, J. J. Lum and A. Jirasek, Raman spectroscopy identifies radiation response in human non-small cell lung cancer xenografts, *Sci. Rep.*, 2016, **6**, 21006.
- 35 S. J. van Nest, L. M. Nicholson, L. Devorkin, A. G. Brolo, J. J. Lum and A. Jirasek, Raman Spectroscopic Signatures Reveal Distinct Biochemical and Temporal Changes in Irradiated Human Breast Adenocarcinoma Xenografts, *Radiat. Res.*, 2018, **189**, 497–504.
- 36 S. J. van Nest, L. M. Nicholson, N. Pavey, M. N. Hindi, A. G. Brolo, A. Jirasek and J. J. Lum, Raman spectroscopy detects metabolic signatures of radiation response and hypoxic fluctuations in non-small cell lung cancer, *BMC Cancer*, 2019, **19**, 474.
- 37 S. K. Paidi, P. M. Diaz, S. Dadgar, S. V. Jenkins, C. M. Quick, R. J. Griffin, R. P. M. Dings, N. Rajaram and I. Barman, Label-Free Raman Spectroscopy Reveals Signatures of Radiation Resistance in the Tumor Microenvironment, *Cancer Res.*, 2019, **79**, 2054–2064.
- 38 J. Pelletier, G. Bellot, P. Gounon, S. Lacas-Gervais, J. Pouysségur and N. M. Mazure, Glycogen synthesis is induced in hypoxia by the hypoxia-inducible factor and promotes cancer cell survival, *Front. Oncol.*, 2012, **2**, 18.
- 39 M. Z. Ibrahim, E. Pascoe, S. Alam and J. Miquel, Glycogen and phosphorylase activity in rat brain during recovery from several forms of hypoxia, *Am. J. Pathol.*, 1970, **60**, 403.
- 40 R. J. Lakshmi, V. B. Kartha, C. M. Krishna, J. G. R. Solomon, G. Ullas and P. U. Devi, Tissue Raman spectroscopy for the study of radiation damage: brain irradiation of mice - PubMed, *Radiat. Res.*, 2002, **157**, 175–182.
- 41 M. S. Vidyasagar, K. Maheedhar, B. M. Vadhiraja, D. J. Fernandes, V. B. Kartha and C. M. Krishna, Prediction of radiotherapy response in cervix cancer by Raman spectroscopy: A pilot study, *Biopolymers*, 2008, **89**, 530–537.
- 42 D. Hanahan and R. A. Weinberg, The Hallmarks of Cancer, *Cell*, 2000, **100**, 57–70.
- 43 D. Hanahan and R. A. Weinberg, Hallmarks of cancer: The next generation, *Cell*, 2011, **144**, 646–674.
- 44 M. H. Barcellos-Hoff, C. Park and E. G. Wright, Radiation and the microenvironment – tumorigenesis and therapy, *Nat. Rev. Cancer*, 2005, **5**, 867–875.
- 45 W. K. Sinclair, Cysteamine: Differential X-ray Protective Effect on Chinese Hamster Cells During the Cell Cycle, *Science*, 1968, **159**, 442–444.
- 46 D. W. Siemann and P. C. Keng, In situ radiation response and oxygen enhancement ratio of KHT sarcoma cells in various phases of the cell cycle, *Br. J. Radiol.*, 1984, **57**, 823–827.
- 47 G. D. Wilson, Radiation and the cell cycle, revisited, *Cancer Metastasis Rev.*, 2004, **23**, 209–225.
- 48 K. Otani, Y. Naito, Y. Sakaguchi, Y. Seo, Y. Takahashi, J. Kikuta, K. Ogawa and M. Ishii, Cell-cycle-controlled radiation therapy was effective for treating a murine malignant melanoma cell line in vitro and in vivo, *Sci. Rep.*, 2016, **6**, 30689.
- 49 J. R. Mourant, K. W. Short, S. Carpenter, N. Kunapareddy, L. Coburn, T. M. Powers and J. P. Freyer, Biochemical differences in tumorigenic and nontumorigenic cells measured by Raman and infrared spectroscopy, *J. Biomed. Opt.*, 2005, **10**, 031106.



- 50 K. W. Short, S. Carpenter, J. P. Freyer and J. R. Mourant, Raman Spectroscopy Detects Biochemical Changes Due to Proliferation in Mammalian Cell Cultures, *Biophys. J.*, 2005, **88**, 4274–4288.
- 51 R. J. Swain, G. Jell and M. M. Stevens, Non-invasive analysis of cell cycle dynamics in single living cells with Raman micro-spectroscopy, *J. Cell. Biochem.*, 2008, **104**, 1427–1438.
- 52 A. G. Brolo, A. Jirasek, J. Lum, Q. Matthews and X. Duan, Variability in Raman Spectra of Single Human Tumor Cells Cultured in Vitro: Correlation with Cell Cycle and Culture Confluency, *Appl. Spectrosc.*, 2010, **64**(8), 871–887.
- 53 S. O. Konorov, H. G. Schulze, J. M. Piret, M. W. Blades and R. F. B. Turner, Label-Free Determination of the Cell Cycle Phase in Human Embryonic Stem Cells by Raman Microspectroscopy, *Anal. Chem.*, 2013, **85**, 8996–9002.
- 54 G. Chen and X. Deng, Cell Synchronization by Double Thymidine Block, *Bio-Protoc.*, 2018, **8**, e2994.
- 55 M. Chen, J. Huang, X. Yang, B. Liu, W. Zhang, L. Huang, F. Deng, J. Ma, Y. Bai, R. Lu, B. Huang, Q. Gao, Y. Zhuo and J. Ge, Serum starvation induced cell cycle synchronization facilitates human somatic cells reprogramming, *PLoS One*, 2012, **7**, e28203.
- 56 R. Khammanit, S. Chantakru, Y. Kitiyanant and J. Saikhun, Effect of serum starvation and chemical inhibitors on cell cycle synchronization of canine dermal fibroblasts, *Theriogenology*, 2008, **70**, 27–34.
- 57 J. Jackman and P. M. O'Connor, Methods for Synchronizing Cells at Specific Stages of the Cell Cycle, *Curr. Protoc. Cell Biol.*, 2001, **8**, 8.3.1–8.3.20.
- 58 M. B. Kastan and J. Bartek, Cell-cycle checkpoints and cancer, *Nature*, 2004, **432**, 316–323.
- 59 R. Visconti, R. Della Monica and D. Grieco, Cell cycle checkpoint in cancer: a therapeutically targetable double-edged sword, *J. Exp. Clin. Cancer Res.*, 2016, **35**, 153.
- 60 Z. Movasaghi, S. Rehman and I. U. Rehman, Raman Spectroscopy of Biological Tissues, *Appl. Spectrosc. Rev.*, 2007, **42**, 493–541.
- 61 S. Akbar, M. I. Majeed, H. Nawaz, N. Rashid, A. Tariq, W. Hameed, S. Shakeel, G. Dastgir, R. Z. A. Bari, M. Iqbal, A. Nawaz and M. Akram, Surface-Enhanced Raman Spectroscopic (SERS) Characterization of Low Molecular Weight Fraction of the Serum of Breast Cancer Patients with Principal Component Analysis (PCA) and Partial Least Square-Discriminant Analysis (PLS-DA), *Anal. Lett.*, 2022, **55**, 1588–1604.
- 62 C. Krafft, L. Neudert, T. Simat and R. Salzer, Near infrared Raman spectra of human brain lipids, *Spectrochim. Acta, Part A*, 2005, **61**, 1529–1535.
- 63 S. Qiu, Y. Weng, Y. Li, Y. Chen, Y. Pan, J. Liu, W. Lin, X. Chen, M. Li, T. Lin, W. Liu, L. Zhang and D. Lin, Raman profile alterations of irradiated human nasopharyngeal cancer cells detected with laser tweezer Raman spectroscopy, *RSC Adv.*, 2020, **10**, 14368–14373.
- 64 J. Gala de Pablo, F. J. Armistead, S. A. Peyman, D. Bonthron, M. Lones, S. Smith and S. D. Evans, Biochemical fingerprint of colorectal cancer cell lines using label-free live single-cell Raman spectroscopy, *J. Raman Spectrosc.*, 2018, **49**, 1323–1332.
- 65 I. Notingher, S. Verrier, S. Haque, J. M. Polak and L. L. Hench, Spectroscopic study of human lung epithelial cells (A549) in culture: Living cells versus dead cells, *Biopolymers*, 2003, **72**, 230–240.
- 66 J. Wang, C. X. Zheng, C. L. Ma, X. X. Zheng, X. Y. Lv, G. D. Lv, J. Tang and G. H. Wu, Raman spectroscopic study of cervical precancerous lesions and cervical cancer, *Lasers Med. Sci.*, 2021, **36**, 1855–1864.
- 67 U. Utzinger, D. L. Heintzelman, A. Mahadevan-Jansen, A. Malpica, M. Follen and R. Richards-Kortum, Near-Infrared Raman Spectroscopy for in Vivo Detection of Cervical Precancers, *Appl. Spectrosc.*, 2001, **55**(8), 955–959.
- 68 C. Krafft, T. Knetschke, A. Siegner, R. H. W. Funk and R. Salzer, Mapping of single cells by near infrared Raman microspectroscopy, *Vib. Spectrosc.*, 2003, **32**, 75–83.
- 69 S. Sato, R. Sekine, H. Kagoshima, K. Kazama, A. Kato, M. Shiozawa and J. Tanaka, All-in-one Raman spectroscopy approach to diagnosis of colorectal cancer: analysis of spectra in the fingerprint regions, *J. Anus, Rectum Colon*, 2019, **3**, 84.
- 70 P. Kesarwani, A. Prabhu, S. Kant, P. Kumar, S. F. Graham, K. L. Buelow, G. D. Wilson, C. R. Miller and P. Chinnaiyan, Tryptophan Metabolism Contributes to Radiation-Induced Immune Checkpoint Reactivation in Glioblastoma, *Clin. Cancer Res.*, 2018, **24**, 3632–3643.
- 71 Y. Akagi, N. Mori, T. Kawamura, Y. Takayama and Y. S. Kida, Non-invasive cell classification using the Paint Raman Express Spectroscopy System (PRESS), *Sci. Rep.*, 2021, **11**, 8818.
- 72 J. Immunol, A. Ramoji, O. Ryabchykov, K. Galler, A. Tannert, R. Markwart, R. P. Requardt, I. Rubio, M. Bauer, T. Bocklitz, J. Popp and U. Neugebauer, Raman Spectroscopy Follows Time-Dependent Changes in T Lymphocytes Isolated from Spleen of Endotoxemic Mice, *Immunohorizons*, 2019, **3**, 45–60.
- 73 K. Milligan, S. J. van Nest, X. Deng, R. Ali-Adeeb, P. Shreeves, S. Punch, N. Costie, N. Pavey, J. M. Crook, D. M. Berman, A. G. Brolo, J. J. Lum, J. L. Andrews and A. Jirasek, Raman spectroscopy and supervised learning as a potential tool to identify high-dose-rate-brachytherapy induced biochemical profiles of prostate cancer, *J. Biophotonics*, 2022, **15**(11), e202200121.
- 74 F. Lussier, V. Thibault, B. Charron, G. Q. Wallace and J. F. Masson, Deep learning and artificial intelligence methods for Raman and surface-enhanced Raman scattering, *TrAC, Trends Anal. Chem.*, 2020, **124**, 115796.
- 75 M. Tollefson, J. Magera, T. Sebo, J. Cohen, A. Drauch, J. Maier and I. Frank, Raman spectral imaging of prostate cancer: can Raman molecular imaging be used to augment standard histopathology?, *BJU Int.*, 2010, **106**, 484–488.
- 76 W. C. Chang, On using principal components before separating a mixture of two multivariate normal distributions, *Appl. Stat.*, 1983, **32**, 267–275.



- 77 N. L. McGinley, *Solid and hollow gold nanoparticles as radiosensitisers in combination with X-ray radiation and targeted radiopharmaceuticals*, University of Strathclyde, 2015.
- 78 G. M. Cooper, in *The Cell: A Molecular Approach*, Sinauer Associates, 2nd edn, 2000.
- 79 D. P. Waterman, J. E. Haber and M. B. Smolka, Checkpoint Responses to DNA Double-strand Breaks, *Annu. Rev. Biochem.*, 2020, **89**, 103.
- 80 S. Burma, B. P. Chen, M. Murphy, A. Kurimasa and D. J. Chen, ATM phosphorylates histone H2AX in response to DNA double-strand breaks, *J. Biol. Chem.*, 2001, **276**, 42462–42467.
- 81 H. Zhao, X. Huang, H. D. Halicka and Z. Darzynkiewicz, Detection of Histone H2AX Phosphorylation on Ser-139 as an Indicator of DNA Damage, *Curr. Protoc. Cytom.*, 2019, **89**, e55.
- 82 S. Burma, B. P. Chen, M. Murphy, A. Kurimasa and D. J. Chen, ATM phosphorylates histone H2AX in response to DNA double-strand breaks, *J. Biol. Chem.*, 2001, **276**, 42462–42467.
- 83 G. Borrego-Soto, R. Ortiz-López and A. Rojas-Martínez, Ionizing radiation-induced DNA injury and damage detection in patients with breast cancer, *Genet. Mol. Biol.*, 2015, **38**, 420.
- 84 M. Boyd, A. Livingstone, L. E. Wilson, E. M. Marshall, A. G. McCluskey, R. J. Mairs and T. E. Wheldon, Dose-response relationship for radiation-induced mutations at micro- and minisatellite loci in human somatic cells in culture, *Int. J. Radiat. Biol.*, 2009, **76**, 169–176.

