

PAPER

[View Article Online](#)
[View Journal](#) | [View Issue](#)
Cite this: *Food Funct.*, 2023, **14**, 112

Anti-inflammatory potential of digested *Brassica* sprout extracts in human macrophage-like HL-60 cells†

Paula Garcia-Ibañez,^{a,b} María A. Núñez-Sánchez,^c Alba Oliva-Bolarín,^{ID c}
 María A. Martínez-Sánchez,^{c,d} Bruno Ramos-Molina,^{ID *c}
 Antonio J. Ruiz-Alcaraz,^{ID *d} and Diego A. Moreno,^{ID a}

Cruciferous vegetables have been reported to be a great source of anti-inflammatory compounds. Specifically, sprouts from the Brassicaceae family stand out for their high content of glucosinolates (and their bioactive derivatives, isothiocyanates), phenolic acids, and anthocyanins. Despite the evident anti-inflammatory activity of certain Brassica phytochemicals such as sulforaphane or phenolic acids, the effect of digested Brassica vegetables on inflammation remains understudied. In this work, we aimed to evaluate the anti-inflammatory potential of the bioaccessible forms of cruciferous bioactives (from red cabbage sprouts (RCS) and red radish sprouts (RRS)) obtained upon *in vitro* gastrointestinal digestion in the HL-60 macrophage-like differentiated human cell line. The study was performed under basal conditions or stimulated with a low dose of LPS for 24 hours as a validated *in vitro* model of chronic inflammation. The cell viability was determined by MTT assay. The gene expression and production of pro-inflammatory cytokines TNF- α , IL-6 and IL-1 β were determined by RT-qPCR and ELISA respectively. Our results revealed no cytotoxicity with any of the treatments in LPS-stimulated macrophage-like HL60 cells. Regarding cytokine production, digestates significantly decreased the production of the three pro-inflammatory cytokines at concentrations of 50 and 100 $\mu\text{g mL}^{-1}$ except for IL-1 β treated with RCS digestates. Furthermore, the RT-qPCR analysis showed a decrease in the relative expression of pro-inflammatory cytokines in LPS-stimulated cells treated with RRS digestates at 100 $\mu\text{g mL}^{-1}$ but not with red cabbage digestates. In conclusion, RRS bioaccessible compounds in the extracts could be used as dietary coadjuncts given their potential anti-inflammatory effect on this *in vitro* model of chronic inflammation.

Received 29th September 2022

Accepted 1st December 2022

DOI: 10.1039/d2fo02914f

rsc.li/food-function

1. Introduction

Nowadays, it is widely accepted that the dietary pattern is a key factor in improving human health. A “healthy diet” is defined as one with a high percentage of vegetables and fiber, an adequate content of proteins, and rich in vitamins and micronutrients.¹ In that sense, in the last few decades a high research

effort has been exerted to demonstrate the beneficial health effects of bioactive compounds present in diverse types of foods, with special attention to those coming from fruits and vegetables. Among them, cruciferous vegetables stand out not only for their nutritional value but also for their diverse content of a wide range of bioactive compounds, such as flavonols, anthocyanins and glucosinolates (GSLs). Furthermore, the dietary intake of these molecules has been related to a reduction in the incidence of non-communicable diseases related to inflammatory processes.²

Due to their short time of production and high concentration of health-promoting biomolecules, cruciferous sprouts have emerged as a great option for incorporation in the daily diet.³ Sprouts obtained from Brassicaceae vegetables are one of the main sources of GSLs, which are the main secondary metabolites present in Brassicaceae species. These molecules present a β -thioglucoside-*N*-hydrosulfate bound to a sulfur β -glucopyranose with a variable side chain depending on the precursor amino acid.⁴ However, the bioactivity of GSLs

^aPhytochemistry and Healthy Food Lab, Department of Food Science Technology, Centro de Edafología y Biología aplicada del Segura (CEBAS-CSIC), Campus de Espinardo, 30100 Murcia, Spain

^bAquaporins Research Group, Centro de Edafología y Biología Aplicada del Segura (CEBAS-CSIC), Campus de Espinardo, 30100 Murcia, Spain

^cObesity and Metabolism Laboratory, Biomedical Research Institute of Murcia (IMIB), 30120 Murcia, Spain. E-mail: bruno.ramos@imib.es; Tel: +34 868 885220

^dDepartment of Biochemistry, Molecular Biology B and Immunology, School of Medicine, University of Murcia, Regional Campus of International Excellence, Spain. E-mail: ajruiz@um.es; Tel: +34 868 889521

†Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2fo02914f>



depends on their hydrolysis products after catalyzation by the myrosinase, mainly isothiocyanates (ITCs) and indoles.⁵ Certain *Brassica* sprouts, such as broccoli, contain high levels of glucoraphanin, of which the corresponding ITC is sulforaphane (SFN). SFN has been widely studied for its anti-tumorigenic, anti-oxidative and anti-inflammatory properties.⁶ In addition, the ITC iberin, derived from the glucoiberin present in some sprouts (e.g. red cabbage), has been also reported to be involved in the induction of several anti-inflammatory pathways.⁷ Glucobrassicin (GBS) is also an abundant indolic GSL present in Brassicaceae, and it mainly produces indole-3-carbinol (I3C). However, I3C usually undergoes dimeric condensation, giving 3,3-diindolylmethane (DIM).⁸ Both molecules have been demonstrated to be agonists of the aryl hydrocarbon receptor (AhR), and therefore increase the rate of detoxification pathways.⁹

Other colored Brassicaceae sprouts, such as red radish, are especially rich in anthocyanins. These compounds belong to the family of flavonoids, showing a C6–C3–C6 carbon skeleton and cyanidin, and they are frequently present in highly glycosylated and acylated forms in Brassicaceae sprouts, particularly in red radish varieties.¹⁰ The interest in anthocyanins has recently increased not only because of their anti-oxidant properties, but also due to their role in reducing the risk of renal and hepatic damage caused by oxidative stress conditions.^{11,12}

In recent decades, low-grade inflammatory processes have been highly related to non-communicable diseases, such as obesity, diabetes and cancer.^{13,14} Inflammation is a coordinated process performed by the immune system, whose main function is to maintain the organism's homeostasis. This process is triggered by both foreign agents and endogenous signals of tissue damage, thus playing a key role in tissue repair.¹⁵ Within this process, a broad variety of molecular mediators are involved, such as pro-inflammatory cytokines such as interleukin 6 (IL-6), interleukin 1 beta (IL-1 β) and tumor necrosis factor alpha (TNF- α), which are produced to a large extent by macrophages.¹⁶ It is noteworthy that an inflammatory process can lead to chronic inflammation if pathogenic agents or tissue damage signals are persistent and the acute inflammation is not correctly resolved. Sustained in time, chronic inflammation stages often are related to non-communicable diseases like type 2 diabetes and cardiovascular and neurodegenerative diseases, among others.¹⁷ Thus, understanding the physiopathology of these types of long-term inflammatory processes is of a great importance for the identification of new targets and strategies against these chronic inflammatory-related diseases.

Currently, the main therapeutic options to treat the above-mentioned inflammatory diseases are non-steroidal anti-inflammatory drugs (NSAIDs), since they act by blocking cyclooxygenases implicated in the synthesis of inflammatory mediators, like prostaglandins.¹⁸ However, NSAIDs present relevant adverse effects, which include cardiovascular, renal, and gastrointestinal toxicity. As a consequence, the need for new strategies and therapeutic agents that could ameliorate the inflammatory condition without these deleterious effects

has become a major research challenge.^{19,20} In this scenario, compounds derived from cruciferous vegetables have emerged as a possible alternative, either as a drug on their own or as a possible nutritional intervention option. For example, SFN has been described to induce the Nrf2 transcription factor, which is implicated in the transcription of diverse genes associated with the anti-inflammatory response.²¹ Furthermore, *in vivo* studies conducted on 40 overweight subjects who consumed 30 g of broccoli sprouts per day in their diet for 10 weeks showed a decrease in IL-6 levels.²² In this line, we previously reported that in macrophage-like HL-60 cells several cruciferous-derived compounds exhibited potential anti-inflammatory activity without showing any cytotoxic effect.²³ Nevertheless, the concentrations used in most *in vitro* assays often differ from the actual concentrations of these compounds that can be found in the circulation after intestinal digestion.²⁴ Thus, the aim of this work was to assess the anti-inflammatory and cytotoxic effect of the resulting product of an *in vitro* digestion process (digestates) of red radish sprout (RRS) and red cabbage sprout (RCS) extracts in a human macrophage-like cell model derived from HL-60.

2. Materials and methods

2.1. Plant material

A total of 50 g of seeds from red cabbage (*Brassica oleracea* L. var. *capitata* f. *rubra*) and red radish (*Raphanus sativus* L. var. *sativus* L. cv. Sango), provided by Intersemillas S.A. (Loriguilla, Valencia, Spain), were induced to germinate after decontamination in 1% bleach in de-ionized water followed by 24 hours of aeration. Subsequently, they were sown on an inert substrate of cellulose and kept in darkness and high relative humidity for 48 hours. Three-day-old germinated seeds were transferred to a growth chamber to grow the sprouts under a 20 W Protect BioLED (SysLed Spain, S.L.) photoperiod of 18 hours light/6 hours darkness, a temperature of 24 °C/18 °C, and a relative humidity of 60%/80%. Once collected, the 8-day sprouts were freeze-dried and ground prior to analysis. The resulting freeze-dried powder was stored at –80 °C.

2.2. Extract preparation and *in vitro* gastrointestinal digestions

Aqueous extracts (1 : 15, w : v) from freeze-dried RRS and RCS powders were elaborated by maceration for 4 hours in darkness and under continuous agitation. The samples were centrifuged at 13500g for 15 minutes. Supernatants were collected and percolated through a qualitative filter. For the *in vitro* gastrointestinal digestion, the INFOGEST standardized protocol was used.²⁵ 1 mL of the extract was mixed with 15 mL of a porcine pepsin solution (2000 units per mL in 100 mM HCl, Sigma-Aldrich, MO, USA) at pH 2 and 37 °C for 3 h. After the gastric digestion phase, 7.5 mL of 0.2 M sodium hydroxide (NaOH) was added to increase the pH to 7. For the intestinal digestion phase, another 7.5 mL of a pancreatin solution in phosphate buffer (2000 unit per mL, pH = 8) was added. For



the intestinal phase, samples were incubated for 12 hours at 37 °C. In order to prevent the contamination of the samples by microorganisms, 75 µL of 0.5 mM merthiolate solution was included in the digestions. After the whole process, samples were kept overnight at 4 °C, then filtered through a 0.22 µm Ø PVDF membrane and freeze-dried for further experiments (Fig. 1). The digestates were labeled as follows: RRS (red radish sprout extract digestates); and (ii) RCS (red cabbage sprout extract digestates).

2.3. Determination of ITCs in digestates

The presence of ITCs in both types of digestate from RRS and RCS extracts was analyzed and quantified using a UHPLC coupled with a 6460 triple quadrupole-MS/MS (Agilent Technologies, Waldbronn, Germany) and a Zorbax Eclipse Plus C18 column (2.1 × 50 mm, 1.8 µm).²⁶ SFN, I3C, and DIM from Santa Cruz Biotech (California, USA) were used as standards. In order to quantify and detect the presence of flavonols and anthocyanins in the digestates, additional analysis with HPLC-DAD was performed and sinapinic acid (Sigma) and cyanidin-3-glucoside (Sigma) were used as external standards.²⁶ Retention times and UV-Vis spectral characteristics, as well as the fragmentation patterns of the compounds of interest, were used in the determinations.

2.4. Cell culture

Digestates from RRS and RCS aqueous extracts were incorporated in the *in vitro* model of macrophage-like cells obtained by the differentiation of the human acute myeloid leukemia cell line HL-60 (ATCC® CCL-240™) as previously described.²⁷ Briefly, the cells were kept undifferentiated and growing at an exponential ratio in complete culture media (CCM), RPMI-1640 (Biowest, Nuaille, France) supplemented with 10% fetal bovine serum (GIBCO Invitrogen, Paisley, UK) and 1%

penicillin/streptomycin (GIBCO) prior to their differentiation. Growing conditions were set at 37 °C and under a 5% CO₂ atmosphere in the incubator. Cells were allowed to differentiate for conduction of the *in vitro* assays, always between passages number 5 and 20.

2.5. Cell viability assays

The cell viability of differentiated HL-60 macrophage-like cells treated with the digestates was measured using the MTT assay.²⁷ Briefly, after cell differentiation with 10 ng mL⁻¹ phorbol myristate acetate (PMA; Sigma Chemical Co., St Louis, MO, USA) in 96-well plates, the cells were pre-incubated with 10, 50, and 100 µg mL⁻¹ of either RRS or RCS digestates, and after 30 minutes the cells were treated with LPS (0.1 µg mL⁻¹) and incubated for 24 hours. MTT (Alfa Aesar, Thermo Fisher, Karlsruhe, Germany) was then added to the final concentration of 483 µM and incubated for 1 hour at 37 °C and under a 5% CO₂ atmosphere. Then, an acidified isopropanol solubilization solution of 0.04 M hydrochloric acid and 0.1% NP-40 detergent was added to each well to lyse the cells and dissolve the purple formazan crystal formed inside the cells, obtaining a purple-colored solution. The absorbance was measured at 550 nm in a plate-reading spectrophotometer (Spectrostar Nano; BMG Labtech, Ortenberg, Germany). The cell viability was determined by comparison with control conditions, setting the latter as 100% viability, and thus 0% cytotoxicity. Control cells were maintained under basal conditions treated only with CCM. Three replicates were performed for each experiment.

2.6. *In vitro* anti-inflammatory assays

The anti-inflammatory potential of the RRS and RCS digestates was examined by determining their ability to modulate the *in vitro* secretion of cytokines under a low-intensity inflamma-




Fig. 1 Schematic presentation of the standardized protocol followed for *in vitro* digestions. This protocol can be divided into a first phase of "gastric digestion", developed at pH = 2 for 3 hours, and followed by a phase of "intestinal digestion" at pH = 7 for 12 hours. The whole protocol was performed at 37 °C. Once digested, the samples were incubated overnight at 4 °C and then kept at -80 °C until analysis.



tory state induced by lipopolysaccharide (LPS).²⁸ To achieve this, HL-60 cells (0.2×10^6 cells per well) were stimulated with 10 ng mL^{-1} phorbol 12-myristate 13-acetate (PMA) in CCM for 24 hours, in order to differentiate them into human macrophage-like cells.²⁷ Afterward, the cells were rested for 24 hours in CCM without PMA. Once the differentiation was complete, the digestates of RRS and RCS aqueous extracts were pre-incubated with different concentrations (10 , 50 , and $100 \text{ } \mu\text{g mL}^{-1}$) at 37°C for 30 minutes. LPS (from *Escherichia coli* 0111.B4, Sigma Chemical Co.) was then added to a final concentration of $0.1 \text{ } \mu\text{g mL}^{-1}$ to induce a low-grade pro-inflammatory state similar to those found in chronic inflammatory diseases. Finally, the plates were incubated at 37°C under a $5\% \text{ CO}_2$ atmosphere for 24 hours. Supernatants were collected and stored at -20°C for further analysis. The cells were washed with PBS and then stored at -80°C until RNA extraction.

2.7. Enzyme-linked immunosorbent assays

Stored supernatants from *in vitro* assays performed on human macrophage-like HL-60 cells were used to determine the levels of the pro-inflammatory cytokines TNF- α , IL-6, and IL-1 β using enzyme-linked immunosorbent assay (ELISA) kits (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA). The absorbance in each well was determined using a microplate reader (Spectrostar Nano; BMG Labtech, Ortenberg, Germany) at 450 nm and corrected at 570 nm . Each cytokine concentration was calculated using the corresponding standard curve. The modulation of cytokines was attained by calculating the level of release inhibition when compared to control conditions. In order to compare the effect of the RRS and RCS digestates from the different experiments ($n \geq 2$), concentrations of the cytokines obtained from enzyme-linked immunosorbent assays (ELISA) (in the ranges $30\text{--}350 \text{ pg mL}^{-1}$ for TNF- α , $250\text{--}800 \text{ pg mL}^{-1}$ for IL-6, and $30\text{--}120 \text{ pg mL}^{-1}$ for IL-1 β) were normalized for each individual assay by setting the negative control of the corresponding experiment as the reference (valued as 1 for control conditions). The effect of the digestates on cytokine levels has been presented as the mean \pm SEM of normalized results.

2.8. Gene expression determination by RT-qPCR

The entire RNA was extracted from macrophage-like differentiated HL-60 cells using the GenElute Mammalian Total RNA Miniprep kit (Sigma) following the manufacturer's instructions. RNA was eluted in RNase-free water and checked for concentration and purity using the Nanodrop spectrophotometer system (ND-100 3.3. Nanodrop Technologies, USA). Only samples with a ratio of Abs260/Abs280 between 1.8 and 2.1 were used for gene expression assays. The generation of cDNA was executed using a high-capacity RNA-to-cDNA kit (Thermo Fisher Scientific, USA), following the manufacturer's instructions. PCR amplification was carried out with a PowerUp SYBR Green PCR Master Mix (Applied Biosystems, USA) using a Fast 7500 real-time instrument (Applied Biosystems, USA). Glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) was used as the reference gene. Primer sequences are presented in the

ESI Table S1.† The relative expression was calculated using the $2^{-\Delta\Delta Ct}$ method. All experiments were performed in triplicate and as three independent experiments.

2.9. Statistical analysis

Statistical analysis was performed using *t*-Student, ANOVA, and Tukey's HSD tests. Values of $p < 0.05$ were considered to be statistically significant. The GraphPad Prism 9 software was used for performing the calculations.

3. Results

3.1. Quantitative composition of secondary metabolites in the digestates

The results of the analysis and quantification of secondary metabolites present in RRS and RCS upon digestion of the aqueous extracts are shown in Table 1. According to ITCs, the SFN concentrations were higher in the digestates of RRS when compared to the digestates of RCS ($p = 0.002$). In addition, sulforaphane (SFE) was only detected in the digestates of RRS. Regarding I3C, the digestates of RCS contained around 3-fold higher I3C amounts compared to the digestates of RRS ($p = 0.015$). No traces of DIM, the naturally formed dimer of I3C, were found in RCS, although it was quantified in RRS. The ITC iberin was detected in digestates of both RRS and RCS in similar contents ($p = 0.075$). Accounting for the total amount of ITCs, the digestates of RCS were significantly higher than the digestates of RRS ($p < 0.05$).

The composition of phenolic acids and anthocyanins in both digestates was also analyzed (Table 1). Among phenolic acids, the characteristic sinapic acid derivatives of these varieties were detected and quantified by HPLC-DAD in similar contents for both types of digestate ($p > 0.05$). However, only the digestates of RRS aqueous extracts presented a quantifiable amount of cyanidins.

Table 1 Composition of GSLs hydrolysis products, ITCs and phenolic compounds (sinapic acid derivatives and anthocyanins) analysed in the digestates obtained from aqueous extracts of RRS and RCS (mM)

(mM)		RRS extract digestate	RCS extract digestate
Isothiocyanates	SFN	$6.60 \pm 0.40\text{a}$	$1.92 \pm 0.18\text{b}$
	SFE	1.68 ± 0.06	—
	I3C	$30.00 \pm 2.50\text{a}$	$74.00 \pm 7.06\text{b}$
	DIM	20.00 ± 2.12	—
	Iberin	$12.94 \pm 0.56\text{a}$	$10.75 \pm 1.07\text{a}$
	Total	$71.19 \pm 3.70\text{a}$	$86.67 \pm 8.25\text{b}$
Phenolic compounds	Sinapic acid derivatives	$11.17 \pm 0.27\text{a}$	$12.56 \pm 0.38\text{a}$
	Anthocyanins (cyanidins)	4.10 ± 0.04	—

Results are represented as the mean \pm SD ($n = 3$). Different letters indicate statistically significant differences between extract digestates in Tukey's HSD test ($p < 0.05$). RCS: red cabbage sprouts, RRS: red radish sprouts, SFN: sulforaphane, SFE: sulforaphane, I3C: indole-3-carbinol, and DIM: 3,3-diindolylmethane.



3.2. The effect of digestates on human macrophage-like HL-60 cell viability

The percentage of cell viability of human macrophage-like HL-60 cells is presented in Fig. 2. The viability of the cells stimulated with LPS did not show any statistically significant difference when compared to those under basal conditions without LPS (control). Viability in terms of the metabolism ratio of the cells was also assessed in the presence of the two tested digestates with different concentrations of 10, 50, and 100 $\mu\text{g mL}^{-1}$ under low-degree inflammation induced by LPS treatment. The RRS extract digestates induced a slight but significant elevation of the cellular metabolism with only a mild concentration of 50 $\mu\text{g mL}^{-1}$ in the presence of LPS, showing an increase of 23% when compared to the control cells ($p = 0.02$), while no differences were found with cells treated with 10 and 100 $\mu\text{g mL}^{-1}$ doses of digestates of RRS and LPS. Regarding RCS extract digestates, increases of 25% and 30% in cell viability were also observed in the cells treated with both 10 and 50 $\mu\text{g mL}^{-1}$ respectively in the presence of LPS when compared to control conditions. No change in the cell viability was observed when they were treated with 100 $\mu\text{g mL}^{-1}$ concentrations of RCS extract digestates and LPS.

3.3. The *in vitro* effect of digestates on pro-inflammatory cytokines production by human macrophage-like HL-60 cells

The results obtained from analyzing the pro-inflammatory cytokine profile (TNF- α , IL-6, and IL-1 β levels) released from human macrophage-like HL-60 cells treated with RRS and RCS




Fig. 2 The effect on cell viability (%) of the macrophage-like HL-60 differentiated cells of red radish sprout (RRS) and red cabbage sprout (RCS) extract digestates (10, 50, and 100 $\mu\text{g mL}^{-1}$) under a low degree of inflammatory conditions (0.1 $\mu\text{g mL}^{-1}$ LPS). *P*-Values were calculated using the Student *t*-test. Values are expressed as percentages referred to a control basal condition without LPS (dash line). Results are presented as the mean \pm SEM from 3 different MTT assays, with three replicates in each assay. * $p < 0.05$ between treatments and LPS-stimulated control. LPS, Lipopolysaccharide.

extract digestates under low degrees of inflammatory conditions are presented in Fig. 3. For TNF- α levels (Fig. 3A), when LPS-stimulated cells were treated with 50 and 100 $\mu\text{g mL}^{-1}$




Fig. 3 ELISA measured relative levels of TNF- α (A), IL-6 (B), and IL-1 β (C) released from macrophage-like HL-60 differentiated cells under LPS-stimulated conditions (0.1 $\mu\text{g mL}^{-1}$ LPS) following treatment with RRS and RCS extract digestates at different concentrations (10, 50 and 100 $\mu\text{g mL}^{-1}$). *P*-Values were calculated using the ANOVA test with a *post hoc* Tukey's HSD test. Values are expressed as a fold-change to control basal conditions without LPS (dash line). Results are represented as the mean \pm SEM from three different ELISA assays with three replicates performed in each assay. * $p < 0.05$, ** $p < 0.01$ between treatments and basal conditions (without LPS). # $p < 0.05$, ## $p < 0.01$ between treatments and the LPS-stimulated control. TNF- α , tumor necrosis factor alpha; IL-6, interleukin 6; IL-1 β , interleukin 1 beta; LPS, lipopolysaccharide.

doses of digestates obtained from RRS extracts, a reduction trend of 50% and 70% was observed respectively when compared with LPS-stimulated cells, while no differences were found for the TNF- α production when the lowest concentration of 10 $\mu\text{g mL}^{-1}$ of the RRS extract digestates was added. Regarding the treatment with RCS extract digestates, a non-significant decrease of 50% in the levels of this pro-inflammatory cytokine was also observed with the highest concentration of 50 $\mu\text{g mL}^{-1}$ when compared to the LPS-stimulated cells. Surprisingly, a further increase in TNF- α levels was observed after treatment with lower doses of 10 and 50 $\mu\text{g mL}^{-1}$ of RCS extract digestates and LPS; these increments were significantly higher compared to those under basal conditions ($p < 0.01$ and $p < 0.05$ respectively for 10 and 50 $\mu\text{g mL}^{-1}$ of RCS extract digestates).

Regarding IL-6 (Fig. 3B), for RRS extract digestates, the highest reduction in this pro-inflammatory cytokine (80%) was achieved with 100 $\mu\text{g mL}^{-1}$ compared to LPS-stimulated levels, reducing this cytokine level even further than those under basal conditions ($p < 0.01$ when compared with LPS-stimulated cells and $p < 0.05$ when compared to those under basal conditions), while the 50 $\mu\text{g mL}^{-1}$ concentration produced a decrease of 46% compared with LPS-stimulated cells ($p > 0.05$). In the case of RCS extract digestates, a similar performance to that observed for TNF- α was also observed for IL-6. When LPS-stimulated macrophage-like cells were exposed to the highest concentration of 100 $\mu\text{g mL}^{-1}$ of RCS extract digestates, a non-significant 30% decrease in IL-6 levels was observed, whilst an increment trend of 70% was found with the 10 $\mu\text{g mL}^{-1}$ concentration. In the case of IL-1 β (Fig. 3C), RRS extract digestates reported 26% and 35% decreasing tendencies respectively for concentrations of 50 and 100 $\mu\text{g mL}^{-1}$, while no difference whatsoever was observed for the lowest concentration of 10 $\mu\text{g mL}^{-1}$. For LPS-stimulated cells treated with RCS extract digestates, no significant differences were

observed between the cells treated with only LPS and those treated with any of the three tested concentrations.

3.4. Effect of digestates on the gene expression of pro-inflammatory cytokines

The gene expression of *TNF*, *IL6*, and *IL1B* was analyzed in the presence of both types of digestate under LPS-stimulated and non-stimulated human macrophage-like HL-60 cells (Fig. 4; ESI Fig. S1†). Assays were performed at the concentration of 100 $\mu\text{g mL}^{-1}$ of both tested digestates since this dose displayed the highest inhibitory effect on *in vitro* pro-inflammatory cytokines release without any cytotoxic effects. For *TNF*, no significant differences were found between the LPS-stimulated control cells and the ones treated with the RRS and RCS extract digestates. However, under control conditions without LPS (a non-inflammatory scenario) the treatment with a 100 $\mu\text{g mL}^{-1}$ concentration of the RCS extract digestate showed a 2-fold increase in the *TNF* relative gene expression when compared to non-treated cells ($p < 0.05$). Regarding the *IL6* expression, HL-60 macrophage-like cells stimulated with LPS showed a 10-times decrease in the relative gene expression after treatment with 100 $\mu\text{g mL}^{-1}$ digestates of the RRS extract ($p < 0.05$). Cells maintained under the same pro-inflammatory conditions treated with the same concentration of RCS extract digestates (100 $\mu\text{g mL}^{-1}$) showed no statistically significant differences in the *IL6* gene expression levels when compared to LPS-treated cells ($p < 0.05$). In the case of unstimulated cells cultured under non-inflammatory conditions, no statistically significant differences in the *IL6* gene relative expression were found between non-treated cells and those treated with any of the digestates. When the *IL1B* gene expression was analyzed in LPS-stimulated cells, a decrease of approximately 50% was observed after treatment with 100 $\mu\text{g mL}^{-1}$ digestates of RRS. No significant differences were found between RCS extract




Fig. 4 mRNA relative expression levels of genes involved in the immune response of human macrophage-like HL-60 differentiated cells exposed to red radish sprout (RRS) or red cabbage sprouts (RCS) extract digestate (100 $\mu\text{g mL}^{-1}$) under low-degree inflammatory conditions (0.1 $\mu\text{g mL}^{-1}$ LPS). *P*-Values were calculated using the ANOVA test with a *post hoc* Tukey's HSD test. Values are expressed as fold-change to the LPS-stimulated control. Results are represented as the mean \pm SEM from three different assays with three replicates performed in each assay. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. TNF, tumor necrosis factor; IL6, interleukin 6; IL1B, interleukin 1 beta; LPS, lipopolysaccharide.



digestates and those with LPS treatment alone. Nonetheless, under unstimulated conditions, a 2-times increase in *IL1B* expression levels was observed after treatment with 100 $\mu\text{g mL}^{-1}$ digestates of the RCS extract ($p < 0.05$).

4. Discussion

In recent years, the need to find alternative agents and/or adjuvants to treat chronic inflammatory diseases has been highlighted. Emerging evidence has proposed the addition of phytochemicals obtained from plant sources with anti-inflammatory properties in the dietary pattern.²⁹ Among those phytochemicals, natural compounds present in plants from the Brassicaceae family stand out because of their reported interaction with key molecular anti-inflammatory pathways.³⁰ However, usually, testing doses used to conduct the corresponding *in vitro* and *in vivo* studies are higher than those concentrations that are used under physiological conditions or else do not represent suitable therapeutic approaches.²⁴ Thus, this work aims to assess the anti-inflammatory potential of bioactive compounds of cruciferous sprouts (red radish and red cabbage) in their bioaccessible form (obtained by the digestion of aqueous extracts), using this simulation of physiological conditions for which the contents of individual compounds were in the concentration range from 0 to 50 μM . This *in vitro* digestion process provides the active fraction of compounds that might reach the gut epithelium and from there reach the lamina propria in human beings after the ingestion of food. To test the anti-inflammatory potential of such digestates we used a well-established *in vitro* inflammation cellular model³¹ consisting of human macrophage-like HL60 cells stimulated with LPS. This model tries to mimic systemic chronic inflammatory conditions present in certain non-communicable diseases such as cardiovascular disease, cancer, and diabetes.

The composition of the digestates of RRS and RCS extracts presented differences with a 20% lower content of total ITCs in RRS than in RCS. However, there was more variability in the compounds present in RRS than in RCS extract digestates, including SFE and DIM, which were exclusively present in RRS. The SFE is a structural analog of SFN, which only differs in one double bond that provides higher stability to this ITC.³² Also, previous studies have reported a high interconversion of GSLs into ITCs in sprouts, making them a great source for obtaining extracts and further digestates.³³ In addition, although both digestates presented similar contents of sinapic acid derivatives, there was also a clear difference between the RCS and RRS, as the presence of cyanidins was only detected in RRS. The highly-acylated and glycosylated cyanidin derivatives of RRS may also interact with other bioactives in the extracts to exert their health-promoting activity in additive or synergistic combination, considering the demonstrated antioxidant and anti-inflammatory activity of the individual compounds, due to the coordination between the immune system and metabolism; developing novel dietary strategies to attenuate inflammation and oxidative stress seems promising and

the combination of different metabolites present in food matrices (e.g. sprout extracts) could exert beneficial effects on metabolic markers and the reduction of inflammatory (cytokines) markers in cells and animals as found individually with anthocyanins^{23,34} and isothiocyanates.^{21,23,35}

As the bioactivity of these digestates was determined *in vitro*, we found that they do not show any level of cytotoxicity with the assessed concentrations. These results were similar to those previously reported by us, in which among a series of diverse bioactive derivatives from Brassica plants, only SFN at the highest tested concentration (50 μM) showed a decrease in the cell viability of this *in vitro* model of HL-60 macrophage-like cells.³¹ In this regard, it is worth noting that the actual concentration of SFN present in the higher tested dose (100 $\mu\text{g mL}^{-1}$) was only 3.72 μM for RRS and 1.08 μM for RCS, concentrations that were shown to be non-cytotoxic in this cellular model. Furthermore, low and medium digestate concentrations even produced a slight increase in the cell viability, in a similar way to that reported in HepG2 cells treated with Bimi® digestates.³⁶ This phenomenon could be due to the presence of sugars and fibers that would favor an increase in the cell metabolism. Since in the extraction by aqueous maceration only hydrophilic molecules are extracted, no specific selection for ITCs has been made, thus, these kinds of molecules could also be present in the sprout extracts obtained before the *in vitro* digestion process.³⁷

The *in vitro* anti-inflammatory potential of the digestates was tested by ELISA assay determination of the production of three pro-inflammatory cytokines TNF- α , IL-6, and IL-1 β . It has been reported that TNF- α molecules exert a pleiotropic effect on diverse cell types, mainly by the activation of caspases, and the induction of intracellular signaling nuclear factor kappa B (NF- κB) and mitogen-activated protein kinases (MAPKs).³⁸ IL-6 was originally defined as a differentiation factor for B cells. Its synthesis is regulated by LPS and TNF- α and shows both pro- and anti-inflammatory effects.³⁹ IL-1 β is also a pro-inflammatory interleukin that is secreted by macrophages in the classical activation pathway.⁴⁰ Our results showed that the treatment of the cells with a low dose of RCS extract digestate (10 $\mu\text{g mL}^{-1}$) unexpectedly tended to increase the TNF- α and IL-6 levels. For this digestate concentration, it might be argued that the ratio between the content of sugars and bioactive compounds could be high enough to activate the cell metabolism by the former (as observed by an increase in the cell metabolism), which would in turn induce the basal macrophage-like cells activity, augmenting thus the expression and release of cytokines in the LPS-dependent low-degree inflammatory scenario established. Furthermore, it has been reported that IL-6 is also a factor involved in glucose homeostasis, but whether its role is beneficial or detrimental is still a controversial topic.⁴¹ Nevertheless, a higher concentration of the 100 $\mu\text{g mL}^{-1}$ RCS extract showed a trend of decreasing both the TNF- α and IL-6 production under LPS-stimulated conditions, and this inhibitory effect was mainly observed at the final protein expression. This activity at higher rates might be related to the inhibitory ability of iberin upon TLRs dimerization, impairing the NF- κB



human macrophage-like HL60 cells maintained in an LPS-stimulated low-grade inflammatory scenario showed a reduction in pro-inflammatory cytokine production with a dose-response trend when treated with RRS extract digestates. Furthermore, this RRS treatment was able to decrease the *IL6* relative gene expression even under non-inflammatory conditions. Therefore, this work highlights the relevance of the precise composition of plant-derived foods, as it is the key factor that can determine the anti-inflammatory potential of complex mixtures of vegetal origin and sheds light on the search for nutritional interventions or adjuvants that might help to improve relevant non-communicable diseases related to low-grade and chronic systemic inflammation.

Author contributions

P. G.-I. performed experiments, analyzed the data, and wrote the manuscript. M. A. N.-S. performed experiments, analyzed the data and wrote the manuscript. A. O.-B. performed experiments. M. A. M.-S. performed experiments and wrote the manuscript. B. R.-M. conceived and designed the experimental plan, performed experiments, analyzed the data, provided the necessary facilities, and wrote the manuscript. A. J. R.-A. conceived, designed and coordinated the experimental plan, performed experiments, analyzed the data, and wrote the manuscript. D. A.-M. conceived and designed the experimental plan, provided plant materials, provided necessary facilities, and wrote the manuscript. All authors read and approved the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This research was funded by the Fundación Séneca – Murcia Regional Agency for Science and Technology (Comunidad Autónoma de la Región de Murcia, CARM), through the grant number 20855/PI/18, D. A. M., and, by the Institute of Health “Carlos III” (ISCIII), and co-funded by the Fondo Europeo de Desarrollo Regional-FEDER (grant number PI20/00505, B. R. M.). P. G.-I., was funded by a grant from the Fundación Séneca-CARM, Spain (21273/FPI/19). M. A. M.-S, was supported by a PFIS contract from the ISCIII (FI21/00003, ISCIII, Spain; co-funded by the Fondo Europeo de Desarrollo Regional-FEDER). B. R.-M., was supported by the “Miguel Servet Type I” program (CP19/00098, ISCIII, Spain; co-funded by the Fondo Europeo de Desarrollo Regional-FEDER).

5. Conclusion

In this work, we were able to establish the anti-inflammatory potential of two different digestates obtained from the aqueous extracts of certain Brassicaceae sprouts. In this way,

References

- 1 L. Galland, Diet and inflammation, *Nutr. Clin. Pract.*, 2010, **25**, 634–640.

- 2 D. Ramirez, A. Abellán-Victorio, V. Beretta, A. Camargo and D. A. Moreno, Functional ingredients from Brassicaceae species: Overview and perspectives, *Int. J. Mol. Sci.*, 2020, **21**, 1998.
- 3 A. Abellán, R. Domínguez-Perles, D. A. Moreno and C. García-Viguera, Sorting out the value of cruciferous sprouts as sources of bioactive compounds for nutrition and health, *Nutrients*, 2019, **11**, 429.
- 4 D. B. Clarke, Glucosinolates, structures and analysis in food, *Anal. Methods*, 2010, **2**, 310–325.
- 5 M. A. Prieto, C. J. López and J. Simal-Gandara, Glucosinolates: Molecular structure, breakdown, genetic, bioavailability, properties and healthy and adverse effects, *Adv. Food Nutr. Res.*, 2019, **90**, 305–350.
- 6 R. Bayat Mokhtari, N. Baluch, T. S. Homayouni, E. Morgatskaya, S. Kumar, P. Kazemi and H. Yeger, The role of sulforaphane in cancer chemoprevention and health benefits: a mini-review, *J. Cell Commun. Signaling*, 2018, **12**, 91–101.
- 7 I. M. A. Ernst, K. Palani, T. Esatbeyoglu, K. Schwarz and G. Rimbach, Synthesis and Nrf2-inducing activity of the isothiocyanates iberiverin, iberin and cheirolin, *Pharmacol. Res.*, 2013, **70**, 155–162.
- 8 K. R. Grose and L. F. Bjeldanes, Oligomerization of indole-3-carbinol in aqueous acid, *Chem. Res. Toxicol.*, 1992, **5**, 188–193.
- 9 T. T. Y. Wang, Q. Pham and Y. S. Kim, Elucidating the role of CD84 and AHR in modulation of LPS-induced cytokines production by cruciferous vegetable-derived compounds indole-3-carbinol and 3,3'-diindolylmethane, *Int. J. Mol. Sci.*, 2018, **19**, 1–16.
- 10 N. Zhang and P. Jing, Anthocyanins in Brassicaceae: Composition, stability, bioavailability and potential health benefits, *Crit. Rev. Food Sci. Nutr.*, 2022, **62**, 2205–2220.
- 11 N. Tena, J. Martín and A. G. Asuero, State of the art of anthocyanins: Antioxidant activity, sources, bioavailability and therapeutic effect in human health, *Antioxidants*, 2020, **9**, 451.
- 12 J. M. Sankhari, M. C. Thounaojam, R. N. Jadeja, R. V. Devkar and A. V. Ramachandran, Anthocyanin-rich red cabbage (*Brassica oleracea* L.) extract attenuates cardiac and hepatic oxidative stress in rats fed an atherogenic diet, *J. Sci. Food Agric.*, 2012, **98**, 1688–1693.
- 13 F. R. Greten and S. I. Grivennikov, Inflammation and cancer: Triggers, mechanisms, and consequences, *Immunity*, 2019, **51**, 27–41.
- 14 S. Galmés, M. Cifre, A. Palou, P. Oliver and F. Serra, A genetic score of predisposition to low-grade inflammation associated with obesity may contribute to discern population at risk for metabolic syndrome, *Nutrients*, 2019, **11**, 298.
- 15 R. Medzhitov, Origin and physiological roles of inflammation, *Nature*, 2008, **454**, 428–435.
- 16 D. R. Germolec, R. P. Frawley and E. Evans, Markers of inflammation, *Methods Mol. Biol.*, 2010, **598**, 53–73.
- 17 R. Scrivo, M. Vasile, I. Bartosiewicz and G. Valesini, Inflammation as 'common soil' of the multifactorial diseases, *Autoimmun. Rev.*, 2011, **10**, 369–374.
- 18 G. A. Green, Understanding NSAIDs: From aspirin to COX-2, *Clin. Cornerstone*, 2001, **3**, 50–59.
- 19 K. M. Knights, A. A. Mangoni and J. O. Miners, Defining the COX inhibitor selectivity of NSAIDs: Implications for understanding toxicity, *Expert Rev. Clin. Pharmacol.*, 2010, **3**, 769–776.
- 20 M. Fine, Quantifying the impact of NSAID-associated adverse events., *Am. J. Manage. Care*, 2013, **19**, 267–272.
- 21 A. J. Greaney, N. K. Maier, S. H. Leppla and M. Moayeri, Sulforaphane inhibits multiple inflammasomes through an Nrf2-independent mechanism, *J. Leukocyte Biol.*, 2016, **99**, 189–199.
- 22 M. T. López-Chillón, C. Carazo-Díaz, D. Prieto-Merino, P. Zafrilla, D. A. Moreno and D. Villano, Effects of long-term consumption of broccoli sprouts on inflammatory markers in overweight subjects, *Clin. Nutr.*, 2018, **38**, 745–752.
- 23 A. J. Ruiz-Alcaraz, M. A. Martínez-Sánchez, P. García-Peñarrubia, M. Martínez-Esparza, B. Ramos-Molina and D. A. Moreno, Analysis of the anti-inflammatory potential of Brassica bioactive compounds in a human macrophage-like cell model derived from HL-60 cells, *Biomed. Pharmacother.*, 2022, **149**, 112804.
- 24 S. Quirante-Moya, P. García-Ibañez, F. Quirante-Moya, D. Villano and D. A. Moreno, The role of *Brassica* bioactives on human health: Are we studying it the right way?, *Molecules*, 2020, **25**, 1591.
- 25 M. Minekus, M. Alminger, P. Alvito, S. Ballance, T. Bohn, C. Bourlieu, F. Carrière, R. Boutrou, M. Corredig, D. Dupont, C. Dufour, L. Egger, M. Golding, S. Karakaya, B. Kirkhus, S. Le Feunteun, U. Lesmes, A. MacIerzanka, A. MacKie, S. Marze, D. J. McClements, O. Ménard, I. Recio, C. N. Santos, R. P. Singh, G. E. Vegarud, M. S. J. Wickham, W. Weitschies and A. Brodtkorb, A standardised static in vitro digestion method suitable for food—an international consensus, *Food Funct.*, 2014, **5**, 1113–1124.
- 26 N. Baenas, C. Suárez-Martínez, C. García-Viguera and D. A. Moreno, Bioavailability and new biomarkers of cruciferous sprouts consumption, *Food Res. Int.*, 2017, **100**, 497–503.
- 27 A. Guirado, J. I. López Sánchez, A. J. Ruiz-Alcaraz, D. Bautista and J. Gálvez, Synthesis and biological evaluation of 4-alkoxy-6,9-dichloro[1,2,4] triazolo[4,3-a]quinoxalines as inhibitors of TNF- α and IL-6, *Eur. J. Med. Chem.*, 2012, **54**, 87–94.
- 28 U. Maitra, H. Deng, T. Glaros, B. Baker, D. G. S. Capelluto, Z. Li and L. Li, Molecular mechanisms responsible for the selective and low-grade induction of proinflammatory mediators in murine macrophages by lipopolysaccharide, *J. Immunol.*, 2012, **189**, 1014–1023.
- 29 A. Durazzo, L. D'Addezio, E. Camilli, R. Piccinelli, A. Turrini, L. Marletta, S. Marconi, M. Lucarini, S. Lisciani, P. Gabrielli, L. Gambelli, A. Aguzzi and S. Sette, From plant compounds to botanicals and back: A current snapshot, *Molecules*, 2018, **23**, 1844.



- 30 P. García-Ibañez, L. Yepes-Molina, A. J. Ruiz-Alcaraz, M. Martínez-Esparza, D. A. Moreno, M. Carvajal and P. García-Peñarrubia, Brassica bioactives could ameliorate the chronic inflammatory condition of endometriosis, *Int. J. Mol. Sci.*, 2020, **21**, 1–20.
- 31 A. J. Ruiz-Alcaraz, M. A. Martínez-Sánchez, P. García-Peñarrubia, M. Martínez-Esparza, B. Ramos-Molina and D. A. Moreno, Analysis of the anti-inflammatory potential of Brassica bioactive compounds in a human macrophage-like cell model derived from HL-60 cells, *Biomed. Pharmacother.*, 2022, **149**, 112804.
- 32 J. Zhang, X. Li, P. Ge, B. Zhang, L. Wen, C. Gu and X. Zhou, Sulforaphene: Formation, stability, separation, purification, determination and biological activities, *Sep. Purif. Rev.*, 2022, **51**(3), 330–339.
- 33 G. R. de Nicola, M. Bagatta, E. Pagnotta, D. Angelino, L. Gennari, P. Ninfali, P. Rollin and R. Iori, Comparison of bioactive phytochemical content and release of isothiocyanates in selected Brassica sprouts, *Food Chem.*, 2013, **141**, 297–303.
- 34 Y. M. Lee, Y. Yoon, H. Yoon, H. M. Park, S. Song and K. J. Yeum, Dietary anthocyanins against obesity and inflammation, *Nutrients*, 2017, **9**, 1–15.
- 35 M. T. López-Chillón, C. Carazo-Díaz, D. Prieto-Merino, P. Zafrilla, D. A. Moreno and D. Villaño, Effects of long-term consumption of broccoli sprouts on inflammatory markers in overweight subjects, *Clin. Nutr.*, 2018, **38**, 745–752.
- 36 P. García-Ibañez, D. A. Moreno and M. Carvajal, Nanoencapsulation of Bimi® extracts increases its bioaccessibility after in vitro digestion and evaluation of its activity in hepatocyte metabolism, *Food Chem.*, 2022, **385**, 132680.
- 37 S. C. Chuo, H. M. Nasir, S. H. Mohd-Setapar, S. F. Mohamed, A. Ahmad, W. A. Wani, M. Muddassir and A. Alarifi, A glimpse into the extraction methods of active compounds from plants, *Crit. Rev. Anal. Chem.*, 2020, **52**, 667–696.
- 38 H. Mitoma, T. Horiuchi, H. Tsukamoto and N. Ueda, Molecular mechanisms of action of anti-TNF- α agents – Comparison among therapeutic TNF- α antagonists, *Cytokine*, 2018, **101**, 56–63.
- 39 M. Akbari and V. Hassan-Zadeh, IL-6 signalling pathways and the development of type 2 diabetes, *Inflammopharmacology*, 2018, **26**, 685–698.
- 40 A. Capobianco and P. Rovere-Querini, Endometriosis, a disease of the macrophage, *Front. Immunol.*, 2013, **4**, 9.
- 41 L. L. Lehrskov and R. H. Christensen, The role of interleukin-6 in glucose homeostasis and lipid metabolism, *Semin. Immunopathol.*, 2019, **41**, 491–499.
- 42 T. Shibata, F. Nakashima, K. Honda, Y. J. Lu, T. Kondo, Y. Ushida, K. Aizawa, H. Suganuma, S. Oe, H. Tanaka, T. Takahashi and K. Uchida, Toll-like receptors as a target of food-derived anti-inflammatory compounds, *J. Biol. Chem.*, 2014, **289**, 32757–32772.
- 43 J. Jiang, T. B. Kang, D. W. Shim, N. H. Oh, T. J. Kim and K. H. Lee, Indole-3-carbinol inhibits LPS-induced inflammatory response by blocking TRIF-dependent signaling pathway in macrophages, *Food Chem. Toxicol.*, 2013, **57**, 256–261.
- 44 H. S. Youn, Y. S. Kim, Z. Y. Park, S. Y. Kim, N. Y. Choi, S. M. Joung, J. A. Seo, K.-M. Lim, M.-K. Kwak, D. H. Hwang and J. Y. Lee, Sulforaphane suppresses oligomerization of TLR4 in a thiol-dependent manner, *J. Immunol.*, 2010, **184**, 305–319.
- 45 J. E. Koo, Z. Y. Park, N. D. Kim and J. Y. Lee, Sulforaphane inhibits the engagement of LPS with TLR4/MD2 complex by preferential binding to Cys133 in MD2, *Biochem. Biophys. Res. Commun.*, 2013, **434**, 600–605.
- 46 E. Heiss, C. Herhaus, K. Klimo, H. Bartsch and C. Gerhäuser, Nuclear factor kappa B is a molecular target for sulforaphane-mediated anti-inflammatory mechanisms, *J. Biol. Chem.*, 2001, **276**, 32008–32015.
- 47 S. Han, Z. Wang, J. Liu and Q. Yuan, Identifying the p65-dependent effect of sulforaphane on esophageal squamous cell carcinoma progression via bioinformatics analysis, *Int. J. Mol. Sci.*, 2020, **22**, 60.
- 48 E. H. Lee, J. H. Shin, S. S. Kim and S. R. Seo, Sinapic acid controls inflammation by suppressing NLRP3 inflammasome activation, *Cells*, 2021, **10**, 2327.
- 49 T. Qiu, Y. Sun, X. Wang, L. Zheng, H. Zhang, L. Jiang, X. Zhu and H. Xiong, Drum drying and extrusion-black rice anthocyanins exert anti-inflammatory effects via suppression of the NF- κ B /MAPKs signaling pathways in LPS-induced RAW 264.7 cells, *Food Biosci.*, 2021, **41**, 100841.

