




Showcasing research from Professor Carell's laboratory,
Faculty of Chemistry and Pharmacy, Ludwig-Maximilians-
University, Munich, Germany.

Gradual evolution of a homo-L-peptide world on
homo-D-configured RNA and DNA

Nucleic acids with hypermodified nucleosides perform
stereoselective RNA/DNA-mediated peptide coupling,
followed by thermal release of the peptide, leading to
temperature-driven one-pot cycles with gradual homochiral
L-peptide enrichment.

As featured in:



See Ewa Węgrzyn, Ivana Mejdrová
and Thomas Carell,
Chem. Sci., 2024, **15**, 14171.

Cite this: *Chem. Sci.*, 2024, 15, 14171

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 23rd May 2024
Accepted 26th July 2024

DOI: 10.1039/d4sc03384a

rsc.li/chemical-science

Gradual evolution of a homo-L-peptide world on homo-D-configured RNA and DNA†

Ewa Węgrzyn,  ‡ Ivana Mejdrová  ‡ and Thomas Carell  *

Modern life requires the translation of genetic information – encoded by nucleic acids – into proteins, which establishes the essential link between genotype and phenotype. During translation, exclusively L-amino acids are loaded onto transfer RNA molecules (tRNA), which are then connected at the ribosome to give homo-L-proteins. In contrast to the homo-L-configuration of amino acids and proteins, the oligonucleotides involved are all D-configured (deoxy)ribosides. Previously, others and us have shown that if peptide synthesis occurs at homo D-configured oligonucleotides, a pronounced L-amino acid selectivity is observed, which reflects the D-sugar/L-amino acid world that evolved in nature. Here we further explore this astonishing selectivity. We show a peptide-synthesis/recapture-cycle that can lead to a gradual enrichment and hence selection of a homo-L-peptide world. We show that even if peptides with a mixed L/D-stereochemistry are formed, they are not competitive against the homo-L-counterparts. We also demonstrate that this selectivity is not limited to RNA but that peptide synthesis on DNA features the same L-amino acid preference. In total, the data bring us a step closer to an understanding of how homochirality on Earth once evolved.

Introduction

Life, as we know it, requires nucleic acid biomolecules to encode genetic information and amino acid-based proteins to catalyse biochemical reactions, that are essential for the maintenance of life. While the genetic code contains the blueprint for the multitude of vital proteins, the proteins in turn are essential for metabolism and required for decoding the sequence information and its replication. This strong interdependence of the genotype established by nucleic acids and the phenotype established by proteins is a hallmark of all life on Earth.¹ The element where the genotype “meets” the phenotype is the ribosome.^{2–4} Consequently, the evolution of the ribosome is a chicken-and-egg conundrum and one of the most challenging mysteries of the origin of life.^{5,6} The translation machinery that is in place today shows an extremely high stereoselectivity towards L-amino acids. First, a specific transfer RNA (tRNA) consisting of D-ribose is selectively loaded with the corresponding L-amino acid.⁷ In the step of translation, these L-amino acids are connected to other L-amino acids in a messenger RNA (mRNA)-based templation process, catalysed by the RNA components (all L-configured) of the ribosome. The exact time-point when homochirality emerged is unknown. Mechanistically it was suggested that it was caused

by a small initial enantiomeric imbalance on the monomer level,⁸ followed by processes of chiral amplification,^{9,10} leading to an enantiomeric induction between homochiral nucleic acids and peptides.^{11,12}

Numerous ideas exist of how the decoding¹³ could have evolved,^{14–16} which are all connected to the RNA world concept.^{17–19} This model predicts that in prebiotic times it was only RNA which encoded information and which also catalysed the essential reactions such as loading itself with specific amino acids.^{20–22} In the RNA world, the process of connecting amino acids was catalysed by RNA.^{22,23} The fact that the peptidyl transferase center of the ribosome is composed of RNA,²⁴ provides strong support for this “RNA-only” idea.^{25,26}

Currently, the idea of an RNA-only world is questioned. Recent studies²⁷ suggest instead the coexistence/coevolution of RNA with other entities like peptides,²⁸ cofactors^{29,30} or DNA.^{31–33} Even a DNA-first theory was formulated.^{33,34} The fact that most known RNA catalysts have rather low turn-over numbers strongly suggests that amino acids, peptides and cofactors might have been early on involved.³⁵ Already in 1976, White pointed out that cofactors alone could be considered as remnants of early life ribozymes,^{29,30} which acted in concert with sulfur-containing modified nucleosides.³⁶ Recently the idea of an RNA-peptide world was brought forward,^{37,38} with the peptides adding stability and catalytic capabilities to the RNA.³⁹ Such an early partnership could have been the foundation for the development of early amino acid-tRNA synthetases.⁴⁰

We recently introduced the idea, that the high number of non-canonical nucleosides present in RNA, are living fossils^{41,42}

Department of Chemistry, Center for Nucleic Acids Therapies at the Institute for Chemical Epigenetics (ICE-M), Ludwig-Maximilians-Universität (LMU), München Butenandtstrasse 5-13, 81377 Munich, Germany. E-mail: thomas.carell@lmu.de

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc03384a>

‡ These authors contributed equally.

of an early RNA/DNA world.⁴³ We could show that these non-canonical nucleosides add functions to RNA that would strongly benefit the evolution of life.⁴⁴ For example, non-canonical nucleosides t⁶A, g⁶A, as well as (m)nm⁵U,^{45–47} enable RNA to grow peptides on itself in a process which can establish an RNA-peptide world (Fig. 1).⁴⁸ The loading with amino acids and peptides and their connection to more complex structures is in principle possible under prebiotically plausible freezing/thawing conditions.^{49,50} It was shown, that this loading reaction exhibits no stereochemical bias, meaning that L- and D-amino acids are loaded onto RNA with no preference.^{12,51} However, when the amino acids are allowed to react with each other to give peptides, a large kinetic advantage was observed for the L-configured amino acids.^{11,52–57}

Based on this observation, we can envision a primitive peptide synthesis cycle as depicted in Fig. 1, that is driven by cold/hot cycles (−20 °C to +90 °C). In this model, amino acids (aa) and small peptides react (*via* nitrosation of N⁶-urea-A) with RNA (step I and II) to give (m⁶)aa⁶A-containing RNA-donor strands ((m⁶)aa⁶A-RNA),⁵¹ which can hybridize to complementary RNA acceptor strands containing the non-canonical

nucleoside (m)nm⁵U (step 0) in the cold. Upon activation and reaction of the amino acids with the (m)nm⁵U nucleoside (step 1), a hairpin structure with a high thermal stability is formed.⁴⁸ During a phase of increased temperature (90 °C), cleavage of the thermally labile urea bond will occur (step 2) together with an immediate dissociation of the strands (step 3). Upon cooling of the solution, a new donor strand can bind (step 4) to continue the hot/cold-governed synthesis cycle.⁵⁸ In the prebiotic context, these temperature variations could have originated from volcanic eruptions, meteorite impact or even day/night-dependent presence of sunlight.^{59,60} Current research reports on one hand cold, even frozen environments on early Earth,^{61,62} and on the other hot temperatures provided by the hydrothermal vents.⁶³

We hypothesise that the initially formed peptides were short and released into the environment upon degradation of the RNA. And such they could have been taken up again by donor strands to continue the process at another RNA “host” strand (step II). Here we show that such a scenario is realistic and we show that homo-L peptides would feature a clear competitive advantage over peptide strands contaminated with D-amino acids.

Results and discussion

In order to establish a stereochemical self-selecting peptide synthesis cycle, we intended to investigate the one-pot growth of a peptide using a racemic mixture of L- and D-Val RNA donor strands (Fig. 2). We selected valine because it is one of the simplest chiral amino acids for which a prebiotic existence is plausible.^{64,65} Valine has also been observed in meteorites^{66,67} and its formation has been demonstrated in Urey–Miller-type spark discharge experiments.^{68–71}




Fig. 1 Cold/hot cycle of amino acid loading onto RNA (steps I and II) and iterative peptide synthesis cycle, that could lead to homochiral peptide formation in an outgrowth process. The steps of the cycle include: annealing of the complementary acceptor and donor strands (step 0), stereoselective peptide coupling (step 1), thermal cleavage of the peptide from the donor strand (step 2), followed by a release of the peptide-RNA acceptor conjugate (step 3), that can then recapture another amino acid-loaded donor strand (step 4), and enter another cycle. R stands for L- or D-Val.




Fig. 2 One-pot synthesis cycle scheme and HPLC-chromatograms. For reasons of simplicity, only the products formed in detectable amounts and relevant for further reactions are indicated in the scheme. Yields were determined by HPLC using the calibration curves of reference compounds (Fig. S2 and S3†).

The reaction mixture was next lyophilized (in analogy to a dry-down step) and the buffer was exchanged (comparable to fresh solution floating in a tidal process). We then added another portion of the equimolar mixture of **ON1_L** and **ON1_D**. After activation, we detected the formation of the hairpin product **ON3_{LLL}** by HPLC and MALDI-ToF, albeit in a low yield of 3% (over three steps). The heterochiral product **ON3_{LID}** was not detected. We then went through another hot-phase and performed the cleavage reaction, which provided in 1% yield (over 4 steps) the RNA strand **ON2_{LLL}** (Fig. 2 and S5; Table S8†). The main reason for the low yield is the large amount of

We noticed during the study that the second amino acid of *e.g.* **ON1_{LL}** undergoes a *ca.* 25% racemization when incubated under the described coupling conditions for 2 h. The mono-Val donors **ON1_L** and **ON1_D**, however, did not racemize when subjected to the same conditions. This phenomenon makes the analysis and interpretation of the described reactions difficult, but the general result is firmly established, that the homo-*L* situation is always winning.



Fig. 3 (A) General scheme for the competitive peptide coupling reactions of equimolar mixture of ON1_x and ON1_y with (B) ON2_L or (C) ON2_D. X and Y can correspond to either LL, LD, DL or DD, while X ≠ Y. The HPLC-chromatograms show the analysed crude reaction mixtures and the tables summarize the results obtained for the competitive coupling reactions, specifically product ratios of (B) [ON3_{xL}]/[ON3_{yL}] and (C) [ON3_{xD}]/[ON3_{yD}]. Reaction conditions: [ON1] = 50 μM; [ON2] = 50 μM; [buffer] = 100 mM; [NaCl] = 100 mM and [activator] = 50 mM. Yields were determined by HPLC analysis using the calibration curve of a reference compound (Fig. S2†).

To complete the set of experiments with dipeptide donors, we also analysed the competition reactions of ON1_{LL}, ON1_{LD}, ON1_{DL} and ON1_{DD} with a 1 : 1 mixture of ON2_L and ON2_D (Fig. 4 and S7†). For this experiment we prepared an equimolar mixture of ON2_L and ON2_D and allowed them to react with the dipeptide RNA donor strand ON1_x under the described coupling conditions. The reaction with ON1_{LL} showed

a selectivity of 66 : 34 in favour of the homochiral _{LLL}-hairpin product ON3_{LLL} over the product ON3_{LDD} (Fig. 4). This diastereoselectivity decreased slightly when we used ON1_{LD} (62 : 38, ON3_{LDD} : ON3_{DDL}). The _{D,D}-Val donor ON1_{DD} showed a reversed selectivity towards the homochiral all-_D product ON3_{DDD} of 71 : 29 compared to ON3_{DDL} (Fig. S7†). The coupling reaction with ON1_{DL} could not be analysed due to a HPLC peak overlap of the product with an impurity.

Finally, we wanted to investigate if the observed tendency to form homo-_L products prevails when one moves from RNA to DNA in light of the recent discussion that life may have started with a DNA-world³³ or a mixed RNA–DNA world.³¹ We therefore synthesized the DNA analogues of the donor strands **dON1_L** and **dON1_D** and the acceptor strands **dON2_L** and **dON2_D** by solid-phase oligonucleotide synthesis to form the hairpin products **dON3**. We conducted the competitive coupling reactions and observed the formation of the **dON3_{LL}** with a high diastereoselectivity of 85 : 15 over the heterochiral product **dON3_{DL}** (Fig. 5a). The reverse competition reaction yielded a 31 : 69 stereoselectivity for **dON3_{LD}** over **dON3_{DD}** (Fig. 5b). These results suggest that DNA will induce a similar diastereoselectivity towards the homo-_L products.

In order to gain more information about the conformation of the RNA duplexes, we finally measured the circular dichroism (CD) spectra of three different solutions: (1) annealed RNA duplex of ON1_L and ON2_L, (2) hairpin duplex ON3_{LL} and (3) the canonical analogue annealed RNA duplex of ON5 and ON6 (Fig. S9†). We observed CD-spectra typical for double-stranded A-form RNA with no significant changes introduced by the




Fig. 4 Competitive peptide coupling reactions of equimolar mixture of ON2_L and ON2_D with ON1_x. X can correspond to either LL, LD, DL or DD. The table summarizes the results obtained for the competitive coupling reactions. Reaction conditions: [ON1] = 50 μM; [ON2] = 50 μM; [buffer] = 100 mM; [NaCl] = 100 mM and [activator] = 50 mM. Yields were determined by HPLC analysis using the calibration curve of a reference compound (Fig. S2†).




Fig. 5 DNA-competitive peptide coupling reactions between dON1_L and dON1_D with (a) dON2_L or (b) dON2_D using EDC/Sulfo-NHS as activator. Yields were determined by HPLC analysis using the calibration curve of a reference compound (Fig. S2†).

nucleobase modification or the hairpin formation between them.

Conclusions

In this work we explored the possibility of a primitive temperature-driven one-pot peptide synthesis cycle, in which stereoselective peptide coupling followed by thermal cleavage can lead to an enriched formation, albeit in low yields, of the homochiral all *L*-peptides.

We investigated how different stereochemical dipeptide combinations, namely the presence of *L*- and *D*-amino acids in the cycle, would influence the peptide-forming chemistry. We found that whatever combination we investigated, in almost all cases we observed faster reactions and higher yields with *L*-amino acids and homo-*L*-peptides. The *L*-situation outcompeted reactions with *D*-amino acids present on the donor or on the acceptor strand, especially when the (*L*-) amino acid was directly attached to the nucleobase and therefore in close contact with the nucleic acid strand. Important is the observation that this stereochemical “preferential handshake” (*D*-ribose and *L*-amino acid) is also true with *D*-deoxyribose, which forms DNA. It shows us that even in a putative prebiotic world in which DNA-only or a mixed DNA–RNA combination were the carrier of genetic information, the stereochemical preference for *L* amino acids prevails.

Data availability

The data supporting this article have been included as part of the ESI.†

Author contributions

E. W. and I. M. were responsible for data curation, formal analysis, investigation and visualisation. T. C. had the research idea and was in charge of funding acquisition, project administration and supervision. All authors were involved in the conceptualization and the writing of the manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We thank the Deutsche Forschungsgemeinschaft for supporting this research through the DFG grants: CA275/11-3 (ID: 326039064), CRC1309 (ID: 325871075, A4), CRC1032 (ID: 201269156, A5) and CRC1361 (ID: 393547839, P2). We thank the Volkswagen Foundation for funding this research (grant EvoRib). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Program under grant agreement No. 741912 (EPiR) and the Marie Skłodowska-Curie grant agreement No. 861381 (Nature-ETN). The research was supported by the BMBF in the framework of the Cluster4Future program (Cluster for Nucleic Acid Therapeutics Munich, CNATM) (Project ID: 03ZU1201AA).

References

- 1 F. Crick, *Nature*, 1970, **227**, 561–563.
- 2 V. Ramakrishnan, *Cell*, 2002, **108**, 557–572.
- 3 T. A. Steitz, *Nat. Rev. Mol. Cell Biol.*, 2008, **9**, 242–253.
- 4 A. Yonath, M. A. Saper, I. Makowski, J. Müssig, J. Piefke, H. D. Bartunik, K. S. Bartels and H. G. Wittmann, *J. Mol. Biol.*, 1986, **187**, 633–636.
- 5 F. H. C. Crick, *Symp. Soc. Exp. Biol.*, 1958, **12**, 138–163.
- 6 F. H. C. Crick, S. Brenner, A. Klug and G. Pieczenik, *Origins Life*, 1976, **7**, 389–397.
- 7 R. Giegé and G. Eriani, in *eLS*, John Wiley & Sons, Ltd, Chichester, 2014, pp. 1007–1030.
- 8 J. E. Hein and D. G. Blackmond, *Acc. Chem. Res.*, 2012, **45**, 2045–2054.
- 9 R. Breslow and M. S. Levine, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 12979–12980.
- 10 R. Breslow and Z.-L. Cheng, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 9144–9146.
- 11 K. Tamura and P. Schimmel, *Science*, 2004, **305**, 1253.
- 12 E. Węgrzyn, I. Mejdrová, F. Müller, M. Nainy, L. Escobar and T. Carell, *Angew. Chem., Int. Ed.*, 2024, **63**, e202319235.
- 13 P. F. Agris, *Nucleic Acids Res.*, 2004, **32**, 223–238.
- 14 X. Y. Guo and M. Su, *Int. J. Mol. Sci.*, 2023, **24**, 197.
- 15 Y. I. Wolf and E. V. Koonin, *Biol. Direct*, 2007, **2**, 14.
- 16 M. Seki, *Genes Genet. Syst.*, 2023, **98**, 9–24.
- 17 W. Gilbert, *Nature*, 1986, **319**, 618.
- 18 G. F. Joyce, *Nature*, 2002, **418**, 214–221.



- 19 J. C. Bowman, N. V. Hud and L. D. Williams, *J. Mol. Evol.*, 2015, **80**, 143–161.
- 20 B. Jash and C. Richert, *Chem. Sci.*, 2020, **11**, 3487–3494.
- 21 B. Jash, P. Tremmel, D. Jovanovic and C. Richert, *Nat. Chem.*, 2021, **13**, 751–757.
- 22 N. Lee, Y. Bessho, K. Wei, J. W. Szostak and H. Suga, *Nat. Struct. Biol.*, 2000, **7**, 28–33.
- 23 I. Agmon, *Int. J. Mol. Sci.*, 2022, **23**, 15756.
- 24 P. Nissen, J. Hansen, N. Ban, P. B. Moore and T. A. Steitz, *Science*, 2000, **289**, 920–930.
- 25 I. Agmon, *Int. J. Mol. Sci.*, 2009, **10**, 2921–2934.
- 26 T. Bose, G. Fridkin, C. Davidovich, M. Krupkin, N. Dinger, A. H. Falkovich, Y. Peleg, I. Agmon, A. Bashan and A. Yonath, *Nucleic Acids Res.*, 2022, **50**, 1815–1828.
- 27 S. Tagami and P. Li, *Dev., Growth Differ.*, 2023, **65**, 167–174.
- 28 D. Kunnev, *Life*, 2020, **10**, 269.
- 29 H. B. White III, *J. Mol. Evol.*, 1976, **7**, 101–104.
- 30 A. D. Goldman and B. Kacar, *J. Mol. Evol.*, 2021, **89**, 127–133.
- 31 J. Xu, V. Chmela, N. J. Green, D. A. Russell, M. J. Janicki, R. W. Góra, R. Szabla, A. D. Bond and J. D. Sutherland, *Nature*, 2020, **582**, 60–66.
- 32 T. Lindahl, *Nature*, 1993, **362**, 709–715.
- 33 J. S. Teichert, F. M. Kruse and O. Trapp, *Angew. Chem., Int. Ed.*, 2019, **58**, 9944–9947.
- 34 J. Oró, S. L. Miller and A. Lazcano, *Annu. Rev. Earth Planet. Sci.*, 1990, **18**, 317–356.
- 35 H. S. Bernhardt, *Biol. Direct*, 2012, **7**, 23.
- 36 J. C. Cochrane and S. A. Strobel, *RNA*, 2008, **14**, 993–1002.
- 37 Y. Kim, K. Opron and Z. F. Burton, *Life*, 2019, **9**, 37.
- 38 A. S. Burton, J. C. Stern, J. E. Elsila, D. P. Glavin and J. P. Dworkin, *Chem. Soc. Rev.*, 2012, **41**, 5459–5472.
- 39 P. T. S. v. d. Gulik, *Life*, 2015, **5**, 1629–1637.
- 40 C. W. Carter Jr, *Nat. Hist.*, 2016, **125**, 28–33.
- 41 M. P. Robertson and S. L. Miller, *Science*, 1995, **268**, 702–705.
- 42 C. Schneider, S. Becker, H. Okamura, A. Crisp, T. Amatov, M. Stadlmeier and T. Carell, *Angew. Chem., Int. Ed.*, 2018, **57**, 5943–5946.
- 43 S. Becker, C. Schneider, H. Okamura, A. Crisp, T. Amatov, M. Dejmek and T. Carell, *Nat. Commun.*, 2018, **9**, 163.
- 44 I. Agmon, *Life*, 2024, **14**, 277.
- 45 T. Carell, C. Brandmayr, A. Hienzsich, M. Müller, D. Pearson, V. Reiter, I. Thoma, P. Thumbs and M. Wagner, *Angew. Chem., Int. Ed.*, 2012, **51**, 7110–7131.
- 46 J.-B. Zhou, E.-D. Wang and X.-L. Zhou, *Cell. Mol. Life Sci.*, 2021, **78**, 7087–7105.
- 47 T. Suzuki, Y. Yashiro, I. Kikuchi, Y. Ishigami, H. Saito, I. Matsuzawa, S. Okada, M. Mito, S. Iwasaki, D. Ma, X. Zhao, K. Asano, H. Lin, Y. Kirino, Y. Sakaguchi and T. Suzuki, *Nat. Commun.*, 2020, **11**, 4269.
- 48 F. Müller, L. Escobar, F. Xu, E. Węgrzyn, M. Nainytė, T. Amatov, C. Y. Chan, A. Pichler and T. Carell, *Nature*, 2022, **605**, 279–284.
- 49 K. Kitada, Y. Suda and N. Takenaka, *J. Phys. Chem. A*, 2017, **121**, 5383–5388.
- 50 C. Menor-Salván and M. R. Marín-Yaseli, *Chem. Soc. Rev.*, 2012, **41**, 5404–5415.
- 51 J. N. Singer, F. M. Müller, E. Węgrzyn, C. Hözl, H. Hurmiz, C. Liu, L. Escobar and T. Carell, *Angew. Chem., Int. Ed.*, 2023, **62**, e202302360.
- 52 J. C. Lacey, R. D. Thomas, M. P. Staves and C. L. Watkins, *Biochim. Biophys. Acta*, 1991, **1076**, 395–400.
- 53 K. Tamura, *BioSystems*, 2008, **92**, 91–98.
- 54 L. F. Wu, M. Su, Z. Liu, S. J. Bjork and J. D. Sutherland, *J. Am. Chem. Soc.*, 2021, **143**, 11836–11842.
- 55 S. J. Roberts, Z. Liu and J. D. Sutherland, *J. Am. Chem. Soc.*, 2022, **144**, 4254–4259.
- 56 M. Su, C. Schmitt, Z. Liu, S. J. Roberts, K. C. Liu, K. Röder, A. Jäschke, D. J. Wales and J. D. Sutherland, *J. Am. Chem. Soc.*, 2023, **145**, 15971–15980.
- 57 O. Doppleb, J. Bremer, M. Bechthold, C. Sánchez Rico, D. Göhringer, H. Griesser and C. Richert, *Chem.–Eur. J.*, 2021, **27**, 13544–13551.
- 58 D. Ross and D. Deamer, *Life*, 2023, **13**, 1749.
- 59 S. Becker, C. Schneider, A. Crisp and T. Carell, *Nat. Commun.*, 2018, **9**, 5174.
- 60 B. Charnay, G. Le Hir, F. Fluteau, F. Forget and D. C. Catling, *Earth Planet. Sci. Lett.*, 2017, **474**, 97–109.
- 61 K. Le Vay, E. Salibi, E. Y. Song and H. Mutschler, *Chem.–Asian J.*, 2020, **15**, 214–230.
- 62 M. J. de Wit and H. Furnes, *Sci. Adv.*, 2016, **2**, e1500368.
- 63 W. Martin, J. Baross, D. Kelley and M. J. Russell, *Nat. Rev. Microbiol.*, 2008, **6**, 805–814.
- 64 L. M. Longo and M. Blaber, *Arch. Biochem. Biophys.*, 2012, **526**, 16–21.
- 65 L. M. Longo, J. Lee and M. Blaber, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 2135–2139.
- 66 J. R. Cronin and C. B. Moore, *Science*, 1971, **172**, 1327–1329.
- 67 A. Shimoyama, C. Ponnampertuma and K. Yanai, *Nature*, 1979, **282**, 394–396.
- 68 S. L. Miller, *Science*, 1953, **117**, 528–529.
- 69 Y. Wolman, W. J. Haverland and S. L. Miller, *Proc. Natl. Acad. Sci. U. S. A.*, 1972, **69**, 809–811.
- 70 A. P. Johnson, H. J. Cleaves, J. P. Dworkin, D. P. Glavin, A. Lazcano and J. L. Bada, *Science*, 2008, **322**, 404.
- 71 E. T. Parker, H. J. Cleaves, J. P. Dworkin, D. P. Glavin, M. Callahan, A. Aubrey, A. Lazcano and J. L. Bada, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 5526–5531.
- 72 L. L. Cummins, S. R. Owens, L. M. Risen, E. A. Lesnik, S. M. Freier, D. McGee, C. J. Guinasso and P. D. Cook, *Nucleic Acids Res.*, 1995, **23**, 2019–2024.
- 73 W. A. Decatur and M. J. Fournier, *Trends Biochem. Sci.*, 2002, **27**, 344–351.

