

REVIEW

View Article Online
View Journal | View Issue



Cite this: *Org. Biomol. Chem.*, 2021, **19**, 4652

Optical sensing of anions by macrocyclic and interlocked hosts

Hui Min Tay and Paul Beer  *

The ubiquity of anions in biological and environmental systems has motivated the development of many novel anion receptors and sensors over the past two decades. Optical anion sensors, which undergo a spectral change in response to anion binding, are particularly desirable due to the technical simplicity and fast response time of such systems. A myriad of macrocyclic host molecules have been shown to be effective anion receptors and present a promising platform for elaboration into optical anion sensors by incorporation of an appropriate fluorogenic or chromogenic group. The enhanced anion binding properties of the three-dimensional binding cavities in mechanically interlocked host molecules have also been exploited to design anion sensors that exhibit remarkable selectivity and sensitivity. This review summarises recent progress in the development of optical anion sensors based on macrocyclic and interlocked hosts. The major classes of macrocyclic receptors possessing neutral, cationic and metal-based binding motifs are examined, followed by a survey of optically-responsive interlocked anion hosts.

Received 29th March 2021,
Accepted 6th May 2021

DOI: 10.1039/d1ob00601k

rsc.li/obc

1. Introduction

Prompted by the pivotal role of anions in multiple biological^{1–3} and environmental^{4–6} processes, the field of anion recognition has grown significantly in recent years. A particularly exciting development is the design of increasingly sophisticated sensors capable of reporting the presence of specific anions in a rapid and sensitive manner.⁷

Macrocyclic hosts are particularly well-suited to act as anion receptors due to the relative ease with which they can be functionalised with multiple anion binding motifs.⁸ This enables the use of a variety of synergistic non-covalent interactions, including hydrogen bonding, electrostatic interactions, metal coordination, halogen bonding and anion- π interactions, to achieve the recognition of a target guest species. By careful choice and placement of these binding groups, it is possible to design cavities that are complementary to specific anions – often a prerequisite for strong and selective recognition. Perhaps unsurprisingly, the earliest anion receptor, reported by Park and Simmons in 1968, was a cryptand-like macrobicyclic host that encapsulated halides between the ammonium groups located at the poles of its cavity.⁹ More recently, mechanically-interlocked molecules, such as catenanes and rotaxanes, have emerged as promising molecular hosts due to their highly pre-organised binding cavities, which have been

shown to enhance the affinity and selectivity for target guests.¹⁰

The modular synthesis of macrocyclic and mechanically interlocked hosts facilitates the incorporation of various reporter groups, opening up possibilities for optical and/or electrochemical sensing of anions.¹¹ The development of colorimetric and fluorescent probes for anions is especially desirable due to the sensitivity, fast response and technical simplicity of such systems.^{12,13} Many examples of optical anion sensors have been reported, including acyclic¹⁴ and metal-based systems¹⁵ which have been comprehensively discussed in other reviews. This review highlights recent developments in macrocyclic and interlocked optical anion sensors, with a particular focus on examples in which the macrocyclic/interlocked nature of the host significantly influences its sensing properties, either by conferring selectivity for specific anions, and/or by facilitating the spectral changes that occur upon anion binding. Macrocycles containing various neutral, cationic and metal-based binding motifs are discussed, followed by a survey of advances in the field of mechanically interlocked anion sensors.

2. Macrocycles with neutral anion-binding motifs

2.1. Calix[*n*]arene-based macrocycles

Calixarenes and their derivatives are versatile scaffolds for guest binding due to their highly pre-organised arene core which can be readily functionalised with a range of binding

Chemistry Research Laboratory, Department of Chemistry, University of Oxford, Mansfield Road, Oxford, OX1 3TA, UK. E-mail: paul.beer@chem.ox.ac.uk



Recently, two further examples of pyrenyl-appended derivatives of calixarenes were developed for optical sensing of halides. A calix[4]arene receptor **4** decorated with *tert*-butyl groups on the upper rim and amide-pyrene groups on the lower rim (Fig. 1) served as a dual cation and anion sensor specific for As(III), Nd(III) and Br⁻. Impressively, **4** exhibited a binding constant of $6.31 \times 10^8 \text{ M}^{-1}$ for Br⁻ in a competitive solvent mixture of acetonitrile/aqueous phosphate buffer (8 : 2, v/v). The formation of hydrogen bonds between Br⁻ and the amide linkers increased the partial negative charge on the pyrenyl-adjacent nitrogen atom, switching on photoinduced



electron transfer (PET) and quenching the pyrenyl fluorescence.²² A hexahomotrioxacalix[3]arene receptor **5** similarly underwent fluorescence quenching upon addition of iodide in THF/H₂O (7 : 3, v/v) (Fig. 1). The authors suggested that iodide was bound by synergistic hydrogen bonding interactions involving the amide groups. The heavy atom effect associated with iodine led to increased intersystem crossing to the non-emissive triplet state of pyrene.²³

The fluorenone-appended calix[4]arene receptor **6** exhibits a high affinity for fluoride, with a binding constant of $1.18 \times 10^6 \text{ M}^{-1}$ in acetonitrile, attributed to the formation of multiple OH and NH hydrogen bonds (Fig. 2). Addition of small quantities of fluoride resulted in significant fluorescence quenching due to an increase in PET from the partially deprotonated amide nitrogen atoms to the fluorenone groups, as well as a 'naked eye' colour change from yellow to orange-red.²⁴ A clear trend towards preferential fluoride binding was observed in a related series of calix[4]arene-based receptors functionalised on the lower rim with various fluorophores or chromophores (Fig. 2), including benzoxadiazole (**7**),²⁵ naphthalene (**8**),²⁶ anthraquinone (**9**),²⁷ nitrophenyl (**10**)²⁸ and naphthol (**11**).²⁹ In these systems, anion binding occurs in the pocket formed by the lower rim and the hydrogen-bonding linker, suggesting that size complementarity exists between this binding site and the small fluoride anion. In a slightly unusual synthesis, the calix[4]arene receptor **12** was functionalised at the upper rim with

thiourea linkers connected to coumarin groups.³⁰ This receptor once again displayed a high affinity for fluoride, with NMR studies in DMSO-*d*₆ suggesting the deprotonation of the thiourea groups and formation of HF₂[−] at high fluoride concentration. This resulted in fluorescence quenching due to enhanced PET. In addition, a new absorption band appeared at 475 nm, causing a colour change from colourless to orange.

A unique bis(bora)calix[4]arene **13** similarly displayed selectively for fluoride (Fig. 3).³¹ Unlike the previous examples, in which hydrogen bonding interactions were responsible for anion binding, **13** binds fluoride in chloroform *via* Lewis acid–


**13**

Fig. 3 Structure of a bis(bora)calix[4]arene for fluoride sensing ($K_a = 2.00 \times 10^6 \text{ M}^{-1}$ in CHCl₃).




Fig. 2 Structures of calix[*n*]arenes **7–12** functionalised with various chromophores/fluorophores highlighted in blue.



base interactions with the boron centres. The authors suggested that the fluoride selectivity arose from *endo* binding of fluoride in the pocket between the two boron centres, which is too small to accommodate other anions. However, a subsequent computational study suggested that cooperative *endo* and *exo* binding modes were responsible.³² The fluorescence of the boron-adjacent phenyl groups was quenched upon fluoride binding, which may be due to increased PET from boron to the phenyl groups or internal conversion to a long-lived 'dark state' that suppressed the 'prompt' fluorescence from the higher excited states.

While many calix[4]arene receptors recognised fluoride selectively, tweaking the nature and position of the binding groups can alter the selectivity for anions. A resin consisting of calix[4]amido-crown-5 receptor **14** attached to 4-benzyloxy benzyl alcohol polymer was found to preferentially bind iodide (Fig. 4). The addition of iodide to a suspension of the resin in $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ (v/v 1 : 1) resulted in an enhancement of fluorescence emission, which was attributed to increased configurational rigidity of the amido-crown macrocycle upon anion binding.³³ A calix[4]arene host **15** was synthesised by Singh and co-workers, which bore long aliphatic linkers ending in urea groups for an enlarged binding pocket that was complementary for iodide (Fig. 4). Unlike the majority of calixarene-based optical sensors, **15** did not contain a fluorophore tag; instead, the receptor was coupled to the surface of gold nanoparticles. Binding of iodide to a suspension of **15** in water induced the aggregation of nanoparticles, as seen by TEM, resulting in a colour change from light pink to blue.³⁴




Fig. 4 Structures of iodide-selective calix[4]arene receptors **14** (where the grey sphere represents a resin of 4-benzyloxybenzyl alcohol polymer) and **15**.




Fig. 5 Structure of an oxacalix[2]arene[2]triazine capable of sensing H_2PO_4^- ($K_a = 7.98 \times 10^3 \text{ M}^{-1}$ in CH_3CN).

An unusual oxacalix[2]arene[2]triazine host **16**, in which two of the benzene rings in the calix[4]arene core were replaced by triazines, was used as a sensor for H_2PO_4^- in acetonitrile (Fig. 5). The binding of the anion was due to a combination of hydrogen bonds with the pendant amide groups as well as anion- π interactions with the electron-poor triazine rings. Binding of H_2PO_4^- to **16** was accompanied by a quenching of the triazine fluorescence at 310 nm and the appearance of a new band at 412 nm.³⁵

2.2. Calix[n]pyrrole-based macrocycles

Calix[n]pyrroles are a well-studied class of macrocycles first synthesised by Baeyer in 1886,³⁶ and subsequently repurposed as an anion receptor by Sessler in 1996.³⁷ To elaborate these anion receptors into functional optical sensors, a natural approach is to attach a fluorophore or chromophore to the macrocyclic ring. This was first demonstrated by Sessler and co-workers, who prepared a series of β -functionalized calix[4]pyrrole anthracene conjugates (**17a-c**), which contain amide linkers of varying length (Fig. 6). These systems formed 1 : 1 complexes with various anions, which were shown by ^1H NMR studies to bind *via* multiple hydrogen bonds in the macrocyclic cavity. The addition of F^- , Cl^- and H_2PO_4^- in dichloromethane resulted in significant fluorescence quenching, attributed to an increase in PET, with electrons being transferred from the pyrrole rings to the anthracene fluorophore. Consistent with this mechanism, the quenching effect was most pronounced in **17a**, which contained the shortest linker.³⁸ Anzenbacher and co-workers recently reported a series of five calix[n]pyrrole-based macrocycles of varying ring sizes and bearing different fluorophores (**18a-e**) (Fig. 6). The macrocycles showed surprisingly high cross-reactivity for different anions, with little size discrimination observed. For example, in acetonitrile, the expanded calix[2]benzo[4]pyrrole **18d** exhibited similar or lower affinities for large carboxylates compared to the smaller calix[4]pyrrole analogue **18a**. Nonetheless, subtle differences in the photophysical responses of the five sensors towards different anions enabled the establishment of a fluorescence-based microarray that successfully classified 18 anions by linear discriminant analysis.³⁹




Fig. 6 Structures of calixpyrrole-based anion receptors 17–18 with chromophores highlighted in blue.

In a bid to improve the binding affinity of calix[4]pyrroles for anions, Lee and co-workers synthesised a bicyclic or 'strapped' calix[4]pyrrole **19**, which bears a dipyrrolylquinoxaline strap that serves the dual roles of providing additional sites for receptor–anion interactions as well as reporting anion binding *via* colour changes in the quinoxaline chromophore (Fig. 7). ^1H NMR anion binding studies in $\text{CD}_3\text{CN}/\text{DMSO}-d_6$ (9 : 1 v/v) showed that fluoride formed hydrogen bonds with the NH groups of the calix[4]pyrrole, but not the pyrrole groups of the strap, which did not undergo any appreciable downfield shift upon fluoride addition. Instead, the β -pyrrolic protons of the pyrrole groups in the strap shifted downfield by up to 0.3 ppm, which the authors attributed to an anion– π interaction between the electron-deficient pyrrole rings and the anion in the cavity. The binding of fluoride and dihydro-

gen phosphate to the receptor in $\text{CH}_3\text{CN}/\text{DMSO}$ (97 : 3 v/v) was accompanied by a 'naked eye' colour change from orange to dark blue.⁴⁰

A novel bis-calix[4]pyrrole-appended phenazine sensor **20**, which uses vibration induced emission (VIE) to differentiate between anions, was designed by Sessler and co-workers (Fig. 8). VIE is a phenomenon in which the fluorophore in question undergoes reversible vibrations between bent and planar configurations, resulting in multiple and tunable emission wavelengths from a single system.⁴¹ The free receptor adopts a planar configuration which has a long π -conjugation pathway and therefore emits orange-red fluorescence. Dubbed a 'molecular calliper' by the authors, **20** binds dicarboxylate anions in a bridging fashion between the two calix[4]pyrrole units, imposing strain on the phenazine core and inducing a




Fig. 7 Structure of 'strapped' calix[4]pyrrole **19** with quinoxaline fluorophore shown in blue. **19** binds F^- with $K_a = 8.97 \times 10^6 \text{ M}^{-1}$ in 97 : 3 $\text{CH}_3\text{CN}/\text{DMSO}$.




Fig. 8 Structure of a bis-calix[4]pyrrole-appended phenazine for anion sensing using VIE, with the phenazine core shown in blue. **20** binds a range of dicarboxylates with K_a values ranging from 10^5 – 10^7 M^{-1} in CH_3CN .



bending vibration that causes the emission to blueshift. In general, the shorter the dicarboxylate anion, the greater the degree of bending and therefore the observed blueshift, allowing structurally-similar anions to be distinguished in acetonitrile based on the colour of the fluorescence signal produced.⁴²

2.3. Amide- and urea-based macrocycles

The incorporation of hydrogen bonding donor groups directly into the core of the macrocycle is a common strategy for designing potent anion receptors, with amides, ureas and thioureas being some of the most widely utilised motifs.¹² An added advantage of the aforementioned groups is their ease of formation, allowing them to be readily constructed during the macrocyclization process, which greatly simplifies synthetic efforts. A diverse range of anion sensors based on amide or urea groups have been reported over the years,^{43–45} with some recent examples highlighted below.

Yoon and co-workers reported a chiral urea-based macrocycle **21** which contains an (*S*)-binaphthol (BINOL) unit that serves both as a chiral recognition site and a chromophore (Fig. 9). **21** bound *D*-Boc-alanine *via* hydrogen bonding interactions between the deprotonated carboxylate of the amino acid and the urea moieties of the macrocycle, resulting in marked enhancements in the UV-Vis absorption at 258 and 370 nm as well as a decrease at 342 nm in DMSO. Addition of *L*-Boc-alanine, however, induced only small changes in the absorption at these wavelengths, allowing discrimination of the two enantiomers.⁴⁶ In the following year, Anzenbacher and co-workers synthesised three chiral macrocycles (**22a–c**) decorated with multiple amide and/or sulfonamide groups (Fig. 9). BINOL was once again chosen as the chiral fluorophore, with both the (*R*)- and (*S*)-enantiomers of each macrocycle prepared. In addition, **22a** also possesses a chromogenic nitrophenyl group that enabled colorimetric detection of anions. The three sensors displayed significant cross-reactivity for simple inorganic ions as well as various chiral nucleoside phosphates,




Fig. 9 Structures of the chiral amide-based anion receptors **21–22**, with BINOL fluorophores shown in blue.







Fig. 10 (Top) Structures of tetraamide macrocyclic anion receptors with azulene units shown in blue. (Bottom) Crystal structure of **23b**·H₂PO₄[−], with hydrogen bonds between the receptor amide groups and H₂PO₄[−] shown as dashed lines. All other hydrogen atoms have been omitted for clarity. *K*_a (H₂PO₄[−]) = 4800 M^{−1} for **23a** and >10⁴ M^{−1} for **23b** in 9 : 1 DMSO/MeOH.

exhibiting moderate enantioselectivities. Promisingly, a multi-sensor array of the six sensors was capable of discriminating seven biologically relevant anions (AMP, ADP, ATP, CMP, GMP, PO₄^{3−} and P₂O₇^{4−}) in DMSO.⁴⁷

Two macrocyclic tetraamide receptors containing one (**23a**) or two (**23b**) azulene units were found to be selective for phosphate-containing anions (H₂PO₄[−] and HP₂O₇^{3−}) (Fig. 10). The addition of these anions to a DMSO solution of the asymmetrical receptor **23a** produced a redshift of the UV-vis absorption, which led to observable colour changes. Crystallographic studies showed that both the free and complexed receptors adopt a bent-sheet conformation that allows all four amides to participate in hydrogen bonding with the anionic guest, displaying a high degree of pre-organisation that favours anion binding. Notably, acyclic analogues of both receptors showed reduced affinity for phosphate, with no detectable spectroscopic changes.⁴⁸

Jolliffe and co-workers synthesised two semi-flexible tetra-thiourea macrocycles **24a–b** as receptors for ditopic anions, in order to study the effects of chelate cooperativity in multitopic anion receptors (Fig. 11). Although not the main focus of the study, fluorescence studies of **24a–b** in H₂O/DMSO (1 : 9 v/v) showed that the emission of the 1,8-diaminocarbazole groups was significantly quenched in the presence of dicarboxylates and sulfate, but not when mono-anions were added. This was attributed to increased PET from the thiourea moieties to the 1,8-diaminocarbazole groups and demonstrates the preference of the receptors for binding ditopic anions.⁴⁹

Amide or urea functionality can also be inserted into the pendant groups of the macrocycle, as demonstrated in a series of hybrid polyether/polyaza macrocycles **25a–d** synthesised by Núñez and co-workers (Fig. 12). Anion recognition in **25a–d**



typically proceeds *via* the formation of hydrogen bonds involving the amide/urea/thiourea groups on the arms in DMSO or acetonitrile. The receptors displayed significant cross-reactivity for a range of anions and also showed affinity for some metal cations, which were coordinated by the lone pairs of the oxygen or nitrogen groups forming the macrocyclic cavity. Binding events were signalled by colour changes associated with changes in the absorption of the nitrophenyl chromophore.^{50–52}

Although the literature on optical anion sensors based on macrocyclic hosts with neutral binding groups has been dominated by the types discussed in the preceding sections, there is certainly scope to design new macrocycles that contain novel anion binding groups or employ unusual sensing mechanisms.

Anion- π interactions, which occur between the HOMO of the anion and the LUMO of an electron-deficient aromatic ring, have been increasingly identified in the crystal structures of small molecules and proteins.⁵³ Recently, several examples of anion receptors based on anion- π interactions have been reported,⁵⁴ including two *s*-tetrazine-based macrocycle examples that possess optical sensing capabilities. Li and co-workers synthesised a tetrazine-containing diamide macrocycle **26**, which bound fluoride selectively through a combination of hydrogen bonding involving the amide NH groups as well as anion- π interactions involving the electron-poor tetrazine (Fig. 13). The addition of fluoride to a solution of **26** dissolved in DMSO led to a rapid colour change from red to green, which the authors attributed to the formation of a tetrazine radical anion due to electron transfer from the fluoride guest. The formation of the radical anion was further confirmed by

Photoswitchable anion receptors that can undergo reversible conformational changes have the potential to demonstrate altered affinities for different guests upon light irradiation. Bandyopadhyay and co-workers reported a fluorescent azobenzene-containing macrocycle **29** which resides predominantly in the ‘open’ *E* form, where it binds preferentially ATP over GTP in aqueous tris buffer (1 mM, pH 6.8) (Fig. 14). This is seen in a 7-fold fluorescence enhancement when ATP is added to **29**, whereas GTP only induces a 3-fold enhancement. Crystallographic and computational studies suggested that the protonated pyridinium group of **29** forms charged interactions with both GTP and ATP, but only the adenosine moiety of ATP contains an appropriately-positioned amine group to form an additional hydrogen bond with the azobenzene lone pair. However, upon irradiation with light at 366 nm, **29** photoisomerises to the ‘closed’ *Z* form, which is unable to form a hydrogen bond with ATP due to geometrical constraints. The

reduced affinity of the Z form for ATP is reflected in the low fluorescence enhancement (1.5-fold), which is slightly lower than the GTP-induced enhancement (1.9-fold).⁵⁸ While the findings are somewhat preliminary, this work demonstrates the potential of photoswitchable macrocycles with tunable affinities for ostensibly similar anionic guests.

Unlike the majority of optical anion sensors, which possess colorimetric and fluorescent signalling units, Pasini and co-workers designed a chiroptical probe **30** that responds to halide binding *via* changes in its circular dichroism spectrum in methylcyclohexane/dichloromethane (9:1, v/v) (Fig. 15). The halide guests docked in the cleft generated by the triazole units and phenyl spacer, forming weak CH-X⁻ hydrogen bonds with the triazole groups and the phenyl ring. The inclusion of halides in the macrocyclic cavity changed the dihedral angle of the BINOL group, which altered the intensity of its characteristic CD signal. Notably, these changes were not observed in the acyclic analogue, which is inherently conformationally flexible.⁵⁹

Another example of atypical anion sensor design comes in the form of four *trans*-homoporphodimethenes **31a–d** synthesised by Ravikanth and co-workers (Fig. 16). These receptors are irreversible chemodosimeter sensors that selectively recognise cyanide *via* covalent bond formation. Addition of cyanide to a THF solution of the **31a** led to significant fluorescence enhancement, with NMR studies suggesting this was due to nucleophilic attack of cyanide on the *meso*-carbon of the dipyrriin moiety, converting it to a cyanodipyrromethane.⁶⁰




Fig. 15 Structure of a chiroptical probe for halide sensing with BINOL shown in blue. The K_a values for binding of halides to **30** range from 8.9–18.1 M⁻¹ in CDCl₃.




Fig. 14 Photoswitchable azobenzene-based anion receptor **29** with photoisomerisable azobenzene groups shown in blue.




Fig. 16 Structure of homoporphodimethenes for selective cyanide sensing with nitrophenyl chromophore highlighted in blue.

The incorporation of cationic groups capable of forming strong electrostatic interactions with anions is another common approach to enhance the potency of macrocyclic anion receptors, particularly when these electrostatic forces are combined with more directional interactions like hydrogen or halogen bonding. Amines, in particular, facilitate reversible guest-binding as they can be protonated in acidic media to ‘turn-on’ the anion recognition capabilities of the host. In some cases, the binding of a basic anionic guest leads to deprotonation of the amine group, which quenches the emission of adjacent fluorophores *via* enhanced PET. Therefore, amines are useful and versatile groups that can assist in binding anions as well as inducing spectral changes in the receptor.

A handful of optical anion sensors rely predominantly on charged interactions established between multiple protonated amines and the anionic guest. Senge and co-workers employed a tetrakis(2-aminophenyl)porphyrin **32** as a selective host for pyrophosphate (Fig. 17). The hexaprotonated form of **32**, formed in the presence of excess TFA, exhibited exceptionally high affinity for pyrophosphate, with an association constant of $2.26 \times 10^9 \text{ M}^{-1}$ in CH_2Cl_2 . Pyrophosphate binding was accompanied by notable changes in the UV-Vis spectrum, including shifts in the Soret band, reduced intensity of the main porphyrin Q band and the appearance of two new Q bands, which resulted in a ‘naked eye’ colour change from yellow-brown to green. Interestingly, upon addition of excess of pyrophosphate (16 equiv.), the increased basicity of the solution causes deprotonation of the amino groups and regeneration of the free receptor. Addition of TFA reverses these changes, demonstrating switchable behaviour that could be repeated for at least eight cycles.⁶¹

Gellini and co-workers synthesised a tetra-aza macrocycle **33** containing an acridine fluorophore (Fig. 17). At low-to-intermediate pH, the free receptor exists predominantly in the +2 or +3 protonation state and can undergo a proton migration from one of the protonated ammonium groups to the acridine moiety, forming an acridinium-like fragment that fluoresces




Fig. 17 Structure of hexaprotonated tetrakis(2-aminophenyl)porphyrin host **32** (K_a ($P_2O_7^{4-}$) = $2.26 \times 10^9 \text{ M}^{-1}$ in CH_2Cl_2) and tetra-aza macrocycle **33** (K_s (F^-) = $3.98 \times 10^9 \text{ M}^{-1}$ and K_s (Cl^-) = $1.58 \times 10^{11} \text{ M}^{-1}$ in water).

intensely at 450 nm. Upon binding of fluoride or chloride in water, the emission is enhanced, with calculations suggesting that the F^- and Cl^- complexes of **33** preferentially undergo the proton transfer upon excitation to the S_2 state due to a reduction of the energy barrier for this reaction. In contrast, addition of bromide or iodide to **33** results in fluorescence quenching, which was attributed to the heavy atom effect.⁶²

Hybrid macrocycles that utilise both electrostatic interactions and hydrogen bonding interactions have been shown to act as potent anion sensors. Kataev and co-workers synthesised a series of hybrid amido-amine macrocycles **34a-c** of varying sizes (Fig. 18). **34a** showed an increase in fluorescence intensity upon addition of fluoride, whereas other halides and oxyanions induced aggregation and fluorescence quenching in aqueous MES buffer (10 mM, pH 6.2). This interesting optical response was attributed to the size of the macrocyclic cavity in **34a**, which was suitable for encapsulating up to three fluoride ions, encouraging protonation of the tertiary amine groups and hindering PET from the amine to the naphthalimide fluorophore. The larger ionic radii of other anions prevent more than one anion from fitting inside the binding cavity,




Fig. 18 Structures of hybrid amine-containing macrocycles **34–36** with various fluorophores shown in blue.

with the second anion bound on the periphery of the macrocycle, thereby resulting in aggregation by intermolecular π - π and electrostatic interactions.⁶³ The same group synthesised a similar hybrid amide-amine macrocycle **35** (Fig. 18), which was able to bind various nucleoside monophosphates, of which cGMP was most strongly bound ($K = 8.1 \times 10^4 \text{ M}^{-1}$ in aqueous acetate buffer with 2% DMSO). The binding of cGMP by the dicationic macrocycle led to fluorescence quenching due to π - π stacking of the guanine base with the pendant naphthalimide groups of the macrocycle.⁶⁴ Subsequently, a cage-like amido-amine receptor **36** was reported, which contains an anthracene fluorophore held in a 'strap' over the macrocyclic cavity (Fig. 18). The receptor formed a 1:2 receptor-anion complex with phosphate in aqueous solution. Initial binding of phosphate leads to fluorescence quenching due to deprotonation of the benzylic amines by the basic phosphate anion, enhancing PET to the anthracene moiety. In excess phosphate, a fluorescence enhancement is observed, attributed to the protonation of the tertiary amines to maintain charge balance when additional anions coordinate to the outside of the receptor cavity.⁶⁵

The versatility of amine groups is further demonstrated in their ability to bind both anionic and cationic guests. Chehri and co-workers synthesised a hybrid macrocycle **37** containing both ether and amine linkers, which served as a dual sensor for Fe(III) and iodide in aqueous solution (Fig. 19). Fe(III) was coordinated by the lone pairs of the ether and amine groups, leading to fluorescence quenching of the naphthalene groups *via* ligand-to-metal charge transfer, which opens a new non-radiative deactivation channel for electrons in the excited state. The addition of I^- to the naked receptor induces a change in the UV-vis absorbance spectrum, with the appearance of a new band at 360 nm, accompanied by a change from colourless to pale-pink. The fluorescence intensity also decreases due to the heavy atom effect.⁶⁶ Another dual sensor for Hg(II) and HSO_4^- consisted of a tetraazamacrocycle attached to a bromosalicylaldehyde pendant arm **38** (Fig. 19). Coordination of Hg(II) led to fluorescence quenching in $\text{CH}_3\text{CN}/\text{HEPES}$ buffer (2:8, v/v, 20 μM , pH 7.4). In contrast, the binding of HSO_4^- in the same solvent mixture, which was postulated to occur *via* multiple hydrogen bonds and/or charged interactions with the amine and hydroxyl groups of **38**, induced chelation-enhanced fluorescence by restricting the free rotation of the sensor.⁶⁷

Ionisable pyridine groups have also been used to enhance the binding of receptors to anions. Johnson and co-workers




Fig. 20 Structures of pyridine-based macrocyclic receptors for chloride sensing.

reported two pyridine-containing macrocycles **39a-b** (Fig. 20), which exhibit chloride-selective binding *via* a combination of hydrogen bonding from the amide/urea groups as well as charge-assisted hydrogen bonding from the protonated pyridinium moiety. The receptors were found to be unstable in aqueous solution, but could be incorporated into a thin film. In the solid state, intermolecular interactions between the fluorophore backbones induced a redshift of the emission peak when the receptors were exposed to an acidic solution of chloride.⁶⁸

3.2. Macrocycles with intrinsically charged groups

Cationic groups such as imidazolium and benzimidazolium units are capable of forming both electrostatic interactions and charge-assisted hydrogen bonds with anionic guests, and have been widely employed in a range of macrocyclic optical anion sensors. The majority of these hosts are cyclophane-like macrocycles consisting of multiple charged binding groups and signalling units connected by methylene spacers. Weakly-associated counterions such as PF_6^- are typically utilised in the preparation of such hosts to minimise competition for binding sites between the counterion and the target anion.

Some early examples of cyclophane-like benzimidazolium receptors were reported by Gao and co-workers.^{69,70} A series of acridine-containing hosts **40a-b** (Fig. 21), containing linkers of varying length and flexibility, were synthesised and found to bind H_2PO_4^- *via* charge-assisted hydrogen bonds between the two benzimidazolium CH groups and the anion. This resulted in a significant redshift of the acridine fluorescence from 430 nm to 556 nm in acetonitrile, which was attributed to the anion-induced assembly of excimers between the acridine rings of two macrocycles. The degree of redshift increased as ring size decreased, which likely indicated differing extents of excimer formation; in smaller rings, there was greater repulsion energy between the benzimidazolium rings in the ground state and largest stabilisation energy in the excited state, leading to the lowest-energy excimer.⁶⁹ Liu and co-workers recently reported a related pair of cyclophanes **41a-b**, possessing two dibenzimidazolium units and an anthraquinone fluorophore (Fig. 21). The smaller macrocycle **41a** exhibited fluorescence enhancement in the presence of acetate while the






Fig. 19 Structures of amine-based dual cation/anion receptors.



42

43



$X = \text{Cl}$ (**46a**)
 $X = \text{Br}$ (**46b**)
 $X = \text{I}$ (**46c**)
 $X = \text{H}$ (**46d**)

This journal is © The Royal Society of Chemistry 2021

and bromide ($K_a = 9.55 \times 10^5 \text{ M}^{-1}$) respectively. The binding of these ions was accompanied by a redshift and enhancement of the naphthalene emission band. Notably, the fluorescence response was observed even in a competitive $\text{CD}_3\text{OD}/\text{D}_2\text{O}$ (9:1) solvent mixture. In contrast, a hydrogen bonding analogue **46d** displayed no fluorescence response in the same solvent mixture, providing evidence for the greater strength and selectivity of halogen bonding for anion recognition in aqueous media.⁷⁷

While imidazolium-based motifs are commonly used in macrocyclic anion receptors, charged pyridinium groups have also been employed. A hybrid macrocycle **47**, containing four cationic pyridinium groups and four amide groups (Fig. 25), was found to preferentially bind 1,4-phenylenediacetate, which exhibits size complementarity for the large cavity. The dicarboxylate binds at the charged pyridinium sites and forms

hydrogen bonds with the nearby amide groups, as well as weak π -stacking interactions with the aromatic anthracene spacer. Fluorescence enhancement was observed upon anion addition to an acetonitrile solution of **47**, attributed to inhibition of PET from the binding sites to the anthracene moieties.⁷⁸ Recently, Cao and co-workers synthesised a octacationic cage **48** based on eight pyridinium moieties and two tetraphenylethylene (TPE) units. **48** was shown to bind the fluorescent dye sulforhodamine 101 (SR101), which contains two sulfonate groups that form electrostatic interactions with the pyridinium sites as well as an aromatic core that can form π -stacking interactions with the TPE moieties. Binding of SR101 to **48** led to quenching of the fluorescence emission at 545 nm as well as the appearance of a sharp emission peak at 621 nm, resulting in a fluorescence colour change from yellow to red under UV illumination.⁷⁹




Fig. 25 Structure of pyridinium-based macrocycles with fluorophores shown in blue.

4. Metallomacrocycles

The cationic and Lewis acidic nature of metal ions, as well as the interesting photophysical properties exhibited by some transition metal and lanthanide ions, has prompted the development of many metal-based systems for anion sensing.¹⁵ Combining metal ions with macrocyclic ligands, usually decorated with additional chromophores or anion-binding sites, can enhance the sensing capabilities of these systems.

Homodimetallic complexes of cryptand-like ligands have been used to selectively encapsulate anions that are of suitable size and geometry to bridge the metal centres. An early example was reported by Ugozzoli and co-workers in 2000, in which three di-copper bis-tren cryptands with linkers of varying length and flexibility (**49a–c**) demonstrated preferential binding for different anions (Fig. 26). **49a** exhibited a preference for linear anions like N_3^- and NCO^- , which are long enough to bridge the two $\text{Cu}(\text{II})$ centres. In contrast, the more flexible ether and 2,5-dimethylfuran spacers in **49b** and **49c** respectively enabled the receptors to rearrange into geometries suitable for binding various halide ions. The coordination of chloride to the copper centres of **49c** in particular gave rise to an intense absorption band at 410 nm in aqueous solution, which was attributed to ligand-to-metal charge transfer.⁸⁰




Fig. 26 Structures of dicopper cryptand-like anion receptors. **50** binds GTP as a 1:2 host–guest complex with $K_1 = 5.01 \times 10^4 \text{ M}^{-1}$ and $K_2 = 1.58 \times 10^4 \text{ M}^{-1}$ in 1:1 $\text{CH}_3\text{OH}/\text{H}_2\text{O}$.



A dicopper complex of an extended cryptand-like ligand **50** was found to bind nucleoside monophosphates in $\text{H}_2\text{O}/\text{CH}_3\text{OH}$ (1 : 1, v/v) (Fig. 26). The anionic guests bridge the two $\text{Cu}(\text{II})$ ions in the cryptand, binding through the phosphate group and a donor atom on the nucleobase. In the absence of a nucleoside monophosphate, **50** could form a non-fluorescent complex with the anionic dye 6-carboxyfluorescein. The dye was displaced upon binding of GMP to the receptor, switching on the yellow fluorescence emission of the free dye.⁸¹ This indicator-displacement method, in which the displacement of a dye (the indicator) from the receptor by the target anion switches on the fluorescence of the free dye, was later used extensively by Hossain and co-workers for the optical sensing of dicarboxylates. Over a series of publications, the authors demonstrated that dinuclear complexes of foldamer-like macrocyclic ligands could form non-fluorescent ion pairs with eosin Y (an anionic dye). The dye was readily displaced by dicarboxylates such as oxalate and citrate, which were bound more strongly by the receptors. Anion binding and the concomitant release of eosin Y was signalled by an increase in the characteristic fluorescence of the free dye.^{82–88}

In a twist on the indicator-displacement method, Zhang and co-workers synthesised a host a 2-(hydroxy)-naphthyl imino-functionalised pillar[5]arene **51** (Fig. 27), which coordinated to $\text{Fe}(\text{III})$ through its imine nitrogen and hydroxyl groups in $\text{H}_2\text{O}/\text{DMSO}$ (1 : 9, v/v). In the presence of $\text{Fe}(\text{III})$, the blue fluorescence of **51** was quenched. However, upon addition of fluoride, $\text{Fe}(\text{III})$ is displaced to form $[\text{FeF}_6]^{3-}$, restoring the fluorescence of the free receptor. Therefore, **51** could be used as either a 'turn-off' fluorescence sensor for $\text{Fe}(\text{III})$ or a 'turn-on' fluorescence sensor for fluoride.⁸⁹

Macrocyclic complexes of zinc have also been explored as anion sensors. A mononuclear $\text{Zn}(\text{II})$ complex of a bipyridine crown ether **52** (Fig. 28) was shown to undergo a 3-fold fluorescence enhancement in the presence of acetate, with smaller enhancements for larger carboxylates in $\text{H}_2\text{O}/\text{CH}_3\text{OH}$ (1 : 4, v/v, buffered at pH 7.4). The carboxylates were postulated to chelate the $\text{Zn}(\text{II})$ centre as well as form hydrophobic interactions with the crown ether ring; interestingly, an acyclic analogue of **52** showed no fluorescence response to carboxylates under the same conditions, suggesting at the importance of the crown ether.⁹⁰ A tetranuclear $\text{Zn}(\text{II})$ -Schiff base complex **53**, which contained multiple imine and amine groups (Fig. 28),




Fig. 27 Structure of pillar[5]arene receptor **51** functionalised with a 2-(hydroxy)-naphthyl imino fluorophore (in blue).




Fig. 28 Structures of Zn -based metallomacrocyclic anion receptors **52** ($K_a(\text{CH}_3\text{COO}^-) = 880 \text{ M}^{-1}$ in 1 : 4 $\text{H}_2\text{O}/\text{MeOH}$) and **53**.

underwent a blueshift of its absorption band upon addition of fluoride or acetate in aqueous methanolic solvent mixtures. Computational studies suggested this was due to a deprotonation of the iminium groups by the basic anions, making the iminium substituents less electron-withdrawing and widening the HOMO–LUMO gap of the adjacent aromatic ring. The spectral changes could be reversed by addition of a protic acid to the receptor–anion solutions.⁹¹

A handful of metallomacrocyclic anion sensors have been synthesised containing metal ions that do not directly coordinate to the anionic guests, but instead serve as fluorogenic or chromogenic signalling units, whilst providing a positively-charged environment to enhance anion binding. $\text{Ru}(\text{II})$ polypyridyl complexes have been widely used in chemosensing due to their favourable photophysical properties, including high photostability, long emission lifetimes and emission in the visible region.⁹² It is therefore unsurprising that Ru complexes are amongst the most common metal-based chromophores used in optical anion sensors.^{93–95} An early example reported in 1996 was a series of macrocyclic and calix[4]arene-based receptors **54a–d** containing bipyridyl units to which ruthenium (II) could be appended (Fig. 29). ^1H NMR studies in $\text{DMSO}-d_6$ indicated that receptors preferably bound H_2PO_4^- over Cl^- via convergent hydrogen bonds. A comparison of the two calix[4]arene receptors showed that **54c**, which contains shorter linkers and therefore a smaller binding cavity, exhibited stronger binding to all anions tested. The interaction of **54c** with H_2PO_4^- resulted in a blueshift of the $\text{Ru}(\text{II})$ MLCT emission band, accompanied by an increase in emission intensity. The latter was attributed to the rigidification of the macrocycle upon anion binding, which decreases the rate of non-radiative decay.⁹⁶ A series of hetero-dinuclear macrocycles containing ruthenium(II) tris(bipyridyl) alongside ferrocene, cobaltocenium and osmium(II) tris(bipyridyl) were subsequently synthesised and found to bind chloride strongly in acetonitrile with a concomitant enhancement in emission intensity.^{96,97}

Mukherjee and co-workers synthesised platinum-based square macrocycles **55a–c** which contain a conjugated Pt -






Fig. 29 Macrocycles incorporating ruthenium(II) bipyridyl complexes (shown in blue) for sensing of various anions.




Fig. 30 Structures of Pt-containing square macrocycles with carbazole fluorophores shown in blue. 55a binds $\text{P}_2\text{O}_7^{4-}$ with $K_a = 2.4 \times 10^4 \text{ M}^{-1}$ in DMF.

ethynyl-carbazole fluorophore and amide groups for anion binding (Fig. 30). The Pt(II) centres were not involved in anion binding; instead, they served the dual purposes of directing

the formation of square macrocycles and facilitating PET from the amide groups to the carbazole, quenching fluorescence from the excited state of the carbazole in the free receptor. Binding of pyrophosphate to the amide groups suppressed electronic transfer, leading to significant fluorescence enhancement in DMF.⁹⁸

5. Mechanically interlocked structures

Mechanically interlocked molecules (MIMs), such as rotaxanes and catenanes, possess topologically unique cavities that mimic the three-dimensional binding sites of proteins. These binding cavities provide a shielded hydrophobic environment to reduce the thermodynamic cost of binding, particularly in competitive solvents. The cavities can also be decorated with multiple pre-organised binding motifs that are complementary in geometry and size to the target guest. Building upon the seminal MIM metal template synthesis work of Sauvage in particular,^{99–103} in the past two decades, significant progress has been made in the development of complementary template-based approaches for MIM synthesis, including anion templation¹⁰⁴ and active metal templation^{105–108} strategies. These powerful synthetic methods have enabled the preparation of complex interlocked structures in greatly improved yields, making MIMs more than synthetic curiosities and unlocking the vast potential of these molecules as platforms for anion sensing.

5.1. MIMs containing organic luminophores

In an analogous manner to macrocyclic anion hosts, optical sensing capabilities can be introduced to MIMs by the incorporation of fluorogenic or chromophoric units that exhibit anion-induced spectral changes. The use of aromatic luminophores such as naphthalene, which also serve as spacer groups to adjust the size of the macrocycles, was a natural starting point.

A halogen bonding [2]catenane 56 was constructed from two identical macrocycles, each of which contained a cationic bromoimidazolium for anion binding as well as naphthalene groups as signalling units (Fig. 31). The [2]catenane selectively bound Br^- ($K_a = 1.48 \times 10^5 \text{ M}^{-1}$) and Cl^- ($K_a = 3.76 \times 10^6 \text{ M}^{-1}$) in acetonitrile solution to form 1 : 1 stoichiometric host-guest complexes, with molecular dynamics (MD) simulations indicating that the halide anions were held in the binding cavity by convergent halogen bonds. Anion binding was accompanied by a decrease in the emission band at 309 nm, alongside the appearance of a new broad band at 445 nm.¹⁰⁹ Another example of a [2]catenane-based anion sensor was the hetero-[2]catenane 57, which bound a range of anions *via* a combination of hydrogen bonding from isophthalamide group of one macrocycle and charge-assisted halogen bonding from the iodotriazolium of the second macrocycle (Fig. 31). Upon addition of anions, in particular acetate and H_2PO_4^- , an enhancement of the naphthalene emission bands was



An indolocarbazole-containing [2]rotaxane **58**, synthesised by active metal templation, displayed an unusual preference for oxyanions (CH_3COO^- and H_2PO_4^-) over halide ions in acetone- d_6 /D $_2$ O (95 : 5, v/v) (Fig. 32). A strong affinity for sulfate was also observed, possibly due to the complementary tetrahedral geometry of the dianion and the arrangement of the hydrogen bond donor groups in the binding cavity of **58**. The addition of anions induced a redshift in the emission band of the indolocarbazole fluorophore. In addition, chloride binding

Higher-order interlocked structures have been shown to produce interesting optical responses upon the binding of anions. A dicationic [3]rotaxane **59** (Fig. 33) containing a naphthalene reporter group was synthesised by chloride anion templation and formed 1:2 host-guest complexes with the monoanions Cl^- , Br^- and AcO^- , producing modest enhancements of fluorescence intensity arising from the increased rigidity of the receptor. In contrast, the addition of sulfate to a $\text{CDCl}_3/\text{CD}_3\text{OD}$ (1:1, v/v) solution of **59** initially resulted in a 1:1 stoichiometric complex with the anion bridging the two macrocycles. The rotary flexibility of the naphthalene linker allowed the rotaxane to fold around the sulfate guest to form a sandwich complex, quenching the naphthalene fluorescence through the formation of close contacts. The second equivalent of sulfate induced a conformational change in **59** to a more open form, forming a 1:1 host-guest complex in which each sulfate is associated with one macrocycle. This conformational change restored the fluorescence of the naphthalene units.¹¹²




Fig. 33 A dicationic [3]rotaxane which displays an unusual optical response to SO_4^{2-} ($K_1 > 10^4 \text{ M}^{-1}$ in 45 : 45 : 10 $\text{CDCl}_3/\text{CD}_3\text{OD}/\text{D}_2\text{O}$).




Fig. 32 A neutral indolocarbazole-containing [2]rotaxane which binds AcO^- ($K_a = 2.36 \times 10^3 \text{ M}^{-1}$) and H_2PO_4^- ($K_a = 2.05 \times 10^3 \text{ M}^{-1}$) in 95:5 acetone- $\text{d}_6/\text{D}_2\text{O}$).




Fig. 34 A chiral (*S*)-BINOL-containing [3]rotaxane with two bis(iodotriazolium) stations, which displays stereoselective binding of (*S*)-/(*R*)-glutamate ($K_S/K_R = 5.7 \pm 0.3$) and fumarate/malonate ($K_{\text{fumarate}}/K_{\text{malonate}} = 4.4 \pm 0.3$) in 60 : 39 : 1 $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{H}_2\text{O}$.

A chiral [3]rotaxane **60**, which possessed a central (*S*)-BINOL unit flanked by two halogen bonding 3,5-bis(iodotriazolium)pyridinium motifs in the axle (Fig. 34), was employed as a sensor for dicarboxylates. The dicarboxylate guests resided in the binding pocket between the two macrocycles, spanning the gap between the halogen bonding sites of the axle to form 1 : 1 stoichiometric sandwich complexes. Significant fluorescence quenching was observed in the presence of dicarboxylates, presumably due to the formation of close contacts between the bridging anion and the (*S*)-BINOL fluorophore, whereas the binding of smaller anions like chloride elicited much weaker responses. **60** exhibited an impressive ability to discriminate between both enantiomers (K_S/K_R for glutamate = 5.7 ± 0.3) and geometric isomers ($K_{\text{fumarate}}/K_{\text{malonate}} = 4.4 \pm 0.3$) in $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (60 : 39 : 1 v/v/v). MD simulations suggested the observed selectivity stemmed from differences in the number of synergistic halogen and hydrogen bonding interactions that could be formed between **60** and the respective stereo- or geometric isomers, alongside important contributions from anion-solvent interactions.¹¹³

5.2. MIMs containing metal-based luminophores

Luminescent metal complexes have also been incorporated into the structure of MIM-based anion receptors to produce effective optical sensors. An early example was a pair of cationic [2]rotaxanes **61a–b** with rhenium(i) or ruthenium(ii) bipyridyl complexes incorporated into the macrocyclic components (Fig. 35). The rotaxanes preferentially bound chloride over larger oxanions, which were unable to penetrate the binding cavity. This preference may have been a consequence of using a chloride template to construct the interlocked structure, producing a cavity that was complementary to the anion template. The addition of chloride or H_2PO_4^- induced an increase in fluorescence intensity in both rotaxanes, with the optical response persisting even in competitive solvent systems containing up to 10% water.¹¹⁴

A ruthenium(ii) bipyridyl complex was once again integrated into the macrocyclic component of a [2]rotaxane **62**, which contained permethylated β -cyclodextrin stoppers for water solubility and exhibited a fluorescence response to




Fig. 35 [2]Rotaxanes with a ruthenium(ii) (**61a**) or rhenium(i) (**61b**) complex integrated into the macrocycle which preferentially bind chloride ($K_a > 10^4 \text{ M}^{-1}$ for **61a** in 9 : 1 acetone/ D_2O and $K_a = 1500 \text{ M}^{-1}$ for **61b** in 7 : 3 acetone/ D_2O).






Fig. 36 A water-soluble ruthenium(II) bipyridyl-functionalised [2]rotaxane which binds iodide ($K_a = 6300 \text{ M}^{-1}$ in water).

anions in pure water (Fig. 36). NMR studies suggested that **62** formed 1 : 1 receptor–anion complexes in which the anionic guest resides in the cavity formed between the macrocycle and axle, where it is held by convergent hydrogen and halogen bonds. The receptor demonstrated a remarkably high affinity for iodide ($K_a = 6300 \text{ M}^{-1}$ in water), as well as moderate affinities for chloride ($K_a = 190 \text{ M}^{-1}$), bromide ($K_a = 1020 \text{ M}^{-1}$) and sulfate ($K_a = 450 \text{ M}^{-1}$). The addition of iodide and sulfate induced modest enhancements of the ruthenium(II) MLCT emission, attributed to rigidification of the receptor upon complexation with the anions, allowing the rotaxane to be employed as a luminescent sensor for these anions in water.¹¹⁵

Recently, luminescent 1,3-di(2-pyridyl)benzene complexes of Pt(II) and Ru(II) were incorporated into a range of acyclic, macrocyclic and interlocked anion hosts (**63a–f**) (Fig. 37). The acyclic Pt receptor **63a** bound anions in a 1 : 1 stoichiometry and underwent significant fluorescence enhancement of 60% upon addition of chloride or sulfate, attributed to rigidifica-

tion of the receptor upon anion coordination. On the other hand, the macrocyclic analogue **63c** showed a smaller fluorescence enhancement due to the higher pre-organisation and rigidity of the free receptor. Promisingly, preliminary luminescent titrations of the cationic [2]rotaxane **63f** with chloride suggested that the interlocked host exhibited a significantly higher affinity for the halide than that of the acyclic or macrocyclic analogues in a competitive solvent mixture of acetone- d_6 /D₂O (7 : 3, v/v).¹¹⁶

An all-halogen bonding [2]rotaxane **64**, which contains a rhenium(I) complex integrated into the macrocyclic component (Fig. 38), was shown to selectively bind halide ions over a range of oxanions. The preference for halides was attributed to complementarity in size and shape between the spherical anions and the binding cavity, with X-ray crystallographic studies confirming the presence of halogen bonds between the chloride anion and the iodotriazole/iodotriazolium groups. Luminescence titrations in solvent systems containing up to 50% water showed an initial increase in the




Fig. 37 Anion receptors incorporating 1,3-di(2-pyridyl)benzene fluorophores **63a–f**.






Fig. 38 [2]rotaxanes with a rhenium(I) complex integrated into the macrocycle. **64** binds various halides while **65** binds chloride preferentially with $K_a = 560 \text{ M}^{-1}$ in CHCl_3 .

intensity of the rhenium(I) MLCT band in the presence of halides, followed by fluorescence quenching when a large excess of the anions was added.¹¹⁷

A related rhenium-containing [2]rotaxane **65** (Fig. 38) was prepared *via* an elegant synthetic route involving an active template Cu-mediated alkyne-azide cycloaddition reaction (AT-CuAAC).¹¹⁸ A macrocycle containing a flexible bis-iodotriazole motif was used to bind the copper(I) template, which promoted a Cu-AAC reaction between the two stopper precursors to yield the interlocked host. After removal of the metal template, introduction of a more sterically-demanding rhenium(I) complex, which coordinated to the bis-iodotriazole group in an exotopic manner, inverted the bis-iodotriazole group so that the halogen bond donors were directed into the macrocyclic cavity. **65** preferentially bound chloride over larger halides and acetates, leading to a decrease in the broad rhenium(I) MLCT absorption band in CHCl_3 .

Surprisingly, despite the useful luminescent properties of lanthanide ions in sensing and imaging,¹¹⁹ the only report of a lanthanide-containing MIM used in optical anion sensing was a europium-containing [2]rotaxane **66**, which was syn-

thesised *via* nitrite anion templation (Fig. 39).¹²⁰ Upon removal of the nitrite template, it was found that fluoride addition caused substantial quenching of the Eu(III) emission in acetone/ H_2O (99 : 1 v/v). The increase in fluorescence lifetime and changes in the intensity of the hypersensitive $\Delta J = 2$ bands in fluoride-bound **66** suggested that fluoride was coordinated directly to the Eu(III) centre. Charge-assisted hydrogen bonds from the axle also enhanced the affinity of **66** for fluoride, as shown by the binding strength of the macrocyclic component alone. Nitrite and acetate led to weaker fluorescence quenching, which was attributed to a different mode of binding, wherein these anions did not directly coordinate to the emissive Eu(III) ion but were instead associated with the organic groups in the binding cavity.

A zinc metalloporphyrin was used as a signalling group in a hetero-[2]catenane **67**, which was synthesised *via* chloride anion templation, with auxiliary contributions from aromatic donor-acceptor interactions and pyridyl coordination to the apical site of Zn(II) centre (Fig. 40). These interactions further




Fig. 39 A [2]rotaxane with a europium(III) complex integrated into the macrocycle which binds fluoride ($K_a = 2.42 \times 10^5 \text{ M}^{-1}$ in 99 : 1 acetone/water).




Fig. 40 A hetero-[2]catenane incorporating a zinc porphyrin chromophore which selectively binds Cl^- ($K_a = 2144 \text{ M}^{-1}$ in DMSO-d_6).

served to pre-organise the interlocked cavity for chloride recognition. Crystallographic studies showed a high degree of complementarity between the binding cavity and the spherical halide, which was bound *via* six convergent CH–Cl and NH–Cl hydrogen bonds in DMSO. The encapsulation of chloride by **67** led to a gradual blueshift of the Soret band absorbance as well as slight increases in emission intensity.¹²¹

5.3. MIMs that undergo anion-induced shuttling

One of the most fascinating and useful properties of mechanically interlocked systems is the ability of the components to undergo reversible co-conformational changes in response to a stimulus. Due to this dynamic behaviour, MIMs are regarded as highly promising candidates for design of molecular machines and switches.^{122,123} A number of MIMs that undergo anion-induced conformational changes have been constructed and show promise as optical anion sensors. The majority of these systems are based on multi-station rotaxanes, which produce a photophysical response when the macrocycles translocate from a photoactive station to an anion binding station.

A simple chloride-selective [2]rotaxane **68** was synthesised by Goldup and co-workers using AT–CuAAC (Fig. 41). In the free receptor, strong intramolecular hydrogen bonds between the urea station of the axle and the bipyridine group of the

macrocycle precluded the binding of anions. However, protonation of the bipyridine group in acidic solutions forced the macrocycle to move away from the urea station and onto the adjacent triazole station. This displacement was inferred from ¹H NMR experiments and confirmed by crystallographic analysis, which additionally showed the formation of a new hydrogen bond between the protonated bipyridine and the triazole as well as π – π interactions between the bipyridine and the naphthalimide stopper. The ‘freed’ urea group in the protonated receptor was able to form hydrogen bonds with various anions, exhibiting a marked preference for chloride ($K_a > 10^4$ in 1 : 1 CDCl₃/CD₃CN) and inducing a fluorescence enhancement in **68**. The authors postulated this was due to a combination of receptor rigidification as well as the formation of π – π interactions with the naphthalimide group in the conformation adopted upon protonation of the [2]rotaxane.¹²⁴

Lin and co-workers prepared a BODIPY-stoppered three-station [2]rotaxane **69**, containing a central dibenzylammonium group and two peripheral triazolium stations (Fig. 42).^{125,126} In acidic solution, the crown ether-based macrocycle preferentially resides over the ammonium group, a co-conformation stabilised by hydrogen bonds between the ammonium NH groups and the oxygen atoms of the crown ether. The addition of OH[–] resulted in deprotonation of the




Fig. 41 (Left) Structure of urea-containing [2]rotaxane **68**; (middle) crystal structure of apo-receptor **68**; (right) crystal structure of **68**·HCl. Hydrogen bonds are shown in dashed lines and all other hydrogen atoms have been omitted for clarity. **68** binds Cl[–] strongly with $K_a > 10^4$ in 1 : 1 CDCl₃/CD₃CN.




Fig. 42 A BODIPY-stoppered [2]rotaxane with a central ammonium station and peripheral imidazolium anion binding stations which binds H₂PO₄[–] ($K = 3.75 \times 10^4 \text{ M}^{-1}$ in CH₃CN).



A hetero-[3]catenane **72**, comprising a large four-station central macrocycle and two smaller isophthalamide-containing macrocycles (Fig. 45), was constructed by chloride anion templation. The central macrocycle contains a tetrachloro-functionalised perylene diimide (PDI) unit as well as two triazolium anion-binding stations. The free receptor exhibited solvent-dependent co-conformational changes; in aprotic solvents like CHCl_3 , the smaller macrocycles reside predominantly on the triazolium stations, while the introduction of a polar protic solvent like CH_3OH induces circumrotatory motion of the small macrocycles to the two naphthalimide subunits of the PDI. This change in co-conformation was accompanied by a naked eye colour change from orange to red. The anion binding properties of **72** were thus investigated in 3 : 1 $\text{CHCl}_3/$

[illegible]

Org. Biomol. Chem., 2021, **19**, 4652–4677 | 4671




Fig. 44 A nitrate-selective [3]rotaxane with iodotriazolium anion binding stations and NDI chromophores. **71** binds NO_3^- with $K_a = 1782 \text{ M}^{-1}$ in 1 : 1 $\text{CDCl}_3/\text{CD}_3\text{OD}$ while the hydrogen bonding analogue exhibits a lower K_a of 653 M^{-1} .




Fig. 45 A hetero-[3]catenane incorporating a perylene diimide chromophore. **72** binds various anions as a 1 : 2 host–guest complex with highest affinity for Cl^- ($K_1 = 1.55 \times 10^4 \text{ M}^{-1}$ and $K_2 = 1.58 \times 10^2 \text{ M}^{-1}$ in 3 : 1 $\text{CHCl}_3/\text{CH}_3\text{OH}$).

CH_3OH , where the small macrocycles preferentially reside on the PDI unit. **72** was found to bind chloride in a 1 : 2 host–guest stoichiometry, with an accompanying colour change from red to orange. This is consistent with the loss of intramolecular interactions between the PDI stations and the hydroquinone groups of the small macrocycles, indicating that chloride binding to the triazolium units induced the concomitant movement of the small macrocycles to the triazolium binding sites. A significant enhancement of the PDI fluorescence was also observed upon chloride binding in a solvent


mixture of 45 : 45 : 10 $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{H}_2\text{O}$, allowing **72** to be used as a fluorescent anion sensor in competitive media.¹²⁹

An exotic fullerene-containing four-station [3]rotaxane **73** was shown to undergo dynamic shuttling and photophysical changes in the presence of chloride (Fig. 46). The axle component comprised a central C_{60} fullerene bis-triazolium motif flanked by two peripheral NDI stations, while the two macrocycles contained isophthalamide binding motifs tethered to ferrocene groups. The [3]rotaxane adopted two distinct co-conformations mediated by anion binding; in the presence of the non-coordinating anion PF_6^- , the ferrocenyl macrocycles reside predominantly on the NDI stations, while addition of chloride induced a translational motion of the macrocycles to the central bis-triazolium stations. In the co-conformation adopted in the PF_6^- salt of **73** in CHCl_3 solution, NDI emission is quenched due to PET from the ferrocenyl groups to the NDI stations. In the chloride-bound [3]rotaxane, the shuttling of the macrocycles to the central fullerene unit results in the formation of a C_{60} fullerene-based charge-separated state and also precludes ferrocenyl-NDI electron transfer, thereby switching on NDI emission.¹³⁰

5.4. MIMs that undergo anion-induced dissociation


The anion-triggered dissociation of MIMs into the respective axle and macrocyclic components can lead to a turn-on of fluorescence arising from the separation of a fluorogenic unit from a quencher. Reversible dissociation was first demonstrated using a pseudorotaxane **74** comprising an NDI-based axle with no stoppers and an isophthalamide naphthohydroquinone-containing macrocycle (Fig. 47). Although **74** is not strictly a MIM due to the lack of mechanical bonds in the structure, in





displaces the NDI axle by formation of hydrogen bonding interactions with the isophthalamide group, resulting in a discoloration of the solution in CHCl_3 .¹³¹

Jasti and co-workers synthesised a self-immolating [2]rotaxane **75** capable of selectively sensing fluoride (Fig. 48).¹³² The macrocyclic component of **75** was a fluorescent carbon ‘nanohoop’ while the asymmetrical axle is stoppered by a fluorescence-quenching 3,5-dinitrobenzyl unit and a fluoride-cleavable triisopropylsilane (TIPS). The intact [2]rotaxane is non-fluorescent in CHCl₃ due to the proximity of the quencher to the macrocyclic fluorophore. The addition of fluoride promoted cleavage of the TIPS stopper, allowing rapid de-threading of the axle from the macrocycle. The separation of the macrocycle from the quencher resulted in a dramatic 123-fold enhancement of fluorescence. A similar concept was employed by Zhu and co-workers using synthesised a [2]rotaxane **76** with a pillar[5]arene-based macrocycle and a silane-stoppered axle (Fig. 48). As before, disassembly of the rotaxane in the presence of fluoride restored the fluorescence of the pillar[5]arene in DMSO/THF (1 : 1, v/v). A simple test kit was made by adsorbing **76** onto strips of filter paper, which exhibited fluorescence turn-on when dipped in THF solutions containing fluoride.¹³³



Notes and references

Mechanically interlocked molecules have emerged as highly promising candidates for optical anion sensing due to the presence of three-dimensional binding cavities that can be exploited to provide shape and size complementarity to target anions. Future developments in MIM-based anion sensors may exploit the three-dimensional nature of the binding cavities to selectively detect environmentally-relevant oxyanions such as nitrate, which are challenging targets due to their high solvation and complex geometries. The propensity of interlocked systems to undergo anion-induced shuttling also provides a novel means of producing an optical response, as exemplified in a series of sophisticated anion sensors based on [3]rotaxanes and [3]catenanes. There is great scope to exploit the switchable nature of MIM-based molecular shuttles in the design of complex sensors and devices, and we foresee exciting developments in this direction.

There are no conflicts to declare.

H. M. T. acknowledges the Clarendon Fund and the Oxford Australia Scholarships Fund for a postgraduate research scholarship.

- 1 M. P. Anderson, R. J. Gregory, S. Thompson, D. W. Souza, S. Paul, R. C. Mulligan, A. E. Smith and M. J. Welsh, *Science*, 1991, **253**, 202.
- 2 F. Delange, *Thyroid*, 1994, **4**, 107–128.
- 3 J. L. Way, *Annu. Rev. Pharmacol. Toxicol.*, 1984, **24**, 451–481.
- 4 V. H. Smith and D. W. Schindler, *Trends Ecol. Evol.*, 2009, **24**, 201–207.
- 5 J. H. P. Watson and D. C. Ellwood, *Nucl. Eng. Des.*, 2003, **226**, 375–385.
- 6 P. K. Dasgupta, J. V. Dyke, A. B. Kirk and W. A. Jackson, *Environ. Sci. Technol.*, 2006, **40**, 6608–6614.
- 7 J. L. Sessler, P. A. Gale and W.-S. Cho, in *Anion Receptor Chemistry*, The Royal Society of Chemistry, 2006, pp. 320–369.
- 8 P. Prados and R. Quesada, *Supramol. Chem.*, 2008, **20**, 201–216.
- 9 C. H. Park and H. E. Simmons, *J. Am. Chem. Soc.*, 1968, **90**, 2431–2432.
- 10 G. T. Spence and P. D. Beer, *Acc. Chem. Res.*, 2013, **46**, 571–586.
- 11 M. J. Chmielewski, J. J. Davis and P. D. Beer, *Org. Biomol. Chem.*, 2009, **7**, 415–424.
- 12 P. A. Gale and C. Caltagirone, *Coord. Chem. Rev.*, 2018, **354**, 2–27.
- 13 D. A. McNaughton, M. Fares, G. Picci, P. A. Gale and C. Caltagirone, *Coord. Chem. Rev.*, 2021, **427**, 213573.
- 14 S. K. Sahoo, G.-D. Kim and H.-J. Choi, *J. Photochem. Photobiol., C*, 2016, **27**, 30–53.
- 15 A. B. Aleffi, D. M. Gillen and T. Gunnlaugsson, *Coord. Chem. Rev.*, 2018, **354**, 98–120.
- 16 R. Kumar, A. Sharma, H. Singh, P. Suating, H. S. Kim, K. Sunwoo, I. Shim, B. C. Gibb and J. S. Kim, *Chem. Rev.*, 2019, **119**, 9657–9721.
- 17 D. M. Homden and C. Redshaw, *Chem. Rev.*, 2008, **108**, 5086–5130.
- 18 F. Sansone, L. Baldini, A. Casnati and R. Ungaro, *New J. Chem.*, 2010, **34**, 2715–2728.
- 19 S. K. Kim, J. H. Bok, R. A. Bartsch, J. Y. Lee and J. S. Kim, *Org. Lett.*, 2005, **7**, 4839–4842.
- 20 E. Brunetti, J.-F. Picon, K. Flidrova, G. Bruylants, K. Bartik and I. Jabin, *J. Org. Chem.*, 2014, **79**, 6179–6188.
- 21 B.-Y. Hou, D.-X. Wang, H.-B. Yang, Q.-Y. Zheng and M.-X. Wang, *J. Org. Chem.*, 2007, **72**, 5218–5226.
- 22 P. G. Sutariya, H. Soni, S. A. Gandhi and A. Pandya, *New J. Chem.*, 2019, **43**, 737–747.
- 23 H.-F. Xie, C. Wu, J. Zou, Y.-X. Yang, H. Xu, Q.-L. Zhang, C. Redshaw and T. Yamato, *Dyes Pigm.*, 2020, **178**, 108340.
- 24 M. Nemati, R. Hosseinzadeh, R. Zadmand and M. Mohadjerani, *Sens. Actuators, B*, 2017, **241**, 690–697.
- 25 B. Uttam, R. Kandi, M. A. Hussain and C. P. Rao, *J. Org. Chem.*, 2018, **83**, 11850.
- 26 A. S. Miranda, P. M. Marcos, J. R. Ascenso, M. N. Berberan-Santos, R. Schurhammer, N. Hickey and S. Geremia, *Molecules*, 2020, **25**, 4708.

- 27 A. A. Vavilova, R. V. Nosov and I. I. Stoikov, *Mendeleev Commun.*, 2016, **26**, 508–510.
- 28 E. Quinlan, S. E. Matthews and T. Gunnlaugsson, *J. Org. Chem.*, 2007, **72**, 7497–7503.
- 29 Y. Chen, X. Liu, H. Huang, W. Wu and Y. Zheng, *Sci. China: Chem.*, 2010, **53**, 569–575.
- 30 Z.-Y. Li, H.-K. Su, H.-X. Tong, Y. Yin, T. Xiao, X.-Q. Sun, J. Jiang and L. Wang, *Spectrochim. Acta, Part A*, 2018, **200**, 307–312.
- 31 S. Arimori, M. G. Davidson, T. M. Fyles, T. G. Hibbert, T. D. James and G. I. Kociok-köhn, *Chem. Commun.*, 2004, **4**, 1640–1641.
- 32 J. Jin, J. Y. Park and Y. S. Lee, *J. Phys. Chem. C*, 2016, **120**, 24324–24334.
- 33 P. R. Sharma, S. Pandey, V. K. Soni, G. Choudhary and R. K. Sharma, *Supramol. Chem.*, 2019, **31**, 634–644.
- 34 M. Singh, N. Singh, J. R. Ascenso and P. M. Marcos, *Supramol. Chem.*, 2019, **31**, 313–321.
- 35 S. Li, D.-X. Wang and M.-X. Wang, *Tetrahedron Lett.*, 2012, **53**, 6226–6229.
- 36 A. Baeyer, *Ber. Dtsch. Chem. Ges.*, 1886, **19**, 2184–2185.
- 37 P. A. Gale, J. L. Sessler, V. Král and V. Lynch, *J. Am. Chem. Soc.*, 1996, **118**, 5140–5141.
- 38 J. Sessler, P. Anzenbacher, H. Miyaji, K. Jursikova, E. Bleasdale and P. A. Gale, *Ind. Eng. Chem. Res.*, 2000, **39**, 3471–3478.
- 39 M. Pushina, P. Koutnik, R. Nishiyabu, T. Minami, P. Savechenkov and P. Anzenbacher, *Chem. – Eur. J.*, 2018, **24**, 4879–4884.
- 40 J. Yoo, M.-S. Kim, S.-J. Hong, J. L. Sessler and C.-H. Lee, *J. Org. Chem.*, 2009, **74**, 1065.
- 41 W. Huang, L. Sun, Z. Zheng, J. Su and H. Tian, *Chem. Commun.*, 2015, **51**, 4462–4464.
- 42 W. Chen, C. Guo, Q. He, X. Chi, V. M. Lynch, Z. Zhang, J. Su, H. Tian and J. L. Sessler, *J. Am. Chem. Soc.*, 2019, **141**, 14798–14806.
- 43 W.-T. Gong, K. Hiratani and S. S. Lee, *Tetrahedron*, 2008, **64**, 11007–11011.
- 44 A. Costero, M. Bañuls, M. Aurell and M. Arellano, *J. Inclusion Phenom. Macrocyclic Chem.*, 2006, **54**, 61–66.
- 45 P. Piatek and J. Jurczak, *Chem. Commun.*, 2002, **2**, 2450–2451.
- 46 F. Wang, R. Nandhakumar, Y. Hu, D. Kim, K. M. Kim and J. Yoon, *J. Org. Chem.*, 2013, **78**, 11571–11576.
- 47 T. Ema, K. Okuda, S. Watanabe, T. Yamasaki, T. Minami, N. A. Esipenko and P. Anzenbacher, *Org. Lett.*, 2014, **16**, 1302.
- 48 D. Lichosyt, P. Dydio and J. Jurczak, *Chem. – Eur. J.*, 2016, **22**, 17673–17680.
- 49 S. N. Berry, L. Qin, W. Lewis and K. A. Jolliffe, *Chem. Sci.*, 2020, **11**, 7015–7022.
- 50 A. Aldrey, V. García, C. Lodeiro, A. Macías, P. Pérez-Lourido, L. Valencia, R. Bastida and C. Núñez, *Tetrahedron*, 2013, **69**, 4578–4585.
- 51 A. Aldrey, C. Núñez, V. García, R. Bastida, C. Lodeiro and A. Macías, *Tetrahedron*, 2010, **66**, 9223–9230.
- 52 C. Núñez, A. Aldrey, V. García, R. Bastida, A. Macías and C. Lodeiro, *Inorg. Chim. Acta*, 2012, **381**, 85–94.
- 53 I. A. Rather, S. A. Wagay and R. Ali, *Coord. Chem. Rev.*, 2020, **415**, 213327.
- 54 P. Molina, F. Zapata and A. Caballero, *Chem. Rev.*, 2017, **117**, 9907–9972.
- 55 Y. Zhao, Y. Li, Z. Qin, R. Jiang, H. Liu and Y. Li, *Dalton Trans.*, 2012, **41**, 13338–13342.
- 56 H.-B. Liu, Q. Zhang and M.-X. Wang, *Angew. Chem., Int. Ed.*, 2018, **57**, 6536–6540.
- 57 D.-H. Tuo, Y.-F. Ao, Q.-Q. Wang and D.-X. Wang, *Org. Lett.*, 2019, **21**, 7158–7162.
- 58 M. S. Hossain, S. A. Rahaman, J. Hatai, M. Saha and S. Bandyopadhyay, *Chem. Commun.*, 2020, **56**, 4172–4175.
- 59 M. Caricato, A. Olmo, C. Gargiulli, G. Gattuso and D. Pasini, *Tetrahedron*, 2012, **68**, 7861–7866.
- 60 P. Isar, M. Rajeswara Rao and M. Ravikanth, *Eur. J. Org. Chem.*, 2018, 3095–3104.
- 61 K. Norvaiša, K. J. Flanagan, D. Gibbons and M. O. Senge, *Angew. Chem., Int. Ed.*, 2019, **58**, 16553–16557.
- 62 R. Chelli, G. Pietrapertzia, A. Bencini, C. Giorgi, V. Lippolis, P. R. Salvi and C. Gellini, *Phys. Chem. Chem. Phys.*, 2015, **17**, 10813–10822.
- 63 A. S. Oshchepkov, T. A. Shumilova, S. R. Namashivaya, O. A. Fedorova, P. V. Dorovatovskii, V. N. Khrustalev and E. A. Kataev, *J. Org. Chem.*, 2018, **83**, 2145–2153.
- 64 A. S. Oshchepkov, T. A. Shumilova, M. Zerson, R. Magerle, V. N. Khrustalev and E. A. Kataev, *J. Org. Chem.*, 2019, **84**, 9034–9043.
- 65 B. S. Morozov, S. S. R. Namashivaya, M. A. Zakharko, A. S. Oshchepkov and E. A. Kataev, *ChemistryOpen*, 2020, **9**, 171–175.
- 66 R. Azadbakht and N. Chehri, *New J. Chem.*, 2018, **42**, 17690–17699.
- 67 M. Sethupathi, G. Muthusankar, V. Thamilarasan, N. Sengottuvelan, G. Gopu, N. M. Vinita, P. Kumar and F. Perdihi, *J. Photochem. Photobiol., B*, 2020, **203**, 111739.
- 68 C. L. Vonnegut, A. M. Shonkwiler, L. N. Zakharov, M. M. Haley and D. W. Johnson, *Chem. Commun.*, 2016, **52**, 9506–9509.
- 69 D. Zhang, X. Jiang, H. Yang, A. Martinez, M. Feng, Z. Dong and G. Gao, *Org. Biomol. Chem.*, 2013, **11**, 3375–3381.
- 70 D. Zhang, X. Jiang, H. Yang, Z. Su, E. Gao, A. Martinez and G. Gao, *Chem. Commun.*, 2013, **49**, 6149–6151.
- 71 Y. Liu, Z. Zhao, R. Huo and Q. Liu, *Sci. Rep.*, 2019, **9**, 502.
- 72 J.-H. Wang, J.-B. Xiong, X. Zhang, S. Song, Z.-H. Zhu and Y.-S. Zheng, *RSC Adv.*, 2015, **5**, 60096–60100.
- 73 J. Yang, C.-C. Dong, X.-L. Chen, X. Sun, J.-Y. Wei, J.-F. Xiang, J. L. Sessler and H.-Y. Gong, *J. Am. Chem. Soc.*, 2019, **141**, 4597–4612.
- 74 H.-T. Niu, Z. Yin, D. Su, D. Niu, J. He and J.-P. Cheng, *Dalton Trans.*, 2008, 3694.
- 75 J. Li, P. Zhang, Y. Xu, Z. Su, Y. Qian, S. Li, T. Yu, P. J. Sadler and H.-K. Liu, *Dalton Trans.*, 2017, **46**, 16205.
- 76 J. Y. C. Lim and P. D. Beer, *Chem*, 2018, **4**, 731–783.



- 77 F. Zapata, A. Caballero, N. G. White, T. D. W. Claridge, P. J. Costa, V. Félix and P. D. Beer, *J. Am. Chem. Soc.*, 2012, **134**, 11533.
- 78 K. Ghosh and A. R. Sarkar, *Tetrahedron Lett.*, 2009, **50**, 85–88.
- 79 H. Duan, Y. Li, Q. Li, P. Wang, X. Liu, L. Cheng, Y. Yu and L. Cao, *Angew. Chem.*, 2020, **59**, 10101–10110.
- 80 V. Amendola, E. Bastianello, L. Fabbrizzi, C. Mangano, P. Pallavicini, A. Perotti, A. Manotti Lanfredi and F. Ugozzoli, *Angew. Chem., Int. Ed.*, 2000, **39**, 2917–2920.
- 81 V. Amendola, G. Bergamaschi, A. Buttafava, L. Fabbrizzi and E. Monzani, *J. Am. Chem. Soc.*, 2010, **132**, 147–156.
- 82 J. S. Mendy, M. A. Saeed, F. R. Fronczek, D. R. Powell and M. A. Hossain, *Inorg. Chem.*, 2010, **49**, 7223–7225.
- 83 M. Rhaman, M. Hasan, A. Alamgir, L. Xu, D. Powell, B. Wong, R. Tandon and M. Hossain, *Sci. Rep.*, 2018, **8**, 286–286.
- 84 M. M. Rhaman, A. Alamgir, B. M. Wong, D. R. Powell and M. A. Hossain, *RSC Adv.*, 2014, **4**, 54263–54267.
- 85 M. M. Rhaman, F. R. Fronczek, D. R. Powell and M. A. Hossain, *Dalton Trans.*, 2014, **43**, 4618–4621.
- 86 M. A. Saeed, D. R. Powell and M. A. Hossain, *Tetrahedron Lett.*, 2010, **51**, 4904–4907.
- 87 M. Hu and G. Feng, *Chem. Commun.*, 2012, **48**, 6951–6953.
- 88 D. Hontz, J. Hensley, K. Hiriyak, J. Lee, J. Luchetta, M. Torsiello, M. Venditto, D. Lucent, W. Terzaghi, D. Mencer, A. Bommarreddy and A. L. Vanwert, *ACS Omega*, 2020, **5**, 19469.
- 89 W. Zhu, H. Fang, J.-X. He, W.-H. Jia, H. Yao, T.-B. Wei, Q. Lin and Y.-M. Zhang, *New J. Chem.*, 2018, **42**, 11548–11554.
- 90 P.-L. Ng, C.-S. Lee, H.-L. Kwong and A. S. C. Chan, *Inorg. Chem. Commun.*, 2005, **8**, 769–772.
- 91 T. Chakraborty, S. Dasgupta, A. Bhattacharyya, E. Zangrando, D. Escudero and D. Das, *New J. Chem.*, 2019, **43**, 13152–13161.
- 92 Q. Zhao, F. Li and C. Huang, *Chem. Soc. Rev.*, 2010, **39**, 3007–3030.
- 93 M. D. Pratt and P. D. Beer, *Tetrahedron*, 2004, **60**, 11227–11238.
- 94 A. Mishra, V. Vajpayee, H. Kim, M. H. Lee, H. Jung, M. Wang, P. J. Stang and K.-w. Chi, *Dalton Trans.*, 2012, **41**, 1195–1201.
- 95 P. De Wolf, P. Waywell, M. Hanson, S. L. Heath, A. J. H. M. Meijer, S. J. Teat and J. A. Thomas, *Chem. – Eur. J.*, 2006, **12**, 2188–2195.
- 96 F. Szemes, D. Hesek, Z. Chen, S. W. Dent, M. G. B. Drew, A. J. Goulden, A. R. Graydon, A. Grieve, R. J. Mortimer, T. Wear, J. S. Weightman and P. D. Beer, *Inorg. Chem.*, 1996, **35**, 5868–5879.
- 97 P. D. Beer, F. Szemes, V. Balzani, C. M. Salà, M. G. B. Drew, S. W. Dent and M. Maestri, *J. Am. Chem. Soc.*, 1997, **119**, 11864–11875.
- 98 S. Shanmugaraju, A. Bar, K. Chi and P. S. Mukherjee, *Organometallics*, 2010, **29**, 2971–2980.
- 99 C. O. Dietrich-Buchecker and J.-P. Sauvage, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 189–192.
- 100 P. Gaviña and J.-P. Sauvage, *Tetrahedron Lett.*, 1997, **38**, 3521–3524.
- 101 M. C. Jiménez, C. Dietrich-Buchecker and J.-P. Sauvage, *Angew. Chem., Int. Ed.*, 2000, **39**, 3284–3287.
- 102 J. F. Nierengarten, C. O. Dietrich-Buchecker and J. P. Sauvage, *J. Am. Chem. Soc.*, 1994, **116**, 375–376.
- 103 N. Solladié, J.-C. Chambron and J.-P. Sauvage, *J. Am. Chem. Soc.*, 1999, **121**, 3684–3692.
- 104 K. M. Bāk, K. Porfyrakis, J. J. Davis and P. D. Beer, *Mater. Chem. Front.*, 2020, **4**, 1052–1073.
- 105 J. E. M. Lewis, P. D. Beer, S. J. Loeb and S. M. Goldup, *Chem. Soc. Rev.*, 2017, **46**, 2577–2591.
- 106 J. E. Beves, B. A. Blight, C. J. Campbell, D. A. Leigh and R. T. McBurney, *Angew. Chem., Int. Ed.*, 2011, **50**, 9260–9327.
- 107 K. D. Hänni and D. A. Leigh, *Chem. Soc. Rev.*, 2010, **39**, 1240–1251.
- 108 J. D. Crowley, S. M. Goldup, A.-L. Lee, D. A. Leigh and R. T. McBurney, *Chem. Soc. Rev.*, 2009, **38**, 1530–1541.
- 109 A. Caballero, F. Zapata, N. G. White, P. J. Costa, V. Félix and P. D. Beer, *Angew. Chem., Int. Ed.*, 2012, **51**, 1876–1880.
- 110 J. M. Mercurio, A. Caballero, J. Cookson and P. D. Beer, *RSC Adv.*, 2015, **20**, 9298–9306.
- 111 A. Brown, T. Lang, K. M. Mullen and P. D. Beer, *Org. Biomol. Chem.*, 2017, **15**, 4587–4594.
- 112 M. J. Langton and P. D. Beer, *Chem. – Eur. J.*, 2012, **18**, 14406–14412.
- 113 J. Y. C. Lim, I. Marques, V. Félix and P. D. Beer, *Chem. Commun.*, 2018, **54**, 10851–10854.
- 114 L. M. Hancock, E. Marchi, P. Ceroni and P. D. Beer, *Chem. – Eur. J.*, 2012, **18**, 11277–11283.
- 115 M. J. Langton, I. Marques, S. W. Robinson, V. Félix and P. D. Beer, *Chem. – Eur. J.*, 2016, **22**, 185–192.
- 116 R. C. Knighton, S. Dapin and P. D. Beer, *Chem. – Eur. J.*, 2020, **26**, 5288–5296.
- 117 B. R. Mullaney, A. L. Thompson and P. D. Beer, *Angew. Chem., Int. Ed.*, 2014, **53**, 11458–11462.
- 118 M. J. Langton, Y. Xiong and P. D. Beer, *Chem. – Eur. J.*, 2015, **21**, 18910–18914.
- 119 C. Allain and S. Faulkner, *Future Med. Chem.*, 2010, **2**, 339–350.
- 120 M. J. Langton, O. A. Blackburn, T. Lang, S. Faulkner and P. D. Beer, *Angew. Chem., Int. Ed.*, 2014, **53**, 11463–11466.
- 121 A. Brown, M. J. Langton, N. L. Kilah, A. L. Thompson and P. D. Beer, *Chem. – Eur. J.*, 2015, **21**, 17664–17675.
- 122 J. F. Stoddart, *Angew. Chem., Int. Ed.*, 2017, **56**, 11094–11125.
- 123 A. W. Heard and S. M. Goldup, *ACS Cent. Sci.*, 2020, **6**, 117–128.
- 124 M. Denis, L. Qin, P. Turner, K. A. Jolliffe and S. M. Goldup, *Angew. Chem., Int. Ed.*, 2018, **57**, 5315–5319.



- 125 R. Arumugaperumal, V. Parthiban, T. Shukla, R. Putikam, R. Singh, S.-P. Wu, M.-C. Lin and H.-C. Lin, *Sens. Actuators, B*, 2018, **270**, 382–395.
- 126 R. Arumugaperumal, V. Srinivasadesikan, M. V. Ramakrishnam Raju, M.-C. Lin, T. Shukla, R. Singh and H.-C. Lin, *ACS Appl. Mater. Interfaces*, 2015, **7**, 26491–26503.
- 127 T. A. Barendt, S. W. Robinson and P. D. Beer, *Chem. Sci.*, 2016, **7**, 5171–5180.
- 128 T. A. Barendt, A. Docker, I. Marques, V. Félix and P. D. Beer, *Angew. Chem., Int. Ed.*, 2016, **55**, 11069–11076.
- 129 T. A. Barendt, L. Ferreira, I. Marques, V. Félix and P. D. Beer, *J. Am. Chem. Soc.*, 2017, **139**, 9026–9037.
- 130 T. A. Barendt, I. Rašović, M. A. Lebedeva, G. A. Farrow, A. Auty, D. Chekulaev, I. V. Sazanovich, J. A. Weinstein, K. Porfyrakis and P. D. Beer, *J. Am. Chem. Soc.*, 2018, **140**, 1924–1936.
- 131 K. M. Mullen, J. J. Davis and P. D. Beer, *New J. Chem.*, 2009, **33**, 769–776.
- 132 J. M. Van Raden, B. M. White, L. N. Zakharov and R. Jasti, *Angew. Chem., Int. Ed.*, 2019, **58**, 7341–7345.
- 133 Q. Li, Y. Wu, Y. Liu, L. Shangguan, B. Shi and H. Zhu, *Org. Lett.*, 2020, **22**, 6662.

