# **RSC Advances**



# **PAPER**

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2020, 10, 42164

# Preferential N-H···:C\(\simeg\) hydrogen bonding involving ditopic *NH*-containing systems and *N*-heterocyclic carbenes†

Zacharias J. Kinney, 📭 Arnold L. Rheingold and John D. Protasiewicz 📭 \*a

Hydrogen bonding plays a critical role in maintaining order and structure in complex biological and synthetic systems. N-heterocyclic carbenes (NHCs) represent one of the most versatile tools in the synthetic chemistry toolbox, yet their potential as neutral carbon hydrogen bond acceptors remains underexplored. This report investigates this capability in a strategic manner, wherein carbene-based hydrogen bonding can be assessed by use of ditopic NH-containing molecules. N-H bonds are unique as there are three established reaction modes with carbenes: non-traditional hydrogen bonding adducts ( $X-H\cdots:C\zeta$ ), salts arising from proton transfer ( $[H-C\zeta]^+[X]^-$ ), or amines from insertion of the carbene into the N-H bond. Yet, there are no established rules to predict product distributions or the strength of these associations. Here we seek to correlate the hydrogen bond strength of symmetric and asymmetric ditopic secondary amines with 1,3-bis(2,6-diisopropylphenyl)imidazol-2-ylidene (IPr, a representative NHC). In symmetric and asymmetric ditopic amine adducts both the solid-state (hydrogen bond lengths, NHC interior angles) and solution-state ( $^1H$   $\Delta\delta$  of NH signals,  $^{13}C$  signals of carbenic carbon) can be related to the  $pK_a$  of the parent amine.

Received 5th October 2020 Accepted 5th November 2020

DOI: 10.1039/d0ra08490e

rsc.li/rsc-advances

# Introduction

Hydrogen bonding interactions play a vital role throughout nature and are central to our understanding of complex biological systems.¹ The biological examples of such systems are incalculable, but prominent examples such as DNA, RNA, and enzymatic processes deserve mention.²-⁴ Non-biological examples are equally diverse, ranging from controlling the reactivity of small molecules,⁵ being central to the unique properties of water,⁶,⁷ to being a key mechanism for generating complex folded structures.⁵,⁶ While the role of hydrogen bonding in complex systems is indisputable, the development of methods to predict and assess the strengths of these interactions are vital to rationally designing novel systems.¹,¹,11

Hydrogen bond adducts X-H···A involve three-center fourelectron bonds formed by interaction of lone pair electrons on a hydrogen bond acceptor (A) with a  $\sigma^*$  orbital of a hydrogen bond donor (X-H).<sup>12</sup> In traditional hydrogen bonding, X is an electronegative atom (such as N, O, S, or F) which sufficiently polarizes the X–H bond leading to a reduction of the  $\sigma^*$  orbitals energy; and A is generally an electronegative atom (N, O, or halogens) having accessible lone pair electrons.<sup>12–14</sup> Nontraditional hydrogen bonds involving carbon centers are known but relatively rare, with association strengths that are generally weaker than conventional hydrogen bonds.<sup>15</sup> Most often these non-traditional species are mediated *via* formally charged carbon centers (such as isonitriles and carbanions) as the hydrogen bond acceptor.<sup>16,17</sup>

Since the first isolation of singlet carbenes their proliferation to all corners of chemistry are readily apparent. 18-22 Though a variety of singlet carbenes are now synthetically accessible, 23,24 N-heterocyclic carbenes (NHCs) based on an imidazole core remain one of the most well-studied and utilized class of carbenes.21,22,25-27 Shortly after the discovery of isolable NHCs, it was postulated that these compounds could serve as strong hydrogen bond acceptors: with their divalent carbon bearing a lone pair (σ-donor) to interact with a myriad of Brønsted acid X-H species. 28,29 Arduengo described the first example of a carhydrogen bond between bene-centered 1,3-bis(2,4,6trimethylphenyl)imidazol-2-ium hexafluorophosphate and 1,3bis(2,4,6-trimethylphenyl)imidazol-2-ylidene (IMes).<sup>30</sup> Clyburne and Davidson later expanded these findings to amine-based non-traditional hydrogen bonding between diphenylamine (DPA) and IMes (DPA···IMes, Fig. 1a), showing these species can be structurally authenticated.31 Movassaghi and Schmidt also identified a MeOH...IMes adduct in their efforts to catalyze

<sup>&</sup>quot;Department of Chemistry, Case Western Reserve University, Cleveland, Ohio 44106, USA. E-mail: protasiewicz@case.edu

<sup>&</sup>lt;sup>b</sup>Department of Chemistry and Biochemistry, University of California, La Jolla, San Diego, California 92093, USA

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental procedures, supplemental figures and analysis, NMR spectra, and crystallographic results. CCDC 2011390, 2011391 and 2011392. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d0ra08490e

Paper **RSC Advances** 

Fig. 1 (a) Representative authenticated X-H···:C( systems; 31-33 (b) general reaction progression elucidated by Bertrand of CAAC inserting into a N-H bond 34

amidation reactions with carbenes, authenticating the adduct both in solid- and solution-state (Fig. 1a).32

In addition to these reports, other computational<sup>35-39</sup> and experimental<sup>33,40-50</sup> studies on non-traditional X-H···:C\( hydrogen bonding have emerged (:C\square is neutral carbene species). The balance between non-traditional hydrogen bonding and proton transfer relies on the weak nucleophilicity but high Lewis basicity of NHCs, which can be further modulated via variation of the substituents on the imidazole-core. 51,52 Comparison of the available solid-state structures of X-H···:C\( hydrogen bonding reveal significant differences can occur. An example of this can be realized in comparing the hydrogen bond lengths (H···:C\( \) distance) in secondary amine-based adducts of carbazole (Cbz) and DPA with common NHCs 1,3bis(2,6-diisopropylphenyl)imidazol-2-ylidene (IPr) and IMes. The hydrogen bond lengths of Cbz···IPr (2.07(3) Å)<sup>33</sup> and DPA··· IMes (2.30(2) Å)<sup>31</sup> suggest a stronger interaction for the carbazole NH than for the DPA NH when employing similar NHCs.53 The interior N-C-N angles of the imidazole-2-ylidene units are slightly distorted as well, where Cbz···IPr = 102.2(2)° and DPA···IMes =  $101.7(1)^{\circ}$ .

These differences might be attributed to the differing steric profiles of IMes and IPr (2,4,6-trimethylphenyl, Mes versus 2,6diisopropylphenyl, Dipp), though more probably these differences are due to the acidities of **DPA** (p $K_a \approx 25.0$  in DMSO)<sup>54</sup> *versus* Cbz (p $K_a \approx 19.9$  in DMSO).<sup>55</sup> Consistent with the latter hypothesis, MeOH···IMes adduct (Fig. 1a) displays a shorter hydrogen bond length (1.96(2) Å) and a more pronounced N-C-N angle of 102.5(1)° (MeOH p $K_a \sim 29.0$  in DMSO)<sup>56,57</sup> than the amine-based species depicted in Fig. 1a. Although making comparisons between the reactivities of different NHCs can be troublesome, relatively minor differences in the basicity of IPr and IMes can be assumed, for the  $pK_a$  of their parent imidazoliums' are within error (p $K_a$  IPr·HCl = 19.29  $\pm$  0.07, p $K_a$ IMes·HCl = 19.40  $\pm$  0.12). 58,59

Though the isolation and authentication of hydrogen bonding adducts of carbenes has thus far been limited to imidazolylidenebased carbenes, hydrogen bonding adducts role in catalysis<sup>60-63</sup> and X-H insertion chemistry has been postulated for some time. Bertrand recently provided computational evidence for these non-traditional hydrogen bonded species as key intermediates for the reversible insertion of cyclic (alkyl)(amino)(carbene)s (CAACs) into the N-H bond of DPA (Fig. 1b).34 The calculations

Chart 1 Ditopic secondary amines.

suggested that the process of a CAAC inserting into an N-H bond proceeds via non-traditional hydrogen bonded and proton transfer intermediates.

Our recent work has established that Cbz and IPr form a hydrogen bonding adduct (Cbz...IPr, Fig. 1a) that has a strength of association in solution  $(C_6D_6)$  of -2.8 kcal mol<sup>-1</sup>.<sup>33</sup> Previous studies of non-traditional X-H···:C\( \) hydrogen bonding (X = N (ref. 31) or O (ref. 32, 47, 48 and 64)) did not quantitate thestrengths of interaction. Interestingly, the corresponding reaction of Cbz with CAAC from this study is reversible and yields the product of insertion of the carbene into the N-H bond of Cbz. The use of more strongly acidic N-H species with CAACs however results in proton transfer to forms salts.33,65 Collectively, these experimental efforts help "visualize" the proposed intermediates for insertion reactions of CAACs into X-H bonds.

While these efforts have proven fruitful in identifying new species with unique hydrogen bonds, there are no steadfast rules to predict reactivity or product distribution. Crystallographic information on H···:C\( \) hydrogen bonds shed some insight into the strength of these associations, but there are no obvious ways to correlate these metrics to solution-state measurements. Computational efforts have been made to determine the hydrogen bond basicity of carbenes, but thus far these measures have been purely theoretical. 66,67

It is of interest therefore to gain further insights into the factors that determine when hydrogen bonding occurs in preference to proton transfer or N-H insertion for reactions between isolable carbenes and NH-containing molecules. An important part of this effort involves delineating the strengths of non-conventional hydrogen bonding. In order to probe this latter question in more detail, a series of symmetric and asymmetric ditopic NH-containing systems consisting of Cbz and DPA units (Chart 1; 3,3'-bicarbazole,68 BC; N,N'-diphenylbenzidine, **DPB**; *N*,*N*′-diphenyl-*p*-phenylenediamine, **DPPD**, 4-(carbazol-3-yl)-N-phenylaniline, CD), which are structurally similar with differing acidities, will be reacted with a single NHC (IPr). These unusual platforms will allow for the gradual tuning of a single component (X-H) while having the carbene (A) remain consistent. This array of hydrogen bonded adducts will be used to assess how a NHC participates in non-traditional hydrogen bonding when presented with multiple reactive sites within single molecules.

# Results and discussion

#### BC…2IPr adduct

Previous studies on a Cbz...IPr hydrogen bonded adduct suggested that IPr could form a di-hydrogen bonded adduct with BC, and that this adduct should be labile and display




Fig. 2 Partial  $^1$ H NMR spectra (500 MHz, THF- $d_8$ ) of BC (50 mM) and BC...2IPr. Dashed lines indicate chemical shift changes upon hydrogen bonding to IPr.

concentration dependent behavior. <sup>1</sup>H NMR analysis of a 1:2 mixture of **BC** and **IPr** in THF- $d_8$  reveals that significant shifts  $(\Delta\delta > 0.05 \text{ ppm})$  occur for all proton resonances of the individual components, as highlighted in Fig. 2. In particular, the drastic downfield shift of  $H_{\alpha}$  ( $\Delta\delta \approx +1.7 \text{ ppm}$ ) is indicative of hydrogen bonding, and is similar to the corresponding shift of  $H_{\alpha}$  during formation of Cbz···IPr. The number of resonances indicate a symmetric structure for the **BC···2IPr** di-hydrogen bonded adduct. Consistent with this formulation, signals 1b and 2b show a substantial upfield shift ( $\Delta\delta \approx -0.9 \text{ ppm}$ ), since upon complexation they are positioned directly into the face of the Dipp groups on **IPr**. Inspection of the <sup>13</sup>C{<sup>1</sup>H} NMR spectrum of **BC···2IPr** reveals a single carbenic resonance ( $\delta$  215.2 ppm) that is noticeably upfield relative to the carbenic resonance of **IPr** ( $\delta$  221.1 ppm in THF- $d_8$ ).

Vapor diffusion of hexanes into a solution of **BC···2IPr** in THF yielded pale yellow crystals suitable for X-ray diffraction. Crystallographic analysis of one such crystal of **BC···2IPr** is portrayed in Fig. 3 (see ESI† for full details). The structure confirms a two-fold hydrogen bonding adduct, with an average H1···C13 distance of 2.10 Å and a N2–C12–N3 angle of 102.2°,




Fig. 3 ORTEP diagram (ellipsoids drawn at 50% probability) of BC··· 2IPr. Non-hydrogen bonding hydrogens and co-crystallized THF are omitted for clarity. Average select bond lengths (Å) and angles (°): H1··· C13, 2.10; N1···C13, 2.96; N2-C13-N3, 102.2; N2-N3-C12-C1, 13.5.

which are nearly identical to corresponding parameters for the mono-functional counterpart Cbz···IPr. Interestingly, the imidazole-2-ylidene  $C_3N_2$  and Cbz ring systems are nearly coplanar as indicated by  $\phi$  N2–N3–C12–C1 = 13.5°, which is in line with the monotopic Cbz···IPr system ( $\phi$  = 11.0°).

#### Model DPA…IPr adduct

Structural data for a DPA···IMes hydrogen bonded adduct were previously reported,31 but for the present study related details on the corresponding DPA···IPr adduct are necessary so that closer comparisons can be made. This material was synthesized by the addition of IPr to a THF solution of DPA. Removal of solvent, followed by recrystallization from toluene at -35 °C affords pure DPA...IPr as white crystals. Hydrogen bond formation is clearly indicted in C<sub>6</sub>D<sub>6</sub> by an upfield shift of the carbenic carbon resonance (218.2 ppm) compared to parent IPr (220.6 ppm in C<sub>6</sub>D<sub>6</sub>).<sup>69</sup> Preliminary <sup>1</sup>H NMR analysis of this material showed concentration dependent behavior. To ascertain the strength of this association a <sup>1</sup>H NMR concentration dependent study of DPA···IPr was performed in C<sub>6</sub>D<sub>6</sub> over a broad concentration range (75-1 mM). Benzene is an ideal solvent in such an analysis as its impact on hydrogen bonding is minimal,<sup>70</sup> while also allowing for comparison to our previous measurements on Cbz...IPr (C6D6, concentration range of 50-1 mM).33 This analysis revealed the association strength of **DPA**···**IPr** to be substantially weaker than Cbz···IPr ( $K_{\rm d} = 105 \pm$ 3  $M^{-1}$ ;  $\Delta G \approx -2.8 \text{ kcal mol}^{-1}$ ), with only the *NH* resonance being significantly influenced upon complexation ( $K_{\rm d}=0.85\pm$ 1.05 M<sup>-1</sup> and  $\Delta G \approx -0.10$  kcal mol<sup>-1</sup>, see ESI† for full details).<sup>71</sup> This drastic preference for the more acidic amine of Cbz was also verified via an exchange reaction (eqn (1)):

$$\mathbf{DPA} \cdots \mathbf{IPr} + \mathbf{Cbz} \rightleftharpoons \mathbf{Cbz} \cdots \mathbf{IPr} + \mathbf{DPA} \tag{1}$$

wherein addition of Cbz to **DPA···IPr** results in near complete conversion to Cbz···IPr and displacement of **DPA**. The dynamic (on the NMR timescale) nature of the mixture is indicated by the fact that a single hydrogen bonding *NH* resonance is observed (albeit not to the extent of the pure adducts, see ESI†) for each of the reaction pairs (**DPA···IPr/DPA** and Cbz···IPr/Cbz).

#### Ditopic DPB···2IPr and DPPD···2IPr adducts

Both ditopic **DPB** and **DPPD** yield 1 : 2 adducts **DPB···2IPr** and **DPPD···2IPr**, respectively, upon reaction with two equivalents of **IPr** (Fig. 4, top) in THF. As previously observed, **DPB···2IPr** and **DPPD···2IPr** are in equilibrium with their individual components, and at higher NMR concentrations these solutions exist predominately as the di-hydrogen bonded species. <sup>1</sup>H NMR analyses (**DPB···2IPr**, **DPPD···2IPr**), as shown in Fig. 4, highlights the subtle chemical shift changes ( $\Delta\delta$  < 0.20 ppm) for resonances compared to **BC···2IPr** ( $\Delta\delta$  > 0.15 ppm). Though these shifts in proton resonances are smaller, the overall directionality matches that observed for **BC···2IPr**. This is best realized in the *NH* signal, H<sub>β</sub>, which shifts by  $\Delta\delta$  = 0.20 ppm (**DPB···2IPr**) and  $\Delta\delta$  = 0.15 ppm (**DPPD···2IPr**), indicative of a significantly weaker hydrogen bond.

Paper

Fig. 4 Partial  $^1$ H NMR spectra (500 MHz, THF- $d_8$ ) of DPB (50 mM), DPB...2IPr, DPPD...2IPr, and DPPD (50 mM). Dashed lines highlight the chemical shift changes upon hydrogen bonding to IPr.

Crystals of **DPPD···2IPr** suitable for a X-ray diffraction studies were obtained by chilling a solution of **DPPD···2IPr** in a 3 : 2 hexanes : THF solvent mixture at  $-35\,^{\circ}$ C for several days. The resulting solid-state structure of **DPPD···2IPr** is portrayed in Fig. 5. The H1···C24 distance of 2.30(2) Å matches that found for the DPA···IMes adduct, as is the interior N2–C13–N3 angle of  $101.9(1)^{\circ}$ . The dihedral between the pseudo-planes of **DPPD···2IPr** ( $\phi$  N2–N3–C2–C4 =  $17^{\circ}$ ) are in agreement with **BC···2IPr**, and slightly less pronounced than DPA···IMes ( $\phi = 38^{\circ}$ ). The N1–H1···C24 bond angle is nearly linear ( $171^{\circ}$ ), albeit quite bent relative to DPA···IMes (N–H····C $< = 179^{\circ}$ ). The hydrogen bond distances are within the sum of the van der Waal radii, and display closer contacts than the hydrogen bond




Fig. 5 ORTEP diagram (ellipsoids drawn at 50% probability) of DPPD··· 2IPr. Non-hydrogen bonding hydrogens are omitted for clarity. Select bond lengths (Å) and angles (°): H1···C24, 2.30(2); N1···C24, 3.174(2); N2-C13-N3, 101.9(1); N2-N3-C4-C2, -17.41(7).

adducts of the Ong group, where carbenes with pendant secondary amines form dimers in the solid-state.<sup>44,45</sup> By contrast, all attempts to crystallize **DPB**···2**IPr** resulted in dissociation of the adduct into individual components. It appears that the poor solubility of **DPB** leads to preferential precipitation/crystallization of **DPB**, which is in facile equilibrium with **IPr** and **DPB**···2**IPr** (vide infra).

Analysis of the metric data obtained for the symmetric ditopic *NH*-bearing adducts in both solid- and solution-state reveals several interesting insights. In the solid-state the N–C–N angles within **IPr** ( $101.5^{\circ}$ )<sup>69</sup> relax upon formation of hydrogen bonded species **DPPD···2IPr** ( $101.9^{\circ}$ ) and **BC···2IPr** ( $102.2^{\circ}$ ). A greater distortion of the N–C–N angle is thus associate with greater interaction, and if taken to an extreme would result in proton transfer (where imidazolium salts typically feature N–C–N angles of  $\sim 108^{\circ}$ ).<sup>30</sup> The length of the hydrogen bonds (H····:C $\leq$  distance; **DPPD···2IPr**, 2.30 Å; **BC···2IPr**, 2.10 Å) also correlate well with the presumed strength of association based on measurements on the monofunctional amine counterparts (*vide supra*).

While  $^1H$  NMR spectra show that the changes in the chemical shifts for protons involved in hydrogen bonding can be predictive of association strengths, the shifts observed in the  $^{13}C\{^1H\}$  NMR spectra for the carbenic carbon are also quite sensitive to these external interactions. As the strength of the hydrogen bond increases the carbenic carbon shifts upfield. $^{30}$ 

Specifically, the  $^{13}$ C{ $^{1}$ H} NMR spectra of **BC**···**2IPr** (215.2 ppm) shows a greater  $\Delta \delta$  (free **IPr**  $\delta$  221.1 ppm) than either **DPB···2IPr** ( $\delta$  220.1 ppm) and **DPPD···2IPr** ( $\delta$  220.3 ppm).  $^{13}$ C spectroscopy might thus serve as a new tool for assessing the strength of the hydrogen bond in non-traditional systems, adding to other experimental measures used to assess the properties of carbenes as ligands (*i.e.* measuring  $\pi$ -acidity *via*  $^{31}$ P or  $^{77}$ Se NMR spectroscopy $^{72}$  and carbene-metal ligand strength assessed *via*  $^{13}$ C spectroscopy $^{73}$ ).

## Internal competitive hydrogen bonding

To probe the competitive nature of these hydrogen bonding adducts mixed carbazole-diphenylamine species CD was synthesized via Suzuki-Miyaura cross-coupling in modest yield (55% isolated, see ESI† for details). Reactions of CD with one and two equivalents of IPr were thus examined (Fig. 6, upper). As expected, formation of the mono- and di-carbene adducts CD...IPr and CD...2IPr is observable by NMR spectroscopy at elevated concentrations. Fig. 6 compares partial <sup>1</sup>H NMR spectra of these products contrasted against CD, highlighting the proton resonances impacted by NHC association. In alignment with the findings above, preferential interaction of the NHC occurs with the carbazole unit of CD upon addition of one equivalent of IPr (black dotted lines in Fig. 6). This assessment is confirmed by the signals for carbazole fragment of CD shifting in the same fashion as the symmetric BC···2IPr, with the caveat that shifts are only  $\sim$ 65% the overall  $\Delta\delta$  of their symmetric counterpart. This may reflect the fact that IPr equilibrates between both carbazole and diphenylamine NH residues to some extent. The perturbations of the proton




Fig. 6 Partial  $^1$ H NMR spectra (500 MHz, THF- $d_8$ ) of CD (50 mM), CD···IPr, and CD···2IPr, highlighting the chemical shift changes upon hydrogen bonding to IPr. CD H $_{\alpha}$  (10.25 ppm), CD···IPr H $_{\alpha}$  (11.30 ppm), and CD···2IPr H $_{\alpha}$  (12.40 ppm) are broadened due to exchange, see ESI† (rings 1 and 2 = black dotted lines, rings 3 and 4 = orange dotted lines).

resonances on rings 3 and 4 (orange dotted lines) are reminiscent of **DPB**···**2IPr** and **DPPD**···**2IPr**, albeit only minor in comparison to rings 1 and 2.

Upon addition of two equivalents of **IPr** to **CD**, the <sup>1</sup>H NMR chemical shifts of CD...2IPr are a near match to the symmetric diamine counterparts, BC ... 2IPr and DPB ... 2IPr respectively. The realignment of the <sup>1</sup>H spectra to mirror the symmetric analogues confirms formation of CD...2IPr. Due to the rapid exchange of **IPr** between both *NH* sites on **CD** the <sup>13</sup>C{<sup>1</sup>H} NMR spectra yield a singular shift for the carbenic resonance. The observed shifts for asymmetric CD···2IPr (δ 217.4 ppm) and **CD···IPr** ( $\delta$  216.4 ppm) fall between their respective symmetric counterparts (BC···2IPr,  $\delta$  215.2 ppm, DPB···2IPr,  $\delta$  220.1 ppm). Unsurprisingly the <sup>13</sup>C{<sup>1</sup>H} NMR signal in CD···IPr resides closer to BC···2IPr, verifying our <sup>1</sup>H measurements which show that IPr preferentially binds to the carbazole side of the ditopic system. When both NH sites are satisfied in CD...2IPr the carbenic carbon resonance is a near perfect average of the observed resonances for the symmetric species (BC···2IPr and DPB···2IPr average resonance carbenic resonance  $\approx 217.6$  ppm).

Unfortunately attempts to obtain single crystals of either CD···IPr and CD···2IPr were thwarted by the preferential crystallization of CD from solutions of these materials (see ESI† for crystallographic details). While this may be due to the reduced solubility of CD relative to IPr, the probable cause is the labile nature of the N-H····:C bond not being conducive to solid-state analysis.

Table 1 Summation of solid- and solution-state metrics<sup>a</sup>

	N-C-N (°)	$^{13}$ C $\delta$ (ppm)	$^{1}$ H $\Delta\delta$ (ppm)	$pK_a$
IPr	$101.5^{b}$	221.1	_	19.3 <sup>c</sup>
BC···2IPr	102.2	215.2	1.66	19.9
DPB…2IPr	_	220.1	0.20	25.0
$DPPD\cdots 2IPr$	101.9	220.3	0.15	25.0
CD···2IPr	_	217.4	$0.99^{d}$	$22.5^{d}$

<sup>&</sup>lt;sup>a</sup> <sup>1</sup>H  $\Delta \delta$  is calculated relative to each adducts parent species at 50 mM in THF- $d_8$ . <sup>b</sup> See ref. 69. <sup>c</sup> p $K_a$  of the conjugate acid (**IPr**·HCl). <sup>58,59</sup> <sup>d</sup> Average values.

# Conclusion

Through a series of symmetric and asymmetric ditopic secondary amines the strength of a X-H···:C\( \) hydrogen bond can be directly assessed via several solid- and solution-state factors, summarized in Table 1. Paramount in predicting the strength of the association is the  $pK_a$  of the parent X-H species, which in these examples correlates well with both solid- and solution-state metrics. In the solid-state limiting the degrees of freedom in the parent molecule is vital, wherein systems with even weak hydrogen bonds can be crystallized in their adduct form (DPPD···2IPr). In BC, DPB, and DPPD systems solution-state analysis is simplified due their high degree of symmetry and rapid equilibrium of the N-H...IPr interaction. In each of these species both the  $\Delta\delta$  of the proton participating in hydrogen bonding and the <sup>13</sup>C resonance of the carbenic carbon are indicators of hydrogen bond strength. 1H NMR spectra show that upon hydrogen bonding with IPr the monomers are rigidified, with proton resonances far removed from area of interaction changing significantly in comparison to the unhindered molecules. When starving the asymmetric CD system of **IPr**, the carbene primarily interacts with the more acidic carbazole residue as evidenced by both <sup>1</sup>H and <sup>13</sup>C spectroscopy. When enough IPr is supplied to satisfy both NH sites the carbenic carbon resonance falls directly between the symmetric <sup>13</sup>C residues, indicating a system that is in rapid equilibrium on the NMR timescale. The reversible nature of these systems and the ability to predict hydrogen bond strength through both solid- and solutionstate metrics could lead to further developments in systems chemistry. Efforts to explore higher functionality NH-containing systems and carbenes are currently underway.

### Conflicts of interest

The authors declare no conflicts of interest.

# Acknowledgements

We thank the National Science Foundation (CHE-1464855 and CHE-1955845) for their support of this work.

# Notes and references

1 G. A. Jeffrey and W. Saenger, *Hydrogen Bonding in Biological Systems*, Springer, Berlin, 1991.

Paper

2 L. McFail-Isom, C. C. Sines and L. D. Williams, DNA Structure: Cations in Charge?, *Curr. Opin. Struct. Biol.*, 1999, **9**, 298–304.

- 3 J. A. Subirana and M. Soler-López, Cations as Hydrogen Bond Donors: A View of Electrostatic Interactions in DNA, *Annu. Rev. Biophys. Biomol. Struct.*, 2003, **32**, 27–45.
- 4 D. Meyer, P. Neumann, R. Ficner and K. Tittmann, Observation of a Stable Carbene at the Active Site of a Thiamin Enzyme, *Nat. Chem. Biol.*, 2013, **9**, 488–490.
- 5 A. G. Doyle and E. N. Jacobsen, Small-Molecule H-Bond Donors in Asymmetric Catalysis, *Chem. Rev.*, 2007, **107**, 5713–5743.
- 6 P. Gallo, K. Amann-Winkel, C. A. Angell, M. A. Anisimov, F. Caupin, C. Chakravarty, E. Lascaris, T. Loerting, A. Z. Panagiotopoulos, J. Russo, J. A. Sellberg, H. E. Stanley, H. Tanaka, C. Vega, L. Xu and L. G. M. Pettersson, Water: A Tale of Two Liquids, *Chem. Rev.*, 2016, 116, 7463-7500.
- 7 F. Cipcigan, V. Sokhan, G. Martyna and J. Crain, Structure and Hydrogen Bonding at the Limits of Liquid Water Stability, *Sci. Rep.*, 2018, **8**, 1–8.
- 8 S. H. Gellman, Foldamers: A Manifesto, *Acc. Chem. Res.*, 1998, **31**, 173–180.
- 9 D. Núñez-Villanueva, G. Iadevaia, A. E. Stross, M. A. Jinks, J. A. Swain and C. A. Hunter, H-Bond Self-Assembly: Folding versus Duplex Formation, *J. Am. Chem. Soc.*, 2017, 139, 6654–6662.
- 10 H. Szatyłowicz, Structural Aspects of the Intermolecular Hydrogen Bond Strength: H-Bonded Complexes of Aniline, Phenol and Pyridine Derivatives, *J. Phys. Org. Chem.*, 2008, 21, 897–914.
- 11 S. J. Pike, E. Lavagnini, L. M. Varley, J. L. Cook and C. A. Hunter, H-Bond Donor Parameters for Cations, *Chem. Sci.*, 2019, **10**, 5943–5951.
- 12 M. D. Joesten, Hydrogen Bonding and Proton Transfer, *J. Chem. Educ.*, 1982, **59**, 362–366.
- 13 J. Emsley, Very Strong Hydrogen Bonding, *Chem. Soc. Rev.*, 1968, 9, 91–124.
- 14 R. H. Crabtree, Hypervalency Secondary Bonding and Hydrogen Bonding: Siblings under the Skin, *Chem. Soc. Rev.*, 2017, **46**, 1720–1729.
- 15 G. R. Desiraju and T. Steiner, *The Weak Hydrogen Bond: In Structural Chemistry and Biology*, 2001.
- 16 J. A. Platts, S. T. Howard and K. Woźniak, Quantum Chemical Evidence for C-H···C Hydrogen Bonding, *Chem. Commun.*, 1996, 1, 63–64.
- 17 A. S. Batsanov, M. G. Davidson, J. A. K. Howard, S. Lamb and C. Lustig, Phosphonium Ylides as Hydrogen Bond Acceptors: Inter Molecular C–H···C Interactions in the Crystal Structure of Triphenylphosphonium Benzylide, *Chem. Commun.*, 1996, **15**, 1791–1792.
- 18 A. J. Arduengo, R. L. Harlow and M. Kline, A Stable Crystalline Carbene, *J. Am. Chem. Soc.*, 1991, 113, 361–363.
- 19 M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, An Overview of *N*-Heterocyclic Carbenes, *Nature*, 2014, **510**, 485–496.

- 20 J. Chen and Y. Huang, Asymmetric Catalysis with *N*-Heterocyclic Carbenes as Non-Covalent Chiral Templates, *Nat. Commun.*, 2014, 5, 1–8.
- 21 V. Nesterov, D. Reiter, P. Bag, P. Frisch, R. Holzner, A. Porzelt and S. Inoue, NHCs in Main Group Chemistry, *Chem. Rev.*, 2018, **118**, 9678–9842.
- 22 E. Peris, Smart *N*-Heterocyclic Carbene Ligands in Catalysis, *Chem. Rev.*, 2018, **118**, 9988–10031.
- 23 M. Soleilhavoup and G. Bertrand, Cyclic (Alkyl)(Amino) Carbenes (CAACs): Stable Carbenes on the Rise, *Acc. Chem. Res.*, 2015, **48**, 256–266.
- 24 M. Melaimi, R. Jazzar, M. Soleilhavoup and G. Bertrand, Cyclic (Alkyl)(Amino)Carbenes (CAACs): Recent Developments, Angew. Chem., Int. Ed., 2017, 56, 10046– 10068
- 25 A. J. Arduengo, Looking for Stable Carbenes: The Difficulty in Starting Anew, *Acc. Chem. Res.*, 1999, **32**, 913–921.
- 26 R. Visbal and M. C. Gimeno, *N*-Heterocyclic Carbene Metal Complexes: Photoluminescence and Applications, *Chem. Soc. Rev.*, 2014, 43, 3551–3574.
- 27 A. Doddi, M. Peters and M. Tamm, N-Heterocyclic Carbene Adducts of Main Group Elements and Their Use as Ligands in Transition Metal Chemistry, Chem. Rev., 2019, 119, 6994–7112.
- 28 I. Alkorta, I. Rozas and J. Elguero, Non-Conventional Hydrogen Bonds, *Chem. Soc. Rev.*, 1998, 27, 163–170.
- 29 J. K. W. Chui, T. Ramnial and J. A. C. Clyburne, N-Heterocyclic Carbenes: Exotic Molecules as Precursors to Unusual Hydrogen Bonds, *Comments Inorg. Chem.*, 2003, 24, 165–187.
- 30 A. J. I. Arduengo, S. F. Gamper, M. Tamm, J. C. Calabrese, F. Davidson and H. A. Craig, A Bis(Carbene)—Proton Complex: Structure of a C—H—C Hydrogen Bond, *J. Am. Chem. Soc.*, 1995, **117**, 572–573.
- 31 J. A. Cowan, J. A. C. Clyburne, M. G. Davidson, R. L. W. Harris, J. A. K. Howard, P. Küpper, M. A. Leech and S. P. Richards, On the Interaction between *N*-Heterocyclic Carbenes and Organic Acids: Structural Authentication of the First N—H···C Hydrogen Bond and Remarkably Short C—H···O Interactions, *Angew. Chem., Int. Ed.*, 2002, 41, 1432–1434.
- 32 M. Movassaghi and M. A. Schmidt, *N*-Heterocyclic Carbene-Catalyzed Amidation of Unactivated Esters with Amino Alcohols, *Org. Lett.*, 2005, 7, 2453–2456.
- 33 J. M. Kieser, Z. J. Kinney, J. R. Gaffen, S. Evariste, A. M. Harrison, A. L. Rheingold and J. D. Protasiewicz, Three Ways Isolable Carbenes Can Modulate Emission of NH-Containing Fluorophores, *J. Am. Chem. Soc.*, 2019, 141, 12055–12063.
- 34 D. Tolentino, S. Neale, C. Isaac, S. A. Macgregor, M. K. Whittlesey, R. Jazzar and G. Bertrand, Reductive Elimination at Carbon under Steric Control, *J. Am. Chem. Soc.*, 2019, 141, 9823–9826.
- 35 I. Alkorta and J. Elguero, Carbenes and Silylenes as Hydrogen Bond Acceptors, *J. Phys. Chem.*, 1996, **100**, 19367–19370.

- 36 A. M. Magill, K. J. Cavell and B. F. Yates, Basicity of Nucleophilic Carbenes in Aqueous and Nonaqueous Solvents—Theoretical Predictions, *J. Am. Chem. Soc.*, 2004, 126, 8717–8724.
- 37 J.-L. Mieusset and U. H. Brinker, The Carbene Reactivity Surface: A Classification, J. Org. Chem., 2008, 73, 1553–1558.
- 38 O. Hollóczki, Unveiling the Peculiar Hydrogen Bonding Behavior of Solvated *N*-Heterocyclic Carbenes, *Phys. Chem. Chem. Phys.*, 2016, **18**, 126–140.
- 39 J. E. Del Bene, I. Alkorta and J. Elguero, Hydrogen-Bonded Complexes with Carbenes as Electron-Pair Donors, *Chem. Phys. Lett.*, 2017, 675, 46–50.
- 40 L. A. Leites, G. I. Magdanurov, S. S. Bukalov and R. West, Intermolecular = C-H····: C Hydrogen Bond in a Crystalline Unsaturated Arduengo-Type Carbene, *Mendeleev Commun.*, 2008, **18**, 14–15.
- 41 M. Kapitein, M. Balmer and C. von Hanisch, A Sterically Encumbered 13/15 Cycle and Its Cleavage with *N*-Heterocyclic Carbenes and Other Lewis Bases, *Z. Anorg. Allg. Chem.*, 2016, **642**, 1275–1281.
- 42 P. A. Petrov, T. S. Sukhikh and M. N. Sokolov, NHC Adducts of Tantalum Amidohalides: The First Example of NHC Abnormally Coordinated to an Early Transition Metal, *Dalton Trans.*, 2017, **46**, 4902–4906.
- 43 H. Jong, B. O. Patrick and M. D. Fryzuk, Amine-Tethered N-Heterocyclic Carbene Complexes of Rhodium(I), *Can. J. Chem.*, 2008, **86**, 803–810.
- 44 W. C. Shih, C. H. Wang, Y. T. Chang, G. P. A. Yap and T. G. Ong, Synthesis and Structure of an Amino-Linked *N*-Heterocyclic Carbene and the Reactivity of Its Aluminum Adduct, *Organometallics*, 2009, **28**, 1060–1067.
- 45 C.-Y. Y. Li, Y.-Y. Y. Kuo, J.-H. H. Tsai, G. P. A. A. Yap and T.-G. G. Ong, Amine-Linked *N*-Heterocyclic Carbenes: The Importance of an Pendant Free-Amine Auxiliary in Assisting the Catalytic Reaction, *Chem.–Asian J.*, 2011, **6**, 1520–1524.
- 46 H. Braunschweig, W. C. Ewing, K. Geetharani and M. Schäfer, The Reactivities of Iminoboranes with Carbenes: BN Isosteres of Carbene-Alkyne Adducts, *Angew. Chem., Int. Ed.*, 2015, 54, 1662–1665.
- 47 M. A. Schmidt, P. Müller and M. Movassaghi, On the Interactions of *N*,*N'*-Bismesitylimidazolin-2-Yl and Alcohols, *Tetrahedron Lett.*, 2008, **49**, 4316–4318.
- 48 N. A. Giffin, M. Makramalla, A. D. Hendsbee, K. N. Robertson, C. Sherren, C. C. Pye, J. D. Masuda and J. A. C. Clyburne, Anhydrous TEMPO-H: Reactions of a Good Hydrogen Atom Donor with Low-Valent Carbon Centres, Org. Biomol. Chem., 2011, 9, 3672–3680.
- 49 A. H. Raut, G. Karir and K. S. Viswanathan, Matrix Isolation Infrared and Ab Initio Study of the Interaction of *N*-Heterocyclic Carbene with Water and Methanol: A Case Study of a Strong Hydrogen Bond, *J. Phys. Chem. A*, 2016, **120**, 9390–9400.
- 50 R. Srivastava, R. Moneuse, J. Petit, P. A. Pavard, V. Dardun, M. Rivat, P. Schiltz, M. Solari, E. Jeanneau, L. Veyre, C. Thieuleux, E. A. Quadrelli and C. Camp, Early/Late Heterobimetallic Tantalum/Rhodium Species Assembled

- Through a Novel Bifunctional NHC-OH Ligand, *Chem.-Eur. J.*, 2018, **24**, 4361–4370.
- 51 B. Maji, M. Breugst and H. Mayr, *N*-Heterocyclic Carbenes: Organocatalysts with Moderate Nucleophilicity but Extraordinarily High Lewis Basicity, *Angew. Chem., Int. Ed.*, 2011, **50**, 6915–6919.
- 52 A. Levens, F. An, M. Breugst, H. Mayr and D. W. Lupton, Influence of the *N*-Substituents on the Nucleophilicity and Lewis Basicity of *N*-Heterocyclic Carbenes, *Org. Lett.*, 2016, **18**, 3566–3569.
- 53 T. Steiner, The Hydrogen Bond in the Solid State, *Angew. Chem.*, *Int. Ed.*, 2002, **41**, 48-76.
- 54 F. G. Bordwell, J. C. Branca, D. L. Hughes and W. N. Olmstead, Equilibriums Involving Organic Anions in Dimethyl Sulfoxide and *N*-Methylpyrrolidin-2-One: Acidities, Ion Pairing, and Hydrogen Bonding, *J. Org. Chem.*, 1980, 45, 3305–3313.
- 55 F. G. Bordwell, G. E. Drucker and H. E. Fried, Acidities of Carbon and Nitrogen Acids: The Aromaticity of the Cyclopentadienyl Anion, *J. Org. Chem.*, 1981, **46**, 632–635.
- 56 P. Ballinger and F. A. Long, Acid Ionization Constants of Alcohols. II. Acidities of Some Substituted Methanols and Related Compounds, J. Am. Chem. Soc., 1960, 82, 795–798.
- 57 W. N. Olmstead, Z. Margolin and F. G. Bordwell, Acidities of Water and Simple Alcohols in Dimethyl Sulfoxide Solution, *J. Org. Chem.*, 1980, 45, 3295–3299.
- 58 M. H. Dunn, N. Konstandaras, M. L. Cole and J. B. Harper, Targeted and Systematic Approach to the Study of pKa Values of Imidazolium Salts in Dimethyl Sulfoxide, *J. Org. Chem.*, 2017, **82**, 7324–7331.
- 59 Z. Wang, X. S. Xue, Y. Fu and P. Ji, Comprehensive Basicity Scales for N-Heterocyclic Carbenes in DMSO: Implications on the Stabilities of N-Heterocyclic Carbene and CO<sub>2</sub> Adducts, Chem.-Asian J., 2020, 15, 169–181.
- 60 S. Santra, A. Porey, B. Jana and J. Guin, N-Heterocyclic Carbenes as Chiral Brønsted Base Catalysts: A Highly Diastereo- and Enantioselective 1,6-Addition Reaction, Chem. Sci., 2018, 9, 6446–6450.
- 61 S. Gehrke and O. Hollóczki, Hydrogen Bonding of *N*-Heterocyclic Carbenes in Solution: Mechanisms of Solvent Reorganization, *Chem.-Eur. J.*, 2018, **24**, 11594–11604.
- 62 O. Hollóczki, The Mechanism of *N*-Heterocyclic Carbene Organocatalysis through a Magnifying Glass, *Chem.–Eur. J.*, 2020, **26**, 4885–4894.
- 63 S. Santra, U. Maji and J. Guin, Enantioselective α-Amination of Acyclic 1,3-Dicarbonyls Catalyzed by *N*-Heterocyclic Carbene, *Org. Lett.*, 2020, 22, 468–473.
- 64 R. Guo, X. Huang, M. Zhao, Y. Lei, Z. Ke and L. Kong, Bifurcated Hydrogen-Bond-Stabilized Boron Analogues of Carboxylic Acids, *Inorg. Chem.*, 2019, 58, 13370–13375.
- 65 C. Pi, X. Yu and W. Zheng, Imidazolium 1,3-Benzazaphospholide Ion Pairs with Strong C-H···N Hydrogen Bonds Synthesis, Structures, and Reactivity, *Eur. J. Inorg. Chem.*, 2015, 2015, 1804–1810.
- 66 J. A. Platts, Theoretical Prediction of Hydrogen Bond Donor Capacity, Phys. Chem. Chem. Phys., 2000, 2, 973–980.

- 67 M. H. Abraham, J. Elguero and I. Alkorta, The Hydrogen-Bond Basicity of Carbenes, Croat. Chem. Acta, 2018, 91, 121-124.
- 68 S. Mallick. S. Maddala. K. Kollimalayan and P. Venkatakrishnan, Oxidative Coupling of Carbazoles: A Substituent-Governed Regioselectivity Profile, J. Org. Chem., 2019, 84, 73-93.
- 69 A. J. Arduengo, H. A. Craig, J. R. Goerlich, W. J. Marshall and M. Unverzagt, Imidazolylidenes, Imidazolinylidenes and Imidazolidines, Tetrahedron, 1999, 55, 14523-14534.
- 70 J. L. Cook, C. A. Hunter, C. M. R. Low, A. Perez-Velasco and J. G. Vinter, Solvent Effects on Hydrogen Bonding, Angew. Chem., Int. Ed., 2007, 46, 3706-3709.
- 71 M. Chu, A. N. Scioneaux and C. S. Hartley, Solution-Phase Dimerization of an Oblong Shape-Persistent Macrocycle, J. Org. Chem., 2014, 79, 9009-9017.
- 72 D. Munz, Pushing Electrons Which Carbene Ligand for Which Application?, Organometallics, 2018, 37, 275-289.
- 73 D. Tapu, D. A. Dixon and C. Roe, <sup>13</sup>C NMR Spectroscopy of "Arduengo-Type" Carbenes and Their Derivatives, Chem. Rev., 2009, 109, 3385-3407.