



Applications of N-heterocyclic imines in main group chemistry

Journal:	<i>Chemical Society Reviews</i>
Manuscript ID	CS-REV-02-2016-000163.R3
Article Type:	Review Article
Date Submitted by the Author:	28-Jun-2016
Complete List of Authors:	Ochiai, Tatsumi; Technische Universität Berlin, Institute of Chemistry Daniel, Franz; Technische Universita Muenschen Inoue, Shigeyoshi; Technische Universitat Munchen



Chemical Society Reviews

ARTICLE

Applications of N-heterocyclic imines in main group chemistry

Tatsumi Ochiai,^a Daniel Franz^b and Shigeyoshi Inoue*^{a,b}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

The imidazolin-2-imino group is an N-heterocyclic **imino functionality** that derives from the compound class of guanidines. The exocyclic nitrogen atom preferably bonds to electrophiles and its electron-donating character is markedly enhanced by efficient delocalization of cationic charge density into the five-membered imidazoline ring. Thus, this imino group is an excellent choice for the thermodynamic stabilization of electron-deficient species. Due to the variety of available imidazolin-based precursors to this ligand, its steric demand can be tailored to meet the requirements for kinetic stabilization of otherwise highly reactive species. Consequently, it does not come as a surprise that the imidazolin-2-iminato ligand has found widespread applications in transition-metal chemistry to furnish pincer complexes or “pogo stick” type compounds. In comparison, the field of main-group metal compounds of this ligand is still in its infancy, however, it has gained growing attention in recent years. A considerable number of electron-poor main-group element species have been described today which are stabilized by **N-heterocyclic iminato** ligands. These include low-valent metal cations and species that are marked by formerly unknown bonding modes. In this article we provide an overview on the present chemistry of main-group element compounds of the imidazolin-2-iminato ligand, as well as selected examples for the related imidazolidin- and benzimidazolin-2-imino system.

Introduction

The imidazolin-2-imino group is a potent electron pair donor

In coordination chemistry nitrogen is particularly recognized for its role as a strong electron-donor atom in ligand systems. Seemingly, this contradicts the fact that this element belongs to the highly electronegative members of the periodic table. However, the trivalent nitrogen atom in amines, as well as in imines has high electron density in the form of a lone pair that it readily shares with various types of hard and soft Lewis

acids.

Tertiary amines and secondary ketimines resemble their nucleophilic properties but, in sharp contrast, the unsaturated carbon atom of the imino functionality is prone to the reaction with nucleophiles or reducing agents whereas the amino carbon atom is inert (Fig. 1). This reactivity results from the π -interaction with the more electronegative nitrogen atom which provides for the higher bond order but also drains electron density from the carbon centre in the σ -, as well as the π -scaffold. Due to the orthogonal orientation of the nitrogen lone pair this is not compensated by π back donation. Interestingly, the electronic properties of the imino-nitrogen atom are, vice versa, stronger affected by the characteristics of the carbon atom than it may be the case for an amino-nitrogen centre. In this regard, the electron-rich π -system of an imidazoline ring not only mitigates the electrophilicity of an

^a Institut für Chemie, Technische Universität Berlin, Straße des 17. Juni 135, 10623 Berlin, Germany.

^b Department of Chemistry, Institute for Silicon Chemistry and Catalysis Research Center, Technische Universität München, Lichtenbergstraße 4, 85748 Garching, Germany.

Email correspondence: s. inoue@tum.de



Tatsumi Ochiai

Tatsumi Ochiai was born in Shizuoka/Japan in 1987. In 2012 he received his MSc degree at the University of Tsukuba under the supervision of Prof. Akira Sekiguchi, for which he received the Dean's Prize. He worked on heteroatom-substituted tetrahedranes. He then moved to Berlin in 2012 starting doctoral work in the research group of Prof. Shigeyoshi Inoue. His research interest mainly focuses on the synthesis of low coordinate heavier group 14 elements, metallylenes and metallyliumylidene ions.



Daniel Franz

Daniel Franz studied chemistry at the Goethe University Frankfurt am Main, Germany where he received his PhD degree under the supervision of Prof. Matthias Wagner in 2012. Following this he moved to Berlin to carry out postdoctoral studies in the group of Prof. Shigeyoshi Inoue. Today he is pursuing his research interests in coordination chemistry of group 13 and 14 elements at the Technische Universität München under the supervision of Prof. Inoue.

imino-carbon atom incorporated at the 2-position of the cycle but also pushes electron density to the exocyclic imino-nitrogen atom (Fig. 1). Notably, phosphoramimines ($R_3P=NR$) resemble the imidazolin-2-imines in the electron-donating character of the imino nitrogen atom (Fig. 1). The resemblance of these two ligand classes is reasoned by the similarities in the electronic properties of the parent phosphine and imidazolin-2-ylidene, respectively. Furthermore, one must recognize the isoelectronic relation between imidazolin-2-ylidenes and N-heterocyclic olefins, which function as strong Lewis bases due to the ylide-like nature of the exocyclic alkene bond (Fig. 1).¹

The allocation of electron density from the five-membered imidazoline ring to the exocyclic nitrogen atom is illustrated by conceivable resonance structures of the anionic imidazolin-2-iminato ligand **A** (Fig. 2).^{2,3} The canonical form **B** in which the nitrogen atom bears two formal anionic charges suggests a significant boost of the electron donating properties of the nitrogen atom as compared to ketimines (Fig. 1, Fig. 2). Form **C** represents the partial N-heterocyclic carbene (NHC) character of the imidazoline moiety (Fig. 2). As apparent from the canonical forms (**A-C**) the imidazolin-2-iminato ligand represents a 2σ electron donor with potential to contribute an additional two or even four π -electrons. Consequently, its metal complexes (**A^M**) may exhibit significant metalla-2-aza-allene (**B^M**) or metalimide (**C^M**) character (Fig. 2). This manifests in an expansion of the imino group's CN distance and shortening of the N–M bond length. Concomitantly, the C–N–M bond angle is widened to approach the angle of 180° in the ideal CCC allene structure motive. As a result of its electron-donating properties, the imidazolin-2-iminato ligand is efficient for the thermodynamic stabilization of electron-poor species. Moreover, the bulkiness of the imidazoline ring can be conveniently modified to meet individual requirements for kinetic stabilization of otherwise elusive compounds.

The scope of this review

In this article we focus on the coordination chemistry of the imidazolin-2-iminato ligand, as well as the strongly related imidazolidine-2-imino group and the benzimidazolin-2-imino group with regard to main group elements. For the two latter only relevant examples will be given. An earlier review of Kuhn, Frenking and co-workers on imidazolin-2-imines dates




Fig. 1 Overview of selected N-donor ligands and the related N-heterocyclic olefin (R = organyl).




Fig. 2 Selected resonance structures for the anionic imidazolin-2-iminato ligand, as well as a model complex with M^+ (R = organyl).

back about 13 years.² The broad spectrum of transition metal complexes that comprise this ligand class and the methods for synthesis of the ligand have recently been reviewed by Tamm and coworkers and will be discussed only in part.³

Moreover, only selected examples will be discussed for compounds of this iminato ligand with the non-metals carbon and nitrogen because this belongs to the field of organic chemistry rather than coordination chemistry.




Shigeyoshi Inoue

Shigeyoshi Inoue studied chemistry at University of Tsukuba where he completed his PhD degree under the supervision of Prof. Akira Sekiguchi in 2008. After being a Humboldt Postdoctoral fellow as well as a JSPS postdoctoral fellow for research abroad with Prof. Matthias Dri   at Technische Universit  t Berlin, he began his independent research career as a Sofia Kovalevskaja Professor in 2010 at same university. Since 2015 he has been W2-Tenure-Track Professor of Silicon Chemistry at the Technische Universit  t M  nchen. His research interests lie on the synthesis and reactivity investigation of low-valent main group compounds with the goal of finding novel applications in synthesis and catalysis.

Group 1 and group 2 element complexes

Background

About 20 years ago Kuhn and co-workers started their pioneering studies on the chemistry of imidazolin-2-imines.⁴ Few alkaline^{5,6} and alkaline earth⁷ compounds of the imino group were reported but not investigated thoroughly probably because of the pronounced polar nature of the N–M bond (M = alkaline or alkaline earth metal). This puts them in the role of a reactive intermediate for ligand transfer via salt metathesis rather than a species with its own follow up reactivity with



Scheme 1 Conversion of the imino lithium species **1** into the imino-stabilized LiOCl aggregate **2** (the lithium chloride derives from the methyl lithium synthesis and peroxide from contaminated solvent). Formation of the bulky imino lithium dimer **3** (Dip = 2,6-diisopropylphenyl).




Fig. 3 Ellipsoid plot (30% level) of the Li₁₂N₈O₂Cl₂ cage in **2** with adjacent imino-carbon atoms and oxygen atoms of coordinated THF.

sustainment of the N–M bond. Accordingly, the chemistry of the group 1 and group 2 imidazolin-2-iminato complexes is only explored to a minor degree to date.

Lithium and potassium complexes


The reaction of L^{Me2}NH (L^{Me2} = 1,3-dimethyl-imidazolin-2-ylidene) with MeLi in Et₂O produces L^{Me2}NLi (**1**) which is the N-lithiated derivative of the imidazolin-2-imine.⁵ The species was characterized by ¹H NMR analysis and according to the reported CHN elemental analysis no solvent was present in the isolated material. If the conversion was carried out in THF/Et₂O with MeLi that was prepared from H₃CCl and elemental lithium without prior separation of lithium chloride, crystals of the unexpected composition [Li₁₂O₂Cl₂(L^{Me2}N)₈(thf)₄]·8THF (**2**) were retrieved in low yield (Scheme 1).

The solid state structure of **2** is marked by a Li₁₂N₈O₂Cl₂ cage that comprises a peroxo moiety in its core (Fig. 3). The authors reasoned that the O₂²⁻ group resulted from contamination of the solvent with traces of peroxide. Crystals of dimeric [L^{Dip}NLi]₂·toluene (**3**·toluene, L^{Dip} = 1,3-bis(2,6-diisopropylphenyl)-imidazolin-2-ylidene) were isolated in good


yield after the reaction of L^{Dip}NH with *n*BuLi in toluene/hexane.⁸ Apparently, the formation of higher aggregates is hampered by the bulkier Dip groups (Dip = 2,6-diisopropylphenyl). Bringing into contact L^{Me2}NH and freshly prepared MeK in Et₂O afforded the heavier alkaline derivative L^{Me2}NK (**4**).⁶ The compound was characterized by elemental analysis and its existence was verified by the synthesis of the dithiocarbimate L^{Me2}NCS₂K (**5**, Scheme 2). Interestingly, the latter shows structural characteristics that account for a bonding situation as represented by resonance structure **5^B** (Scheme 2) with the C–N_{imino} bond length significantly elongated (a range from 1.369(16) Å to 1.379(18) Å is observed in the solid state structure; cf. **2**: C–N_{imino} = 1.260(4)–1.263(4) Å; **3**: C–N_{imino} = 1.241(3) Å, 1.242(4) Å). Accordingly, the C–S distances in **5** (1.733(13)–1.755(13) Å) resemble typical CS single bond lengths.

Magnesium complexes

As rare examples for N-heterocyclic iminato complexes of the group 2 metals the magnesium compounds (L^{Me2}NH)₄MgI₂ (**6**[I]₂), as well as {L^{Me2}NMgI}_n (**7**) and {(L^{Me2}N)₂Mg}_n (**8**) were reported by Kuhn and coworkers (n ≥ 1).⁷ They are accessed through L^{Me2}NH via conversion with 0.25 MgI₂, MeMgI and (*n*Bu)₂Mg, respectively (Scheme 3). Single crystal XRD (X-ray diffraction) data were obtained for **6**[I]₂ (Fig. 4) while the degree of aggregation (n) of **7** and **8** was not elucidated by structural analysis. Notably, **6**[I]₂ also formed if less than four equiv of the imine were reacted with MgI₂. The authors attributed this observation to the high basicity of the ligand.



Scheme 2 Conversion of potassium imide **4** with carbon disulfide to the thiocarbimate **5** (represented by resonance structures **5^A** and **5^B**).



Scheme 3 Synthesis of the imino-magnesium compounds **6**[I]₂, **7** and **8**.




Fig. 4 Ball&stick representation of $(L^{\text{Me}_2\text{NH}})_4\text{Mg}^{2+}$ (6^{2+}) as derived from XRD analysis (non-N-bonded hydrogen atoms have been omitted).

Group 13 element complexes


Background

Of the group 13 elements only few aluminium complexes with an N-heterocyclic iminato ligand had been reported until respective research was resumed by our group.⁶ Reports^{8,9,10} in the year 2014 were the first to describe imidazolin-2-imino complexes of boron. In contrast, the coordination chemistry of related phosphoranimes of boron¹¹ and aluminium^{11a,11c,11d,12} is thoroughly studied.

Boron complexes


The Lewis acid base adducts $\text{LNH}(\text{BH}_3)$ (**9**, Scheme 4) between $\text{L}^{\text{Dip}}\text{NH}$, as well as $\text{L}^{\text{Mes}}\text{NH}$ ($\text{L}^{\text{Mes}} = 1,3\text{-dimesityl-imidazolin-2-ylidene}$) and the parent borane were isolated after conversion of the imine with $\text{Me}_2\text{S}\cdot\text{BH}_3$ in toluene.⁹ When treated with imidazolin-2-ylidenes (L) dihydrogen is abstracted from the $\text{HN}\text{--}\text{BH}$ moiety of these imine-borane compounds and along with $\text{L}(\text{H}_2)$ (hydrogenated at the formerly carbenoid centre) the NHC-adducts of respective iminato boron dihydrides ($\text{LN}(\text{BH}_2)\text{L}$) are formed (**10**, Scheme 4).⁹ These NHC-adducts undergo hydride-mediated ring-expansion reaction, that is the boron atom transfers its two hydrides to the adjacent carbon atom and inserts into the $\text{C}_{\text{carbenoid}}\text{N}$ bond of the NHC (**11**, Scheme 4).⁹ Presumably, the interaction between the lone pair at the imino nitrogen atom and the unoccupied p-orbital at the boron centre supports the trigonalization of the metalloid atom. Interestingly, this insertion occurs at higher temperatures with more sterically hindered substituents at the boron atom. Moreover, $\text{H}_2\text{L}^{\text{Mes}}$ (1,3-dimesityl-imidazolidin-2-ylidene, NHC saturated at the ligand backbone) is subject to ring-expansion reaction at significantly lower temperatures than its congener of very similar sterical encumbrance L^{Mes} (NHC unsaturated at the ligand backbone). It was reasoned that the conjugated ring system in L^{Mes} is more efficient for the delocalization of positive charge density and, thus, stabilizes the boron dihydride form (LNBH_2L).

Similarly, ring-activation and expansion reaction of L^{Dip} took place by heating the amido-substituted hydridoborane $\text{L}^{\text{Dip}}(\text{BH}_2)\text{HN}^{\text{Dip}}$ reported by Rivard and coworkers.¹³



Scheme 4 Conversion of the imine-borane adduct **9** with NHC to iminoboron dihydride NHC complex **10** and the formation of **11** via ring expansion reaction. $\text{R}^1 = \text{Mes}$ or Dip , $\text{R}^2 = \text{Me}$ (for $\text{R}^3 = \text{Me}$), Mes or Dip (both for $\text{R}^3 = \text{H}$); $\text{R}^2 = \text{Mes}$ and $\text{R}^3 = \text{H}$ for unsaturated backbone; not all combinations of imine and NHC are viable for **10** and **11**.⁹


Conversion of the bis(iminiumtosylate) $\mathbf{12}[\text{OTs}]_2$ with $\text{Li}[\text{BH}_4]$ furnishes the boronium salt $\mathbf{13}[\text{OTs}]$ (Scheme 5).¹⁰ The compound reacts with yellow sulfur to give a rare example of a cationic thioxoborane $\mathbf{14}[\text{OTs}]$ that was structurally characterized (Scheme 5).¹⁰ The B–S bond length (1.710(5) Å) in $\mathbf{14}^+$ is the shortest that had been reported to date for a molecular complex. Notably, the B–N bond lengths significantly decrease upon transformation of tetrahedral $\mathbf{13}^+$ into trigonal-planar $\mathbf{14}^+$ ($\mathbf{13}^+$: 1.573(5) Å, 1.577(5) Å; $\mathbf{14}^+$:




Scheme 5 Reaction of the bis(iminiumtosylate) $\mathbf{12}[\text{OTs}]_2$ with lithium borohydride to the boronium salt $\mathbf{13}[\text{OTs}]$ (Ts = tosyl) and its conversion to the thioxoborane salt $\mathbf{14}[\text{OTs}]$.

1.483(5) Å, 1.493(5) Å) and, consequently, a partial double bond character can be attributed to the boron-nitrogen interactions in **14**⁺. Concomitantly, the C–N distances of the imino groups increase (**13**⁺: 1.317(5) Å, 1.318(4) Å; **14**⁺: 1.359(4) Å, 1.363(4) Å) which is in accordance with the formulation of resonance structure **14**^{A+} (Scheme 5) that represents the delocalization of positive charge density into the **N-heterocycles** and the polarization of the BS bond towards the sulfur atom (NBO charge at S = –0.58). DFT (density functional theory) calculations supported the interpretation of the remarkably short B–S distance in terms of a boron sulfur double bond. For example, the HOMO (highest occupied molecular orbital) shows mainly the sulfur lone pair and the HOMO–1 reveals the B=S π-bonding orbital. The NBO charge at the boron centre of **14**⁺ was calculated to +0.63 which accounts for the boron cation character of the complex as it is illustrated by the canonical form **14**^{B+} (Scheme 5).

Very recently, Rivard and coworkers described the conversion of the imidazolin-2-imino trimethylsilane **15** to the imino boron dichloride **16** and its organyl derivative **17** by reaction of **15** with BCl₃ and PhBCl₂, respectively (Scheme 6).¹⁴ The bisimino boron monochloride **18** was furnished in a reaction between **15** and **16** (Scheme 6). Moreover, the synthesis of the diphenyl congener **19** was accomplished by conversion of **17** with phenylmagnesium bromide (Scheme 6). The solid state structure of the dihalide **16** hints toward the significant bora-2-aza-allene properties of the CNB moiety (type **B**^M, Fig. 2) as concluded from the C–N–B bond angle of 180° and the short B–N bond length (1.302(6) Å) which implies high boron-nitrogen double bond character. Remarkably, **16**, **17** and **19** react with amine-boranes (R_nH_{3–n}N·BH₃; R = H, Me; n = 1, 2) to produce the respective dihydrogenated imino boron compounds L^{Dip}NH(B(H)R'R'') **20–22** (R' = R'' = Cl for **20**; R' = Cl, R'' = Ph for **21**; R' = R'' = Ph for **22**) and a mixture of the



Scheme 6 Conversion of the imino trimethylsilane **15** to the imino chloroboranes **16–18** and the imino diphenylborane **19**. Formation of the imine-borane adducts via abstraction of dihydrogene from amine-borane adducts.



Scheme 7 Conversion of the trimethylsilylimine **25** to the aluminium trichloride imine complex **26** and its transformation into trimeric **27**.


amine-borane dehydrogenation products (Scheme 6).¹⁴ The authors conclude that this imino boron compound acts as an intramolecular frustrated Lewis acid base pair. It should be noted that compound **17** displays catalytic activity in dehydrocoupling of MeNH₂·BH₃ to yield [MeNBH]₃ along with oligomeric aminoboranes, which shows its great potential with respect to further application in metal-free catalysis for the dehydrocoupling of amine-boranes and related species.

Aluminium complexes

The bis(imino) aluminium complexes (L^{Me2}NSiMe₃)₂AlMe₂[Cl] (**23**) and (L^{Me2}NH)₂AlMe₂[Cl] (**24**) were synthesized by conversion of L^{Me2}NSiMe₃ (**25**) and L^{Me2}NH, respectively, with 0.5 equiv AlMe₂Cl.⁷ The ion-separated forms were postulated on the basis of NMR spectroscopic data. The related Lewis acid base adduct L^{Me2}NSiMe₃·AlCl₃ (**26**) releases Me₃SiCl upon heating the neat compound to 180 °C and converts into {L^{Me2}NAICl₂}₃ (**27**, Scheme 7).⁷ An X-ray crystallographic analysis of compound **27** verified its trimeric structure with a six-membered Al₃N₃ cycle.

The imino aluminium dihydride {L^{Dip}NAIH₂}₂ (**28**) results from the reaction of L^{Dip}NH with Me₃N·AlH₃ (Scheme 7).⁸ From the dihydride derive the dihalides {L^{Dip}NAIX₂}₂ (**29–31**, X = Cl, Br, I) by conversion with BX₃ (two equiv) which were described to form dimers in the solid state, as well as in solution (Scheme 8).⁷ Obviously, the bulkier iminato ligand in **29** leads to the formation of a four-membered Al₂N₂ ring with smaller N–Al–N angles (87.8(1)°, 92.3(1)°) in comparison to the six-membered ring in **27** with larger angles (108–110°). Interestingly, the sterically hindered phosphoranimo aluminium dihydride and -dichloride form dimers with four-membered Al₂N₂ rings [(μ-R₃PN)AlX₂]₂ (R = *i*Pr, *t*Bu; X = H, Cl), as well.^{11c,12a} Furthermore, the dihydride **28** reacts with the electrophiles Me₂S·BH₃ (four equiv) and Me₃SiOTf (two equiv, Tf = triflyl) to the aluminium borohydride **32** and the aluminium monohydride triflate **33**, respectively (Scheme 8).⁸ Notably, the substitution of both aluminium bonded hydrides in **33** for triflate substituents could not be accomplished by use of a larger excess of

Me_3SiOTf , even at elevated temperature. In contrast, the conversion of **28** with only two equiv of $\text{Me}_2\text{S}\cdot\text{BH}_3$ does not



Scheme 8 Overview on syntheses of imino aluminium compounds derived from the aluminium dihydride **28**: Reaction of **28** to the dihalides **29–31**, the borohydride **32** and the triflate **33** ($X = \text{Cl}, \text{Br}, \text{I}$). Synthesis of the aluminium mono- and bis(hydrogensulfides) **34**, as well as **35**. Conversion of **28** to the ditelluride **36** and its monotopic aluminium telluride offspring **37** ($L^{\text{Et}} = 1,3\text{-diethyl-4,5-dimethyl-imidazolin-2-ylidene}$). Transformation of **37** to ditopic **39** via the presumed intermediate **38**.

afford an expected aluminium monohydride borohydride as a product but yields mixtures of **28** and **32**. Obviously, the electron withdrawing triflyl groups in **33** mitigate the hydride-donor strength of the remaining AlH functionality. Accordingly, only aluminium monohydride triflates of the related phosphoraniminato or the 1,3-diketiminato ligand have been reported.^{11c,15} The conversion of the aluminium dihydride **28** with yellow sulfur affords a rare example of an aluminium hydride hydrogensulfide complex (**34**) by insertion of a sulfur atom into the AlH bond (Scheme 8).¹⁶ Similar to **33** the remaining hydride functionalities at the aluminium centres in **34** are less reactive than in the parent compound. However, the transformation with S_8 to form the bis(hydrogensulfide) **35** can be forced onto the system by heating (90 °C for four days, Scheme 8).¹⁶ As apparent from the XRD study the Al–S distances in **35** (2.231(1)–2.240(1) Å) are slightly shorter than the respective distances in the monohydrogensulfide **34** (2.250(1) Å and 2.252(1) Å).

In order to furnish a heavier aluminium chalcogenide of the imidazolin-2-iminato ligand **28** was converted with the tellurium atom transfer reagent $n\text{Bu}_3\text{PTe}$ (two equiv).¹⁷ This conversion yields ditopic aluminium ditelluride **36** as a rare example of an electron-precise aluminium complex with the chalcogen in the oxidation state –1 (Scheme 8). The hydrides left at the aluminium centres in **36** do not react further with

excess $n\text{Bu}_3\text{PTe}$. However, the compound converts with NHC (L^{Et} , 5 equiv, $L^{\text{Et}} = 1,3\text{-diethyl-4,5-dimethyl-imidazolin-2-ylidene}$) in a dehydrogenative redox process to form the monotopic aluminium telluride **37** (Fig. 5, Scheme 8) with the chalcogen in the oxidation state –2 along with dihydrogenated NHC ($L^{\text{Et}}(\text{H}_2)$).¹⁷




Fig. 5 Ellipsoid plot (30% level) of the aluminium telluride **37** (hydrogen atoms, *isopropyl* groups and non-N-bonded methyl groups have been omitted).

The structural study of **37** revealed a remarkably short Al–Te distance of 2.5130(14) Å and DFT calculations determined an enhanced aluminium-tellurium interaction ($WBI_{AlTe} = 1.20$; NPA charges: Al = +1.24, Te = –0.95; WBI = Wiberg Bond Index, NPA = Natural Population Analysis). It has to be pointed out that the terminal position of the tellurium atom is a very scant structure motif as group 16 atoms commonly assume bridging positions in aluminium chalcogenides. Upon heating a benzene solution of **37** to 80 °C one of the two L^{Et} ligands is released and the putative intermediate $L^{Dip}N(AlTe)L^{Et}$ (**38**) aggregates to **39** (Scheme 8).

The reaction pathway via **38** was suggested by DFT calculations, however, the isolation of a bulkier congener to this elusive species was not accomplished by the use of more sterically hindered NHC. The structural investigation of **39** revealed significantly elongated Al–Te distances (2.6143(14) Å, 2.6211(15) Å) and a decreased bond order for the AlTe interaction ($WBI_{AlTe} = 0.75$; NPA charges: Al = +1.21, Te = –0.79) with respect to **37**.¹⁷ It is of note that in ditopic **39** the aluminium centres are bridged via the tellurium atoms. Notably, **37** and **39** contrast the other given examples for aluminium complexes of the imidazolin-2-iminato ligand in that the aluminium centres are not connected via the nitrogen atoms of the imino groups. Taking into account the marked changes in the Al–Te distances and the values for the WBI_{AlTe} upon transformation of **37** into **39** the nature of the AlTe interaction in **37** was presumed to possess high Al=Te double bond character.

Group 14 element complexes


Background

In initial reports on the chemistry of **N-heterocyclic iminato ligands** Kuhn and coworkers described the imino trimethylsilane **25** (Scheme 7) which was used as an alternate transmetallation reagent to the alkaline metal salts mentioned above.^{4,7} Presumably, the bulkier $L^{Mes}NSiMe_3$ is formed as an intermediate in the synthesis of $L^{Mes}NH$ via a Staudinger-type reaction described by Cameron, Jenkins, Clyburne and coworkers in 2001.¹⁸ Tamm and coworkers established the general method for the preparation of trimethylsilyl-functionalized bulkier imidazolin-2-iminato ligands such as $L^{Mes}NSiMe_3$ and $L^{Dip}NSiMe_3$ in 2004.¹⁹ This method has tremendous advantages for the convenient and high-yield synthesis of various imidazolin-2-imines. Moreover, a silicon atom was incorporated into the spacer group between the imino- and the arene moiety in oligodentate ligands reported by Tamm and coworkers.²⁰ However, it played a rather passive role in the chemistry of the transition metal complexes derived from this ligand system. As outlined in the following section a

considerable time elapsed from Kuhn's initial report until the coordination chemistry of the imidazolin-2-iminato ligand with tetrel atoms was thoroughly investigated.

Silicon complexes

Our group commenced work on main group element complexes of the imidazolin-2-imino ligand a few years ago




Scheme 9 Synthesis of Cp*-substituted iminosilylene **40** and its borane adduct **42**, as well as the dibromide precursor **41**. Silylene-nitrene formulation **40^A** and sila-2-aza-allene canonical structure **42^A**.

and described its complex with a silicon(II) centre in 2012.²¹ High interest for molecular low-valent silicon compounds originates from their various applications for catalysis and bond activation.^{22,23} Conversion of $L^{Dip}NLI$ with $Cp^*Si(B(C_6F_5)_4)_2$ (Cp^* = pentamethylcyclopentadienyl) as a source of silicon(II) afforded the pentamethylcyclopentadienyl imino silylene **40** with η^2 coordination of the silicon centre by the organyl ligand (Scheme 9). As an alternative synthetic route **40** can be accessed via reaction of $L^{Dip}NLI$ with Cp^*SiBr_3 followed by reductive dehalogenation of the $L^{Dip}NSi(Br)_2Cp^*$ (**41**). Unfortunately, this method grants only very poor yields of the silylene. DFT calculations on **40** show some π bonding interaction between the imino nitrogen lone pair and the unoccupied p-orbital at the silicon centre. The WBI_{SiN} of 0.80 and the Si–N bond length (1.691(5) Å) imply single bond character. Thus, multiple bond interaction as illustrated by the general canonical structures **B^M** and **C^M** (Fig. 2) cannot be concluded for **40**. A key motivation of the study was to explore potential silylene-nitrene character of complexes between a low-valent silicon atom and the imidazolin-2-iminato ligand as represented by the canonical structure **40^A** (Scheme 9). However, structural and theoretical investigation verified the imino-substituted silylene formulation **40** with no relevant silylene-nitrene character (**40^A**). Conversion of **40** with tris(pentafluorophenyl)borane furnished the silylene-borane adduct **42**. It is interesting to note that the Cp^* ligand is coordinated in a η^1 -mode with one σ bond to the silicon atom in sharp contrast to the precursor **40**, in which η^2 -mode Si– Cp^* bonding is observed. As compared to **40** the Si–N bond length

is considerably reduced to 1.605(3) Å and its W_{SiN} rises to 0.90 which accounts for partial SiN double bond character. The C–N–Si angle of 158.7(3)° in **42** is wider than in **40** (136.6(4)°). Accordingly, relevant 1-sila-2-azallene nature (cf. **B^M**, Fig. 2) can be attributed to **42** as represented by resonance structure **42^A** (Scheme 9).

In order to exploit the strongly electron-donating properties of an N-heterocyclic iminato ligand for tuning the reactivity of low-valent silicon species Rivard and




Scheme 10 Reduction of dibromosilane **43** with KC_8 to form the unexpected anionic compound **45** instead of intended **46**. The siloxane **47**, as well as the tribromide **44**.

coworkers attempted the synthesis of a hypothetical bisiminosilylene. Access to the bisiminodibromosilane precursor **43** is granted by conversion of $\text{L}^{\text{Dip}}\text{NSiMe}_3$ (**15**) with SiBr_4 in appropriate stoichiometry (Scheme 10).²⁴ The monoimino derivative $\text{L}^{\text{Dip}}\text{NSiBr}_3$ (**44**) is synthesized in similar fashion (Scheme 10).²⁴ The reductive dehalogenation of **43** with KC_8 yielded the potassium salt **45** instead of the desired silylene ($\text{L}^{\text{Dip}}\text{N})_2\text{Si}$ (**46**, Scheme 10).²⁴ This product (**45**) was presumed to result from an intermediate potassium silanide via migration of a Dip group. The formation of minor amounts of the siloxane **47** was reasoned by the presence of silicon grease in the reaction mixture (Scheme 10).

Germanium complexes

The reductive dehalogenation of the bulky bisiminodichlorogermane **48** with sodium naphthalene affords the bisiminogermylene **49** as reported by Rivard and coworkers (Scheme 11).²⁴ Notably, the related monoiminotrichlorogermane **50** was also described (Scheme 11). In the solid state the germanium(II) compound (**49**) exhibits longer Ge–N distances (both: 1.8194(15) Å) and a decreased N–Ge–N bond angle (99.48(10)°) with respect to its

halogenated precursor **48** (Ge–N = 1.7528(14) Å, 1.7582(14) Å; N–Ge–N = 106.33(7)°; Fig. 6). These structural features were interpreted by the authors in terms of a higher p-character of the Ge–N bond in **49** as compared to **48**. Theoretical calculations indicated a low singlet-triplet gap of 45.8 kcal/mol for the bisiminogermylene **49**, which is a similar value to that of the elusive bisiminosilylene **46** (44.5 kcal/mol). This computational study suggests high inclination for the sterically hindered metal centre to insert into element-element bonds of small substrate molecules. However, upon conversion of **49** with dihydrogen Rivard and coworkers observed the formation of $\text{L}^{\text{Dip}}\text{NH}$ as the only soluble species instead of the expected $(\text{L}^{\text{Dip}}\text{N})_2\text{GeH}_2$.²⁴ This may account for the pronounced proton affinity of the imidazolin-2-imino group. Interestingly, the bisiminogermene is not formed in the reaction of **48** with hydride transfer reagents as $\text{K}[\text{HB}^t\text{Bu}_3]$ or potassium hydride.²⁴ Another synthetic approach to germanium(II) complexes of the imidazolin-2-iminato ligand uses the Lappert's germylene $(\text{Me}_3\text{Si})_2\text{N}_2\text{Ge}$ as a low-valent metal source. Its conversion with one equiv of $\text{L}^{\text{Dip}}\text{NH}$ at 60 °C furnishes the amino(imino)germylene **51** in the form of a viscous liquid (Scheme 12).²⁵ It acts as a ligand towards iron carbonyls as demonstrated by the formation of the germylene complex **52** after reaction of **51** with diironnonacarbonyl (Scheme 12).²⁵ The XRD analysis of **52** reveals a Ge–N_{imino} distance of 1.755(2) Å which is significantly shorter than the Ge–N_{amino} bond length of 1.839(2) Å and also reduced with respect to the free bisiminogermylene **49** (vide supra). Considering the WBIs of the Ge–N bonds in **52** ($\text{Ge–N}_{\text{imino}} = 0.86$, $\text{Ge–N}_{\text{amino}} = 0.60$) it is reasonable to assume that the bulky imidazolin-2-iminato ligand bonds stronger to the germanium(II) centre than the bis(trimethylsilyl)amino group; presumably as a result of the iminato ligand's higher electron donating character.



Scheme 11 Reductive dehalogenation of the dichlorogermane **48** to the bisiminogermylene **49**. The trichlorogermane **50**.






Fig. 6 Ellipsoid plot (30% level) of the bis(imino)germylene **49** (hydrogen atoms and isopropyl groups have been omitted).



Scheme 12 Preparation of amino(imino)metallylenes **51** and **54** and the iron carbonyl **52**, as well as the dmap adduct **55** (dmap = 4-dimethylamino-pyridine). Conversion of **54** to dimeric stannylene azide **57**.



Scheme 13 Synthesis of four-membered metallyliumylidenes **53⁺** and **56⁺** by methyl-abstraction from the amino(imino)metallylenes **51** and **54**.

If treated with tris(pentafluorophenyl)borane compound **51** undergoes a methyl abstraction and ring closing reaction to form the cyclic germyliumylidene **53**[MeB(C₆F₅)₃] as an example for a cationic complex of germanium(II) (Scheme 13, Fig. 7).²⁵ The bonding situation in the **53⁺** is found to be suitably described as an amino-bonded cationic germanium(II) atom that is stabilized by via dative bond type interaction with an intramolecularly tethered imino group. This is indicated by a weaker interaction between the Ge(II) centre and the N_{imino} atom (Ge–N_{imino} = 1.9694(14) Å, WBI_{GeN} = 0.48) and a stronger bond between the Ge(II) centre and the N_{amino} atom (Ge–N_{amino} = 1.839(2) Å, WBI_{GeN} = 0.73). This bonding situation between the metal centre and the N atoms appears inverse to the uncharged congeners **51** and **52** (vide supra). Moreover, the C–N_{imino} distance of 1.335(2) Å in **53⁺** is elongated in comparison to **51** (C–N_{imino} = 1.296(3) Å) and hints towards delocalization of cationic charge into the imidazolin ring system similar to the observations reported for **5**, as well as **14⁺**. This accounts for the pronounced ability of the imidazolin-2-iminato ligand to stabilize cationic species which was verified yet again by a very recent report on the isolation of bifunctional germylene-germyliumylidenes.^{25b}

Tin complexes

In 2015 the formation of the amino(imino)stannylene **54** was reported that proceeds in similar fashion to the lighter congener **51** via reaction of ((Me₃Si)₂)₂Sn with L^{Dip}NH at 60 °C (Scheme 12).²⁶ Notably, the ¹¹⁹Sn NMR chemical shift of –208 ppm for **54** (C₆D₆) is considerably shifted to higher field with respect to the precursor (767 ppm, C₆D₆) which was accredited to an aggregated species in solution with a higher coordinate tin(II) centre. The compound (**54**) was obtained as an oil and reacted with 4-dimethylamino-pyridine (dmap) to give the solid tin(II) adduct **55** that exhibited a resonance at –3 ppm in the ¹¹⁹Sn NMR spectroscopic analysis (Scheme 12).²⁶ The XRD study of **55** shows a shorter Sn–N_{imino} contact (2.0588(13) Å) and a longer Sn–N_{amino} distance (2.1647(12) Å). This was interpreted in terms of a stronger bond of the metal centre to the iminato ligand and a weaker interaction with the amino group as it was described for the germanium congener **52**, as well (vide supra).^{25,26} The reaction of **54** with tris(pentafluorophenyl)borane affords the stannylidene salt **56**[MeB(C₆F₅)₃] in a methyl abstraction and ring closing reaction similar to the process that afforded the germanium analogue **53**[MeB(C₆F₅)₃] (Scheme 13, vide supra). The bonding situations in **56⁺** and in **53⁺** resemble, that is an amino bonded metallylium ylidene cation which is stabilized by a dative bond to the imino group. Accordingly, the Sn–N_{imino} distance of 2.197(2) Å in **56⁺** is longer than the Sn–N_{amino} bond length of 2.062(2) Å which is an observation on the contrary to the uncharged congener **55** that possesses a shorter Sn–N_{imino} contact. The isolation of **56**[MeB(C₆F₅)₃] is another example for the high potential of N-heterocyclic imino systems to stabilize cationic species. Interestingly, the amino(imino)stannylene **54** converts with azidotrimethylsilane to the dimeric iminostannylene azide **57** (Scheme 12).²⁶ Apparently, an

expected stannaimine of the type $(L^{\text{Dip}}\text{N})((\text{Me}_3\text{Si})_2\text{N})\text{Sn}(\text{NSiMe}_3)$ is not formed but ligand exchange results in the liberation of $(\text{Me}_3\text{Si})_3\text{N}$ from the system. Interestingly, in solution (THF- d_8) dimeric **57** ($\delta(^{119}\text{Sn}) = -285$ ppm) exists in equilibrium with a monomeric species ($\delta(^{119}\text{Sn}) = 39$ ppm).





Fig. 7 Ellipsoid plot (30% level) of the germyliumylidene **53*** (hydrogen atoms and isopropyl groups have been omitted).



Scheme 14 Synthesis of bisiminostannylene **58** and reactivity with I_2 , MeI, ClSiMe_3 and MeLi to products **59–62**.

Very recently, our group described the bisiminochlorostannate **58** which forms by the reaction of $L^{\text{Dip}}\text{NLi}$ with half an equivalent of $\text{SnCl}_2 \cdot \text{diox}$ (Scheme 14).²⁷ The ^{119}Sn NMR spectrum of **58** (thf- d_8) shows a resonance at -18 ppm that is shifted to lower field in comparison to common monomeric trigonal pyramidal-coordinate 1,3-diketiminato tin(II) chlorides (-118 ppm to -337 ppm).²⁸ The XRD analysis of **58** reveals a butterfly-shaped four-membered SnN_2Li stannacycle with no bonding interaction between the Sn atom and the Li atom (Fig. 8). The $\text{Sn}-\text{N}_{\text{imino}}$ distances of 2.143(5) Å and 2.179(4) Å are elongated compared with the dmap adduct **55**. The $\text{Li}-\text{N}_{\text{imino}}$ bond lengths of 1.946(9) Å and 2.004(9) Å are comparable to those reported for the imino lithium dimer $[\text{L}^{\text{Dip}}\text{NLi}]_2 \cdot \text{toluene}$ (**3**·toluene).⁸

Compound **58** reacts with electrophiles such as I_2 and MeI to form the oxidative addition products **59** and **60** which demonstrates its stannylene character. For the bulky substrate Me_3SiCl formation of the stannane $(\text{Me}_3\text{Si})\text{ClSn}(L^{\text{Dip}})_2$ is suppressed and $L^{\text{Dip}}\text{SiMe}_3$ is formed along with the dimeric chlorostannylene $[\text{L}^{\text{Dip}}\text{SnCl}]_2$ (**61**). The bisiminochlorostannate (**58**) may also act as an electrophile. Its conversion with MeLi leads to the formation of Me-substituted stannate **62** that exhibited a planar four-membered LiN_2Sn ring. Theoretical calculations on **58** revealed the high single-bond character of the $\text{Sn}-\text{N}_{\text{imino}}$ interactions as concluded from the comparison of the WBI values of **58** (0.43 and 0.44) with the ones in **61** (0.24 and 0.24) which mark considerable dative-bond character for the latter. Moreover, the computational study of the natural population analysis (NPA) charge distribution in **58** and **61** shows that the Sn atom in **58** is less positively polarized (+1.22) than that in **61** (+1.42). These theoretical results account for the stannyl anion character of **58** as illustrated by the resonance structure **58^A** (Scheme 14). The ambiphilic reactivity of the tin(II) centre in **58**, that is it functions as a nucleophile in the synthesis of **59** and **60** and as an electrophile in the conversion to **62** has to be pointed out. It allows for the conclusion that the compound (**58**) possesses high stannylene character and, thus represents a heavier congener of carbenoids.

Miscellaneous: survey of carbon chemistry

In the field of coordination chemistry the tethering of the exocyclic imino-nitrogen atom of an **N-heterocyclic imino** group to a carbon atom mostly serves the creation of Taylor made ligand systems. These synthetic methods have been reviewed elsewhere.^{2,3} They can be complemented by a report of Tamm and coworkers in 2014 on a modified synthesis of the bisimine 1,2- $(L^{\text{iPr}_2\text{Me}_2}\text{N})_2\text{-C}_2\text{H}_4$ ($L^{\text{iPr}_2\text{Me}_2} = 1,3$ -diisopropyl-4,5-dimethyl-imidazolin-2-ylidene, Scheme 15); a chelate-fashioned ligand system which had been described before in the year 2007.^{29,28}





Fig. 8 Ellipsoid plot (30% level) of the Stannyleneid **58** (hydrogen atoms and *isopropyl* groups have been omitted).



Scheme 15 Synthesis of the bisimine compound 1,2-(L^{*iPr*2Me2}N)₂-C₂H₄ (L^{*iPr*2Me2} = 1,3-diisopropyl-4,5-dimethyl-imidazol-2-ylidene) from an imidazolium salt precursor.


Group 15 element complexes

Background

For the pnictogen family compounds of phosphorus with the imidazolidin-2-imino group (saturated in the ligand backbone) dominate the field. As outlined in the following this ligand system is implemented for the stabilization of phosphorus-centred radicals and suits the requirements for the isolation of cationic species similar to the strongly related imidazolin-2-imino group (unsaturated in the ligand backbone). Most notably, pioneering work on imidazolin-2-imino-substituted phosphanes was reported by Kuhn and coworkers in 1996 and 1998.³¹ Also, compounds with the imidazolin-2-imino structural motif adjacent to a nitrogen atom are abundant. Examples include common types of organic compounds as azines of cyclic ureas, cyclic bisguanidines, as well as triazenes and diazotates with the corresponding C₃N₂ five-membered ring backbone. These will be discussed in the miscellaneous section of this review. Interestingly, the respective chemistry of the heavier pnictogens remains largely unexplored to date.

Phosphorus compounds

Phosphorus mononitrides and phosphinonitrenes have developed into an established subgenre of the iminato ligand-stabilized phosphorus chemistry and respective research was sparked by Bertrand and coworkers in 2010. They reported the use of the imidazolidin-2-imino lithium reagent (H₂)L^{Dip}NLi for the synthesis of the phosphorus dichloride **63** which undergoes reductive dehalogenation with magnesium in the presence of cyclic alkylamino carbene (CAAC) to afford **64** (Scheme 16, Fig. 9).³² The authors demonstrated that this compound (**64**) can be regarded as a molecular congener to phosphorus mononitride stabilized by a CAAC as a ligand to the phosphorus atom and an NHC at the P-bonded nitrogen atom (**64^A**, Scheme 16). This resonance structure (**64^A**) is reminiscent of the diphosphorus compounds **65** and **66** that bear two NHC ligands or two CAACs, respectively (Scheme 16).^{33,34}



Scheme 16 Synthesis of phosphazabutadiene **64** and its conversion to the radical cation **64***. Phosphorus mononitride formulation **64^A**. NHC- and CAAC-stabilized diphosphorus complexes **65** and **66**.





Fig. 9 Ball&stick representation of the phosphazabutadiene **64** as derived from XRD analysis (hydrogen atoms and *isopropyl* groups have been omitted).

The formulation **64** represents the phosphazabutadiene character of the compound. In the ^{31}P NMR spectroscopic analysis the chemical shift of **64** is observed at 134 ppm which is shifted to lower field with respect to the heavier congeners **65** and **66** (range: from 59 ppm to -74 ppm). As derived from XRD analysis the geometry of **64** (Fig. 9) was described as trans-bent with a short P-C_{CAAC} bond (1.719(2) Å), as well as a N-C_{NHC} distance (1.282(3) Å) that is in the range of C-N bond lengths of imino groups (*vide supra*). The P-N distance of 1.7085(16) Å is similar to typical P-N single bonds. Oxidation of **64** with $\text{Ph}_3\text{C}[\text{B}(\text{C}_6\text{F}_5)_4]$ (trityl tetrakis(pentafluorophenylborate)) afforded the radical cation **64^{•+}** (Scheme 16).³² This process is reversible as was shown by the regeneration of uncharged **64** via reduction with potassium graphite (KC_8 , Scheme 16). In the theoretical analysis of **64** and **64^{•+}** the shape of the HOMO and the SOMO (singly occupied molecular orbital), respectively, are very similar. They majorly comprise a π^* orbital of the PN group that shows bonding interaction with a p-type orbital at the carbenic atom of the NHC-, as well as the CAAC ligand.³² The EPR study of **64^{•+}** in frozen fluorobenzene at 100 K revealed g-tensors of $g_x = 2.0028$, $g_y = 2.0087$ and $g_z = 2.0052$ which is comparable to the respective values in **65^{•+}** and **66^{•+}**.³²⁻³⁴

The scope of applications of N-heterocyclic imines in phosphorus chemistry was extended in 2011 when Bertrand and coworkers reported the reduction of the bisimino compound **67**[OTf] (Tf = triflyl) with KC_8 to the uncharged phosphinyl radical **67[•]** (Scheme 16, Fig. 10).³⁵ Notably, the synthesis of **67**[OTf] proceeds via the chloride salt **67**[Cl] that could not be isolated in analytically pure form at that time but is purified in the course of the anion exchange (chloride vs triflate, Scheme 16).³⁵ The paramagnetic nature of **67[•]** was verified by its EPR study at 100 K in frozen THF from which the g-tensors $g_x = 2.0074$, $g_y = 2.0062$ and $g_z = 2.0024$ derived. A comparison with hyper fine coupling constants of atomic phosphorus revealed that an unpaired electron is primarily localized on the 3p(P) orbital (62%) with small contribution of 3s(P) orbital (2%).



Scheme 17 Conversion of the bisimino phosphonium salt **67**[OTf] to the phosphinyl radical **67[•]**. Synthesis of the imino(vanadyl)phosphinyl radical **69[•]** via chlorophosphin **68**. Vanadium-centres formulation **69^{•+}**.




Fig. 10 Ellipsoid plot (30% level) of the bis(imino)phosphinyl radical **67[•]** (hydrogen atoms and *isopropyl* groups have been omitted).

Bertrand and coworkers extended their investigation of imino-substituted phosphinyl radicals: The phosphorus dichloride **63** served as a precursor to the nitridovanadium-functionalized phosphorus monochloride **68** (Scheme 17).³⁵ In analogous fashion as for **67[•]** the reduction of **68** with KC_8 furnished the phosphinyl radical **69[•]** (Scheme 16).³⁵ From the EPR study of **69[•]** g-tensors of $g_x = 1.9726$, $g_y = 2.0048$ and $g_z = 1.9583$ were determined. Taking into account the hyper fine coupling constants to ^{51}V as well as ^{31}P it was concluded that the spin density in **69[•]** mainly resides on the vanadium centre (67%) and is only localized to a minor degree on the phosphorus atom and the NHC moiety. In contrast, the spin density of the bisimino derivative **67[•]** was found to reside with 62% in the 3p(P) orbital and with 2% in the 3s(P) orbital. In line with the structural and theoretical analysis the authors concluded that the bisimino radical **67[•]** is a phosphorus centred radical with little spin delocalization over the iminato ligands. The nitridovanadium congener **69[•]**, however, is best represented by the canonical structure **69^{•+}**, that is a vanadium(IV) complex with a phosphinimide ligand.

In 2012 Bertrand and coworkers reported the remarkable transformation of the azido bisimino phosphane **70** to the phosphinonitrene species **71** via irradiation at 254 nm (Scheme 18).³⁶ As a starting material the bisimino phosphonium salt **67**[Cl] was used that had also been implemented in the synthesis of the phosphinyl radical **67[•]** (Scheme 17).^{35,36} The theoretical analysis of the nitrene (**71**) suggested that the back donation of a nitrogen lone-pairs into accessible σ^* orbitals at the phosphorus atom significantly contributes to the thermodynamic stability of the compound. The phosphorus

atom in **71** is the centre of a trigonal-plane and the sum of the angles around the P atom amounts to 360° (Fig. 11). Notably, the P–N_{nitrene} bond length of 1.457(8) Å in **71** is significantly shorter than the P–N_{imino} distances (1.618(8) Å, 1.629(8) Å), as well as the P–N_{azido} distance of 1.895(11) Å in the precursor (**70**). This is in good agreement with the upfield shift in the ³¹P NMR spectrum of **71** (8 ppm) in comparison to **70** (111 ppm) which indicates the multiple-bond character of the PN functionality. The phosphininirene (**71**) reacts with *isopropyl isonitrile* (*i*PrNC) to yield the carbodiimide **72** that was not structurally characterized (Scheme 16).³⁶ In consequence, the created NCN*i*Pr group can be abstracted from the phosphorus atom implementing *isopropyltriflate* as an alkylating agent. In the outcome the starting material **67**⁺ is generated in the form of the triflate salt (**67**[OTf], Scheme 18).

Bertrand and coworkers described the transformation of the phosphininirene **71** to iminophosphonium triflates in 2013.³⁷ The methylation or protonation of **71** using methyltriflate or triflic acid, respectively, furnished **73**[OTf] or **74**[OTf] (Scheme 19). The P=N bond length of the phosphoranimine functionality in **74**⁺ amounts to 1.526(2) Å which is longer than the distance of these atoms in the precursor (vide supra, note that the structural parameters of **73**⁺ are not discussed due to poor data quality). Vice versa, the distance in between these atoms are decreased to 1.553(2) Å and 1.559(2) Å as compared to **71** (vide supra). This suggests a stronger interaction between the P-centre and the imino nitrogen atoms and accounts for the potential of **N-heterocyclic imino systems** in stabilizing cationic species. The addition of water to **73**[OTf] or **74**[OTf] yielded the cationic phosphine oxides **75**[OTf] or **76**[OTf], respectively (Scheme 19).³⁷ The expected electrophilic properties of **74**⁺ were verified by its conversion with dmap that generated the Lewis acid base adduct **79**[OTf] (Scheme 19).³⁷ Notably, the ³¹P NMR chemical shift of **79**⁺ was observed at significantly higher field (–1 ppm) than the precursor **74**⁺ (73 ppm).






Fig. 11 Ball&stick representation of the phosphininirene **71** as derived from XRD analysis (hydrogen atoms and *isopropyl* groups have been omitted).




Scheme 19 Methylation and protonation of **71** to **73**[OTf] and **74**[OTf] and their reactivity towards H₂O and dmap to produce (**75–77**)[OTf].




Scheme 18 Synthesis of phosphininirene **71** by photoirradiation of azidophosphate **70** and conversion to carbodiimide **72**.

In 2014 the literature on the phosphininirene species (**71**) was enriched by Bertrand and coworkers with their investigation of coinage metal-nitrene compounds.³⁸ By conversion of **71** with a corresponding equivalent of copper- or silver triflate (MOTf) the respective complexes with bridging or terminal phosphininirene ligands are generated (**78–83**, Scheme 20). The reaction of **71** with two equiv of MOTf furnished the bimetallic complexes **78** or **79** with a bridging nitrenic atom. These showed similar structural features in the solid state, that is a planar coordination environment of the phosphorus atom and significantly elongated P–N_{nitrenic} bond lengths (1.510(5) Å for **78** and 1.528(3) Å for **79**) with respect to **71**. Furthermore, the P–N_{imino} bond lengths are shortened (1.573(3) Å for **78** and 1.561(3) Å for **79**) and, vice versa, the C–N_{imino} distances are lengthened (range of 1.31–1.36 Å for **78** and **79**) which indicates the stronger allocation of electron density from the imidazolidin-2-imino system to the phosphorus atom than in the precursor. After conversion of two equiv **71** with MOTf the linear complexes **82** and **83** with

terminal bis(phosphinonitrene) ligands were obtained.³⁸ Notably, the M–N_{nitrenic} bond lengths (1.801(2) Å, 1.807(3) Å for **82** and 2.017(3) Å, 2.029(4) Å for **83**) in the linear complexes are decreased in comparison to the bimetallic systems (1.817(3) Å for **78** and 2.080(3) Å, 2.086(3) Å for **79**). Interestingly, for the conversions of the phosphinonitrene and MOTf in a one to one ratio it was concluded on the generation of **80** and **81**, respectively, by NMR spectroscopic analysis. However, these compounds were found in a dynamic equilibrium with their bridging and terminal congeners (**78**, **82** for **80** and **79**, **83** for **81**).³⁸



Scheme 20 The formations of metal-nitrene complexes **78–83** from phosphinonitrene **71**.




Scheme 21 Thermal conversion of phosphinonitrene **71** to **84** and reactivity with small molecules to imino complexes **85–91**.

A thorough study on the reactivity of the phosphinonitrene **71** was published in 2015.³⁹ The authors described its thermal transformation to the iminophosphorane **84**, as well as several conversions with typical small molecule substrates (Scheme 21). At elevated temperature quantitative rearrangement of **71** was observed. The nitrenic atom inserts into a tertiary carbon CH bond of an *isopropyl* side chain followed by migration of the Dip moiety to the phosphorus centre to create the five membered PNC₃ ring in **84**. The addition of an excess amount of acetonitrile to the phosphinonitrene (**71**) afforded a mixture (16:1) of the ketenimine **85** and the diazaphosphate **86** (Scheme 21).³⁹ Notably, the ketenimine is transformed into the diazaphosphate at elevated temperature (90 °C). This process was reasoned by the initial deprotonation of acetonitrile by the nitrenic centre and nucleophilic attack of the generated cyanomethylanion at the phosphorus atom to afford **85**. Subsequent cyclization and proton migration leads to formation of **86**. Reaction of the phosphinonitrene (**71**) with carbon dioxide or carbon disulfide yields the isocyanate **87** or the isothiocyanate **88**, respectively (Scheme 21).³⁹ One should point out the cleavage of the thermodynamically stable C=E double bond (E = O, S) in this process. Compound **71** activates elemental sulfur (S₈), as well as white phosphorus (P₄).³⁹ The reaction with S₈ furnishes the phosphine sulfide **89** that bears a thiosulfinylamino group at the phosphorus atom (Scheme 21). The conversion of **71** with P₄ affords phosphorus enriched **90** with a unique P₅N moiety via insertion of the PN_{nitrene} fragment into a P–P single bond of the P₄ cluster (Scheme 21). With a slight excess of water the phosphinonitrene (**71**) reacted to the aminophosphine oxide **91** as the product of the addition of H₂O to the PN_{nitrene} bond (Scheme 21).³⁹


Implementing the imidazolin-2-imino trimethylsilane **92** Vidović and coworkers synthesized the imino phosphorus dichloride **93** that was reacted with carbodiphosphorane to yield the phosphonium salt **94**[Cl] (Scheme 22).⁴⁰ The latter was subjected to chloride abstraction with two equiv silver hexafluoroantimonate and in the outcome the dicationic phosphinimine **95**[SbF₆]₂ was formed (Scheme 22).⁴⁰ The dication assumes a trans-bent structure motif and the P–N distance of 1.594(6) Å is significantly shorter than the respective bond lengths in the CAAC congener **64** (1.7085(16) Å) and its radical cation **64^{•+}** (1.645(4) Å). This suggests relevant double bond character for the PN fragment in **95²⁺**. As concluded from the theoretical analysis of the dication the authors attributed the increased PN interaction to the removal of electrons from the HOMO which majorly comprises the PN π* antibonding orbital. Taking into account structural parameters as the comparably long C–N_{imino} bond (1.367(8) Å) and bond polarizations derived from the NBO analysis it was presumed that **95²⁺** possesses dicationic phosphorus mononitride character to a minor degree (**95^{A2+}**, Scheme 22).³⁶ Regardless of the dominant resonance structure of **95²⁺** its isolation affirms the potential of the imidazolin-2-imino ligand for stabilizing cationic species.

The application of (benz)imidazolin-2-imino substituents as supporting groups for P-based ligands has recently emerged as a subgenre of the field of phosphorus compounds of the


iminato ligand. The chemistry relies on pioneering work of Kuhn and coworkers who converted $L^{Me_2}SiMe_3$ (**25**) to the imino dichlorophosphane **96** (Scheme 7, Scheme 23).^{31a} If treated with $AlCl_3$ this compound reacts to the ditopic phosphonium salt **97** $[AlCl_4]_2$ the structural formulation of which was based on NMR spectroscopic characteristics (Scheme 23).^{31a} The authors described that in solution **97**²⁺ is in equilibrium with **96** in dependence of the nucleophilic properties of the solvent. Moreover, Kuhn and coworkers described the iminophosphanes **98** and **99**, as well as the conversion of **99** to the iminophosphorane **100** (Scheme 23).^{31b}




Scheme 22 Reaction of imino phosphorus dichloride to the phosphonium salt **94** $[Cl]$ and its conversion to the dicationic phosphinimine **95**²⁺. Phosphorus mononitride formulation **95**⁴²⁺.



Scheme 23 Preparation of the imino phosphorus dichloride **96** from the imino trimethylsilane **25**. Reaction to the dimeric phosphonium cation **97**²⁺. The iminophosphanes **98** and **99**, as well as the iminophosphorane **100**.



Scheme 24 Reaction of the imino diphenylphosphine **101** to the borane-adduct **102**, as well as the phosphorus chalcogenids **103**–**106**.




Scheme 25 Synthesis of benzimidazolin-2-imino-substituted phosphines **108**–**113**, as well as the imidazolin-2-imino-substituted phosphines **115** and **116**. (a) 1.) $nBuLi$, THF, $-78\text{ }^\circ\text{C}$; 2.) chlorophosphine, room temperature. (b) 1.) $nBuLi$, THF, $-78\text{ }^\circ\text{C}$; 2.) $[Fe(C_5H_4PCl_2)_2]$. (c) **115**: iPr_2PCl , THF; **116**: 1.) PCl_3 , THF, $-78\text{ }^\circ\text{C}$; 2.) $iPrMgCl$, THF, $-78\text{ }^\circ\text{C}$.

In 2015 Mallik, Panda and coworkers reported the imino diphenylphosphine **101** that was converted to the borane adduct **102**, as well as the phosphorus chalcogenides **103**–**106** (Scheme 24).⁴¹

Also in 2015 Dielmann and coworkers established the use of imidazolin-2-imino-substituted phosphines as electron-rich ligands to transition metals.⁴² By lithiation of the benzimidazolin-2-imine $BL^{iPr}NH$ (**107**, $BL^{iPr} = 1,3$ -diisopropylbenzimidazolin-2-ylidene) and reaction with corresponding chlorophosphines the synthesis of the iminophosphines **108**–**113** was accomplished (Scheme 25).⁴² In addition, the conversion of the bulkier $L^{Me_2Mes_2}NSiMe_3$ (**114**, $L^{Me_2Mes_2} = 1,3$ -dimesityl-4,5-dimethyl-imidazolin-2-ylidene) with respective chlorophosphines lead to the iminophosphines **115** and **116** in the outcome (Scheme 25).⁴² To assess the electron-donor strength of these phosphorus-based ligands the Tolman Electronic Parameters (TEP) of their nickel tricarbonyl complexes were determined.⁴³ Moreover, they were evaluated according to the Huynh's method. The ^{13}C NMR-spectroscopic shift of the carbene carbon of the BL^{iPr} group in the $trans$ - $\{PdBr_2(BL^{iPr})\}$ ligand is sensitive to the donor abilities of the ligand, in which the carbene resonance of the BL^{iPr} is downfield shifted with increasing donor strength of the ligand $trans$ to the BL^{iPr} group.⁴⁴ The Huynh's parameters of the iminophosphines show same qualitative trend as the TEP analysis. As a result, many of the iminophosphines were found to be more potent electron pair donors than most electron-rich trialkylphosphines. Remarkably, the authors concluded that the iminophosphines **111** and **112**, as well as **115** and **116** are stronger donor ligands than classical NHCs. In addition, the bisimine **116** was presumed to be a more potent donor ligand than the very strongly electron donating abnormal NHCs exceeding the capability of monoimine **115**, as well as the bis- and the trisimine **111** and **112**. Consequently, it was reasoned

that the imidazolin-2-imino ligand is a stronger π -electron donor than the related benzimidazolin-2-imino ligand.

Uncharged organosuperbases that comprise the imidazolidin-2-imino fragment as a chiral bis(guanidine)iminophosphorane were described by Takeda and Terada in 2013.⁴⁵ The respective iminophosphonium salts **117**·HCl, **118**·HCl, **119**·HCl and **120**·HBr were synthesized by conversion of respective aminoguanidinium halides with phosphorus pentachloride in the presence of base followed by acidic work-up (Scheme 26).⁴⁵ The stability of the iminophosphonium hydrohalide salt and, thus the high Brønsted basicity of the uncharged compounds relies on the properties of the iminophosphorane as an electron-rich oligodentate ligand. The free base was not characterized but generated by reaction with potassium *tert*butoxide and used in situ for the assessment of catalytic activity in the electrophilic amination of tetralones with azodicarboxylate.⁴⁵ Notably, no particular reason for the use of the imidazolidin-2-imino group instead of acyclic guanidino groups was pointed out by the authors. We assume that its implementation rather follows synthetic applicability for furnishing the chiral bis(guanidine)iminophosphorane species. Notably, the scope of catalytic applications of this compound as the chiral uncharged organosuperbase was expanded in recent years.⁴⁶




Scheme 26 Preparation of the chiral imino phosphonium halides (*M*)-**117**·HCl–(*M*)-**120**·HBr and the stereoisomer (*P*)-**117**·HCl.


Miscellaneous: survey of related nitrogen compounds

Compounds in which the exocyclic nitrogen atom of an **N-heterocyclic imino** fragment bonds to another nitrogen atom are abundant in the literature but may be accounted to the field of classic organic chemistry rather than inorganic or organometallic coordination chemistry which is the focus of this review. They can be categorized into triazenes⁴⁷ (representative example: **121**, Scheme 27), azines^{18,48} (**122**, subcategory: bisguanidines, Scheme 27), as well as diazotates⁴⁹ (**123**, Scheme 28) and their azoimidazolium⁵⁰ derivatives (**124**[BPh₄], Scheme 28).

Imidazolyl triazenes (**121**) release dinitrogen under thermal conditions which leads to the formation of imino organyls whereas their exposure to acidic condition generates a diazonium species along with the imine. The latter accounts for the pronounced proton affinity of the imidazolin-2-imino group. Aside of applications in organic chemistry azines (**122**) and diazotates (**123**), as well as their azoimidazolium spin-offs (**124**[BPh₄]) are found to be employed as ligands to main group elements or transition metals in rare instances. The mechanism for the formation of **124**[BPh₄] is proposed to involve AlCl₃-mediated oxygen abstraction to afford the dicationic diazonium compound, followed by its azo coupling with mesitylene. As an interesting difference in their bonding modes the C–N_{imino} distance in azines is generally shorter than in the reported diazotates. This suggests high CN double bond character for the former and considerable single bond character along with delocalization of positive charge into the imidazoline ring for the latter. Similarly, the azoimidazolium cation **124**⁺ exhibits a comparably long C–N_{imino} bond length of 1.386(2) Å which indicates that the positive charge is majorly distributed among the atoms of the five-membered ring (Scheme 28, Fig. 12).



Scheme 27 Synthesis of representative examples for the compounds classes: triazenes (**121**) and azines (**122**).



Scheme 28 Synthesis of representative examples for the compound classes: diazotates (**123**) and azoimidazolium salts (**124**[BPh₄]).




Fig. 12 Ellipsoid plot (30% level) of the azoimidazolium cation **124**⁺ (hydrogen atoms have been omitted).

Conclusions

This survey of the coordination chemistry of the main group element complexes with N-heterocyclic imines shows that these ligands are suitable for the isolation of otherwise elusive species (e.g. Al=Te double bond, stannyleneid, phosphinonitrene). In particular, they have proven valuable for the thermodynamic stabilization of electron-deficient central atoms, and thus enabled the isolation of rare types of low-coordinate cationic metal complexes (e.g. cationic thioxoborane, gerymliumylidene). The strongly electron-donating character of the imidazolin-2-iminato ligand derives from the efficient delocalization of cationic charge density into the five-membered ring system. The lengthening of the C–N_{imino} distance is an indicator for the allocation of electron density by the ligand as it is often observed upon transformation of an uncharged species into a cationic offspring. The exploration of the phosphorus chemistry of this imino ligand demonstrates its applicability for the stabilization of charged, as well as uncharged phosphorus-centred radicals. Moreover, the electron-rich nature of the imidazolin-2-imino group has resulted in a new class of phosphines that bear supporting imino groups and act as highly electron donating phosphorus-centred ligands.

The chemistry of N-heterocyclic imine complexes of main group elements is still in its infancy as compared to the wider spread field of metal amides.⁵¹ However, its evolvement in recent years underlines the usefulness as ancillary ligands and distinguishes it from other classes of nitrogen-based ligand systems. Future work should study yet unexplored complexes of the imidazoli(di)n-2-imino group with heavier main group metals and focus on catalytic applications of respective systems.

Acknowledgements

Financial support of the WACKER Chemie AG, as well as the European Research Council (SILION 637394) is gratefully acknowledged.

Notes and references

- (a) N. Kuhn, H. Bohnen, J. Kreutzberg, D. Bläser and R. Boese, *J. Chem. Soc., Chem. Commun.*, 1993, 1136; (b) S. M. Ibrahim Al-Rafia, A. C. Malcolm, S. K. Liew, M. J. Ferguson, R. McDonald and E. Rivard, *Chem. Commun.*, 2011, **47**, 6987; (c) S. M. Ibrahim Al-Rafia, M. J. Ferguson and E. Rivard, *Inorg. Chem.*, 2011, **50**, 10543; (d) S. Kronig, P. G. Jones and M. Tamm, *Eur. J. Inorg. Chem.*, 2013, 2301; (e) Y. Wang, M. Y. Abraham, R. J. Gilliard, Jr., D. R. Sexton, P. Wei and G. H. Robinson, *Organometallics*, 2013, **32**, 6639.
- N. Kuhn, M. Göhner, M. Grathwohl, J. Wiethoff, G. Frenking and Y. Chen, *Z. Anorg. Allg. Chem.*, 2003, **629**, 793.
- (a) A. G. Trambitas, T. K. Panda and M. Tamm, *Z. Anorg. Allg. Chem.*, 2010, **636**, 2156; (b) Y. Wu and M. Tamm, *Coord. Chem. Rev.*, 2014, **260**, 116.
- N. Kuhn, R. Fawzi, M. Steimann, J. Wiethoff, D. Bläser and R. Boese, *Z. Naturforsch.*, 1995, **50b**, 1779.
- N. Kuhn, U. Abram, C. Maichle-Mößmer and J. Wiethoff, *Z. Anorg. Allg. Chem.*, 1997, **623**, 1121.
- N. Kuhn, R. Fawzi, M. Steimann, J. Wiethoff and G. Henkel, *Z. Anorg. Allg. Chem.*, 1997, **623**, 1577.
- N. Kuhn, R. Fawzi, M. Steimann and J. Wiethoff, *Z. Anorg. Allg. Chem.*, 1997, **623**, 554.
- D. Franz, E. Irran and S. Inoue, *Dalton Trans.*, 2014, **43**, 4451.
- D. Franz and S. Inoue, *Chem. – Asian J.*, 2014, **9**, 2083.
- D. Franz, E. Irran and S. Inoue, *Angew. Chem., Int. Ed.*, 2014, **53**, 14264.
- (a) K. Dehnicke and F. Weller, *Coord. Chem. Rev.*, 1997, **158**, 103; (b) S. Courtenay, J. Y. Mutus, R. W. Schurko and D. W. Stephan, *Angew. Chem., Int. Ed.*, 2002, **41**, 498; (c) S. Courtenay, D. Walsh, S. Hawkeswood, P. Wei, A. K. Das and D. Stephan, *Inorg. Chem.*, 2007, **46**, 3623; (d) O. Alhomainan, E. Hollink and D. W. Stephan, *Organometallics*, 2007, **26**, 3041; (e) M. H. Holthausen, I. Mallov and D. W. Stephan, *Dalton Trans.*, 2014, **43**, 15201; (f) K. Spannhoff, R. Rojas, R. Fröhlich, G. Kehr and E. Erker, *Organometallics*, 2011, **30**, 2377; (g) K. Jaiswal, B. Prashanth, S. Ravi, K. R. Shamasundar and S. Singh, *Dalton Trans.*, 2015, **44**, 15779.
- (a) C. M. Ong, P. McKarns and D. W. Stephan, *Organometallics*, 1999, **18**, 4197; (b) K. Aparna, R. McDonald, M. Ferguson and R. G. Cavell, *Organometallics*, 1999, **18**, 4241; (c) Z.-X. Wang and Y.-X. Li, *Organometallics*, 2003, **22**, 4900; (d) J. Guo, J.-S. Lee, M.-C. Foo, K.-C. Lau, H.-W. Xi, K. H. Lim and C.-W. So, *Organometallics*, 2010, **29**, 939; (e) C. V. Cárdenas, M. Á. M. Hernández and J.-M. Grévy, *Dalton Trans.*, 2010, **39**, 6441.
- S. M. I. Al-Rafia, R. McDonald, M. J. Ferguson and E. Rivard, *Chem. – Eur. J.*, 2012, **18**, 13810.
- M. W. Lui, N. R. Paisley, R. McDonald, M. J. Ferguson and E. Rivard, *Chem. – Eur. J.*, 2016, **22**, 2134.
- Z. Yang, M. Zhong, X. Ma, S. De, C. Anusha, P. Parameswaran and H. W. Roesky, *Angew. Chem., Int. Ed.*, 2015, **54**, 10225.
- D. Franz and S. Inoue, *Chem. – Eur. J.*, 2014, **20**, 10645.
- (a) D. Franz, T. Szilvási, E. Irran and S. Inoue, *Nat. Commun.*, 2015, **6**, 10037; (b) D. Franz and S. Inoue, *Dalton Trans.* 2016, **45**, 9385.
- J. M. Hopkins, M. Bowdridge, K. N. Robertson, T. S. Cameron, H. A. Jenkins and J. A. C. Clyburne, *J. Org. Chem.*, 2001, **66**, 5713.

- 19 (a) M. Tamm, S. Randoll, T. Bannenberg and E. Herdtweck, *Chem. Commun.*, 2004, 876; (b) M. Tamm, D. Petrovic, S. Randoll, S. Beer, T. Bannenberg, P. G. Jones and J. Grunenberg, *Org. Biomol. Chem.*, 2007, **5**, 523.
- 20 S. Randoll, P. G. Jones and M. Tamm, *Organometallics*, 2008, **27**, 3232.
- 21 S. Inoue and K. Leszczyńska, *Angew. Chem., Int. Ed.*, 2012, **51**, 8589.
- 22 M. Asay, C. Jones and M. Driess, *Chem. Rev.*, 2011, **111**, 354.
- 23 G.-H. Lee, R. West and T. Müller, *J. Am. Chem. Soc.*, 2003, **125**, 8114.
- 24 M. W. Lui, C. Merten, M. J. Ferguson, R. McDonald, Y. Xu and E. Rivard, *Inorg. Chem.*, 2015, **54**, 2040.
- 25 (a) T. Ochiai, D. Franz, X.-N. Wu and S. Inoue, *Dalton Trans.*, 2015, **44**, 10952; (b) T. Ochiai, T. Szilvási, D. Franz, E. Irran and S. Inoue, *Angew. Chem., Int. Ed.*, 2016, DOI: [10.1002/anie.201605636](https://doi.org/10.1002/anie.201605636).
- 26 (a) T. Ochiai, D. Franz, E. Irran and S. Inoue, *Chem. – Eur. J.*, 2015, **21**, 6704; (b) T. Ochiai and S. Inoue, *Phosphorus Sulfur Silicon Relat. Elem.*, 2016, **191**, 624.
- 27 T. Ochiai, D. Franz, X.-N. Wu, E. Irran and S. Inoue, *Angew. Chem., Int. Ed.*, 2016, **55**, 6983.
- 28 (a) A. Akkari, J. J. Byrne, I. Saur, G. Rima, H. Gornitzka and J. Barrau, *J. Organomet. Chem.*, 2001, **622**, 190; (b) Y. Ding, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt and P. P. Power, *Organometallics*, 2001, **20**, 1190; (c) P. B. Hitchcock, J. Hu, M. F. Lappert and J. R. Severn, *Dalton Trans.*, 2004, 14193; (d) A. P. Dove, V. C. Gibson, E. L. Marshall, H. S. Rzepa, A. J. P. White and D. J. Williams, *J. Am. Chem. Soc.*, 2006, **128**, 9834. (e) S. L. Choong, C. Schenk, A. Stasch, D. Dange and C. Jones, *Chem. Commun.*, 2012, **48**, 2504. (f) R. Olejnik, Z. Padělková, R. Mundil, J. Merna and A. Růžička, *Appl. Organometal. Chem.*, 2014, **28**, 405.
- 29 J. Volbeda, P. G. Jones and M. Tamm, *Inorg. Chim. Acta*, 2014, **422**, 158.
- 30 D. Petrovic, T. Glöge, T. Bannenberg, C. G. Hrib, S. Randoll, P. G. Jones and M. Tamm, *Eur. J. Inorg. Chem.*, 2007, 3472.
- 31 (a) N. Kuhn, R. Fawzi, M. Steimann and J. Wiethoff, *Chem. Ber.*, 1996, **129**, 479; (b) N. Kuhn, H. Kotowski and J. Wiethoff, *Phosphorus Sulfur Silicon Relat. Elem.*, 1998, **133**, 237.
- 32 R. Kinjo, B. Donnadieu and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 5930.
- 33 Y. Wang, Y. Xie, P. Wie, R. B. King, H. F. Schaefer, III, P. v. R. Schleyer and G. H. Robinson, *J. Am. Chem. Soc.*, 2008, **130**, 14970.
- 34 O. Back, B. Donnadieu, P. Parameswaran, G. Frenking and G. Bertrand, *Nat. Chem.*, 2010, **2**, 369.
- 35 O. Back, B. Donnadieu, M. v. Hopffgarten, S. Klein, R. Tonner, G. Frenking and G. Bertrand, *Chem. Sci.*, 2011, **2**, 858.
- 36 F. Dielmann, O. Back, M. Henry-Ellinger, P. Jerabek, G. Frenking and G. Bertrand, *Science*, 2012, **337**, 1526.
- 37 F. Dielmann, C. E. Moore, A. Rheingold and G. Bertrand, *J. Am. Chem. Soc.*, 2013, **135**, 14071.
- 38 F. Dielmann, D. M. Andrada, G. Frenking and G. Bertrand, *J. Am. Chem. Soc.*, 2014, **136**, 3800.
- 39 F. Dielmann and G. Bertrand, *Chem. – Eur. J.*, 2015, **21**, 191.
- 40 Y. K. Loh, C. Gurnani, R. Ganguly and D. Vidović, *Inorg. Chem.*, 2015, **54**, 3087.
- 41 K. Naktode, S. D. Gupta, A. Kundu, S. K. Jana, H. P. Nayek, B. S. Mallik and T. K. Panda, *Austr. J. Chem.*, 2015, **68**, 127.
- 42 M. A. Wünsche, P. Mehlmann, T. Witteler, F. Buß, P. Rathmann and F. Dielmann, *Angew. Chem., Int. Ed.*, 2015, **54**, 11857.
- 43 C. A. Tolman, *Chem. Rev.*, 1977, **77**, 313.
- 44 H. V. Huynh, Y. Han, R. Jothibas and J. A. Yang, *Organometallics*, 2009, **28**, 5395.
- 45 T. Takeda and M. Terada, *J. Am. Chem. Soc.*, 2013, **135**, 15306.
- 46 (a) T. Takeda and M. Terada, *Aust. J. Chem.*, 2014, **67**, 1124; (b) A. Kondoh, M. Oishi, T. Takeda and M. Terada, *Angew. Chem., Int. Ed.*, 2015, **54**, 15836; (c) T. Takeda, A. Kondoh and M. Terada, *Angew. Chem., Int. Ed.*, 2016, **55**, 4734.
- 47 (a) D. M. Khramov and C. W. Bielawski, *Chem. Commun.*, 2005, 4958; (b) D. M. Khramov and C. W. Bielawski, *J. Org. Chem.*, 2007, **72**, 9407; (c) D. J. Coady, D. M. Khramov, B. C. Norris, A. G. Tennyson and C. W. Bielawski, *Angew. Chem., Int. Ed.*, 2009, **48**, 5187; (d) A. G. Tennyson, D. M. Khramov, C. D. Varnado Jr., P. T. Creswell, J. W. Kamplain, V. M. Lynch and C. W. Bielawski, *Organometallics*, 2009, **28**, 5142; (e) A. G. Tennyson, E. J. Moorhead, B. L. Madison, J. A. V. Er, V. M. Lynch and C. W. Bielawski, *Eur. J. Org. Chem.*, 2010, 6277; (f) R. J. Ono, Y. Suzuki, D. M. Khramov, M. Ueda, J. L. Sessler and C. W. Bielawski, *J. Org. Chem.*, 2011, **76**, 3239; (g) D. Jishkariani, C. D. Hall, A. Demircan, B. J. Tomlin, P. J. Steel and A. R. Katritzky, *J. Org. Chem.*, 2013, **78**, 3349; (h) S. Patil, K. White and A. Bugarin, *Tetrahedron Lett.*, 2014, **55**, 4826; (i) F. W. Kimani and J. C. Jewett, *Angew. Chem., Int. Ed.*, 2015, **54**, 4051.
- 48 (a) B. Bildstein, M. Malaun, H. Kopačka, K.-H. Ongania and K. Wurst, *J. Organomet. Chem.*, 1999, **572**, 177; (b) M. Reinmuth, C. Neuhäuser, P. Walter, M. Enders, E. Kaifer and H.-J. Himmel, *Eur. J. Inorg. Chem.*, 2011, 83; (c) J. Tauchman, K. Hladíková, F. Uhlík, I. Císařová and Štěpnička, *New J. Chem.*, 2013, **37**, 2019; (d) H. Herrmann, M. Reinmuth, S. Wiesner, O. Hübner, E. Kaifer, H. Wadepl and H.-J. Himmel, *Eur. J. Inorg. Chem.*, 2015, 2345.
- 49 (a) A. G. Tskhovrebov, E. Solari, M. D. Wodrich, R. Scopelliti and K. Severin, *Angew. Chem., Int. Ed.*, 2012, **51**, 232; (b) A. G. Tskhovrebov, E. Solari, M. D. Wodrich, R. Scopelliti and K. Severin, *J. Am. Chem. Soc.*, 2012, **134**, 1471; (c) A. G. Tskhovrebov, B. Vuichoud, E. Solari, R. Scopelliti and K. Severin, *J. Am. Chem. Soc.*, 2013, **135**, 9486.
- 50 A. G. Tskhovrebov, L. C. E. Naested, E. Solari, R. Scopelliti and K. Severin, *Angew. Chem., Int. Ed.*, 2015, **54**, 1289.
- 51 (a) M. F. Lappert, P. P. Power, A. V. Protchenko and A. L. Seeber, *Metal Amide Chemistry*, Wiley-VCH, Weinheim, 2009; (b) D. L. Kays, *Chem. Soc. Rev.*, 2016, **45**, 1004.